



Ann. Funct. Anal. 4 (2013), no. 2, 87–96

ANNALS OF FUNCTIONAL ANALYSIS

ISSN: 2008-8752 (electronic)

URL: www.emis.de/journals/AFA/

OPTIMAL BUNDLES IN NORMED SPACES

H. H. CUENYA, F. E. LEVIS* AND C. N. RODRIGUEZ

Communicated by J. Soria

ABSTRACT. In this paper, we prove existence of optimal bundles for a countable set of data in a broad class of normed spaces, which extend previous known results for a finite data set in a Hilbert space. In addition, we study the behavior of deviations and diameters for an increasing sequence of data sets.

DEPARTAMENTO DE MATEMÁTICA, FCEFQYN, UNIVERSIDAD NACIONAL DE RÍO CUARTO, RUTA 36 KM 601, RÍO CUARTO, ARGENTINA.

E-mail address: hcuenya@exa.unrc.edu.ar

E-mail address: flevis@exa.unrc.edu.ar

E-mail address: crodriguez@exa.unrc.edu.ar

Date: Received: 4 September 2012; Revised: 18 December 2012; Accepted: 5 January 2013.

* Corresponding author.

2010 *Mathematics Subject Classification.* Primary 41A65; Secondary 41A28.

Key words and phrases. Optimal bundles, deviation, diameter, normed space.