

**THE CANONICAL TENSOR FIELDS  
OF TYPE (1, 1) ON  $(J^r(\odot^2 T^*))^*$**

PAWEŁ MICHAŁEC

ABSTRACT. We prove that every natural affnor on  $(J^r(\odot^2 T^*))^*(M)$  is proportional to the identity affnor if  $\dim M \geq 3$ .

0. INTRODUCTION

For every  $n$ -dimensional manifold  $M$  we have the vector bundle

$$J^r(\odot^2 T^*)(M) = \{j_x^r \tau \mid \tau \text{ is a symmetric tensor field type } (0, 2) \text{ on } M, x \in M\}.$$

Every local diffeomorphism  $\varphi : M \rightarrow N$  between  $n$ -manifolds gives a vector bundle homomorphism  $J^r(\odot^2 T^*)(\varphi) : J^r(\odot^2 T^*)(M) \rightarrow J^r(\odot^2 T^*)(N)$ ,  $j_x^r \tau \rightarrow j_{\varphi(x)}^r(\varphi_* \tau)$ . Functor  $J^r(\odot^2 T^*) : \mathcal{M}f_n \rightarrow \mathcal{VB}$  is a vector natural bundle over  $n$ -manifolds in the sense of [5]. Let  $(J^r(\odot^2 T^*))^* : \mathcal{M}f_n \rightarrow \mathcal{VB}$  be the dual vector bundle,  $(J^r(\odot^2 T^*))^*(M) = (J^r(\odot^2 T^*)(M))^*$ ,  $(J^r(\odot^2 T^*))^*(\varphi) = (J^r(\odot^2 T^*)(\varphi^{-1}))^*$  for  $M$  and  $\varphi$  as above.

An affnor on a manifold  $M$  is a tensor field of type (1, 1) on  $M$ .

A natural affnor  $Q$  on  $(J^r(\odot^2 T^*))^*$  is a system of affnors

$$Q : T(J^r(\odot^2 T^*))^*(M) \rightarrow T(J^r(\odot^2 T^*))^*(M)$$

on  $(J^r(\odot^2 T^*))^*(M)$  for every  $n$ -manifold  $M$  satisfying the naturality condition  $T(J^r(\odot^2 T^*))^*(\varphi) \circ Q = Q \circ T(J^r(\odot^2 T^*))^*(\varphi)$  for every local diffeomorphism  $\varphi : M \rightarrow N$  between  $n$ -manifolds.

In this paper we prove, that every natural affnor  $Q$  on  $(J^r(\odot^2 T^*))^*$  over  $n$ -manifolds is proportional to the identity affnor if  $n \geq 3$ .

The proof of the classification theorem is based on the method from paper [7], where there are determined the natural affnors on  $(J^r(\wedge^2 T^*))^*$ . However the proof is different, because the tensor field  $dx^1 \odot dx^1$  on  $\mathbf{R}^n$  is non-zero, in contrast to  $dx^1 \wedge dx^1$ .

---

2000 *Mathematics Subject Classification*: 58A20.

*Key words and phrases*: natural affnor, natural bundle, natural transformation.

Received December 1, 2001.

Natural affinors on some natural bundle  $F$  can be used to study torsions  $[Q, \Gamma]$  of a connection  $\Gamma$  of  $F$ . That is why, the natural affinors have been study in many papers, [1] ... [11], e.t.c.

The usual coordinates on  $\mathbf{R}^n$  are denoted by  $x^i$ . The canonical vector fields on  $\mathbf{R}^n$  are denoted by  $\partial_i = \frac{\partial}{\partial x^i}$ .

All manifolds are assumed to be finite dimensional and smooth, i.e. of class  $C^\infty$ . Mappings between manifolds are assumed to be smooth.

### 1. THE LINEAR NATURAL TRANSFORMATIONS $T(J^r(\odot^2 T^*))^* \rightarrow (J^r(\odot^2 T^*))^*$

A natural transformation  $T(J^r(\odot^2 T^*))^* \rightarrow (J^r(\odot^2 T^*))^*$  over  $n$ -manifolds is a system of fibred maps

$$A : T(J^r(\odot^2 T^*))^*(M) \rightarrow (J^r(\odot^2 T^*))^*(M)$$

over  $\text{id}_M$  for every  $n$ -manifold  $M$  such that

$$(J^r(\odot^2 T^*))^*(f) \circ A = A \circ T(J^r(\odot^2 T^*))^*(f)$$

for every local diffeomorphism  $f : M \rightarrow N$  between  $n$ -manifolds.

A natural transformation  $A : T(J^r(\odot^2 T^*))^* \rightarrow (J^r(\odot^2 T^*))^*$  is called linear if  $A$  gives a linear map  $T_y(J^r(\odot^2 T^*))^*(M) \rightarrow ((J^r(\odot^2 T^*))^*(M))_x$  for any  $y \in ((J^r(\odot^2 T^*))^*(M))_x$ ,  $x \in M$ .

**Theorem 1.** *If  $n \geq 3$  and  $r$  are natural numbers, then every linear natural transformation  $A : T(J^r(\odot^2 T^*))^* \rightarrow (J^r(\odot^2 T^*))^*$  over  $n$ -manifolds is equal to 0.*

The proof of Theorem 1 will occupy Sections 2 – 6.

### 2. THE REDUCIBILITY PROPOSITIONS

Every element from the fibre  $((J^r(\odot^2 T^*))^*(\mathbf{R}^n))_0$  is a linear combination of all elements  $(j_0^r(x^\alpha dx^i \odot dx^j))^*$ , where  $\alpha \in (\mathbf{N} \cup \{0\})^n$ ,  $|\alpha| \leq r$ ,  $i \leq j$ ,  $i, j = 1, \dots, n$ . The elements  $(j_0^r(x^\alpha dx^i \odot dx^j))^*$  are dual basis to the basis  $j_0^r(x^\alpha dx^i \odot dx^j)$  of  $(J^r(\odot^2 T^*)(\mathbf{R}^n))_0$ .

Consider a linear natural transformation  $A : T(J^r(\odot^2 T^*))^* \rightarrow (J^r(\odot^2 T^*))^*$ .

**Lemma 1.** *Suppose  $A$  satisfies*

$$\langle A(u), j_0^r(x^\alpha dx^i \odot dx^j) \rangle = 0$$

for every  $u \in (T(J^r(\odot^2 T^*))^*(\mathbf{R}^n))_0$ ,  $\alpha \in (\mathbf{N} \cup \{0\})^n$ ,  $|\alpha| \leq r$ ,  $i \leq j$ ,  $i, j = 1, \dots, n$ . Then  $A = 0$ .

**Proof.** If assumptions of Lemma 1 meets, then  $A(u) = 0$  for every  $u \in (T(J^r(\odot^2 T^*))^*(\mathbf{R}^n))_0$ . Let  $w \in (T(J^r(\odot^2 T^*))^*(M))_x$ ,  $x \in M$ . There exists a chart  $\varphi : M \supset U \rightarrow \mathbf{R}^n$  such that  $\varphi(x) = 0$  and  $U$  is open subset including  $x$ . Since  $A$  is invariant with respect to  $\varphi$ , we have  $A(w) = T(J^r(\odot^2 T^*))^*(\varphi^{-1})(A(u))$ , where  $u = T(J^r(\odot^2 T^*))^*(\varphi)(w) \in (T(J^r(\odot^2 T^*))^*(\mathbf{R}^n))_0$ . Then  $A(w) = 0$ , because  $A(u) = 0$ . That is why  $A = 0$ . The lemma is proved.  $\square$

**Lemma 2.** *Suppose that*

$$\langle A(u), j_0^r(x^\alpha dx^1 \odot dx^1) \rangle = \langle A(u), j_0^r(x^\alpha dx^1 \odot dx^2) \rangle = 0$$

for every  $u \in (T(J^r(\odot^2 T^*))^*(\mathbf{R}^n))_0$ ,  $\alpha \in (\mathbf{N} \cup \{0\})^n$ ,  $|\alpha| \leq r$ ,  $i \leq j$ ,  $i, j = 1, \dots, n$ . Then  $A = 0$ .

**Proof.** Let  $u \in (T(J^r(\odot^2 T^*))^*(\mathbf{R}^n))_0$ ,  $\alpha \in (\mathbf{N} \cup \{0\})^n$ ,  $|\alpha| \leq r$ ,  $i \leq j$ ,  $i, j = 1, \dots, n$ . It is enough to prove, that  $\langle A(u), j_0^r(x^\alpha dx^i \odot dx^j) \rangle = 0$ .

Consider two cases

a)  $i = j$ . Let  $\varphi : \mathbf{R}^n \rightarrow \mathbf{R}^n$  be a diffeomorphism transforming  $x^i$  into  $x^1$  and  $x^\alpha$  into  $x^{\tilde{\alpha}}$  for some  $\tilde{\alpha} \in (\mathbf{N} \cup \{0\})^n$ ,  $|\tilde{\alpha}| \leq r$ . From the invariance of  $A$  with respect to  $\varphi$  and the assumption of Lemma 2, we have  $\langle A(u), j_0^r(x^\alpha dx^i \odot dx^i) \rangle = \langle A(\tilde{u}), j_0^r(x^{\tilde{\alpha}} dx^1 \odot dx^1) \rangle = 0$ , where  $\tilde{u} = T(J^r(\odot^2 T^*))^*(\varphi)(u)$

b)  $i \neq j$ . Let  $\varphi : \mathbf{R}^n \rightarrow \mathbf{R}^n$  be a diffeomorphism transforming  $x^i$  in  $x^1$ ,  $x^j$  in  $x^2$  and  $x^\alpha$  in  $x^{\tilde{\alpha}}$  for some  $\tilde{\alpha} \in (\mathbf{N} \cup \{0\})^n$ ,  $|\tilde{\alpha}| \leq r$ . From invariance of  $A$  with respect to  $\varphi$  and the assumption of Lemma 2, we have  $\langle A(u), j_0^r(x^\alpha dx^i \odot dx^j) \rangle = \langle A(\tilde{u}), j_0^r(x^{\tilde{\alpha}} dx^1 \odot dx^2) \rangle = 0$ , where  $\tilde{u} = T(J^r(\odot^2 T^*))^*(\varphi)(u)$ .  $\square$

**Lemma 3.** *Suppose  $A$  satisfies*

$$\begin{aligned} \langle A(u), j_0^r(dx^1 \odot dx^1) \rangle &= \langle A(u), j_0^r(x^3 dx^1 \odot dx^1) \rangle \\ &= \langle A(u), j_0^r(dx^1 \odot dx^2) \rangle = \langle A(u), j_0^r(x^3 dx^1 \odot dx^2) \rangle = 0 \end{aligned}$$

for every  $u \in (T(J^r(\odot^2 T^*))^*(\mathbf{R}^n))_0$ ,  $\alpha \in (\mathbf{N} \cup \{0\})^n$ ,  $|\alpha| \leq r$ ,  $i \leq j$ ,  $i, j = 1, \dots, n$ . Then  $A = 0$ .

**Proof.** Let  $\alpha \in (\mathbf{N} \cup \{0\})^n$ ,  $|\alpha| \leq r$ ,  $u \in (T(J^r(\odot^2 T^*))^*(\mathbf{R}^n))_0$ ,  $\alpha \neq e_3 = (0, 0, 1, 0, \dots, 0) \in (\mathbf{N} \cup \{0\})^n$ .

On the strength of Lemma 2 it is enough to prove that

$$\langle A(u), j_0^r(x^\alpha dx^1 \odot dx^1) \rangle = \langle A(u), j_0^r(x^\alpha dx^1 \odot dx^2) \rangle = 0.$$

We can set that  $\alpha \neq 0$ . Let  $\varphi : \mathbf{R}^n \rightarrow \mathbf{R}^n$  be a diffeomorphism transforming  $x^1$  in  $x^1$ ,  $x^2$  in  $x^2$  and  $x^3 + x^\alpha$  in  $x^3$ . From the invariance of  $A$  with respect to  $\varphi$  and the assumption of Lemma 3, we have

$$\begin{aligned} \langle A(u), j_0^r(x^\alpha dx^1 \odot dx^1) \rangle &= \langle A(u), j_0^r(x^3 dx^1 \odot dx^1) \rangle + \langle A(u), j_0^r(x^\alpha dx^1 \odot dx^1) \rangle \\ &= \langle A(u), j_0^r((x^3 + x^\alpha) dx^1 \odot dx^1) \rangle \\ &= \langle A(\tilde{u}), j_0^r(x^3 dx^1 \odot dx^1) \rangle = 0 \end{aligned}$$

where  $\tilde{u} = T(J^r(\odot^2 T^*))^*(\varphi)(u)$ .

Similarly  $\langle A(u), j_0^r(x^\alpha dx^1 \odot dx^2) \rangle = 0$ .  $\square$

**Lemma 4.** *Suppose that*

$$\langle A(u), dx^1 \odot dx^2 \rangle = \langle A(u), j_0^r(x^3 dx^1 \odot dx^2) \rangle = 0$$

for every  $u \in (T(J^r(\odot^2 T^*))^*(\mathbf{R}^n))_0$ . Then  $A = 0$ .

**Proof.** By Lemma 3 it is sufficient to show that

$$\langle A(u), dx^1 \odot dx^1 \rangle = \langle A(u), j_0^r(x^3 dx^1 \odot dx^1) \rangle = 0$$

for every  $u \in (T(J^r(\odot^2 T^*))^*(\mathbf{R}^n))_0$ .

Let  $u \in (T(J^r(\odot^2 T^*))^*(\mathbf{R}^n))_0$ . Consider a diffeomorphism  $\varphi : \mathbf{R}^n \rightarrow \mathbf{R}^n$  transforming  $x^1$  in  $x^1$ ,  $x^2$  in  $x^1 + x^2$  and  $x^3$  in  $x^3$ . Then from the invariance of  $A$  with respect to  $\varphi$  and the assumption of lemma, we have

$$\begin{aligned} 0 &= \langle A(\tilde{u}), j_0^r(dx^1 \odot dx^2) \rangle \\ &= \langle A(u), j_0^r(dx^1 \odot (dx^1 + dx^2)) \rangle \\ &= \langle A(u), j_0^r(dx^1 \odot dx^1) \rangle + \langle A(u), j_0^r(dx^1 \odot dx^2) \rangle, \end{aligned}$$

where  $\tilde{u} = T(J^r(\odot^2 T^*))^*(\varphi^{-1})(u)$ . So  $\langle A(u), j_0^r(dx^1 \odot dx^1) \rangle = 0$ .

Similarly  $\langle A(u), j_0^r(x^3 dx^1 \odot dx^1) \rangle = 0$ .  $\square$

Using Lemma 4 we see that Theorem 1 will be proved after proving the following two propositions.

**Proposition 1.** *We have*

$$\langle A(u), j_0^r(dx^1 \odot dx^2) \rangle = 0$$

for every  $u \in (T(J^r(\odot^2 T^*))^*(\mathbf{R}^n))_0$ .

**Proposition 2.** *We have*

$$\langle A(u), j_0^r(x^3 dx^1 \odot dx^2) \rangle = 0$$

for every  $u \in (T(J^r(\odot^2 T^*))^*(\mathbf{R}^n))_0$ .

### 3. SOME NOTATIONS

We have the obvious trivialization

$$(T(J^r(\odot^2 T^*))^*(\mathbf{R}^n))_0 \cong \mathbf{R}^n \times ((J^r(\odot^2 T^*))^*(\mathbf{R}^n))_0 \times ((J^r(\odot^2 T^*))^*(\mathbf{R}^n))_0$$

given by  $(u_1, u_2, u_3) \rightarrow (\tilde{u}_1)^C(u_2) + \frac{d}{dt}|_{t=0}(u_2 + tu_3)$ , where  $\tilde{u}_1$  is the constant vector field on  $\mathbf{R}^n$  such that  $\tilde{u}_1|_0 = u_1 \in \mathbf{R}^n \cong T_0\mathbf{R}^n$  and  $(\tilde{u}_1)^C$  is the complete lift of  $\tilde{u}_1$  to  $(J^r(\odot^2 T^*))^*$ .

Each  $u_\tau \in ((J^r(\odot^2 T^*))^*(\mathbf{R}^n))_0$ ,  $\tau = 2, 3$  can be expressed in the form

$$u_\tau = \sum u_{\tau,\alpha,i,j} (j_0^r(x^\alpha dx^i \odot dx^j))^*,$$

where the sum is over all  $\alpha \in (\mathbf{N} \cup \{0\})^n$ ,  $|\alpha| \leq r$ ,  $i \leq j$ ,  $i, j = 1, \dots, n$ .

It defines  $u_{\tau,\alpha,i,j}$  for each  $u_\tau$  as above.

### 4. PROOF OF PROPOSITION 1

We start with the following lemma.

**Lemma 5.** *There exists the number  $\lambda \in \mathbf{R}$  such that*

$$\langle A(u), j_0^r(dx^1 \odot dx^2) \rangle = \lambda u_{3,(0),1,2}$$

for every  $u = (u_1, u_2, u_3) \in (T(J^r(\odot^2 T^*))^*(\mathbf{R}^n))_0$ .

**Proof.** Let  $\Phi : \mathbf{R}^n \times ((J^r(\odot^2 T^*))^*(\mathbf{R}^n))_0 \times ((J^r(\odot^2 T^*))^*(\mathbf{R}^n))_0 \rightarrow \mathbf{R}$  be such that

$$\Phi(u_1, u_2, u_3) = \langle A(u), j_0^r(dx^1 \odot dx^2) \rangle,$$

where  $u = (u_1, u_2, u_3)$ ,  $u_1 = (u_1^\iota) \in \mathbf{R}^n$ ,  $\iota = 1, \dots, n$ ,  $u_2 \in ((J^r(\odot^2 T^*))^*(\mathbf{R}^n))_0$ ,  $u_3 \in ((J^r(\odot^2 T^*))^*(\mathbf{R}^n))_0$ .

The invariance of  $A$  with respect to the homotheties  $a_t = (t^1 x^1, \dots, t^n x^n)$  for  $t = (t^1, \dots, t^n) \in \mathbf{R}_+^n$  gives the homogeneous condition

$$\Phi(T(J^r(\odot^2 T^*))^*(a_t)(u)) = t^1 t^2 \Phi(u).$$

Then from the homogeneous function theorem, [5], it follows that  $\Phi(u)$  is the linear combination of monomials in  $u_1^\iota$ ,  $u_{\tau, \alpha, i, j}$  of weight  $t^1 t^2$ . Moreover  $\Phi(u_1, u_2, u_3)$  is linear in  $u_1, u_3$  for  $u_2$ , since  $A$  is linear. It implies the lemma.  $\square$

In particular from Lemma 5 it follows that

$$(*) \quad \langle A(\partial_{1|w}^C), j_0^r(dx^1 \odot dx^2) \rangle = \langle A(e_1, w, 0), j_0^r(dx^1 \odot dx^2) \rangle = 0$$

for every  $w \in ((J^r(\odot^2 T^*))^*(\mathbf{R}^n))_0$ , where  $\partial_1 = \frac{\partial}{\partial x^1}$  and  $()^C$  is the complete lift to  $(J^r(\odot^2 T^*))^*$ .

We are now in position to prove Proposition 1. Let  $\lambda$  be from Lemma 5. It is enough to prove that  $\lambda$  is equal to 0.

We see that  $\lambda = \langle A(0, 0, (j_0^r(dx^1 \odot dx^2))^*), j_0^r(dx^1 \odot dx^2) \rangle$ .

We have

$$\begin{aligned} 0 &= \langle A((x^1)^{r+1} \partial_1)|_w^C, j_0^r(dx^1 \odot dx^2) \rangle \\ (**) \quad &= (r+1) \langle A(0, w, (j_0^r(dx^1 \odot dx^2))^* + \dots), j_0^r(dx^1 \odot dx^2) \rangle \\ &= (r+1) \langle A(0, 0, (j_0^r(dx^1 \odot dx^2))^*), j_0^r(dx^1 \odot dx^2) \rangle, \end{aligned}$$

where  $w = (j_0^r((x^1)^r dx^1 \odot dx^2))^*$  and the dots is a linear combination of the  $(j_0^r(x^\alpha dx^i \odot dx^j))^*$  with  $(j_0^r(x^\alpha dx^i \odot dx^j))^* \neq (j_0^r(dx^1 \odot dx^2))^*$ .

It remains to explain (\*\*).

At first we show the second equality in (\*\*). Let  $\varphi_t$  be the flow of  $(x^1)^{r+1} \partial_1$ . We have the following sequences of equalities

$$\begin{aligned} \langle (x^1)^{r+1} \partial_1|_w^C, j_0^r(dx^1 \odot dx^2) \rangle &= \left\langle \frac{d}{dt} \Big|_{t=0} (J^r(\odot^2 T^*))_0^*(\varphi_t)(w), j_0^r(dx^1 \odot dx^2) \right\rangle \\ &= \frac{d}{dt} \Big|_{t=0} \langle (J^r(\odot^2 T^*))_0^*(\varphi_t)(w), j_0^r(dx^1 \odot dx^2) \rangle \\ &= \frac{d}{dt} \Big|_{t=0} \langle w, j_0^r((\varphi_{-t})_* dx^1 \odot dx^2) \rangle \\ &= \langle w, j_0^r\left(\frac{d}{dt} \Big|_{t=0} (\varphi_{-t})_* dx^1 \odot dx^2\right) \rangle \\ &= \langle w, j_0^r(L_{(x^1)^{r+1} \partial_1}(dx^1 \odot dx^2)) \rangle \\ &= (r+1) \langle w, j_0^r((x^1)^r dx^1 \odot dx^2) \rangle = r+1. \end{aligned}$$

Then  $((x^1)^{r+1}\partial_1)|_w^C = (r+1)(j_0^r(dx^1 \odot dx^2))^* + \dots$  under the canonical isomorphism  $V_w((J^r(\odot^2 T^*))^*(\mathbf{R}^n)) \cong ((J^r(\odot^2 T^*))^*(\mathbf{R}^n))_0$ . So we have the second equality in (\*\*).

The last equality in (\*\*) is clear because of Lemma 5.

We can prove the first equality in (\*\*) as follows. Vector fields  $\partial_1 + (x^1)^{r+1}\partial_1$  and  $\partial_1$  have the same  $r$ -jets at  $0 \in \mathbf{R}^n$ . Then, by [12], there exists a diffeomorphism  $\varphi : \mathbf{R}^n \rightarrow \mathbf{R}^n$  such that  $j_0^{r+1}\varphi = \text{id}$  and  $\varphi_*\partial_1 = \partial_1 + (x^1)^{r+1}\partial_1$  in a certain neighborhood of 0. Obviously,  $\varphi$  preserves  $j_0^r(dx^1 \odot dx^2)$  that is  $j_0^r(dx^1 \odot dx^2) = J_0^r(\odot^2 T^*)(\varphi)(j_0^r(dx^1 \odot dx^2))$  because  $j_0^{r+1}\varphi = \text{id}$ . Then, using the invariance of  $A$  with respect to  $\varphi$ , from (\*) it follows that  $\langle A(\partial_1 + (x^1)^{r+1}\partial_1)|_w^C, j_0^r(dx^1 \odot dx^2) \rangle = \langle A(\partial_1)|_w^C, j_0^r(dx^1 \odot dx^2) \rangle = 0$  for every  $w \in ((J^r(\odot^2 T^*))^*(\mathbf{R}^n))_0$ . Now, using the linearity of  $A$ , we end the proof of the first equality of (\*\*).

The proof of Proposition 1 is complete. □

### 5. PROOF OF PROPOSITION 2

The proof of Proposition 2 is similar to the proof of Proposition 1. We start with the following lemma.

**Lemma 6.** *For every  $u = (u^1, u^2, u^3) \in (T(J^r(\odot^2 T^*))^*(\mathbf{R}^n))_0$  we have*

$$\begin{aligned} \langle A(u), j_0^r(x^3 dx^1 \odot dx^2) \rangle &= au_1^1 u_{2,(0),2,3} + bu_1^2 u_{2,(0),1,3} + cu_1^3 u_{2,(0),1,2} \\ &\quad + eu_{3,e_2,2,3} + fu_{3,e_2,1,3} + gu_{3,e_3,1,2} \end{aligned}$$

where  $e_i = (0, 0, \dots, 1, 0, \dots, 0) \in (\mathbf{N} \cup \{0\})^n$ , 1 in  $i$ -position.

**Proof.** We will use the similar arguments as in the proof of Lemma 5.

Let  $\Phi : \mathbf{R}^n \times ((J^r(\odot^2 T^*))^*(\mathbf{R}^n))_0 \times ((J^r(\odot^2 T^*))^*(\mathbf{R}^n))_0 \rightarrow \mathbf{R}$  such that

$$\Phi(u_1, u_2, u_3) = \langle A(u), j_0^r(x^3 dx^1 \odot dx^2) \rangle,$$

$u = (u_1, u_2, u_3)$ ,  $u_1 = (u_1^\iota) \in \mathbf{R}^n$ ,  $\iota = 1, \dots, n$ ,  $u_2 \in ((J^r(\odot^2 T^*))^*(\mathbf{R}^n))_0$ ,  $u_3 \in ((J^r(\odot^2 T^*))^*(\mathbf{R}^n))_0$ . The invariance of  $A$  with respect to the homotheties  $a_t = (t^1 x^1, \dots, t^n x^n)$  for  $t = (t^1, \dots, t^n) \in \mathbf{R}_+^n$  gives the homogeneous condition

$$\Phi(T(J^r(\odot^2 T^*))^*(a_t)(u)) = t^1 t^2 t^3 \Phi(u).$$

Then from the homogeneous function theorem, [5], it follows that  $\Phi(u)$  is the linear combination of monomials in  $u_1^\iota, u_{\tau,\alpha,i,j}$  of weight  $t^1 t^2 t^3$ . Moreover  $\Phi(u_1, u_2, u_3)$  is linear in  $u_1$  and  $u_3$  for  $u_2$ , since  $A$  is linear. It implies the lemma. □

To prove Proposition 2 we have to show that  $a = b = c = e = f = g = 0$ . We need the following lemmas.

**Lemma 7.** *For every  $u \in (T(J^r(\odot^2 T^*))^*(\mathbf{R}^n))_0$  we have*

$$\langle A(u), j_0^r(x^3 dx^1 \odot dx^2) \rangle = -\langle A(u'), j_0^r(x^3 dx^1 \odot dx^2) \rangle,$$

where  $u'$  is the image of  $u$  by  $(x^2, x^3, x^1) \times \text{id}_{\mathbf{R}^{n-3}}$ .

**Proof.** Consider  $u \in (T(J^r(\odot^2 T^*))^*(\mathbf{R}^n))_0$ . Let  $\tilde{u}$  be the image of  $u$  by  $\varphi = (x^1 + x^1 x^3, x^2, \dots, x^n)$ . From Proposition 1 we have

$$\langle A(\tilde{u}), j_0^r(dx^1 \odot dx^2) \rangle = \langle A(u), j_0^r(dx^1 \odot dx^2) \rangle = 0.$$

Using the invariance of  $A$  with respect to  $\varphi^{-1}$  we have

$$\begin{aligned} 0 &= \langle A(u), j_0^r(dx^1 \odot dx^2) \rangle \\ &= \langle A(u), j_0^r(x^3 dx^1 \odot dx^2) \rangle + \langle A(u), j_0^r(x^1 dx^2 \odot dx^3) \rangle \end{aligned}$$

because  $\varphi^{-1}$  preserves  $A$ , it transforms  $\tilde{u}$  in  $u$  and  $j_0^r(dx^1 \odot dx^2)$  in  $j_0^r(dx^1 \odot dx^2) + j_0^r(x^3 dx^1 \odot dx^2) + j_0^r(x^1 dx^2 \odot dx^3)$ . So,  $\langle A(u), j_0^r(x^3 dx^1 \odot dx^2) \rangle = -\langle A(u), j_0^r(x^1 dx^2 \odot dx^3) \rangle$ . Hence we have the lemma because  $(x^2, x^3, x^1) \times \mathbf{R}^{n-3}$  sends  $u$  in  $u'$  and  $j_0^r(x^1 dx^2 \odot dx^3)$  in  $j_0^r(x^3 dx^1 \odot dx^2)$ .  $\square$

**Lemma 8.** *We have  $g = f = e = 0$ .*

**Proof.** Obviously

$$g = \langle A(0, 0, (j_0^r(x^3 dx^1 \odot dx^2))^*), j_0^r(x^3 dx^1 \odot dx^2) \rangle$$

by Lemma 6. Similarly

$$\begin{aligned} f &= \langle A(0, 0, (j_0^r(x^2 dx^1 \odot dx^3))^*), j_0^r(x^3 dx^1 \odot dx^2) \rangle, \\ e &= \langle A(0, 0, (j_0^r(x^1 dx^2 \odot dx^3))^*), j_0^r(x^3 dx^1 \odot dx^2) \rangle. \end{aligned}$$

So, to prove Lemma 8 we have to show

$$\begin{aligned} &\langle A(0, 0, (j_0^r(x^3 dx^1 \odot dx^2))^*), j_0^r(x^3 dx^1 \odot dx^2) \rangle \\ &= \langle A(0, 0, (j_0^r(x^2 dx^1 \odot dx^3))^*), j_0^r(x^3 dx^1 \odot dx^2) \rangle \\ &= \langle A(0, 0, (j_0^r(x^1 dx^2 \odot dx^3))^*), j_0^r(x^3 dx^1 \odot dx^2) \rangle = 0. \end{aligned}$$

We can see that  $(x^2, x^3, x^1) \times \text{id}_{\mathbf{R}^{n-3}}$  sends  $(j_0^r(x^3 dx^1 \odot dx^2))^*$  in  $(j_0^r(x^2 dx^1 \odot dx^3))^*$  and  $(j_0^r(x^2 dx^1 \odot dx^3))^*$  in  $(j_0^r(x^1 dx^2 \odot dx^3))^*$ . Then using Lemma 7 it is enough to verify that  $\langle A(0, 0, (j_0^r(x^3 dx^1 \odot dx^2))^*), j_0^r(x^3 dx^1 \odot dx^2) \rangle = 0$ . So, it is enough to prove the sequence of equalities:

$$\begin{aligned} 0 &= \langle A((x^1)^r \partial_1)|_w^C, j_0^r(x^3 dx^1 \odot dx^2) \rangle \\ (***) &= r \langle A(0, w, (j_0^r(x^3 dx^1 \odot dx^2))^*), j_0^r(x^3 dx^1 \odot dx^2) \rangle \\ &= r \langle A(0, 0, (j_0^r(x^3 dx^1 \odot dx^2))^*), j_0^r(x^3 dx^1 \odot dx^2) \rangle, \end{aligned}$$

where  $w = (j_0^r(x^3(x^1)^{r-1} dx^1 \odot dx^2))^* \in ((J^r(\odot^2 T^*))^*(\mathbf{R}^n))_0$ .

The third equality in (\*\*\*) is clear on the basis of Lemma 6.

Let us explain the first equality in (\*\*\*). Vector fields  $\partial_1 + (x^1)^r \partial_1$  and  $\partial_1$  have the same  $(r-1)$ -jets at  $0 \in \mathbf{R}^n$ . Then, by [12] there exist a diffeomorphism  $\varphi = \varphi_1 \times \text{id}_{\mathbf{R}^{n-1}} : \mathbf{R}^n = \mathbf{R} \times \mathbf{R}^{n-1} \rightarrow \mathbf{R}^n = \mathbf{R} \times \mathbf{R}^{n-1}$  such that  $\varphi_1 : \mathbf{R} \rightarrow \mathbf{R}$ ,  $j_0^r \varphi = \text{id}$  and  $\varphi_* \partial_1 = \partial_1 + (x^1)^r \partial_1$  in a certain neighborhood of  $0 \in \mathbf{R}^n$ . Let  $\varphi^{-1}$  sends  $\omega$  in  $\tilde{\omega}$ . Then  $\tilde{\omega}$  is a linear combination of the elements  $(j_0^r(x^\alpha dx^i \odot dx^j))^* \in ((J^r(\odot^2 T^*))^*(\mathbf{R}^n))_0$  for  $r \geq |\alpha| \geq 1$ ,  $i, j = 1, \dots, n$ ,  $i \leq j$ . (For  $\langle \tilde{\omega}, j_0^r(dx^i \odot dx^j) \rangle = \langle \omega, j_0^r(d(x^i \circ \varphi^{-1}) \odot d(x^j \circ \varphi^{-1})) \rangle = 0$ .) Then, by Lemma 6,  $\langle A(\partial_1|_{\tilde{\omega}}^C), j_0^r(x^3 dx^1 \odot dx^2) \rangle = \langle A(e_1, \tilde{\omega}, 0), j_0^r(x^3 dx^1 \odot dx^2) \rangle$ .

$dx^2\rangle = 0$  (as  $j_0^r\varphi = \text{id}$ ). Then from naturality of  $A$  with respect to  $\varphi$  we obtain  $\langle A((\partial_1 + (x^1)^r\partial_1)|_\omega^C), j_0^r(x^3 dx^1 \odot dx^2)) \rangle = 0$ . Now, using the linearity of  $A$  we have  $\langle A(((x^1)^r\partial_1)|_\omega^C), j_0^r(x^3 dx^1 \odot dx^2)) \rangle = 0$ . This ends the proof of the first equality in (\*\*\*)).

Let us explain the second equality in (\*\*). Analysing the flow of vector field  $(x^1)^r\partial_1$  and taking  $\omega = (j_0^r(x^3(x^1)^{r-1} dx^1 \odot dx^2))^* \in ((J^r(\odot^2 T^*))^*(\mathbf{R}^n))_0$  we have (similarly as in the justification of the second equality of (\*\*))

$$\begin{aligned} \langle ((x^1)^r\partial_1)|_\omega^C, j_0^r(\alpha dx^i \odot dx^j) \rangle &= \langle \omega, j_0^r(L_{(x^1)^r\partial_1}(x^\alpha dx^i \odot dx^j)) \rangle \\ &= \langle \omega, \alpha_1 j_0^r((x^1)^{r-1} x^\alpha dx^i \odot dx^j) \rangle \\ &\quad + \langle \omega, j_0^r(x^\alpha \delta_1^i r (x^1)^{r-1} dx^1 \odot dx^j) \rangle, \end{aligned}$$

where  $\delta_1^i$  is the Kronecker delta.

Since  $\omega = (j_0^r(x^3(x^1)^{r-1} dx^1 \odot dx^2))^*$  the last sum is equal to  $r$  if  $\alpha = e_3$  and  $(i, j) = (1, 2)$ , and 0 in the other cases. Then  $(x^1)^r\partial_1|_\omega^C = r(j_0^r(x^3 dx^1 \odot dx^2))^*$ .

This ends the proof of the second equality of (\*\*). □

The proof of Lemma 8 is complete. □

**Lemma 9.** *We have  $a = b = c = 0$ .*

**Proof.** Using Lemma 7 (similarly as for  $g = f = e$ ) it is sufficient to prove that  $c = 0$ , i.e.  $\langle A(\partial_3|_{(j_0^r(dx^1 \odot dx^2))^*}, j_0^r(x^3 dx^1 \odot dx^2)) \rangle = 0$ .

But we have

$$\begin{aligned} 0 &= \langle A(\partial_3|_{(j_0^r((x^1)^r dx^1 \odot dx^2))^*}, j_0^r(x^3 dx^1 \odot dx^2)) \rangle \\ (***) &= \langle A(\partial_3|_{(j_0^r(dx^1 \odot dx^2))^* + \dots}, j_0^r(x^3 dx^1 \odot dx^2)) \rangle \\ &= \langle A(\partial_3|_{(j_0^r(dx^1 \odot dx^2))^*}, j_0^r(x^3 dx^1 \odot dx^2)) \rangle, \end{aligned}$$

where the dots is the linear combination of elements  $(j_0^r(x^\alpha dx^i \odot dx^j))^* \neq (j_0^r(dx^1 \odot dx^2))^*$ ,  $\alpha \in (\mathbf{N} \cup \{0\})^n$ ,  $|\alpha| \leq r$ ,  $i \leq j$ ,  $i, j = 1, \dots, n$ .

Equalities first and third are clear because of Lemma 6.

Let us explain the second equality. Consider the local diffeomorphism  $\varphi = (x^1 + \frac{1}{r+1}(x^1)^{r+1}, x^2, \dots, x^n)^{-1}$ . We see that  $\varphi^{-1}$  preserves  $j_0^r(x^3 dx^1 \odot dx^2)$  and  $\partial_3$ . Moreover  $\varphi^{-1}$  sends  $(j_0^r((x^1)^r dx^1 \odot dx^2))^*$  in  $(j_0^r(dx^1 \odot dx^2))^* + \dots$ , where the dots is as above. Now, by the invariance of  $A$  with respect to  $\varphi^{-1}$  we get the second equality in(\*\*\*\*).

The proof of Lemma 9 is complete. □

The proof of Proposition 2 is complete. □

The proof of Theorem 1 is complete. □

## 7. THE NATURAL AFFINORS ON $(J^r(\odot^2 T^*))^*$ OF VERTICAL TYPE

A natural affnor  $Q : T(J^r(\odot^2 T^*))^* \rightarrow T(J^r(\odot^2 T^*))^*$  on  $(J^r(\odot^2 T^*))^*$  is of *vertical type* if the image of  $Q$  is in the vertical space  $V(J^r(\odot^2 T^*))^*(M)$  for every  $n$ -manifolds  $M$ .



We have the natural isomorphism

$$V(J^r(\odot^2 T^*))^*(M) \cong (J^r(\odot^2 T^*))^*(M) \times (J^r(\odot^2 T^*))^*(M)$$

given by  $(u, w) = \frac{d}{dt}|_{t=0}(u + tv)$ ,  $u, v \in (J^r(\odot^2 T^*))^*_x(M)$ ,  $x \in M$ , and the natural projection  $pr_2 : V(J^r(\odot^2 T^*))^*M \rightarrow (J^r(\odot^2 T^*))^*M$  on the second factor.

Let  $Q : T(J^r(\odot^2 T^*))^* \rightarrow T(J^r(\odot^2 T^*))^*$  on  $(J^r(\odot^2 T^*))^*$  be a natural affinator of vertical type. Composing  $Q$  with  $pr_2$  we get a natural linear transformation  $pr_2 \circ Q : T(J^r(\odot^2 T^*))^* \rightarrow (J^r(\odot^2 T^*))^*$  over  $n$ -manifolds. It is equal to 0 because of Theorem 1. So, we have the following corollary.

**Corollary 1.** *Let  $n \geq 3$ ,  $r$  be natural numbers. Every natural affinator  $Q$  of vertical type on  $(J^r(\odot^2 T^*))^*$  over  $n$ -manifolds is equal to 0.*

### 8. THE LINEAR NATURAL TRANSFORMATIONS $T(J^r(\odot^2 T^*))^* \rightarrow T$

Let  $\pi$  be the projection of natural bundle  $(J^r(\odot^2 T^*))^*$ . Then the tangent map  $T\pi_M : T(J^r(\odot^2 T^*))^*(M) \rightarrow TM$  defines a linear natural transformation  $T\pi : T(J^r(\odot^2 T^*))^* \rightarrow T$ . ( The definition of a linear natural transformation  $T(J^r(\odot^2 T^*))^* \rightarrow T$  over  $n$ -manifolds is similar to the one in Section 1.)

**Theorem 2.** *Let  $n$  and  $r$  be natural numbers. Every linear natural transformation  $B : T(J^r(\odot^2 T^*))^* \rightarrow T$  over  $n$ -manifolds is proportional to  $T\pi$ .*

### 9. PROOF OF THEOREM 2

Consider a linear natural transformation  $B : T(J^r(\odot^2 T^*))^* \rightarrow T$ . We have

**Lemma 10.** *If  $\langle B(u), d_0x^1 \rangle = 0$  for every  $u \in (T(J^r(\odot^2 T^*))^*(\mathbf{R}^n))_0$  then  $B = 0$ .*

**Proof.** The proof of Lemma 10 is similar to the proofs of Lemmas 1 – 4. From the invariance of  $B$  with respect to the coordinate permutation we see that  $\langle B(u), d_0x^i \rangle = 0$  for  $i = 1, \dots, n$  and  $u \in (T(J^r(\odot^2 T^*))^*(\mathbf{R}^n))_0$ . So  $B(u) = 0$  for every  $u \in (T(J^r(\odot^2 T^*))^*(\mathbf{R}^n))_0$ . Then using the invariance of  $B$  with respect to the charts we obtain that  $B = 0$ . □

**Lemma 11.** *We have  $\langle B(u), d_0x^1 \rangle = \lambda u_1^1$  for some  $\lambda \in \mathbf{R}$ , where  $u = (u_1, u_2, u_3)$ ,  $u_1 = (u_1^i) \in \mathbf{R}^n$ ,  $i = 1, \dots, n$ , and  $u_2, u_3 \in ((J^r(\odot^2 T^*))^*(\mathbf{R}^n))_0$ .*

**Proof.** The proof of Lemma 11 is similar to the proof of Lemma 5. □

Lemma 11 shows that  $\langle (B - \lambda T\pi)(u), d_0x^1 \rangle = 0$  for every  $u \in (T(J^r(\odot^2 T^*))^*(\mathbf{R}^n))_0$ . Then  $B - \lambda T\pi = 0$  by Lemma 10, i.e.  $B = \lambda T\pi$ .

The proof of Theorem 2 is complete. □

## 10. THE MAIN RESULT

The main result of the present paper is the following theorem.

**Theorem 3.** *Let  $n \geq 3$  and  $r$  be natural numbers. Every natural affinor  $Q : T(J^r(\odot^2 T^*))^* \rightarrow T(J^r(\odot^2 T^*))^*$  on  $(J^r(\odot^2 T^*))^*$  over  $n$ -manifolds is proportional to the identity affinor.*

**Proof.** The composition  $T\pi \circ Q : T(J^r(\odot^2 T^*))^* \rightarrow T$  is a linear natural transformation. Hence, by Theorem 2,  $T\pi \circ Q = \lambda T\pi$  for some  $\lambda \in \mathbf{R}$ . Then  $Q - \lambda \text{id} : T(J^r(\odot^2 T^*))^* \rightarrow T(J^r(\odot^2 T^*))^*$  is a natural affinor of vertical type, because  $T\pi \circ (Q - \lambda \text{id}) = T\pi \circ Q - \lambda T\pi = 0$ . From Corollary 1 we obtain that  $Q - \lambda \text{id} = 0$ . Thus  $Q = \lambda \text{id}$ . The proof of Theorem 3 is complete.  $\square$

## REFERENCES

- [1] Doupovec, M., Kolář, I., *Natural affinors on time-dependent Weil bundles*, Arch. Math. (Brno) 27 (1991), 205-209.
- [2] Doupovec, M., Kurek, J., *Torsions of connections of higher order cotangent bundles*, Czech. Math. J. (to appear).
- [3] Gancarzewicz, J., Kolář, I., *Natural affinors on the extended  $r$ -th order tangent bundles*, Suppl. Rendiconti Circolo Mat. Palermo, 1993, 95-100.
- [4] Kolář, I., Modugno, M., *Torsion of connections on some natural bundles*, Diff. Geom. and Appl. 2(1992), 1-16.
- [5] Kolář, I., Michor, P. W., Slovák, J., *Natural Operations in Differential Geometry*, Springer-Verlag, Berlin 1993.
- [6] Kurek, J., *Natural affinors on higher order cotangent bundles*, Arch. Math. (Brno) 28 (1992), 175-180.
- [7] Mikulski, W. M., *The natural affinors on dual  $r$ -jet prolongation of bundles of 2-forms*, Ann. UMCS Lublin 2002, (to appear).
- [8] Mikulski, W. M., *Natural affinors on  $r$ -jet prolongation of the tangent bundle*, Arch. Math. (Brno) 34 (2) (1998). 321-328.
- [9] Mikulski, W. M., *The natural affinors on  $\otimes^k T^{(k)}$* , Note di Matematica vol. 19-n. 2. (1999), 269-274.
- [10] Mikulski, W. M., *The natural affinors on generalized higher order tangent bundles*, Rend. Mat. Roma vol. 21. (2001). (to appear).
- [11] Mikulski, W. M., *Natural affinors on  $(J^{r,s,q}(\cdot, \mathbf{R}^{1,1})_0)^*$* , Coment. Math. Carolinae 42 (2001), (to appear).
- [12] Zajtz, A., *On the order of natural operators and liftings*, Ann. Polon. Math. 49 (1988), 169-178.

INSTITUTE OF MATHEMATICS, CRACOW UNIVERSITY OF TECHNOLOGY  
 31-155 KRAKÓW, UL. WARSZAWSKA 24, POLAND  
*E-mail:* pmichale@usk.pk.edu.pl