

**ON LOCALLY LIPSCHITZ LOCALLY COMPACT
TRANSFORMATION GROUPS OF MANIFOLDS**

GEORGE MICHAEL, A. A.

ABSTRACT. In this paper we show that a “locally Lipschitz” locally compact transformation group acting continuously and effectively on a connected paracompact locally Euclidean topological manifold is a Lie group. This is a contribution to the proof of the Hilbert-Smith conjecture. It generalizes the classical Bochner-Montgomery-Kuranishi Theorem [1, 9] and also the Repovš-Ščepin Theorem [17] which holds only for Riemannian manifolds.

A topological group G acting continuously and effectively (i.e. every non-trivial element of G acts non-trivially) on an n -dimensional topological manifold M is said to be locally Lipschitz if $\forall a \in M, \exists (U_a, \varphi_a)$ chart of M where $a \in U_a$ open $\subseteq M$, $\varphi_a: U_a \rightarrow \mathbf{R}^n$ an open embedding and $1 \in V$ open $\subseteq G$, $a \in U$ open $\subseteq U_a$ such that $V(U) \subseteq U_a$ and $\forall g \in V, d(gx, gy) \leq c_g d(x, y) \forall x, y \in U$ for some $c_g \in \mathbf{R}$ where d is the transported Euclidean distance from \mathbf{R}^n via the map φ_a . This definition generalizes the corresponding one given in [8].

In this paper we generalize the results of [8] and we show that a locally Lipschitz locally compact transformation group acting continuously and effectively on a connected paracompact locally Euclidean manifold is a Lie group, Theorem 2. This provides a contribution to the proof of the Hilbert-Smith conjecture. It generalizes the classical Bochner-Montgomery-Kuranishi Theorem [1, 9] where the group acts by C^1 diffeomorphisms on a smooth manifold and also the Repovš-Ščepin Theorem [17] which holds only for Riemannian manifolds.

The following theorem is all what we need to establish our result. It replaces the Hausdorff dimension argument in [17] by an elementary theorem in dimension theorem from Nagata book [16, p.83]. This will enable us to obtain the required generalization from Riemannian manifolds to topological manifolds.

2000 *Mathematics Subject Classification*: Primary 57S05, Secondary 54H15.

Key words and phrases: locally Lipschitz transformation group, Hilbert-Smith conjecture.

Received May 2, 2006, revised March, 2007.

Theorem 1. *Let G be a zero-dimensional compact topological group acting continuously on a paracompact n -dimensional topological manifold M such that G is locally Lipschitz then $\dim M/G = \dim M = n$ ($\dim =$ covering dimension).*

Proof. Let $a \in M$ and let (U_a, φ_a) chart of M where $a \in U_a$ open $\subseteq M$, $\varphi_a: U_a \rightarrow \mathbf{R}^n$ an open embedding. Let $a \in U$ open $\subseteq U^-$ compact $\subseteq U_a$ and H open normal subgroup of G such that $H(U^-) \subseteq U_a$ [1, TGIII.36] such that $\forall g \in H$, $d(gx, gy) \leq c_g d(x, y) \forall x, y \in H(U^-)$ for some $c_g \in \mathbf{R}$, where d is the transported Euclidean distance from \mathbf{R}^n via the map φ_a .

Since $H = \cup_{k \geq 1} \{g \in H : d(gx, gy) \leq k d(x, y) \forall x, y \in H(U^-)\}$, we may assume by a Baire category argument that $c_g = c \forall g \in H$.

Let β be the invariant Haar measure on H of total mass 1, then

$$d^-(x, y) = \int_H d(gx, gy) d\beta(g), \quad \forall x, y \in H(U^-)$$

defines an equivalent H -invariant metric on $H(U^-)$ and if $\pi_1: M \rightarrow M/H$ is the canonical map, then $d^*(\pi_1 x, \pi_1 y) = d^-(Hx, Hy)$ defines an admissible metric on $H(U^-)/H$.

Note that if $A \subseteq H(U^-)$, then $\text{diam}(\pi_1, d^*) \leq \text{diam}(A, d^-) \leq c \text{diam}(A, d)$ ($\text{diam}(B, \delta) =$ diameter of set B under the metric δ) so that if

$$m_n(A, d) = \sup_{\varepsilon > 0} \inf \left\{ \sum_{i \geq 1} (\text{diam}(A_i, d))^n : A = \bigcup_{i=1}^n A_i, \text{diam}(A_i, d) < \varepsilon, \forall i \geq 1 \right\}$$

[16, p.81], and if we identify $H(U^-)$ by its image in \mathbf{R}^n under φ_a then $m_n(A, d) = \left(\frac{4}{\pi}\right)^{\frac{n}{2}} \Gamma\left(1 + \frac{n}{2}\right) \lambda_n^*(A)$ where λ_n^* (resp. λ_n) is the Lebesgue outer measure (resp. measure) on \mathbf{R}^n [6, p.157].

We have $m_{n+1}(H(U^-), d) = 0$ since $\lambda_{n+1}(H(U^-)) = 0$ and

$$m_{n+1}(H(U^-)/H, d^*) = 0$$

so that $\dim H(U^-)/H = \dim \pi_1(U^-) \leq n$ [16, p.83].

Now, if $\pi: M \rightarrow M/G$ and $\pi_2: M/H \rightarrow (M/H)/(G/H)$ are the canonical maps, then

$$\begin{aligned} \dim \pi(U^-) &= \dim \pi_2(G/H\pi_1(U^-)) && [2, \text{TG III.15}] \\ &= \dim G/H\pi_1(U^-) && [13, \text{Theorem 4.1}] \\ &= \dim \pi_1(U^-) && [15, \text{p.53}] \\ &\leq n \end{aligned}$$

hence $n = \dim M \leq \dim M/G \leq n$ [15, p.62+p.130]. \square

Now we can establish our main theorem which, in view of [10, Lemma 2.1], generalizes Repovš-Ščepin Theorem [17] which holds only for Riemannian manifolds.

Theorem 2 (Main Theorem). *Let G be a locally compact topological group acting continuously and effectively on a connected paracompact n -dimensional topological manifold M such that G is locally Lipschitz, then G is a Lie group.*

Proof. Note that G is metrizable [2, TG III 36 + TG III 76] and $\dim G < \infty$ [11], so that by the main approximation theorem [12, p.175] if G is not a Lie group, then G contains a zero-dimensional torsion free compact group by [2, TG III 60], [15, p.161], [14, Theorem 2.1] and [5, p.206] hence it contains the p -adic group $\mathbf{Z}_p = \varprojlim \mathbf{Z}/p^n \mathbf{Z}$ for some prime number p [7, p.426] and

$$\begin{aligned} n &= \dim M = \dim M/\mathbf{Z}_p && \text{by Theorem 1} \\ &= \dim_{\mathbf{Z}} M/\mathbf{Z}_p && [15, \text{p.206, p.210}] \\ &= n + 2 && [3, 18] \end{aligned}$$

($\dim_{\mathbf{Z}}$ = cohomological dimension over \mathbf{Z}) which is absurd. □

Corollary 1 (Bochner-Montgomery-Kuranishi [1, 9]). *Let G be a locally compact topological group acting continuously and effectively on a connected paracompact n -dimensional C^1 manifold M such that G acts on M by C^1 diffeomorphisms, then G is a Lie group.*

Proof. Clearly by the mean value theorem [4, p.155] G is locally Lipschitz, hence our assertion follows from Theorem 2. □

REFERENCES

- [1] Bochner, S., Montgomery, D., *Locally compact groups of differentiable transformations*, Ann. of Math. (2) **47** (1946), 639–653.
- [2] Bourbaki, N., *Topologie générale, Chap. 1-4*, Hermann, Paris 1971.
- [3] Bredon, G. E., Raymond, F., Williams, R. F., *p -Adic transformation groups*, Trans. Amer. Math. Soc. **99** (1961), 488–498.
- [4] Dieudonné, J., *Foundations of modern analysis*, Academic Press, New York–London 1960.
- [5] Dress, A., *Newman’s theorems on transformation groups*, Topology, **8** (1969), 203–207.
- [6] Federer, H., *Geometric measure theory*, Springer-Verlag, Berlin–Heidelberg–New York, N.Y., 1969.
- [7] Hofmann, K. H., Morris, S. A., *The structure of compact groups*, de Gruyter Stud. Math. **25** (1998).
- [8] Karube, T., *Transformation groups satisfying some local metric conditions*, J. Math. Soc. Japan **18**, No. 1 (1966), 45–50.
- [9] Kuranishi, M., *On conditions of differentiability of locally compact groups*, Nagoya Math. J. **1** (1950), 71–81.
- [10] Michael, G., *On the smoothing problem*, Tsukuba J. Math. **25**, No. 1 (2001), 13–45.
- [11] Montgomery, D., *Finite dimensionality of certain transformation groups*, Illinois J. Math. **1** (1957), 28–35.
- [12] Montgomery, D., Zippin, L., *Topological transformation groups*, Interscience Publishers, New York, 1955.
- [13] Nagami, K. R., *Mappings of finite order and dimension theory*, Japan J. Math. **30** (1960), 25–54.
- [14] Nagami, K. R., *Dimension-theoretical structure of locally compact groups*, J. Math. Soc. Japan **14**, No. 4 (1962), 379–396.
- [15] Nagami, K. R., *Dimension theory*, Academic Press, New York, 1970.

- [16] Nagata, J., *Modern dimension theory* Sigma Ser. Pure Math. **2** (1983).
- [17] Repovš, D., Ščepin, E. V., *A proof of the Hilbert-Smith conjecture for actions by Lipschitz maps*, Math. Ann. **308** (1997), 361–364.
- [18] Yang, C. T., *p -adic transformation groups*, Michigan Math. J. **7** (1960), 201–218.

DEPARTMENT OF MATHEMATICS, VOORHEES COLLEGE
DENMARK, SC 29042, U.S.A.
E-mail: adelgeorge1@yahoo.com