

STABLY CAYLEY SEMISIMPLE GROUPS

MIKHAIL BOROVoi AND BORIS KUNYAVSKIĬ

Received: July 9, 2014

ABSTRACT. A linear algebraic group G over a field k is called a Cayley group if it admits a Cayley map, i.e., a G -equivariant birational isomorphism over k between the group variety G and its Lie algebra $\mathrm{Lie}(G)$. A prototypical example is the classical ‘‘Cayley transform’’ for the special orthogonal group \mathbf{SO}_n defined by Arthur Cayley in 1846. A linear algebraic group G is called stably Cayley if $G \times S$ is Cayley for some split k -torus S . We classify stably Cayley semisimple groups over an arbitrary field k of characteristic 0.

2010 Mathematics Subject Classification: 20G15, 20C10.

Keywords and Phrases: Linear algebraic group, stably Cayley group, quasi-permutation lattice.

To Alexander Merkurjev on the occasion of his 60th birthday

0 INTRODUCTION

Let k be a field of characteristic 0 and \bar{k} a fixed algebraic closure of k . Let G be a connected linear algebraic k -group. A birational isomorphism $\phi: G \xrightarrow{\sim} \mathrm{Lie}(G)$ is called a *Cayley map* if it is equivariant with respect to the conjugation action of G on itself and the adjoint action of G on its Lie algebra $\mathrm{Lie}(G)$, respectively. A linear algebraic k -group G is called *Cayley* if it admits a Cayley map, and *stably Cayley* if $G \times_k (\mathbb{G}_{m,k})^r$ is Cayley for some $r \geq 0$. Here $\mathbb{G}_{m,k}$ denotes the multiplicative group over k . These notions were introduced by Lemire, Popov and Reichstein [LPR]; for a more detailed discussion and numerous classical examples we refer the reader to [LPR, Introduction]. The main results of [LPR] are the classifications of Cayley and stably Cayley *simple* groups over an algebraically closed field k of characteristic 0. Over an arbitrary field k of characteristic 0 stably Cayley *simple* k -groups, stably Cayley *simply connected* semisimple k -groups and stably Cayley *adjoint* semisimple k -groups were classified in the paper [BKLR] of Borovoi, Kunyavskiĭ, Lemire and Reichstein. In

the present paper, building on results of [LPR] and [BKLR], we classify all stably Cayley *semisimple* k -groups (not necessarily simple, or simply connected, or adjoint) over an arbitrary field k of characteristic 0.

By a semisimple (or reductive) k -group we always mean a *connected* semisimple (or reductive) k -group. We shall need the following result of [BKLR] extending [LPR, Theorem 1.28].

THEOREM 0.1 ([BKLR, Theorem 1.4]). *Let k be a field of characteristic 0 and G an absolutely simple k -group. Then the following conditions are equivalent:*

- (a) G is stably Cayley over k ;
- (b) G is an arbitrary k -form of one of the following groups:

$$\mathbf{SL}_3, \mathbf{PGL}_2, \mathbf{PGL}_{2n+1} \ (n \geq 1), \mathbf{SO}_n \ (n \geq 5), \mathbf{Sp}_{2n} \ (n \geq 1), \mathbf{G}_2,$$

or an inner k -form of \mathbf{PGL}_{2n} ($n \geq 2$).

In this paper we classify stably Cayley semisimple groups over an *algebraically closed* field k of characteristic 0 (Theorem 0.2) and, more generally, over an *arbitrary* field k of characteristic 0 (Theorem 0.3). Note that Theorem 0.2 was conjectured in [BKLR, Remark 9.3].

THEOREM 0.2. *Let k be an algebraically closed field of characteristic 0 and G a semisimple k -group. Then G is stably Cayley if and only if G decomposes into a direct product $G_1 \times_k \cdots \times_k G_s$ of its normal subgroups, where each G_i ($i = 1, \dots, s$) either is a stably Cayley simple k -group (i.e., isomorphic to one of the groups listed in Theorem 0.1) or is isomorphic to the stably Cayley semisimple k -group \mathbf{SO}_4 .*

THEOREM 0.3. *Let G be a semisimple k -group over a field k of characteristic 0 (not necessarily algebraically closed). Then G is stably Cayley over k if and only if G decomposes into a direct product $G_1 \times_k \cdots \times_k G_s$ of its normal k -subgroups, where each G_i ($i = 1, \dots, s$) is isomorphic to the Weil restriction $R_{l_i/k} G_{i,l_i}$ for some finite field extension l_i/k , and each G_{i,l_i} is either a stably Cayley absolutely simple group over l_i (i.e., one of the groups listed in Theorem 0.1) or an l_i -form of the semisimple group \mathbf{SO}_4 (which is always stably Cayley, but is not absolutely simple and can be not l_i -simple).*

Note that the “if” assertions in Theorems 0.2 and 0.3 follow immediately from the definitions.

The rest of the paper is structured as follows. In Section 1 we recall the definition of a quasi-permutation lattice and state some known results, in particular, an assertion from [LPR, Theorem 1.27] that reduces Theorem 0.2 to an assertion on lattices. In Sections 2 and 3 we construct certain families of non-quasi-permutation lattices. In particular, we correct an inaccuracy in [BKLR]; see Remark 2.5. In Section 4 we prove (in the language of lattices) Theorem

0.2 in the special case when G is isogenous to a direct product of simple groups of type \mathbf{A}_{n-1} with $n \geq 3$. In Section 5 we prove (again in the language of lattices) Theorem 0.2 in the general case. In Section 6 we deduce Theorem 0.3 from Theorem 0.2. In Appendix A we prove in terms of lattices only, that certain quasi-permutation lattices are indeed quasi-permutation.

1 PRELIMINARIES ON QUASI-PERMUTATION GROUPS AND ON CHARACTER LATTICES

In this section we gather definitions and known results concerning quasi-permutation lattices, quasi-invertible lattices and character lattices that we need for the proofs of Theorems 0.2 and 0.3. For details see [BKLR, Sections 2 and 10] and [LPR, Introduction].

1.1. By a *lattice* we mean a pair (Γ, L) where Γ is a finite group acting on a finitely generated free abelian group L . We say also that L is a Γ -lattice. A Γ -lattice L is called a *permutation* lattice if it has a \mathbb{Z} -basis permuted by Γ . Following Colliot-Thélène and Sansuc [CTS], we say that two Γ -lattices L and L' are *equivalent*, and write $L \sim L'$, if there exist short exact sequences

$$0 \rightarrow L \rightarrow E \rightarrow P \rightarrow 0 \quad \text{and} \quad 0 \rightarrow L' \rightarrow E \rightarrow P' \rightarrow 0$$

with the same Γ -lattice E , where P and P' are permutation Γ -lattices. For a proof that this is indeed an equivalence relation see [CTS, Lemma 8, p. 182] or [Sw, Section 8]. Note that if there exists a short exact sequence of Γ -lattices

$$0 \rightarrow L \rightarrow L' \rightarrow Q \rightarrow 0$$

where Q is a permutation Γ -lattice, then, taking in account the trivial short exact sequence

$$0 \rightarrow L' \rightarrow L' \rightarrow 0 \rightarrow 0,$$

we obtain that $L \sim L'$. If Γ -lattices L, L', M, M' satisfy $L \sim L'$ and $M \sim M'$, then clearly $L \oplus M \sim L' \oplus M'$.

Definition 1.2. A Γ -lattice L is called a *quasi-permutation* lattice if there exists a short exact sequence

$$0 \rightarrow L \rightarrow P \rightarrow P' \rightarrow 0, \tag{1.1}$$

where both P and P' are permutation Γ -lattices.

LEMMA 1.3 (well-known). *A Γ -lattice L is quasi-permutation if and only if $L \sim 0$.*

Proof. If L is quasi-permutation, then sequence (1.1) together with the trivial short exact sequence

$$0 \rightarrow 0 \rightarrow P \rightarrow P \rightarrow 0$$

shows that $L \sim 0$. Conversely, if $L \sim 0$, then there are short exact sequences

$$0 \rightarrow L \rightarrow E \rightarrow P \rightarrow 0 \quad \text{and} \quad 0 \rightarrow 0 \rightarrow E \rightarrow P' \rightarrow 0,$$

where P and P' are permutation lattices. From the second exact sequence we have $E \cong P'$, hence E is a permutation lattice, and then the first exact sequence shows that L is a quasi-permutation lattice. \square

Definition 1.4. A Γ -lattice L is called *quasi-invertible* if it is a direct summand of a quasi-permutation Γ -lattice.

Note that if a Γ -lattice L is not quasi-invertible, then it is not quasi-permutation.

LEMMA 1.5 (well-known). *If a Γ -lattice L is quasi-permutation (resp., quasi-invertible) and $L' \sim L$, then L' is quasi-permutation (resp., quasi-invertible) as well.*

Proof. If L is quasi-permutation, then using Lemma 1.3 we see that $L' \sim L \sim 0$, hence L' is quasi-permutation. If L is quasi-invertible, then $L \oplus M$ is quasi-permutation for some Γ -lattice M , and by Lemma 1.3 we have $L \oplus M \sim 0$. We see that $L' \oplus M \sim L \oplus M \sim 0$, and by Lemma 1.3 we obtain that $L' \oplus M$ is quasi-permutation, hence L' is quasi-invertible. \square

Let $\mathbb{Z}[\Gamma]$ denote the group ring of a finite group Γ . We define the Γ -lattice J_Γ by the exact sequence

$$0 \rightarrow \mathbb{Z} \xrightarrow{N} \mathbb{Z}[\Gamma] \rightarrow J_\Gamma \rightarrow 0,$$

where N is the norm map, see [BKLR, before Lemma 10.4]. We refer to [BKLR, Proposition 10.6] for a proof of the following result, due to Voskresenskii [Vo1, Corollary of Theorem 7]:

PROPOSITION 1.6. *Let $\Gamma = \mathbb{Z}/p\mathbb{Z} \times \mathbb{Z}/p\mathbb{Z}$, where p is a prime. Then the Γ -lattice J_Γ is not quasi-invertible.*

Note that if $\Gamma = \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$, then $\text{rank } J_\Gamma = 3$.

We shall use the following lemma from [BKLR]:

LEMMA 1.7 ([BKLR, Lemma 2.8]). *Let W_1, \dots, W_m be finite groups. For each $i = 1, \dots, m$, let V_i be a finite-dimensional \mathbb{Q} -representation of W_i . Set $V := V_1 \oplus \dots \oplus V_m$. Suppose $L \subset V$ is a free abelian subgroup, invariant under $W := W_1 \times \dots \times W_m$. If L is a quasi-permutation W -lattice, then for each $i = 1, \dots, m$ the intersection $L_i := L \cap V_i$ is a quasi-permutation W_i -lattice.*

We shall need the notion, due to [LPR] and [BKLR], of the character lattice of a reductive k -group G over a field k . Let \bar{k} be a separable closure of k . Let $T \subset G$ be a maximal torus (defined over k). Set $\bar{T} = T \times_k \bar{k}$, $\bar{G} = G \times_k \bar{k}$. Let $X(\bar{T})$

denote the character group of $\bar{T} := T \times_k \bar{k}$. Let $W = W(\bar{G}, \bar{T}) := \mathcal{N}_G(\bar{T})/\bar{T}$ denote the Weyl group, it acts on $X(\bar{T})$. Consider the canonical Galois action on $X(\bar{T})$, it defines a homomorphism $\text{Gal}(\bar{k}/k) \rightarrow \text{Aut } X(\bar{T})$. The image $\text{im } \rho \subset \text{Aut } X(\bar{T})$ normalizes W , hence $\text{im } \rho \cdot W$ is a subgroup of $\text{Aut } X(\bar{T})$. By the character lattice of G we mean the pair $\mathcal{X}(G) := (\text{im } \rho \cdot W, X(\bar{T}))$ (up to an isomorphism it does not depend on the choice of T). In particular, if k is algebraically closed, then $\mathcal{X}(G) = (W, X(T))$.

We shall reduce Theorem 0.2 to an assertion about quasi-permutation lattices using the following result due to [LPR]:

PROPOSITION 1.8 ([LPR, Theorem 1.27], see also [BKLR, Theorem 1.3]). *A reductive group G over an algebraically closed field k of characteristic 0 is stably Cayley if and only if its character lattice $\mathcal{X}(G)$ is quasi-permutation, i.e., $X(T)$ is a quasi-permutation $W(G, T)$ -lattice.*

We shall use the following result due to Cortella and Kunyavskii [CK] and to Lemire, Popov and Reichstein [LPR].

PROPOSITION 1.9 ([CK], [LPR]). *Let D be a connected Dynkin diagram. Let $R = R(D)$ denote the corresponding root system, $W = W(D)$ denote the Weyl group, $Q = Q(D)$ denote the root lattice, and $P = P(D)$ denote the weight lattice. We say that L is an intermediate lattice between Q and P if $Q \subset L \subset P$ (note that $L = Q$ and $L = P$ are possible). Then the following list gives (up to an isomorphism) all the pairs (D, L) such that L is a quasi-permutation intermediate $W(D)$ -lattice between $Q(D)$ and $P(D)$:*

$$Q(\mathbf{A}_n), Q(\mathbf{B}_n), P(\mathbf{C}_n), \mathcal{X}(\mathbf{SO}_{2n}) \text{ (then } D = \mathbf{D}_n),$$

or D is any connected Dynkin diagram of rank 1 or 2 (i.e. $\mathbf{A}_1, \mathbf{A}_2, \mathbf{B}_2$, or \mathbf{G}_2) and L is any lattice between $Q(D)$ and $P(D)$, (i.e., either $L = P(D)$ or $L = Q(D)$).

Proof. The positive result (the assertion that the lattices in the list are indeed quasi-permutation) follows from the assertion that the corresponding groups are stably Cayley (or that their generic tori are stably rational), see the references in [CK], Section 3. See Appendix A below for a proof of this positive result in terms of lattices only. The difficult part of Proposition 1.9 is the negative result (the assertion that all the other lattices are not quasi-permutation). This was proved in [CK, Theorem 0.1] in the cases when $L = Q$ or $L = P$, and in [LPR, Propositions 5.1 and 5.2] in the cases when $Q \subsetneq L \subsetneq P$ (this can happen only when $D = \mathbf{A}_n$ or $D = \mathbf{D}_n$). \square

Remark 1.10. It follows from Proposition 1.9 that, in particular, the following lattices are quasi-permutation: $Q(\mathbf{A}_1), P(\mathbf{A}_1), P(\mathbf{A}_2), P(\mathbf{B}_2), Q(\mathbf{C}_2), Q(\mathbf{G}_2) = P(\mathbf{G}_2), Q(\mathbf{D}_3) = Q(\mathbf{A}_3), \mathcal{X}(\mathbf{SL}_4/\mu_2) = \mathcal{X}(\mathbf{SO}_6)$.

2 A FAMILY OF NON-QUASI-PERMUTATION LATTICES

In this section we construct a family of non-quasi-permutation (even non-quasi-invertible) lattices.

2.1. We consider a Dynkin diagram $D \sqcup \Delta$ (disjoint union). We assume that $D = \bigsqcup_{i \in I} D_i$ (a finite disjoint union), where each D_i is of type \mathbf{B}_{l_i} ($l_i \geq 1$) or \mathbf{D}_{l_i} ($l_i \geq 2$) (and where $\mathbf{B}_1 = \mathbf{A}_1$, $\mathbf{B}_2 = \mathbf{C}_2$, $\mathbf{D}_2 = \mathbf{A}_1 \sqcup \mathbf{A}_1$, and $\mathbf{D}_3 = \mathbf{A}_3$ are permitted). We denote by m the cardinality of the finite index set I . We assume that $\Delta = \bigsqcup_{\iota=1}^{\mu} \Delta_{\iota}$ (disjoint union), where Δ_{ι} is of type $\mathbf{A}_{2n_{\iota}-1}$, $n_{\iota} \geq 2$ ($\mathbf{A}_3 = \mathbf{D}_3$ is permitted). We assume that $m \geq 1$ and $\mu \geq 0$ (in the case $\mu = 0$ the diagram Δ is empty).

For each $i \in I$ we realize the root system $R(D_i)$ of type \mathbf{B}_{l_i} or \mathbf{D}_{l_i} in the standard way in the space $V_i := \mathbb{R}^{l_i}$ with basis $(e_s)_{s \in S_i}$ where S_i is an index set consisting of l_i elements; cf. [Bou, Planche II] for \mathbf{B}_l ($l \geq 2$) (the relevant formulas for \mathbf{B}_1 are similar) and [Bou, Planche IV] for \mathbf{D}_l ($l \geq 3$) (again, the relevant formulas for \mathbf{D}_2 are similar). Let $M_i \subset V_i$ denote the lattice generated by the basis vectors $(e_s)_{s \in S_i}$. Let $P_i \supset M_i$ denote the weight lattice of the root system D_i . Set $S = \bigsqcup_i S_i$ (disjoint union). Consider the vector space $V = \bigoplus_i V_i$ with basis $(e_s)_{s \in S}$. Let $M_D \subset V$ denote the lattice generated by the basis vectors $(e_s)_{s \in S}$, then $M_D = \bigoplus_i M_i$. Set $P_D = \bigoplus_i P_i$.

For each $\iota = 1, \dots, \mu$ we realize the root system $R(\Delta_{\iota})$ of type $\mathbf{A}_{2n_{\iota}-1}$ in the standard way in the subspace V_{ι} of vectors with zero sum of the coordinates in the space $\mathbb{R}^{2n_{\iota}}$ with basis $\varepsilon_{\iota,1}, \dots, \varepsilon_{\iota,2n_{\iota}}$; cf. [Bou, Planche I]. Let Q_{ι} be the root lattice of $R(\Delta_{\iota})$ with basis $\varepsilon_{\iota,1} - \varepsilon_{\iota,2}, \varepsilon_{\iota,2} - \varepsilon_{\iota,3}, \dots, \varepsilon_{\iota,2n_{\iota}-1} - \varepsilon_{\iota,2n_{\iota}}$, and let $P_{\iota} \supset Q_{\iota}$ be the weight lattice of $R(\Delta_{\iota})$. Set $Q_{\Delta} = \bigoplus_{\iota} Q_{\iota}$, $P_{\Delta} = \bigoplus_{\iota} P_{\iota}$.

Set

$$W := \prod_{i \in I} W(D_i) \times \prod_{\iota=1}^{\mu} W(\Delta_{\iota}), \quad L' = M_D \oplus Q_{\Delta} = \bigoplus_{i \in I} M_i \oplus \bigoplus_{\iota=1}^{\mu} Q_{\iota},$$

then W acts on L' and on $L' \otimes_{\mathbb{Z}} \mathbb{R}$. For each i consider the vector

$$x_i = \sum_{s \in S_i} e_s \in M_i,$$

then $\frac{1}{2}x_i \in P_i$. For each ι consider the vector

$$\xi_{\iota} = \varepsilon_{\iota,1} - \varepsilon_{\iota,2} + \varepsilon_{\iota,3} - \varepsilon_{\iota,4} + \dots + \varepsilon_{\iota,2n_{\iota}-1} - \varepsilon_{\iota,2n_{\iota}} \in Q_{\iota},$$

then $\frac{1}{2}\xi_{\iota} \in P_{\iota}$; see [Bou, Planche I]. Write

$$\xi'_{\iota} = \varepsilon_{\iota,1} - \varepsilon_{\iota,2}, \quad \xi''_{\iota} = \varepsilon_{\iota,3} - \varepsilon_{\iota,4} + \dots + \varepsilon_{\iota,2n_{\iota}-1} - \varepsilon_{\iota,2n_{\iota}},$$

then $\xi_{\iota} = \xi'_{\iota} + \xi''_{\iota}$. Consider the vector

$$v = \frac{1}{2} \sum_{i \in I} x_i + \frac{1}{2} \sum_{\iota=1}^{\mu} \xi_{\iota} = \frac{1}{2} \sum_{s \in S} e_s + \frac{1}{2} \sum_{\iota=1}^{\mu} \xi_{\iota} \in P_D \oplus P_{\Delta}.$$

Set

$$L = \langle L', v \rangle, \tag{2.1}$$

then $[L : L'] = 2$ because $v \in \frac{1}{2}L' \setminus L'$. Note that the sublattice $L \subset P_D \oplus P_\Delta$ is W -invariant. Indeed, the group W acts on $(P_D \oplus P_\Delta)/(M_D \oplus Q_\Delta)$ trivially.

PROPOSITION 2.2. *We assume that $m \geq 1$, $m + \mu \geq 2$. If $\mu = 0$, we assume that not all of D_i are of types \mathbf{B}_1 or \mathbf{D}_2 . Then the W -lattice L as in (2.1) is not quasi-invertible, hence not quasi-permutation.*

Proof. We consider a group $\Gamma = \{e, \gamma_1, \gamma_2, \gamma_3\}$ of order 4, where $\gamma_1, \gamma_2, \gamma_3$ are of order 2. The idea of our proof is to construct an embedding

$$j: \Gamma \rightarrow W$$

in such a way that L , viewed as a Γ -lattice, is equivalent to its Γ -sublattice L_1 , and L_1 is isomorphic to a direct sum of a Γ -sublattice $L_0 \simeq J_\Gamma$ of rank 3 and a number of Γ -lattices of rank 1. Since by Proposition 1.6 J_Γ is not quasi-invertible, this will imply that L_1 and L are not quasi-invertible Γ -lattices, and hence L is not quasi-invertible as a W -lattice. We shall now fill in the details of this argument in four steps.

Step 1. We begin by partitioning each S_i for $i \in I$ into three (non-overlapping) subsets $S_{i,1}$, $S_{i,2}$ and $S_{i,3}$, subject to the requirement that

$$|S_{i,1}| \equiv |S_{i,2}| \equiv |S_{i,3}| \equiv l_i \pmod{2} \text{ if } D_i \text{ is of type } \mathbf{D}_{l_i}. \tag{2.2}$$

We then set U_1 to be the union of the $S_{i,1}$, U_2 to be the union of the $S_{i,2}$, and U_3 to be the union of the $S_{i,3}$, as i runs over I .

LEMMA 2.3. (i) *If $\mu \geq 1$, the subsets $S_{i,1}$, $S_{i,2}$ and $S_{i,3}$ of S_i can be chosen, subject to (2.2), so that $U_1 \neq \emptyset$.*

(ii) *If $\mu = 0$ (and $m \geq 2$), the subsets $S_{i,1}$, $S_{i,2}$ and $S_{i,3}$ of S_i can be chosen, subject to (2.2), so that $U_1, U_2, U_3 \neq \emptyset$.*

To prove the lemma, first consider case (i). For all i such that D_i is of type \mathbf{D}_{l_i} with *odd* l_i , we partition S_i into three non-empty subsets of odd cardinalities. For all the other i we take $S_{i,1} = S_i$, $S_{i,2} = S_{i,3} = \emptyset$. Then $U_1 \neq \emptyset$ (note that $m \geq 1$) and (2.2) is satisfied.

In case (ii), if one of the D_i is of type \mathbf{D}_{l_i} where $l_i \geq 3$ is *odd*, then we partition S_i for each such D_i into three non-empty subsets of odd cardinalities. We partition all the other S_i as follows:

$$S_{i,1} = S_{i,2} = \emptyset \text{ and } S_{i,3} = S_i. \tag{2.3}$$

Clearly $U_1, U_2, U_3 \neq \emptyset$ and (2.2) is satisfied.

If there is no D_i of type \mathbf{D}_{l_i} with odd $l_i \geq 3$, but one of the D_i , say for $i = i_0$, is \mathbf{D}_l with *even* $l \geq 4$, then we partition S_{i_0} into two non-empty subsets $S_{i_0,1}$ and

$S_{i_0,2}$ of even cardinalities, and set $S_{i_0,3} = \emptyset$. We partition the sets S_i for $i \neq i_0$ as in (2.3) (note that by our assumption $m \geq 2$). Once again, $U_1, U_2, U_3 \neq \emptyset$ and (2.2) is satisfied.

If there is no D_i of type \mathbf{D}_{l_i} with $l_i \geq 3$ (odd or even), but one of the D_i , say for $i = i_0$, is of type \mathbf{B}_l with $l \geq 2$, we partition S_{i_0} into two non-empty subsets $S_{i_0,1}$ and $S_{i_0,2}$, and set $S_{i_0,3} = \emptyset$. We partition the sets S_i for $i \neq i_0$ as in (2.3) (again, note that $m \geq 2$). Once again, $U_1, U_2, U_3 \neq \emptyset$ and (2.2) is satisfied.

Since by our assumption not all of D_i are of type \mathbf{B}_1 or \mathbf{D}_2 , we have exhausted all the cases. This completes the proof of Lemma 2.3. \square

Step 2. We continue proving Proposition 2.2. We construct an embedding $\Gamma \hookrightarrow W$.

For $s \in S$ we denote by c_s the automorphism of L taking the basis vector e_s to $-e_s$ and fixing all the other basis vectors. For $\iota = 1, \dots, \mu$ we define $\tau_\iota^{(12)} = \text{Transp}((\iota, 1), (\iota, 2)) \in W_\iota$ (the transposition of the basis vectors $\varepsilon_{\iota,1}$ and $\varepsilon_{\iota,2}$). Set

$$\tau_\iota^{>2} = \text{Transp}((\iota, 3), (\iota, 4)) \cdot \dots \cdot \text{Transp}((\iota, 2n_\iota - 1), (\iota, 2n_\iota)) \in W_\iota.$$

Write $\Gamma = \{e, \gamma_1, \gamma_2, \gamma_3\}$ and define an embedding $j: \Gamma \hookrightarrow W$ as follows:

$$\begin{aligned} j(\gamma_1) &= \prod_{s \in S \setminus U_1} c_s \cdot \prod_{\iota=1}^{\mu} \tau_\iota^{(12)} \tau_\iota^{>2}; \\ j(\gamma_2) &= \prod_{s \in S \setminus U_2} c_s \cdot \prod_{\iota=1}^{\mu} \tau_\iota^{(12)}; \\ j(\gamma_3) &= \prod_{s \in S \setminus U_3} c_s \cdot \prod_{\iota=1}^{\mu} \tau_\iota^{>2}. \end{aligned}$$

Note that if D_i is of type \mathbf{D}_{l_i} , then by (2.2) for $\varkappa = 1, 2, 3$ the cardinality $\#(S_i \setminus S_{i,\varkappa})$ is even, hence the product of c_s over $s \in S_i \setminus S_{i,\varkappa}$ is contained in $W(D_i)$ for all such i , and therefore, $j(\gamma_\varkappa) \in W$. Since $j(\gamma_1), j(\gamma_2)$ and $j(\gamma_3)$ commute, are of order 2, and $j(\gamma_1)j(\gamma_2) = j(\gamma_3)$, we see that j is a homomorphism. If $\mu \geq 1$, then, since $2n_1 \geq 4$, clearly $j(\gamma_\varkappa) \neq 1$ for $\varkappa = 1, 2, 3$, hence j is an embedding. If $\mu = 0$, then the sets $S \setminus U_1, S \setminus U_2$ and $S \setminus U_3$ are nonempty, and again $j(\gamma_\varkappa) \neq 1$ for $\varkappa = 1, 2, 3$, hence j is an embedding.

Step 3. We construct a Γ -sublattice L_0 of rank 3. Write a vector $\mathbf{x} \in L$ as

$$\mathbf{x} = \sum_{s \in S} b_s e_s + \sum_{\iota=1}^{\mu} \sum_{\nu=1}^{2n_\iota} \beta_{\iota,\nu} \varepsilon_{\iota,\nu},$$

where $b_s, \beta_{\iota,\nu} \in \frac{1}{2}\mathbb{Z}$. Set $n' = \sum_{\iota=1}^{\mu} (n_\iota - 1)$. Define a Γ -equivariant homomorphism

$$\phi: L \rightarrow \mathbb{Z}^{n'}, \quad \mathbf{x} \mapsto (\beta_{\iota,2\lambda-1} + \beta_{\iota,2\lambda})_{\iota=1,\dots,\mu, \lambda=2,\dots,n_\iota}$$

(we skip $\lambda = 1$). We obtain a short exact sequence of Γ -lattices

$$0 \rightarrow L_1 \rightarrow L \xrightarrow{\phi} \mathbb{Z}^{n'} \rightarrow 0,$$

where $L_1 := \ker \phi$. Since Γ acts trivially on $\mathbb{Z}^{n'}$, we have $L_1 \sim L$. Therefore, it suffices to show that L_1 is not quasi-invertible.

Recall that

$$v = \frac{1}{2} \sum_{s \in S} e_s + \frac{1}{2} \sum_{\iota=1}^{\mu} \xi_{\iota}.$$

Set $v_1 = \gamma_1 \cdot v$, $v_2 = \gamma_2 \cdot v$, $v_3 = \gamma_3 \cdot v$. Set

$$L_0 = \langle v, v_1, v_2, v_3 \rangle.$$

We have

$$v_1 = \frac{1}{2} \sum_{s \in U_1} e_s - \frac{1}{2} \sum_{s \in U_2 \cup U_3} e_s - \frac{1}{2} \sum_{\iota=1}^{\mu} \xi_{\iota},$$

whence

$$v + v_1 = \sum_{s \in U_1} e_s. \tag{2.4}$$

We have

$$v_2 = \frac{1}{2} \sum_{s \in U_2} e_s - \frac{1}{2} \sum_{s \in U_1 \cup U_3} e_s + \frac{1}{2} \sum_{\iota=1}^{\mu} (-\xi'_{\iota} + \xi''_{\iota}),$$

whence

$$v + v_2 = \sum_{s \in U_2} e_s + \sum_{\iota=1}^{\mu} \xi''_{\iota}. \tag{2.5}$$

We have

$$v_3 = \frac{1}{2} \sum_{s \in U_3} e_s - \frac{1}{2} \sum_{s \in U_1 \cup U_2} e_s + \frac{1}{2} \sum_{\iota=1}^{\mu} (\xi'_{\iota} - \xi''_{\iota}),$$

whence

$$v + v_3 = \sum_{s \in U_3} e_s + \sum_{\iota=1}^{\mu} \xi'_{\iota}. \tag{2.6}$$

Clearly, we have

$$v + v_1 + v_2 + v_3 = 0.$$

Since the set $\{v, v_1, v_2, v_3\}$ is the orbit of v under Γ , the sublattice $L_0 = \langle v, v_1, v_2, v_3 \rangle \subset L$ is Γ -invariant. If $\mu \geq 1$, then $U_1 \neq \emptyset$, and we see from (2.4), (2.5) and (2.6) that $\text{rank } L_0 \geq 3$. If $\mu = 0$, then $U_1, U_2, U_3 \neq \emptyset$, and again we see from (2.4), (2.5) and (2.6) that $\text{rank } L_0 \geq 3$. Thus $\text{rank } L_0 = 3$ and $L_0 \simeq J_{\Gamma}$, whence by Proposition 1.6 L_0 is not quasi-invertible.

Step 4. We show that L_0 is a direct summand of L_1 . Set $m' = |S|$.

First assume that $\mu \geq 1$. Choose $u_1 \in U_1 \subset S$. Set $S' = S \setminus \{u_1\}$. For each $s \in S'$ (i.e., $s \neq u_1$) consider the one-dimensional (i.e., of rank 1) lattice $X_s = \langle e_s \rangle$. We obtain $m' - 1$ Γ -invariant one-dimensional sublattices of L_1 .

Denote by Υ the set of pairs (ι, λ) such that $1 \leq \iota \leq \mu$, $1 \leq \lambda \leq n_\iota$, and if $\iota = 1$, then $\lambda \neq 1, 2$. For each $(\iota, \lambda) \in \Upsilon$ consider the one-dimensional lattice

$$\Xi_{\iota, \lambda} = \langle \varepsilon_{\iota, 2\lambda-1} - \varepsilon_{\iota, 2\lambda} \rangle.$$

We obtain $-2 + \sum_{\iota=1}^{\mu} n_\iota$ one-dimensional Γ -invariant sublattices of L_1 .

We show that

$$L_1 = L_0 \oplus \bigoplus_{s \in S'} X_s \oplus \bigoplus_{(\iota, \lambda) \in \Upsilon} \Xi_{\iota, \lambda}. \tag{2.7}$$

Set $L'_1 = \langle L_0, (X_s)_{s \neq u_1}, (\Xi_{\iota, \lambda})_{(\iota, \lambda) \in \Upsilon} \rangle$, then

$$\text{rank } L'_1 \leq 3 + (m' - 1) - 2 + \sum_{\iota=1}^{\mu} n_\iota = m' + \sum_{\iota=1}^{\mu} (2n_\iota - 1) - \sum_{\iota=1}^{\mu} (n_\iota - 1) = \text{rank } L_1. \tag{2.8}$$

Therefore, it suffices to check that $L'_1 \supset L_1$. The set

$$\{v\} \cup \{e_s \mid s \in S\} \cup \{\varepsilon_{\iota, 2\lambda-1} - \varepsilon_{\iota, 2\lambda} \mid 1 \leq \iota \leq \mu, 1 \leq \lambda \leq n_\iota\}$$

is a set of generators of L_1 . By construction $v, v_1, v_2, v_3 \in L_0 \subset L'_1$. We have $e_s \in X_s \subset L'_1$ for $s \neq u_1$. By (2.4) $\sum_{s \in U_1} e_s \in L'_1$, hence $e_{u_1} \in L'_1$. By construction

$$\varepsilon_{\iota, 2\lambda-1} - \varepsilon_{\iota, 2\lambda} \in L'_1, \text{ for all } (\iota, \lambda) \neq (1, 1), (1, 2).$$

From (2.6) and (2.5) we see that

$$\sum_{\iota=1}^{\mu} (\varepsilon_{\iota, 1} - \varepsilon_{\iota, 2}) \in L'_1, \quad \sum_{\iota=1}^{\mu} \xi''_\iota \in L'_1.$$

Thus

$$\varepsilon_{1, 1} - \varepsilon_{1, 2} \in L'_1, \quad \varepsilon_{1, 3} - \varepsilon_{1, 4} \in L'_1.$$

We conclude that $L'_1 \supset L_1$, hence $L_1 = L'_1$. From dimension count (2.8) we see that (2.7) holds.

Now assume that $\mu = 0$. Then for each $\varkappa = 1, 2, 3$ we choose an element $u_\varkappa \in U_\varkappa$ and set $U'_\varkappa = U_\varkappa \setminus \{u_\varkappa\}$. We set $S' = U'_1 \cup U'_2 \cup U'_3 = S \setminus \{u_1, u_2, u_3\}$. Again for $s \in S'$ (i.e., $s \neq u_1, u_2, u_3$) consider the one-dimensional lattice $X_s = \langle e_s \rangle$. We obtain $m' - 3$ one-dimensional Γ -invariant sublattices of $L_1 = L$. We show that

$$L_1 = L_0 \oplus \bigoplus_{s \in S'} X_s. \tag{2.9}$$

Set $L'_1 = \langle L_0, (X_s)_{s \in S'} \rangle$, then

$$\text{rank } L'_1 \leq 3 + m' - 3 = m' = \text{rank } L_1. \tag{2.10}$$

Therefore, it suffices to check that $L'_1 \supset L_1$. The set $\{v\} \cup \{e_s \mid s \in S\}$ is a set of generators of $L_1 = L$. By construction $v, v_1, v_2, v_3 \in L'_1$ and $e_s \in L'_1$ for $s \neq u_1, u_2, u_3$. We see from (2.4), (2.5), (2.6) that $e_s \in L'_1$ also for $s = u_1, u_2, u_3$. Thus $L'_1 \supset L_1$, hence $L'_1 = L_1$. From dimension count (2.10) we see that (2.9) holds.

We see that in both cases $\mu \geq 1$ and $\mu = 0$, the sublattice L_0 is a direct summand of L_1 . Since by Proposition 1.6 L_0 is not quasi-invertible as a Γ -lattice, it follows that L_1 and L are not quasi-invertible as Γ -lattices. Thus L is not quasi-invertible as a W -lattice. This completes the proof of Proposition 2.2. \square

Remark 2.4. Since $\text{III}^2(\Gamma, J_\Gamma) \cong \mathbb{Z}/2\mathbb{Z}$ (Voskresenskiĭ, see [BKLR, Section 10] for the notation and the result), our argument shows that $\text{III}^2(\Gamma, L) \cong \mathbb{Z}/2\mathbb{Z}$.

Remark 2.5. The proof of [BKLR, Lemma 12.3] (which is a version with $\mu = 0$ of Lemma 2.3 above) contains an inaccuracy, though the lemma as stated is correct. Namely, in [BKLR] we write that if there exists i such that Δ_i is of type \mathbf{D}_{l_i} where $l_i \geq 3$ is odd, then we partition S_i for *one* such i into three non-empty subsets $S_{i,1}, S_{i,2}$ and $S_{i,3}$ of odd cardinalities, and we partition all the other S_i as in [BKLR, (12.4)]. However, this partitioning of the sets S_i into three subsets does not satisfy [BKLR, (12.3)] for *other* i such that Δ_i is of type \mathbf{D}_{l_i} with odd l_i . This inaccuracy can be easily corrected: we should partition S_i for *each* i such that Δ_i is of type \mathbf{D}_{l_i} with odd l_i into three non-empty subsets of odd cardinalities.

3 MORE NON-QUASI-PERMUTATION LATTICES

In this section we construct another family of non-quasi-permutation lattices.

3.1. For $i = 1, \dots, r$ let $Q_i = \mathbb{Z}\mathbf{A}_{n_i-1}$ and $P_i = \Lambda_{n_i}$ denote the root lattice and the weight lattice of \mathbf{SL}_{n_i} , respectively, and let $W_i = \mathfrak{S}_{n_i}$ denote the corresponding Weyl group (the symmetric group on n_i letters) acting on P_i and Q_i . Set $F_i = P_i/Q_i$, then W_i acts trivially on F_i . Set

$$Q = \bigoplus_{i=1}^r Q_i, \quad P = \bigoplus_{i=1}^r P_i, \quad W = \prod_{i=1}^r W_i,$$

then $Q \subset P$ and the Weyl group W acts on Q and P . Set

$$F = P/Q = \bigoplus_{i=1}^r F_i,$$

then W acts trivially on F .

We regard $Q_i = \mathbb{Z}\mathbf{A}_{n_i-1}$ and $P_i = \Lambda_{n_i}$ as the lattices described in Bourbaki [Bou, Planche I]. Then we have an isomorphism $F_i \cong \mathbb{Z}/n_i\mathbb{Z}$. Note that for each $1 \leq i \leq r$, the set $\{\alpha_{\varkappa,i} \mid 1 \leq \varkappa \leq n_i - 1\}$ is a \mathbb{Z} -basis of Q_i .

Set $c = \gcd(n_1, \dots, n_r)$; we assume that $c > 1$. Let $d > 1$ be a divisor of c . For each $i = 1, \dots, r$, let $\nu_i \in \mathbb{Z}$ be such that $1 \leq \nu_i < d$, $\gcd(\nu_i, d) = 1$, and assume that $\nu_1 = 1$. We write $\boldsymbol{\nu} = (\nu_i)_{i=1}^r \in \mathbb{Z}^r$. Let $\overline{\boldsymbol{\nu}}$ denote the image of $\boldsymbol{\nu}$ in $(\mathbb{Z}/d\mathbb{Z})^r$. Let $S_{\boldsymbol{\nu}} \subset (\mathbb{Z}/d\mathbb{Z})^r \subset \bigoplus_{i=1}^r \mathbb{Z}/n_i\mathbb{Z} = F$ denote the cyclic subgroup of order d generated by $\overline{\boldsymbol{\nu}}$. Let $L_{\boldsymbol{\nu}}$ denote the preimage of $S_{\boldsymbol{\nu}} \subset F$ in P under the canonical epimorphism $P \twoheadrightarrow F$, then $Q \subset L_{\boldsymbol{\nu}} \subset P$.

PROPOSITION 3.2. *Let W and the W -lattice $L_{\boldsymbol{\nu}}$ be as in Subsection 3.1. In the case $d = 2^s$ we assume that $\sum n_i > 4$. Then $L_{\boldsymbol{\nu}}$ is not quasi-permutation.*

This proposition follows from Lemmas 3.3 and 3.8 below.

LEMMA 3.3. *Let $p|d$ be a prime. Then for any subgroup $\Gamma \subset W$ isomorphic to $(\mathbb{Z}/p\mathbb{Z})^m$ for some natural m , the Γ -lattices $L_{\boldsymbol{\nu}}$ and $L_{\mathbf{1}} := L_{(1, \dots, 1)}$ are equivalent for any $\boldsymbol{\nu} = (\nu_1, \dots, \nu_r)$ as above (in particular, we assume that $\nu_1 = 1$).*

Note that this lemma is trivial when $d = 2$.

3.4. We compute the lattice $L_{\boldsymbol{\nu}}$ explicitly. First let $r = 1$. We have $Q = Q_1$, $P = P_1$. Then P_1 is generated by Q_1 and an element $\omega \in P_1$ whose image in P_1/Q_1 is of order n_1 . We may take

$$\omega = \frac{1}{n_1}[(n_1 - 1)\alpha_1 + (n_1 - 2)\alpha_2 + \dots + 2\alpha_{n_1-2} + \alpha_{n_1-1}],$$

where $\alpha_1, \dots, \alpha_{n_1-1}$ are the simple roots, see [Bou, Planche I]. There exists exactly one intermediate lattice L between Q_1 and P_1 such that $[L : Q_1] = d$, and it is generated by Q_1 and the element

$$w = \frac{n_1}{d}\omega = \frac{1}{d}[(n_1 - 1)\alpha_1 + (n_1 - 2)\alpha_2 + \dots + 2\alpha_{n_1-2} + \alpha_{n_1-1}].$$

Now for any natural r , the lattice $L_{\boldsymbol{\nu}}$ is generated by Q and the element

$$w_{\boldsymbol{\nu}} = \frac{1}{d} \sum_{i=1}^r \nu_i [(n_i - 1)\alpha_{1,i} + (n_i - 2)\alpha_{2,i} + \dots + 2\alpha_{n_i-2,i} + \alpha_{n_i-1,i}].$$

In particular, $L_{\mathbf{1}}$ is generated by Q and

$$w_{\mathbf{1}} = \frac{1}{d} \sum_{i=1}^r [(n_i - 1)\alpha_{1,i} + (n_i - 2)\alpha_{2,i} + \dots + 2\alpha_{n_i-2,i} + \alpha_{n_i-1,i}].$$

3.5. *Proof of Lemma 3.3.* Recall that $L_\nu = \langle Q, w_\nu \rangle$ with

$$Q = \langle \alpha_{\varkappa,i} \rangle, \quad \text{where } i = 1, \dots, r, \varkappa = 1, \dots, n_i - 1.$$

Set $Q_\nu = \langle \nu_i \alpha_{\varkappa,i} \rangle$. Denote by \mathfrak{T}_ν the endomorphism of Q that acts on Q_i by multiplication by ν_i . We have $Q_1 = Q$, $Q_\nu = \mathfrak{T}_\nu Q_1$, $w_\nu = \mathfrak{T}_\nu w_1$. Consider

$$\mathfrak{T}_\nu L_1 = \langle Q_\nu, w_\nu \rangle.$$

Clearly the W -lattices L_1 and $\mathfrak{T}_\nu L_1$ are isomorphic. We have an embedding of W -lattices $Q \hookrightarrow L_\nu$, in particular, an embedding $Q \hookrightarrow L_1$, which induces an embedding $\mathfrak{T}_\nu Q \hookrightarrow \mathfrak{T}_\nu L_1$. Set $M_\nu = L_\nu / \mathfrak{T}_\nu L_1$, then we obtain a homomorphism of W -modules $Q / \mathfrak{T}_\nu Q \rightarrow M_\nu$, which is an isomorphism by Lemma 3.6 below.

Now let $p|d$ be a prime. Let $\Gamma \subset W$ be a subgroup isomorphic to $(\mathbb{Z}/p\mathbb{Z})^m$ for some natural m . As in [LPR, Proof of Proposition 2.10], we use Roiter's version [Ro, Proposition 2] of Schanuel's lemma. We have exact sequences of Γ -modules

$$\begin{aligned} 0 \rightarrow \mathfrak{T}_\nu L_1 \rightarrow L_\nu \rightarrow M_\nu \rightarrow 0, \\ 0 \rightarrow Q \xrightarrow{\mathfrak{T}_\nu} Q \rightarrow M_\nu \rightarrow 0. \end{aligned}$$

Since all ν_i are prime to p , we have $|\Gamma| \cdot M_\nu = p^m M_\nu = M_\nu$, and by [Ro, Corollary of Proposition 3] the morphisms of $\mathbb{Z}[\Gamma]$ -modules $L_\nu \rightarrow M_\nu$ and $Q \rightarrow M_\nu$ are projective in the sense of [Ro, §1]. Now by [Ro, Proposition 2] there exists an isomorphism of Γ -lattices $L_\nu \oplus Q \simeq \mathfrak{T}_\nu L_1 \oplus Q$. Since Q is a quasi-permutation W -lattice, it is a quasi-permutation Γ -lattice, and by Lemma 3.7 below, $L_\nu \sim \mathfrak{T}_\nu L_1$ as Γ -lattices. Since $\mathfrak{T}_\nu L_1 \simeq L_1$, we conclude that $L_\nu \sim L_1$. \square

LEMMA 3.6. *With the above notation $L_\nu / \mathfrak{T}_\nu L_1 \simeq Q / \mathfrak{T}_\nu Q = \bigoplus_{i=2}^r Q_i / \nu_i Q_i$.*

Proof. We have $\mathfrak{T}_\nu L_1 = \langle S_\nu \rangle$, where $S_\nu = \{ \nu_i \alpha_{\varkappa,i} \}_{i,\varkappa} \cup \{ w_\nu \}$. Note that

$$dw_\nu = \sum_{i=1}^r \nu_i [(n_i - 1)\alpha_{1,i} + (n_i - 2)\alpha_{2,i} + \dots + 2\alpha_{n_i-2,i} + \alpha_{n_i-1,i}].$$

We see that dw_ν is a linear combination with integer coefficients of $\nu_i \alpha_{\varkappa,i}$ and that $\alpha_{n_1-1,1}$ appears in this linear combination with coefficient 1 (because $\nu_1 = 1$). Set $B'_\nu = S_\nu \setminus \{ \alpha_{n_1-1,1} \}$, then $\langle B'_\nu \rangle \ni \alpha_{n_1-1,1}$, hence $\langle B'_\nu \rangle = \langle S_\nu \rangle = \mathfrak{T}_\nu L_1$, thus B'_ν is a basis of $\mathfrak{T}_\nu L_1$. Similarly, the set $B_\nu := \{ \alpha_{\varkappa,i} \}_{i,\varkappa} \cup \{ w_\nu \} \setminus \{ \alpha_{n_1-1,1} \}$ is a basis of L_ν . Both bases B_ν and B'_ν contain $\alpha_{1,1}, \dots, \alpha_{n_1-2,1}$ and w_ν . For all $i = 2, \dots, r$ and all $\varkappa = 1, \dots, n_i - 1$, the basis B_ν contains $\alpha_{\varkappa,i}$, while B'_ν contains $\nu_i \alpha_{\varkappa,i}$. We see that the homomorphism of W -modules $Q / \mathfrak{T}_\nu Q = \bigoplus_{i=2}^r Q_i / \nu_i Q_i \rightarrow L_\nu / \mathfrak{T}_\nu L_1$ is an isomorphism. \square

LEMMA 3.7. *Let Γ be a finite group, A and A' be Γ -lattices. If $A \oplus B \sim A' \oplus B'$, where B and B' are quasi-permutation Γ -lattices, then $A \sim A'$.*

Proof. Since B and B' are quasi-permutation, by Lemma 1.3 they are equivalent to 0, and we have

$$A = A \oplus 0 \sim A \oplus B \sim A' \oplus B' \sim A' \oplus 0 = A'.$$

This completes the proof of Lemma 3.7 and hence of Lemma 3.3. \square

To complete the proof of Proposition 3.2 it suffices to prove the next lemma.

LEMMA 3.8. *Let $p|d$ be a prime. Then there exists a subgroup $\Gamma \subset W$ isomorphic to $(\mathbb{Z}/p\mathbb{Z})^m$ for some natural m such that the Γ -lattice $L_1 := L_{(1,\dots,1)}$ is not quasi-permutation.*

3.9. Denote by U_i the space \mathbb{R}^{n_i} with canonical basis $\varepsilon_{1,i}, \varepsilon_{2,i}, \dots, \varepsilon_{n_i,i}$. Denote by V_i the subspace of codimension 1 in U_i consisting of vectors with zero sum of the coordinates. The group $W_i = \mathfrak{S}_{n_i}$ (the symmetric group) permutes the basis vectors $\varepsilon_{1,i}, \varepsilon_{2,i}, \dots, \varepsilon_{n_i,i}$ and thus acts on U_i and V_i . Consider the homomorphism of vector spaces

$$\chi_i: U_i \rightarrow \mathbb{R}, \quad \sum_{\lambda=1}^{n_i} \beta_{\lambda,i} \varepsilon_{\lambda,i} \mapsto \sum_{\lambda=1}^{n_i} \beta_{\lambda,i}$$

taking a vector to the sum of its coordinates. Clearly this homomorphism is W_i -equivariant, where W_i acts trivially on \mathbb{R} . We have short exact sequences

$$0 \rightarrow V_i \rightarrow U_i \xrightarrow{\chi_i} \mathbb{R} \rightarrow 0.$$

Set $U = \bigoplus_{i=1}^r U_i$, $V = \bigoplus_{i=1}^r V_i$. The group $W = \prod_{i=1}^r W_i$ naturally acts on U and V , and we have an exact sequence of W -spaces

$$0 \rightarrow V \rightarrow U \xrightarrow{\chi} \mathbb{R}^r \rightarrow 0, \quad (3.1)$$

where $\chi = (\chi_i)_{i=1,\dots,r}$ and W acts trivially on \mathbb{R}^r .

Set $n = \sum_{i=1}^r n_i$. Consider the vector space $\bar{U} := \mathbb{R}^n$ with canonical basis $\bar{\varepsilon}_1, \bar{\varepsilon}_2, \dots, \bar{\varepsilon}_n$. Consider the natural isomorphism

$$\varphi: U = \bigoplus_i U_i \xrightarrow{\sim} \bar{U}$$

that takes $\varepsilon_{1,1}, \varepsilon_{2,1}, \dots, \varepsilon_{n_1,1}$ to $\bar{\varepsilon}_1, \bar{\varepsilon}_2, \dots, \bar{\varepsilon}_{n_1}$, takes $\varepsilon_{1,2}, \varepsilon_{2,2}, \dots, \varepsilon_{n_2,2}$ to $\bar{\varepsilon}_{n_1+1}, \bar{\varepsilon}_{n_1+2}, \dots, \bar{\varepsilon}_{n_1+n_2}$, and so on. Let \bar{V} denote the subspace of codimension 1 in \bar{U} consisting of vectors with zero sum of the coordinates. Sequence (3.1) induces an exact sequence of W -spaces

$$0 \rightarrow \varphi(V) \rightarrow \bar{V} \xrightarrow{\psi} \mathbb{R}^r \xrightarrow{\Sigma} \mathbb{R} \rightarrow 0. \quad (3.2)$$

Here $\psi = (\psi_i)_{i=1,\dots,r}$, where ψ_i takes a vector $\sum_{j=1}^n \beta_j \bar{\varepsilon}_j \in \bar{V}$ to $\sum_{\lambda=1}^{n_i} \beta_{n_1+\dots+n_{i-1}+\lambda}$, and the map Σ takes a vector in \mathbb{R}^r to the sum of its coordinates. Note that W acts trivially on \mathbb{R}^r and \mathbb{R} .

We have a lattice $Q_i \subset V_i$ for each $i = 1, \dots, r$, a lattice $Q = \bigoplus_i Q_i \subset \bigoplus_i V_i$, and a lattice $\bar{Q} := \mathbb{Z}\mathbf{A}_{n-1}$ in \bar{V} with basis $\bar{\varepsilon}_1 - \bar{\varepsilon}_2, \dots, \bar{\varepsilon}_{n-1} - \bar{\varepsilon}_n$. The isomorphism φ induces an embedding of $Q = \bigoplus_i Q_i$ into \bar{Q} . Under this embedding

$$\begin{aligned} \alpha_{1,1} &\mapsto \bar{\alpha}_1, \alpha_{2,1} \mapsto \bar{\alpha}_2, \dots, \alpha_{n_1-1,1} \mapsto \bar{\alpha}_{n_1-1}, \\ \alpha_{1,2} &\mapsto \bar{\alpha}_{n_1+1}, \alpha_{2,2} \mapsto \bar{\alpha}_{n_1+2}, \dots, \alpha_{n_2-1,2} \mapsto \bar{\alpha}_{n_1+n_2-1}, \\ &\dots\dots\dots \\ \alpha_{1,r} &\mapsto \bar{\alpha}_{n_1+n_2+\dots+n_{r-1}+1}, \dots, \alpha_{n_{r-1},r} \mapsto \bar{\alpha}_{n-1}, \end{aligned}$$

while $\bar{\alpha}_{n_1}, \bar{\alpha}_{n_1+n_2}, \dots, \bar{\alpha}_{n_1+n_2+\dots+n_{r-1}}$ are skipped.

3.10. We write L for L_1 and w for $w_1 \in \frac{1}{d}Q$, where $Q = \bigoplus_i Q_i$. Then

$$w = \sum_{i=1}^r w_i, \quad w_i = \frac{1}{d}[(n_i - 1)\alpha_{1,i} + \dots + \alpha_{n_i-1,i}].$$

Recall that

$$Q_i = \mathbb{Z}\mathbf{A}_{n_i-1} = \{(a_j) \in \mathbb{Z}^{n_i} \mid \sum_{j=1}^{n_i} a_j = 0\}.$$

Set

$$\bar{w} = \frac{1}{d} \sum_{j=1}^{n-1} (n-j)\bar{\alpha}_j.$$

Set $\Lambda_n(d) = \langle \bar{Q}, \bar{w} \rangle$. Note that $\Lambda_n(d) = Q_n(n/d)$ with the notation of [LPR, Subsection 6.1]. Set

$$N = \varphi(Q \otimes_{\mathbb{Z}} \mathbb{R}) \cap \Lambda_n(d) = \varphi(V) \cap \Lambda_n(d).$$

LEMMA 3.11. $\varphi(L) = N$.

Proof. Write $j_1 = n_1, j_2 = n_1 + n_2, \dots, j_{r-1} = n_1 + \dots + n_{r-1}$. Set $J = \{1, 2, \dots, n-1\} \setminus \{j_1, j_2, \dots, j_{r-1}\}$. Set

$$\mu = \frac{1}{d} \sum_{j \in J} (n-j)\bar{\alpha}_j = \bar{w} - \sum_{i=1}^{r-1} \frac{n-j_i}{d} \bar{\alpha}_{j_i}.$$

Note that $d|n$ and $d|j_i$ for all i , hence the coefficients $(n-j_i)/d$ are integral, and therefore $\mu \in \Lambda_n(d)$. Since also $\mu \in \varphi(Q \otimes_{\mathbb{Z}} \mathbb{R})$, we see that $\mu \in N$.

Let $y \in N$. Then

$$y = b\bar{w} + \sum_{j=1}^{n-1} a_j \bar{\alpha}_j$$

where $b, a_j \in \mathbb{Z}$, because $y \in \Lambda_n(d)$. We see that in the basis $\bar{\alpha}_1, \dots, \bar{\alpha}_{n-1}$ of $\Lambda_n(d) \otimes_{\mathbb{Z}} \mathbb{R}$, the element y contains $\bar{\alpha}_{j_i}$ with coefficient

$$b \frac{n - j_i}{d} + a_{j_i}.$$

Since $y \in \varphi(Q \otimes_{\mathbb{Z}} \mathbb{R})$, this coefficient must be 0:

$$b \frac{n - j_i}{d} + a_{j_i} = 0.$$

Consider

$$\begin{aligned} y - b\mu &= y - b \left(\bar{w} - \sum_{i=1}^{r-1} \frac{n - j_i}{d} \bar{\alpha}_{j_i} \right) = y - b\bar{w} + \sum_{i=1}^{r-1} \frac{b(n - j_i)}{d} \bar{\alpha}_{j_i} \\ &= \sum_{j=1}^{n-1} a_j \bar{\alpha}_j + \sum_{i=1}^{r-1} \frac{b(n - j_i)}{d} \bar{\alpha}_{j_i} = \sum_{j \in J} a_j \bar{\alpha}_j, \end{aligned}$$

where $a_j \in \mathbb{Z}$. We see that $y \in \langle \bar{\alpha}_j (j \in J), \mu \rangle$ for any $y \in N$, hence $N \subset \langle \bar{\alpha}_j (j \in J), \mu \rangle$. Conversely, $\mu \in N$ and $\bar{\alpha}_j \in N$ for $j \in J$, hence $\langle \bar{\alpha}_j (j \in J), \mu \rangle \subset N$, thus

$$N = \langle \bar{\alpha}_j (j \in J), \mu \rangle. \tag{3.3}$$

Now

$$\varphi(w) = \frac{1}{d} \left[\sum_{j=1}^{n_1-1} (n_1 - j) \bar{\alpha}_j + \sum_{j=1}^{n_2-1} (n_2 - j) \bar{\alpha}_{n_1+j} + \dots + \sum_{j=1}^{n_r-1} (n_r - j) \bar{\alpha}_{j_{r-1}+j} \right]$$

while

$$\mu = \frac{1}{d} \left[\sum_{j=1}^{n_1-1} (n - j) \bar{\alpha}_j + \sum_{j=1}^{n_2-1} (n - n_1 - j) \bar{\alpha}_{n_1+j} + \dots + \sum_{j=1}^{n_r-1} (n_r - j) \bar{\alpha}_{j_{r-1}+j} \right].$$

Thus

$$\mu = \varphi(w) + \frac{n - n_1}{d} \sum_{j=1}^{n_1-1} \bar{\alpha}_j + \frac{n - n_1 - n_2}{d} \sum_{j=1}^{n_2-1} \bar{\alpha}_{n_1+j} + \dots + \frac{n_r}{d} \sum_{j=1}^{n_r-1} \bar{\alpha}_{j_{r-1}+j},$$

where the coefficients

$$\frac{n - n_1}{d}, \quad \frac{n - n_1 - n_2}{d}, \quad \dots, \quad \frac{n_r}{d}$$

are integral. We see that

$$\langle \bar{\alpha}_j (j \in J), \mu \rangle = \langle \bar{\alpha}_j (j \in J), \varphi(w) \rangle. \tag{3.4}$$

From (3.3) and (3.4) we obtain that

$$N = \langle \bar{\alpha}_j (j \in J), \mu \rangle = \langle \bar{\alpha}_j (j \in J), \varphi(w) \rangle = \varphi(L). \quad \square$$

3.12. Now let $p \mid \gcd(n_1, \dots, n_r)$. Recall that $W = \prod_{i=1}^r \mathfrak{S}_{n_i}$. Since $p \mid n_i$ for all i , we can naturally embed $(\mathfrak{S}_p)^{n_i/p}$ into \mathfrak{S}_{n_i} . We obtain a natural embedding

$$\Gamma := (\mathbb{Z}/p\mathbb{Z})^{n/p} \hookrightarrow (\mathfrak{S}_p)^{n/p} \hookrightarrow W.$$

In order to prove Lemma 3.8, it suffices to prove the next Lemma 3.13. Indeed, if n has an odd prime factor p , then by Lemma 3.13 L is not quasi-permutation. If $n = 2^s$, then we take $p = 2$. By the assumptions of Proposition 3.2, $n > 4 = 2^2$, and again by Lemma 3.13 L is not quasi-permutation. This proves Lemma 3.8.

LEMMA 3.13. *If either p odd or $n > p^2$, then L is not quasi-permutation as a Γ -lattice.*

Proof. By Lemma 3.11 it suffices to show that N is not quasi-permutation. Since $N = \Lambda_n(d) \cap \varphi(V)$, we have an embedding

$$\Lambda_n(d)/N \hookrightarrow \overline{V}/\varphi(V).$$

By (3.2) $\overline{V}/\varphi(V) \simeq \mathbb{R}^{r-1}$ and W acts on $\overline{V}/\varphi(V)$ trivially. Thus $\Lambda_n(d)/N \simeq \mathbb{Z}^{r-1}$ and W acts on \mathbb{Z}^{r-1} trivially. We have an exact sequence of W -lattices

$$0 \rightarrow N \rightarrow \Lambda_n(d) \rightarrow \mathbb{Z}^{r-1} \rightarrow 0,$$

with trivial action of W on \mathbb{Z}^{r-1} . We obtain that $N \sim \Lambda_n(d)$ as a W -lattice, and hence, as a Γ -lattice. Therefore, it suffices to show that $\Lambda_n(d) = Q_n(n/d)$ is not quasi-permutation as a Γ -lattice if either p is odd or $n > p^2$. This is done in [LPR] in the proofs of Propositions 7.4 and 7.8. This completes the proof of Lemma 3.13 and hence those of Lemma 3.8 and Proposition 3.2. \square

4 QUASI-PERMUTATION LATTICES – CASE \mathbf{A}_{n-1}

In this section we prove Theorem 0.2 in the special case when G is isogenous to a direct product of groups of type \mathbf{A}_{n-1} for $n \geq 3$.

We maintain the notation of Subsection 3.1. Let L be an intermediate lattice between Q and P , i.e., $Q \subset L \subset P$ ($L = Q$ or $L = P$ are possible). Let S denote the image of L in F , then L is the preimage of $S \subset F$ in P . Since W acts trivially on F , the subgroup $S \subset F$ is W -invariant, and therefore, the sublattice $L \subset P$ is W -invariant.

THEOREM 4.1. *With the notation of Subsection 3.1 assume that $n_i \geq 3$ for all $i = 1, 2, \dots, r$. Let L between Q and P be an intermediate lattice, and assume that $L \cap P_i = Q_i$ for all i such that $n_i = 3$ or $n_i = 4$. If L is a quasi-permutation W -lattice, then $L = Q$.*

Proof. We prove the theorem by induction on r . The case $r = 1$ follows from our assumptions if $n_1 = 3$ or $n_1 = 4$, and from Proposition 1.9 if $n_1 > 4$.

We assume that $r > 1$ and that the assertion is true for $r - 1$. We prove it for r .

For i between 1 and r we set

$$Q'_i = \bigoplus_{j \neq i} Q_j, \quad P'_i = \bigoplus_{j \neq i} P_j, \quad F'_i = \bigoplus_{j \neq i} F_j, \quad W'_i = \prod_{j \neq i} W_j,$$

then $Q'_i \subset Q$, $P'_i \subset P$, $F'_i \subset F$ and $W'_i \subset W$. If L is a quasi-permutation W -lattice, then by Lemma 1.7 $L \cap P'_i$ is a quasi-permutation W'_i -lattice, and by the induction hypothesis $L \cap P'_i = Q'_i$.

Now let $Q \subset L \subset P$, and assume that $L \cap P'_i = Q'_i$ for all $i = 1, \dots, r$. We shall show that if $L \neq Q$ then L is not a quasi-permutation W -lattice. This will prove Theorem 4.1.

Assume that $L \neq Q$. Set $S = L/Q \subset F$, then $S \neq 0$. We first show that $(L \cap P'_i)/Q'_i = S \cap F'_i$. Indeed, clearly $(L \cap P'_i)/Q'_i \subset L/Q \cap P'_i/Q'_i = S \cap F'_i$. Conversely, let $f \in S \cap F'_i$, then f can be represented by some $l \in L$ and by some $p \in P'_i$, and $q := l - p \in Q$. Since $L \supset Q$, we see that $p = l - q \in L \cap P'_i$, hence $f \in (L \cap P'_i)/Q'_i$, and therefore $S \cap F'_i \subset (L \cap P'_i)/Q'_i$. Thus $(L \cap P'_i)/Q'_i = S \cap F'_i$.

By assumption we have $L \cap P'_i = Q'_i$, and we obtain that $S \cap F'_i = 0$ for all $i = 1, \dots, r$. Let $S_{(i)}$ denote the image of S under the projection $F \rightarrow F_i$. We have a canonical epimorphism $p_i: S \rightarrow S_{(i)}$ with kernel $S \cap F'_i$. Since $S \cap F'_i = 0$, we see that $p_i: S \rightarrow S_{(i)}$ is an isomorphism. Set $q_i = p_i \circ p_1^{-1}: S_{(1)} \rightarrow S_{(i)}$, it is an isomorphism.

We regard $Q_i = \mathbb{Z}\mathbf{A}_{n_i-1}$ and $P_i = \Lambda_{n_i}$ as the lattices described in [Bou, Planche I]. Then we have an isomorphism $F_i \cong \mathbb{Z}/n_i\mathbb{Z}$. Since $S_{(i)}$ is a subgroup of the cyclic group $F_i \cong \mathbb{Z}/n_i\mathbb{Z}$ and $S \cong S_{(i)}$, we see that S is a cyclic group, and we see also that $|S|$ divides n_i for all i , hence $d := |S|$ divides $c := \gcd(n_1, \dots, n_r)$.

We describe all subgroups S of order d in $\bigoplus_{i=1}^r \mathbb{Z}/n_i\mathbb{Z}$ such that $S \cap (\bigoplus_{j \neq i} \mathbb{Z}/n_j\mathbb{Z}) = 0$ for all i . The element $a_i := n_i/d + n_i\mathbb{Z}$ is a generator of $S_{(i)} \subset F_i = \mathbb{Z}/n_i\mathbb{Z}$. Set $b_i = q_i(a_1)$. Since b_i is a generator of $S_{(i)}$, we have $b_i = \bar{\nu}_i a_i$ for some $\bar{\nu}_i \in (\mathbb{Z}/d\mathbb{Z})^\times$. Let $\nu_i \in \mathbb{Z}$ be a representative of $\bar{\nu}_i$ such that $1 \leq \nu_i < d$, then $\gcd(\nu_i, d) = 1$. Moreover, since $q_1 = \text{id}$, we have $b_1 = a_1$, hence $\bar{\nu}_1 = 1$ and $\nu_1 = 1$. We obtain an element $\nu = (\nu_1, \dots, \nu_r)$. With the notation of Subsection 3.1 we have $S = S_\nu$ and $L = L_\nu$.

By Proposition 3.2 L_ν is not a quasi-permutation W -lattice. Thus L is not quasi-permutation, which completes the proof of Theorem 4.1. □

5 PROOF OF THEOREM 0.2

LEMMA 5.1 (well-known). *Let P_1 and P_2 be abelian groups. Set $P = P_1 \oplus P_2 = P_1 \times P_2$, and let $\pi_1: P \rightarrow P_1$ denote the canonical projection. Let $L \subset P$ be a*

subgroup. If $\pi_1(L) = L \cap P_1$, then

$$L = (L \cap P_1) \oplus (L \cap P_2).$$

Proof. Let $x \in L$. Set $x_1 = \pi_1(x) \in \pi_1(L)$. Since $\pi_1(L) = L \cap P_1$, we have $x_1 \in L \cap P_1$. Set $x_2 = x - x_1$, then $x_2 \in L \cap P_2$. We have $x = x_1 + x_2$. This completes the proof of Lemma 5.1. \square

5.2. Let I be a finite set. For any $i \in I$ let D_i be a connected Dynkin diagram. Let $D = \bigsqcup_i D_i$ (disjoint union). Let Q_i and P_i be the root and weight lattices of D_i , respectively, and W_i be the Weyl group of D_i . Set

$$Q = \bigoplus_{i \in I} Q_i, \quad P = \bigoplus_{i \in I} P_i, \quad W = \prod_{i \in I} W_i.$$

5.3. We construct certain quasi-permutation lattices L such that $Q \subset L \subset P$.

Let $\{\{i_1, j_1\}, \dots, \{i_s, j_s\}\}$ be a set of non-ordered pairs in I such that D_{i_l} and D_{j_l} for all $l = 1, \dots, s$ are of type $\mathbf{B}_1 = \mathbf{A}_1$ and all the indices $i_1, j_1, \dots, i_s, j_s$ are distinct. Fix such an l . We write $\{i, j\}$ for $\{i_l, j_l\}$ and we set $D_{i,j} := D_i \sqcup D_j$, $Q_{i,j} := Q_i \oplus Q_j$, $P_{i,j} := P_i \oplus P_j$. We regard $D_{i,j}$ as a Dynkin diagram of type \mathbf{D}_2 , and we denote by $M_{i,j}$ the intermediate lattice between $Q_{i,j}$ and $P_{i,j}$ isomorphic to $\mathcal{X}(\mathbf{SO}_4)$, the character lattice of the group \mathbf{SO}_4 ; see Section 1, after Lemma 1.7. Let f_i be a generator of the lattice Q_i of rank 1, and let f_j be a generator of Q_j , then $P_i = \langle \frac{1}{2}f_i \rangle$ and $P_j = \langle \frac{1}{2}f_j \rangle$. Set $e_1^{(l)} = \frac{1}{2}(f_i + f_j)$, $e_2^{(l)} = \frac{1}{2}(f_i - f_j)$, then $\{e_1^{(l)}, e_2^{(l)}\}$ is a basis of $M_{i,j}$, and

$$M_{i,j} = \left\langle Q_{i,j}, e_1^{(l)} \right\rangle, \quad P_{i,j} = \left\langle M_{i,j}, \frac{1}{2}(e_1^{(l)} + e_2^{(l)}) \right\rangle. \tag{5.1}$$

We have $M_{i,j} \cap P_i = Q_i$, $M_{i,j} \cap P_j = Q_j$, and $[M_{i,j} : Q_{i,j}] = 2$. Concerning the Weyl group, we have

$$W(D_{i,j}) = W(D_i) \times W(D_j) = W(\mathbf{D}_2) = \mathfrak{S}_2 \times \{\pm 1\},$$

where the symmetric group \mathfrak{S}_2 permutes the basis vectors $e_1^{(l)}$ and $e_2^{(l)}$ of $M_{i,j}$, while the group $\{\pm 1\}$ acts on $M_{i,j}$ by multiplication by scalars. We say that $M_{i,j}$ is an *indecomposable quasi-permutation lattice* (it corresponds to the semisimple Cayley group \mathbf{SO}_4 which does not decompose into a direct product of its normal subgroups).

Set $I' = I \setminus \bigcup_{l=1}^s \{i_l, j_l\}$. For $i \in I'$ let M_i be any quasi-permutation intermediate lattice between Q_i and P_i (such an intermediate lattice exists if and only if D_i is of one of the types $\mathbf{A}_n, \mathbf{B}_n, \mathbf{C}_n, \mathbf{D}_n, \mathbf{G}_2$, see Proposition 1.9). We say that M_i is a *simple quasi-permutation lattice* (it corresponds to a stably Cayley simple group). We set

$$L = \bigoplus_{l=1}^s M_{i_l, j_l} \oplus \bigoplus_{i \in I'} M_i. \tag{5.2}$$

We say that a lattice L as in (5.2) is a *direct sum of indecomposable quasi-permutation lattices and simple quasi-permutation lattices*. Clearly L is a quasi-permutation W -lattice.

THEOREM 5.4. *Let D, Q, P, W be as in Subsection 5.2. Let L be an intermediate lattice between Q and P , i.e., $Q \subset L \subset P$ (where $L = Q$ and $L = P$ are possible). If L is a quasi-permutation W -lattice, then L is as in (5.2). Namely, then L is a direct sum of indecomposable quasi-permutation lattices $M_{i,j}$ for some set of pairs $\{\{i_1, j_1\}, \dots, \{i_s, j_s\}\}$ and some family of simple quasi-permutation intermediate lattices M_i between Q_i and P_i for $i \in I'$.*

Remark 5.5. The set of pairs $\{\{i_1, j_1\}, \dots, \{i_s, j_s\}\}$ in Theorem 5.4 is uniquely determined by L . Namely, a pair $\{i, j\}$ belongs to this set if and only if the Dynkin diagrams D_i and D_j are of type $\mathbf{B}_1 = \mathbf{A}_1$ and

$$L \cap P_i = Q_i, \quad L \cap P_j = Q_j, \quad \text{while } L \cap (P_i \oplus P_j) \neq Q_i \oplus Q_j.$$

Proof of Theorem 5.4. We prove the theorem by induction on $m = |I|$, where I is as in Subsection 5.2. The case $m = 1$ is trivial.

We assume that $m \geq 2$ and that the theorem is proved for all $m' < m$. We prove it for m . First we consider three special cases.

Split case. Assume that for some subset $A \subset I$, $A \neq I$, $A \neq \emptyset$, we have $\pi_A(L) = L \cap P_A$, where $P_A = \bigoplus_{i \in A} P_i$ and $\pi_A: P \rightarrow P_A$ is the canonical projection. Then by Lemma 5.1 we have $L = (L \cap P_A) \oplus (L \cap P_{A'})$, where $A' = I \setminus A$. By Lemma 1.7 $L \cap P_A$ is a quasi-permutation W_A -lattice, where $W_A = \prod_{i \in A} W_i$. By the induction hypothesis the lattice $L \cap P_A$ is a direct sum of indecomposable quasi-permutation lattices and simple quasi-permutation lattices. Similarly, $L \cap P_{A'}$ is such a direct sum. We conclude that $L = (L \cap P_A) \oplus (L \cap P_{A'})$ is such a direct sum, and we are done.

\mathbf{A}_{n-1} -case. Assume that all D_i are of type \mathbf{A}_{n_i-1} , where $n_i \geq 3$ (so \mathbf{A}_1 is not permitted). We assume also that when $n_i = 3$ and when $n_i = 4$ (that is, for \mathbf{A}_2 and for $\mathbf{A}_3 = \mathbf{D}_3$) we have $L \cap P_i = Q_i$ (for $n_i > 4$ this is automatic because $L \cap P_i$ is a quasi-permutation W_i -lattice, see Proposition 1.9). In this case by Theorem 4.1 we have $L = Q = \bigoplus Q_i$, hence L is a direct sum of simple quasi-permutation lattices, and we are done.

\mathbf{A}_1 -case. Assume that all D_i are of type \mathbf{A}_1 . Then by [BKLR, Theorem 18.1] the lattice L is a direct sum of indecomposable quasi-permutation lattices and simple quasi-permutation lattices, and we are done.

Now we shall show that these three special cases exhaust all the quasi-permutation lattices. In other words, we shall show that if $Q \subset L \subset P$ and L is not as in one of these three cases, then L is not quasi-permutation. This will complete the proof of the theorem.

For the sake of contradiction, let us assume that $Q \subset L \subset P$, that L is not in one of the three special cases above, and that L is a quasi-permutation W -lattice.

We shall show in three steps that L is as in Proposition 2.2. By Proposition 2.2, L is not quasi-permutation, which contradicts our assumptions. This contradiction will prove the theorem.

Step 1. For $i \in I$ consider the intersection $L \cap P_i$, it is a quasi-permutation W_i -lattice (by Lemma 1.7), hence D_i is of one of the types $\mathbf{A}_{n-1}, \mathbf{B}_n, \mathbf{C}_n, \mathbf{D}_n, \mathbf{G}_2$ (by Proposition 1.9). Note that $\pi_i(L) \neq L \cap P_i$ (otherwise we are in the split case).

Now assume that for some $i \in I$, the Dynkin diagram D_i is of type \mathbf{G}_2 or \mathbf{C}_n for some $n \geq 3$, or D_i is of type \mathbf{A}_2 and $L \cap P_i \neq Q_i$. Then $L \cap P_i$ is a quasi-permutation W_i -lattice (by Lemma 1.7), hence $L \cap P_i = P_i$ (by Proposition 1.9). Since $P_i \supset \pi_i(L) \supset L \cap P_i$, we obtain that $\pi_i(L) = L \cap P_i$, which is impossible. Thus no D_i can be of type \mathbf{G}_2 or $\mathbf{C}_n, n \geq 3$, and if D_i is of type \mathbf{A}_2 for some i , then $L \cap P_i = Q_i$.

Thus all D_i are of types $\mathbf{A}_{n-1}, \mathbf{B}_n$ or \mathbf{D}_n , and if D_i is of type \mathbf{A}_2 for some $i \in I$, then $L \cap P_i = Q_i$. Since L is not as in the \mathbf{A}_{n-1} -case, we may assume that one of the D_i , say D_1 , is of type \mathbf{B}_n for some $n \geq 1$ ($\mathbf{B}_1 = \mathbf{A}_1$ is permitted), or of type \mathbf{D}_n for some $n \geq 4$, or of type \mathbf{D}_3 with $L \cap P_1 \neq Q_1$. Indeed, otherwise all D_i are of type \mathbf{A}_{n_i-1} for $n_i \geq 3$, and in the cases \mathbf{A}_2 ($n_i = 3$) and \mathbf{A}_3 ($n_i = 4$) we have $L \cap P_i = Q_i$, i.e., we are in the \mathbf{A}_{n-1} -case, which contradicts our assumptions.

Step 2. In this step, using the Dynkin diagram D_1 of type \mathbf{B}_n or \mathbf{D}_n from the previous step, we construct a quasi-permutation sublattice $L' \subset L$ of index 2 such that L' is as in (5.2). First we consider the cases \mathbf{B}_n and \mathbf{D}_n separately.

Assume that D_1 is of type \mathbf{B}_n for some $n \geq 1$ ($\mathbf{B}_1 = \mathbf{A}_1$ is permitted). We have $[P_1 : Q_1] = 2$. Since $P_1 \supset \pi_1(L) \supsetneq L \cap P_1 \supset Q_1$, we see that $\pi_1(L) = P_1$ and $L \cap P_1 = Q_1$. Set $M_1 = Q_1$. We have $\pi_1(L) = P_1, L \cap P_1 = M_1$, and $[P_1 : M_1] = 2$.

Now assume that D_1 is of type \mathbf{D}_n for some $n \geq 4$, or of type \mathbf{D}_3 with $L \cap P_1 \neq Q_1$. Set $M_1 = L \cap P_1$, then M_1 is a quasi-permutation W_1 -lattice by Lemma 1.7, and it follows from Proposition 1.9 that $(W_1, M_1) \simeq \mathcal{X}(\mathbf{SO}_{2n})$, where $\mathcal{X}(\mathbf{SO}_{2n})$ denotes the character lattice of \mathbf{SO}_{2n} ; see Section 1, after Lemma 1.7. It follows that $[M_1 : Q_1] = 2$ and $[P_1 : M_1] = 2$. Since $P_1 \supset \pi_1(L) \supsetneq L \cap P_1 = M_1$, we see that $\pi_1(L) = P_1$. Again we have $\pi_1(L) = P_1, L \cap P_1 = M_1$, and $[P_1 : M_1] = 2$.

Now we consider together the cases when D_1 is of type \mathbf{B}_n for some $n \geq 1$ and when D_1 is of type \mathbf{D}_n for some $n \geq 3$, where in the case \mathbf{D}_3 we have $L \cap P_1 \neq Q_1$. Set

$$L' := \ker[L \xrightarrow{\pi_1} P_1 \rightarrow P_1/M_1].$$

Since $\pi_1(L) = P_1$, and $[P_1 : M_1] = 2$, we have $[L : L'] = 2$. Clearly we have $\pi_1(L') = M_1$. Set

$$L_1^\dagger := \ker[\pi_1 : L \rightarrow P_1] = L \cap P_1',$$

where $P'_1 = \bigoplus_{i \neq 1} P_i$. Since L is a quasi-permutation W -lattice, by Lemma 1.7 the lattice L^\dagger_1 is a quasi-permutation W'_1 -lattice, where $W'_1 = \prod_{i \neq 1} W_i$. By the induction hypothesis, L^\dagger_1 is a direct sum of indecomposable quasi-permutation lattices and simple quasi-permutation lattices as in (5.2). Since $M_1 = L \cap P_1$, we have $M_1 \subset L' \cap P_1$, and $L' \cap P_1 \subset L \cap P_1 = M_1$, hence $L' \cap P_1 = M_1 = \pi_1(L')$, and by Lemma 5.1 we have $L' = M_1 \oplus L^\dagger_1$. Since M_1 is a simple quasi-permutation lattice, we conclude that L' is a direct sum of indecomposable quasi-permutation lattices and simple quasi-permutation lattices as in (5.2), and $[L : L'] = 2$.

Step 3. In this step we show that L is as in Proposition 2.2. We write

$$L' = \bigoplus_{l=1}^s (L' \cap P_{i_l, j_l}) \oplus \bigoplus_{i \in I'} (L' \cap P_i),$$

where $P_{i_l, j_l} = P_{i_l} \oplus P_{j_l}$, the Dynkin diagrams D_{i_l} and D_{j_l} are of type $\mathbf{A}_1 = \mathbf{B}_1$, and $L' \cap P_{i_l, j_l} = M_{i_l, j_l}$ as in (5.1). For any $i \in I'$, we have $[\pi_i(L) : \pi_i(L')] \leq 2$, because $[L : L'] = 2$. Furthermore, for $i \in I'$ we have

$$\pi_i(L') = L' \cap P_i \subset L \cap P_i \subsetneq \pi_i(L),$$

hence $[\pi_i(L) : (L \cap P_i)] = 2$ and $L' \cap P_i = L \cap P_i$. Similarly, for any $l = 1, \dots, s$, if we write $i = i_l, j = j_l$, then we have

$$M_{i_l, j_l} = L' \cap P_{i_l, j_l} \subset L \cap P_{i_l, j_l} \subsetneq \pi_{i_l, j_l}(L) \subset P_{i_l, j_l}, \quad [P_{i_l, j_l} : M_{i_l, j_l}] = 2,$$

whence $\pi_{i_l, j_l}(L) = P_{i_l, j_l}$, $L \cap P_{i_l, j_l} = M_{i_l, j_l}$, and therefore $[\pi_{i_l, j_l}(L) : (L \cap P_{i_l, j_l})] = [P_{i_l, j_l} : M_{i_l, j_l}] = 2$ and $L' \cap P_{i_l, j_l} = M_{i_l, j_l} = L \cap P_{i_l, j_l}$.

We view the Dynkin diagram $D_{i_l} \sqcup D_{j_l}$ of type $\mathbf{A}_1 \sqcup \mathbf{A}_1$ corresponding to the pair $\{i_l, j_l\}$ ($l = 1, \dots, s$) as a Dynkin diagram of type \mathbf{D}_2 . Thus we view L' as a direct sum of indecomposable quasi-permutation lattices and simple quasi-permutation lattices corresponding to Dynkin diagrams of type $\mathbf{B}_n, \mathbf{D}_n$ and \mathbf{A}_n .

We wish to show that L is as in Proposition 2.2. We change our notation in order to make it closer to that of Proposition 2.2.

As in Subsection 2.1, we now write D_i for Dynkin diagrams of types \mathbf{B}_{l_i} and \mathbf{D}_{l_i} only, appearing in L' , where $\mathbf{B}_1 = \mathbf{A}_1, \mathbf{B}_2 = \mathbf{C}_2, \mathbf{D}_2 = \mathbf{A}_1 \sqcup \mathbf{A}_1$ and $\mathbf{D}_3 = \mathbf{A}_3$ are permitted, but for \mathbf{D}_{l_i} with $l_i = 2, 3$ we require that

$$L \cap P_i = M_i := \mathcal{X}(\mathbf{SO}_{2l_i}).$$

We write $L'_i := L \cap P_i = L' \cap P_i$. We have $[\pi_i(L) : L'_i] = 2$, hence $[P_i : L'_i] \geq 2$. If D_i is of type \mathbf{B}_{l_i} , then $[P_i : L'_i] = 2$. If D_i is of type \mathbf{D}_{l_i} , then $L'_i = L \cap P_i \neq Q_i$, for \mathbf{D}_2 and \mathbf{D}_3 by our assumption and for \mathbf{D}_{l_i} with $l_i \geq 4$ because $L \cap P_i$ is a quasi-permutation W_i -lattice (see Proposition 1.9); again we have $[P_i : L'_i] = 2$.

We see that for all i we have $[P_i : L'_i] = 2$, $\pi_i(L) = P_i$, and the lattice $L'_i = M_i$ is as in Subsection 2.1. We realize the root system $R(D_i)$ of type \mathbf{B}_{l_i} or \mathbf{D}_{l_i} in the standard way (cf. [Bou, Planches II, IV]) in the space $V_i := \mathbb{R}^{l_i}$ with basis $(e_s)_{s \in S_i}$, then L'_i is the lattice generated by the basis vectors $(e_s)_{s \in S_i}$ of V_i , and we have $P_i = \langle L'_i, \frac{1}{2}x_i \rangle$, where

$$x_i = \sum_{s \in S_i} e_s \in L'_i.$$

In particular, when D_i is of type \mathbf{D}_2 we have $x_i = e_1^{(l)} + e_2^{(l)}$ with the notation of formula (5.1).

As in Subsection 2.1, we write Δ_ι for Dynkin diagrams of type $\mathbf{A}_{n'_\iota-1}$ appearing in L' , where $n'_\iota \geq 3$ and for $\mathbf{A}_3 = \mathbf{D}_3$ we require that $L \cap P_\iota = Q_\iota$. We write $L'_\iota := L \cap P_\iota = L' \cap P_\iota$. Then $L'_\iota = Q_\iota$ for all ι , for \mathbf{A}_2 by Step 1, for \mathbf{A}_3 by our assumption, and for other $\mathbf{A}_{n'_\iota-1}$ because L'_ι is a quasi-permutation W_ι -lattice; see Proposition 1.9. We have $\pi_\iota(L) \supseteq L \cap P_\iota = L'_\iota$ and $[\pi_\iota(L) : L'_\iota] = [\pi_\iota(L) : \pi_\iota(L')] \leq 2$ (because $[L : L'] = 2$). It follows that $[\pi_\iota(L) : L'_\iota] = 2$, i.e., $[\pi_\iota(L) : Q_\iota] = 2$. We know that P_ι/Q_ι is a cyclic group of order n'_ι . Since it has a subgroup $\pi_\iota(L)/Q_\iota$ of order 2, we conclude that n'_ι is even, $n'_\iota = 2n_\iota$ (where $2n_\iota \geq 4$), and $\pi_\iota(L)/Q_\iota$ is the unique subgroup of order 2 of the cyclic group P_ι/Q_ι of order $2n_\iota$. As in Subsection 2.1, we realize the root system Δ_ι of type $\mathbf{A}_{2n_\iota-1}$ in the standard way (cf. [Bou, Planche I]) in the subspace V_ι of vectors with zero sum of the coordinates in the space \mathbb{R}^{2n_ι} with basis $\varepsilon_{\iota,1}, \dots, \varepsilon_{\iota,2n_\iota}$. We set

$$\xi_\iota = \varepsilon_{\iota,1} - \varepsilon_{\iota,2} + \varepsilon_{\iota,3} - \varepsilon_{\iota,4} + \dots + \varepsilon_{\iota,2n_\iota-1} - \varepsilon_{\iota,2n_\iota},$$

then $\xi_\iota \in L'_\iota$ and $\frac{1}{2}\xi_\iota \in \pi_\iota(L) \setminus L'_\iota$ (cf. [Bou, Planche I, formula (VI)]), hence $\pi_\iota(L) = \langle L'_\iota, \frac{1}{2}\xi_\iota \rangle$.

Now we set

$$v = \frac{1}{2} \sum_{i \in I} x_i + \frac{1}{2} \sum_{\iota=1}^{\mu} \xi_\iota.$$

We claim that

$$L = \langle L', v \rangle.$$

Proof of the claim. Let $w \in L \setminus L'$, then $L = \langle L', w \rangle$, because $[L : L'] = 2$. Set $z_i = \frac{1}{2}x_i - \pi_i(w)$, then $z_i \in L'_i \subset L'$, because $\frac{1}{2}x_i, \pi_i(w) \in \pi_i(L) \setminus L'_i$. Similarly, we set $\zeta_\iota = \frac{1}{2}\xi_\iota - \pi_\iota(w)$, then $\zeta_\iota \in L'_\iota \subset L'$. We see that

$$v = w + \sum_i z_i + \sum_\iota \zeta_\iota,$$

where $\sum_i z_i + \sum_\iota \zeta_\iota \in L'$, and the claim follows. □

It follows from the claim that L is as in Proposition 2.2 (we use the assumption that we are not in the \mathbf{A}_1 -case). Now by Proposition 2.2 L is not quasi-invertible, hence not quasi-permutation, which contradicts our assumptions. This contradiction proves Theorem 5.4. \square

Proof of Theorem 0.2. Theorem 0.2 follows immediately from Theorem 5.4 by virtue of Proposition 1.8. \square

6 PROOF OF THEOREM 0.3

In this section we deduce Theorem 0.3 from Theorem 0.2.

Let G be a stably Cayley semisimple k -group. Then $\overline{G} := G \times_k \bar{k}$ is stably Cayley over an algebraic closure \bar{k} of k . By Theorem 0.2, $G_{\bar{k}} = \prod_{j \in J} G_{j, \bar{k}}$ for some finite index set J , where each $G_{j, \bar{k}}$ is either a stably Cayley simple group or is isomorphic to $\mathbf{SO}_{4, \bar{k}}$. (Recall that $\mathbf{SO}_{4, \bar{k}}$ is stably Cayley and semisimple, but is not simple.) Here we write $G_{j, \bar{k}}$ for the factors in order to emphasize that they are defined over \bar{k} . By Remark 5.5 the collection of direct factors $G_{j, \bar{k}}$ is determined uniquely by \overline{G} . The Galois group $\text{Gal}(\bar{k}/k)$ acts on $G_{\bar{k}}$, hence on J . Let Ω denote the set of orbits of $\text{Gal}(\bar{k}/k)$ in J . For $\omega \in \Omega$ set $G_{\bar{k}}^\omega = \prod_{j \in \omega} G_{j, \bar{k}}$, then $\overline{G} = \prod_{\omega \in \Omega} G_{\bar{k}}^\omega$. Each $G_{\bar{k}}^\omega$ is $\text{Gal}(\bar{k}/k)$ -invariant, hence it defines a k -form G_k^ω of $G_{\bar{k}}^\omega$. We have $G = \prod_{\omega \in \Omega} G_k^\omega$.

For each $\omega \in \Omega$ choose $j = j_\omega \in \omega$. Let l_j/k denote the Galois extension in \bar{k} corresponding to the stabilizer of j in $\text{Gal}(\bar{k}/k)$. The subgroup $G_{j, \bar{k}}$ is $\text{Gal}(\bar{k}/l_j)$ -invariant, hence it comes from an l_j -form G_{j, l_j} . By the definition of Weil's restriction of scalars (see e.g. [Vo2, Subsection 3.12]) $G_{\bar{k}}^\omega \cong R_{l_j/k} G_{j, l_j}$, hence $G \cong \prod_{\omega \in \Omega} R_{l_j/k} G_{j, l_j}$. Each G_{j, l_j} is either absolutely simple or an l_j -form of \mathbf{SO}_4 .

We complete the proof using an argument from [BKLR, Proof of Lemma 11.1]. We show that G_{j, l_j} is a direct factor of $G_{l_j} := G \times_k l_j$. It is clear from the definition that $G_{j, \bar{k}}$ is a direct factor of $G_{\bar{k}}$ with complement $G'_{\bar{k}} = \prod_{i \in J \setminus \{j\}} G_{i, \bar{k}}$. Then $G'_{\bar{k}}$ is $\text{Gal}(\bar{k}/l_j)$ -invariant, hence it comes from some l_j -group G'_{l_j} . We have $G_{l_j} = G_{j, l_j} \times_{l_j} G'_{l_j}$, hence G_{j, l_j} is a direct factor of G_{l_j} .

Recall that G_{j, l_j} is either a form of \mathbf{SO}_4 or absolutely simple. If it is a form of \mathbf{SO}_4 , then clearly it is stably Cayley over l_j . It remains to show that if G_{j, l_j} is absolutely simple, then G_{j, l_j} is stably Cayley over l_j . The group $G_{\bar{k}}$ is stably Cayley over \bar{k} . Since $G_{j, \bar{k}}$ is a direct factor of the stably Cayley k -group $G_{\bar{k}}$ over the algebraically closed field \bar{k} , by [LPR, Lemma 4.7] $G_{j, \bar{k}}$ is stably Cayley over \bar{k} . Comparing [LPR, Theorem 1.28] and [BKLR, Theorem 1.4], we see that G_{j, l_j} is either stably Cayley over l_j (in which case we are done) or an outer form of \mathbf{PGL}_{2n} for some $n \geq 2$. Thus assume by the way of contradiction that G_{j, l_j} is an outer form of \mathbf{PGL}_{2n} for some $n \geq 2$. Then by [BKLR, Example 10.7] the character lattice of G_{j, l_j} is not quasi-invertible,

and by [BKLR, Proposition 10.8] the group G_{j,l_j} cannot be a direct factor of a stably Cayley l_j -group. This contradicts the fact that G_{j,l_j} is a direct factor of the stably Cayley l_j -group G_{l_j} . We conclude that G_{j,l_j} cannot be an outer form of \mathbf{PGL}_{2n} for any $n \geq 2$. Thus G_{j,l_j} is stably Cayley over l_j , as desired. \square

A APPENDIX: SOME QUASI-PERMUTATION CHARACTER LATTICES

The positive assertion of Proposition 1.9 above is well known. It is contained in [CK, Theorem 0.1] and in [BKLR, Theorem 1.4]. However, [BKLR] refers to [CK, Theorem 0.1], and [CK] refers to a series of results on rationality (rather than only stable rationality) of corresponding generic tori. In this appendix for the reader's convenience we provide a proof of the following positive result in terms of lattices only.

PROPOSITION A.1. *Let G be any form of one of the following groups*

$$\mathbf{SL}_3, \mathbf{PGL}_n \text{ (} n \text{ odd)}, \mathbf{SO}_n \text{ (} n \geq 3), \mathbf{Sp}_{2n}, \mathbf{G}_2$$

or an inner form of \mathbf{PGL}_n (n even). Then the character lattice of G is quasi-permutation.

Proof. \mathbf{SO}_{2n+1} . Let L be the character lattice of \mathbf{SO}_{2n+1} (including \mathbf{SO}_3). Then the Dynkin diagram is $D = \mathbf{B}_n$. The Weyl group is $W = \mathfrak{S}_n \ltimes (\mathbb{Z}/2\mathbb{Z})^n$. Then $L = \mathbb{Z}^n$ with the standard basis e_1, \dots, e_n . The group \mathfrak{S}_n naturally permutes e_1, \dots, e_n , while $(\mathbb{Z}/2\mathbb{Z})^n$ acts by sign changes. Since W permutes the basis up to \pm sign, the W -lattice L is quasi-permutation, see [Lo, §2.8].

\mathbf{SO}_{2n} , *any form, inner or outer.* Let L be the character lattice of \mathbf{SO}_{2n} (including \mathbf{SO}_4). Then the Dynkin diagram is $D = \mathbf{D}_n$, with root system $R = R(D)$. We consider the pair (A, L) where $A = \text{Aut}(R, L)$, then (A, L) is isomorphic to the character lattice of \mathbf{SO}_{2n+1} , hence is quasi-permutation.

\mathbf{Sp}_{2n} . The character lattice of \mathbf{Sp}_{2n} is isomorphic to the character lattice of \mathbf{SO}_{2n+1} , hence is quasi-permutation.

\mathbf{PGL}_n , *inner form.* The character lattice of \mathbf{PGL}_n is the root lattice $L = Q$ of \mathbf{A}_{n-1} . It is a quasi-permutation \mathfrak{S}_n -lattice, cf. [Lo, Example 2.8.1].

\mathbf{PGL}_n , *outer form, n odd.* Let P be the *weight* lattice of \mathbf{A}_{n-1} , where $n \geq 3$ is odd. Then P is generated by elements e_1, \dots, e_n subject to the relation

$$e_1 + \dots + e_n = 0.$$

The automorphism group $A = \text{Aut}(\mathbf{A}_{n-1})$ is the product of \mathfrak{S}_n and \mathfrak{S}_2 . The group A acts on P as follows: \mathfrak{S}_n permutes e_1, \dots, e_n , and the nontrivial element of \mathfrak{S}_2 takes each e_i to $-e_i$.

We denote by M the A -lattice of rank $2n+1$ with basis $s_1, \dots, s_n, t_1, \dots, t_n, u$. The group \mathfrak{S}_n permutes s_i and permutes t_i ($i = 1, \dots, n$), and the nontrivial

element of \mathfrak{S}_2 permutes s_i and t_i for each i . The group A acts trivially on u . Clearly M is a permutation lattice.

We define an A -epimorphism $\pi: M \rightarrow P$ as follows:

$$\pi: \quad s_i \mapsto e_i, \quad t_i \mapsto -e_i, \quad u \mapsto 0.$$

Set $M' = \ker \pi$, it is an A -lattice of rank $n+2$. We show that it is a permutation lattice. We write down a set of $n+3$ generators of M' :

$$\rho_i = s_i + t_i, \quad \sigma = s_1 + \cdots + s_n, \quad \tau = t_1 + \cdots + t_n, \quad u.$$

There is a relation

$$\rho_1 + \cdots + \rho_n = \sigma + \tau.$$

We define a new set of $n+2$ generators:

$$\tilde{\rho}_i = \rho_i + u, \quad \tilde{\sigma} = \sigma + \frac{n-1}{2}u, \quad \tilde{\tau} = \tau + \frac{n-1}{2}u,$$

where $\frac{n-1}{2}$ is integral because n is odd. We have

$$\tilde{\rho}_1 + \cdots + \tilde{\rho}_n - \tilde{\sigma} - \tilde{\tau} = u,$$

hence this new set indeed generates M' , hence it is a basis. The group \mathfrak{S}_n permutes $\tilde{\rho}_1, \dots, \tilde{\rho}_n$, while \mathfrak{S}_2 permutes $\tilde{\sigma}$ and $\tilde{\tau}$. Thus A permutes our basis, and therefore M' is a permutation lattice. We have constructed a left resolution of P :

$$0 \rightarrow M' \rightarrow M \rightarrow P \rightarrow 0,$$

(with permutation lattices M and M'), which by duality gives a right resolution of the root lattice $Q \cong P^\vee$ of \mathbf{A}_{n-1} :

$$0 \rightarrow Q \rightarrow M^\vee \rightarrow (M')^\vee \rightarrow 0$$

with permutation lattices M^\vee and $(M')^\vee$. Thus the character lattice Q of \mathbf{PGL}_n is a quasi-permutation A -lattice for odd n .

The assertion that the character lattice of G is quasi-permutation in the remaining cases \mathbf{SL}_3 and \mathbf{G}_2 follows from the next Lemma A.2.

LEMMA A.2 ([BKLR, Lemma 2.5]). *Let Γ be a finite group and L be any Γ -lattice of rank $r = 1$ or 2 . Then L is quasi-permutation.*

This lemma, which is a version of [Vo2, § 4.9, Examples 6 and 7], was stated in [BKLR] without proof. For the sake of completeness we supply a short proof here.

We may assume that Γ is a maximal finite subgroup of $\mathbf{GL}_r(\mathbb{Z})$. If $r = 1$, then $\mathbf{GL}_1(\mathbb{Z}) = \{\pm 1\}$, and the lemma reduces to the case of the character lattice of \mathbf{SO}_3 treated above.

Now let $r = 2$. Up to conjugation there are two maximal finite subgroups of $\mathbf{GL}_2(\mathbb{Z})$, they are isomorphic to the dihedral groups D_8 (of order 8) and to D_{12} (of order 12), resp., see e.g. [Lo, § 1.10.1, Table 1.2]. The group D_8 is the group of symmetries of a square, and in this case it suffices to show that the character lattice of \mathbf{SO}_5 is quasi-permutation, which we have done above. The group D_{12} is the group of symmetries of a regular hexagon, and in this case it suffices to show that the character lattice of \mathbf{PGL}_3 (outer form) is quasi-permutation, which we have done above as well. This completes the proofs of Lemma A.2 and Proposition A.1. \square

ACKNOWLEDGEMENTS. The authors are very grateful to the anonymous referee for prompt and thorough refereeing the paper and for noticing a (correctable) error in Theorem 4.1 and in the proof of Theorem 5.4. The authors thank Rony A. Bitan for his help in proving Lemma 3.8. The first-named author was supported in part by the Hermann Minkowski Center for Geometry. The second-named author was supported in part by the Israel Science Foundation, grant 1207/12, and by the Minerva Foundation through the Emmy Noether Institute for Mathematics.

REFERENCES

- [BKLR] M. Borovoi, B. Konyavskii, N. Lemire and Z. Reichstein, Stably Cayley groups in characteristic 0, *Int. Math. Res. Not.* 2014, no. 19, 5340–5397.
- [Bou] N. Bourbaki, *Groupes et algèbres de Lie. Chap. IV-VI*, Hermann, Paris, 1968.
- [CTS] J.-L. Colliot-Thélène et J.-J. Sansuc, La R -équivalence sur les tores, *Ann. Sci. École Norm. Sup.* (4) 10 (1977), 175–229.
- [CK] A. Cortella and B. Konyavskii, Rationality problem for generic tori in simple groups, *J. Algebra* 225 (2000), 771–793.
- [LPR] N. Lemire, V.L. Popov and Z. Reichstein, Cayley groups, *J. Amer. Math. Soc.* 19 (2006), 921–967.
- [Lo] M. Lorenz, *Multiplicative Invariant Theory*, Encyclopaedia of Mathematical Sciences, 135, Invariant Theory and Algebraic Transformation Groups, VI, Springer-Verlag, Berlin, 2005.
- [Ro] A. V. Roiter, On integral representations belonging to one genus, *Izv. Akad. Nauk SSSR Ser. Mat.* 30 (1966), 1315–1324; English transl.: *Amer. Math. Soc. Transl.* (2) 71 (1968), 49–59.

- [Sw] R. G. Swan, Noether's problem in Galois theory, in: *Emmy Noether in Bryn Mawr (Bryn Mawr, Pa., 1982)*, 21–40, Springer, New York, 1983.
- [Vo1] V. E. Voskresenskiĭ, Birational properties of linear algebraic groups, *Izv. Akad. Nauk SSSR Ser. Mat.* 34 (1970), 3–19; English transl.: *Math. USSR Izv.* 4 (1970), 1–17.
- [Vo2] V. E. Voskresenskiĭ, *Algebraic Groups and Their Birational Invariants*, Transl. Math. Monographs, vol. 179, Amer. Math. Soc., Providence, RI, 1998.

Mikhail Borovoi
Raymond and Beverly
Sackler School of
Mathematical Sciences,
Tel Aviv University
6997801 Tel Aviv
Israel
borovoi@post.tau.ac.il

Boris Kunyavskiĭ
Department of Mathematics
Bar-Ilan University
5290002 Ramat Gan
Israel
kunyav@macs.biu.ac.il