

Research Article

Differentiability Properties of the Pre-Image Pressure

Kesong Yan,^{1,2} Fanping Zeng,^{2,3} and Gengrong Zhang³

¹ Department of Mathematics, University of Science and Technology of China, Hefei, Anhui 230026, China

² Department of Mathematics and Computer Science, Liuzhou Teachers College, Liuzhou, Guangxi 545004, China

³ Institute of Mathematics, Guangxi University, Nanning, Guangxi 530004, China

Correspondence should be addressed to Kesong Yan, ksyang@mail.ustc.edu.cn

Received 8 March 2012; Accepted 26 March 2012

Academic Editor: M. De la Sen

Copyright © 2012 Kesong Yan et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We study the differentiability properties of the pre-image pressure. For a TDS (X, T) with finite topological pre-image entropy, we prove the pre-image pressure function $P_{\text{pre}}(T, \bullet)$ is Gateaux differentiable at $f \in C(X, \mathbb{R})$ if and only if $P_{\text{pre}}(T, \bullet)$ has a unique tangent functional at f . Also, we obtain some equivalent conditions for $P_{\text{pre}}(T, \bullet)$ to be Fréchet differentiable at f .

1. Introduction

By a topological dynamical system (for short TDS), we mean a pair (X, T) where X is a compact metric space and $T : X \rightarrow X$ is a continuous surjection from X to itself. Entropies are fundamental to our current understanding of dynamical systems. The classical measure-theoretic entropy for an invariant measure and the topological entropy were introduced in [1, 2], respectively, and the classical variational principle was completed in [3, 4]. Topological entropy measures the maximal exponential growth rate of orbits for arbitrary topological dynamical systems, and measure-theoretic entropy measures the maximal loss of information for the iteration of finite partitions in a measure-preserving transformation.

Topological pressure is a generalization of topological entropy for a dynamical system. The notion was first introduced by Ruelle [5] in 1973 for an expansive dynamical system and later by Walters [6] for the general case. The theory related to the topological pressure, variational principle, and equilibrium states plays a fundamental role in statistical mechanics, ergodic theory, and dynamical systems (see, e.g., the books [7–12]). Since the works of Bowen [13] and Ruelle [14], the topological pressure has become a basic tool in the dimension theory

related to dynamical systems. One of the basic questions of physical interest in that of differentiability of the pressure. The differentiability of the pressure was considered by many people (see, e.g., [15–18]).

Recently, the pre-image structure of maps has become deeply characterized via entropies. In several papers (see [19–24]), some important pre-image entropy invariants of dynamical systems have been introduced and their relationships with topological entropy have been established. In a certain sense, these new invariants give a quantitative estimate of how “noninvertible” a system is. In [25], we defined the topological pre-image pressure of topological dynamical systems, which is a generalization of the Cheng-Newhouse pre-image entropy (see [19]), and proved a variational principle for it. We gave some applications of the pre-image pressure to equilibrium states (see [25, 26]). Under the assumption that $h_{\text{pre}}(T) < \infty$ and the metric pre-image entropy function $h_{\{\text{pre}, \bullet\}}(T)$ is upper semicontinuous, we obtained a way to describe a kind of continuous dependence of equilibrium states. Also, we proved that the set of all continuous functions with unique equilibrium states is a dense G_δ -set of $C(X, \mathbb{R})$, and for any finite collection of ergodic measures, we can find some continuous function such that its set of equilibrium states contains the given set (see [26]).

The purpose of this paper is to study the differentiability properties of the pre-image pressure of the TDS (X, T) with finite topological pre-image entropy. In Section 2, we concentrate on reviewing some basic definitions and give some basic properties of tangent functionals to the pre-image pressure.

In Section 3, the Gateaux differentiability of the pre-image pressure is discussed. We show that the pre-image pressure function $P_{\text{pre}}(T, \bullet)$ is Gateaux differentiable at $f \in C(X, \mathbb{R})$ if and only if it has a unique tangent functional at f .

In Section 4, we discuss the Fréchet differentiability of the pre-image pressure. We obtain some equivalent conditions for $P_{\text{pre}}(T, \bullet)$ to be Fréchet differentiable at f . Also, we show that the pre-image function $P_{\text{pre}}(T, \bullet)$ is Fréchet differentiable if and only if (X, T) is uniquely ergodic, and hence the pre-pressure is linear.

2. Preliminaries

Throughout the paper, let (X, T) be a TDS with finite topological pre-image entropy $h_{\text{pre}}(T)$ (see [19] for definition). In this section, we will recall some basic definitions and give some useful properties.

Let (X, T) be a TDS and let $\mathcal{B}(X)$ be the collection of all Borel subsets of X . Recall that a *cover* of X is a family of Borel subsets of X whose union is X . An *open cover* is one that consists of open sets. A *partition* of X is a cover of X consisting of pairwise disjoint sets. We denote the set of finite covers, finite open covers, and finite partition of X by \mathcal{C}_X , \mathcal{C}_X^o , and \mathcal{P}_X , respectively. Given two covers \mathcal{U}, \mathcal{V} , \mathcal{U} is said to be *finer* than \mathcal{V} (denoted by $\mathcal{U} \geq \mathcal{V}$) if each element of \mathcal{U} is contained in some element of \mathcal{V} . We set $\mathcal{U} \vee \mathcal{V} = \{U \cap V : U \in \mathcal{U}, V \in \mathcal{V}\}$.

Denote by $C(X, \mathbb{R})$ the Banach space of all continuous, real-valued functions on X endowed with the supremum norm. For $f \in C(X, \mathbb{R})$ and $n \in \mathbb{N}$, we denote $\sum_{i=0}^{n-1} f(T^i(x))$ by $(S_n f)(x)$.

2.1. Topological Pre-Image Pressure

In an early paper with Zeng et al. [25], following the idea of topological pressure (see Chapter 9, [12]), we defined a new notion of pre-image pressure, which extends the Cheng-Newhouse pre-image entropy $h_{\text{pre}}(T)$ [19]. For a given TDS (X, T) , the pre-image pressure of

T is a map $P_{\text{pre}}(T, \bullet) : C(X, \mathbb{R}) \rightarrow \mathbb{R}$ which is convex, Lipschitz continuous, increasing, with $P_{\text{pre}}(T, 0) = h_{\text{pre}}(T)$. More precisely, let $\mathcal{U} \in \mathcal{C}_X^o$. For $x \in X$ and $k \in \mathbb{N}$, we put

$$P_n(T, f, \mathcal{U}, T^{-k}(x)) := \inf_{\mathcal{U}} \sum_{B \in \mathcal{U}} \sup_{y \in B} e^{(S_n f)(y)}, \quad (2.1)$$

where the infimum is taken over all finite subcovers \mathcal{U} of $\bigvee_{i=0}^{n-1} T^{-i} \mathcal{U}$ with respect to $T^{-k}(x)$. We define the *pre-image pressure of T related to \mathcal{U} at f* as

$$P_{\text{pre}}(T, f, \mathcal{U}) := \lim_{n \rightarrow \infty} \frac{1}{n} \log P_{\text{pre}, n}(T, f, \mathcal{U}), \quad (2.2)$$

where $P_{\text{pre}, n}(T, f, \mathcal{U}) := \sup_{x \in X, k \geq n} P_n(T, f, \mathcal{U}, T^{-k}(x))$. The *pre-image pressure of T at f* is defined by

$$P_{\text{pre}}(T, f) := \sup_{\mathcal{U} \in \mathcal{C}_X^o} P_{\text{pre}}(T, f, \mathcal{U}). \quad (2.3)$$

It is clear that $P_{\text{pre}}(T, f) \leq P(T, f)$ (topological pressure, see [12]) and $P_{\text{pre}}(T, 0) = h_{\text{pre}}(T)$. $P_{\text{pre}}(T, f) \leq \|f\|$ if T is a homeomorphism.

2.2. Measure-Theoretic Pre-Image Entropy

Denote by $\mathcal{M}(X)$, $\mathcal{M}(X, T)$, and $\mathcal{M}^e(X, T)$ the set of all Borel probability measures, T -invariant Borel probability measures and T -invariant ergodic measures, on X , respectively. Note that $\mathcal{M}(X, T) \supseteq \mathcal{M}^e(X, T) \neq \emptyset$, and both $\mathcal{M}(X)$ and $\mathcal{M}(X, T)$ are convex compact metric spaces when endowed with the weak*-topology; $\mathcal{M}^e(X, T)$ is a G_δ subset of $\mathcal{M}(X, T)$ (see [12, Chapter 6]). Beside the weak*-topology on $\mathcal{M}(X, T)$, we also have the norm topology arising from the metric:

$$\|\mu - \nu\| := \sup \left\{ \left| \int g d\mu - \int g d\nu \right| : g \in C(X, \mathbb{R}), \|g\| \leq 1 \right\}. \quad (2.4)$$

Note that $\|\mu - \nu\| = 2$ if $\mu \neq \nu \in \mathcal{M}^e(X, T)$.

Given $\alpha \in \mathcal{D}_X$, $\mu \in \mathcal{M}(X)$ and a sub- σ -algebra $\mathcal{A} \subseteq \mathcal{B}(X)$, define

$$H_\mu(\alpha | \mathcal{A}) := \sum_{A \in \alpha} \int_X -\mathbb{E}(1_A | \mathcal{A}) \log \mathbb{E}(1_A | \mathcal{A}) d\mu, \quad (2.5)$$

where $\mathbb{E}(1_A | \mathcal{A})$ is the conditional expectation of 1_A with respect to \mathcal{A} . It is a standard fact that $H_\mu(\alpha | \mathcal{A})$ increases with respect to α and decreases with respect to \mathcal{A} . Note that $\beta \in \mathcal{D}_X$ naturally generates a sub- σ -algebra $\mathcal{F}(\beta)$ of $\mathcal{B}(X)$; where there is no ambiguity, we write $\mathcal{F}(\beta)$ as β . It is easy to check, for $\alpha, \beta \in \mathcal{D}_X$, that $H_\mu(\alpha | \beta) = H_\mu(\alpha \vee \beta) - H_\mu(\beta)$. More generally, for a sub- σ -algebra $\mathcal{A} \subseteq \mathcal{B}(X)$, we have

$$H_\mu(\alpha \vee \beta | \mathcal{A}) = H_\mu(\beta | \mathcal{A}) + H_\mu(\alpha | \beta \vee \mathcal{A}). \quad (2.6)$$

When $\mu \in \mathcal{M}(X, T)$ and $\mathcal{A} \subseteq \mathcal{B}(X)$ is a T -invariant sub- σ -algebra, that is, $T^{-1}\mathcal{A} = \mathcal{A}$ (up to μ -null sets), it is not hard to see that $a_n = H_\mu(\bigvee_{i=0}^{n-1} T^{-i}\alpha \mid \mathcal{A})$ is a nonnegative subadditive sequence for a given $\alpha \in \mathcal{D}_X$, that is, $a_{n+m} \leq a_n + a_m$ for all positive integers n and m . It is well known that

$$\lim_{n \rightarrow \infty} \frac{a_n}{n} = \inf_{n \geq 1} \frac{a_n}{n}. \quad (2.7)$$

The conditional entropy of α with respect to \mathcal{A} is then defined by

$$h_\mu(T, \alpha \mid \mathcal{A}) := \lim_{n \rightarrow \infty} \frac{1}{n} H_\mu \left(\bigvee_{i=0}^{n-1} T^{-i}\alpha \mid \mathcal{A} \right) = \inf_{n \geq 1} \frac{1}{n} H_\mu \left(\bigvee_{i=0}^{n-1} T^{-i}\alpha \mid \mathcal{A} \right). \quad (2.8)$$

Moreover, the metric conditional entropy of (X, T) with respect to \mathcal{A} is defined by

$$h_\mu(T, X \mid \mathcal{A}) = \sup_{\alpha \in \mathcal{D}_X} h_\mu(T, \alpha \mid \mathcal{A}). \quad (2.9)$$

Note that if $\mathcal{N} = \{\emptyset, X\}$ is a trivial sub- σ -algebra, we recover the metric entropy, and we write $h_\mu(T, \alpha \mid \mathcal{N})$ and $h_\mu(T, X \mid \mathcal{N})$ simple by $h_\mu(T, \alpha)$ and $h_\mu(T)$.

Particularly, set $\mathcal{B}^- = \bigcap_{n=0}^{\infty} T^{-n}\mathcal{B}(X)$, then \mathcal{B}^- is a T -invariant sub- σ algebra. We call \mathcal{B}^- the infinite past σ -algebra related to $\mathcal{B}(X)$. We define the measure-theoretic (or metric) pre-image entropy of α with respect to (X, T) by

$$h_{\text{pre}, \mu}(T, \alpha) := h_\mu(T, \alpha \mid \mathcal{B}^-) = \lim_{n \rightarrow \infty} \frac{1}{n} H_\mu \left(\bigvee_{i=0}^{n-1} T^{-i}\alpha \mid \mathcal{B}^- \right). \quad (2.10)$$

Moreover, we define the metric pre-image entropy of (X, T) by

$$h_{\text{pre}, \mu}(T) := \sup_{\alpha \in \mathcal{D}_X} h_{\text{pre}, \mu}(T, \alpha). \quad (2.11)$$

2.3. A Variational Principle for Pre-Image Pressure

The following variational relationship for topological pre-image pressure and measure-theoretic pre-image entropy is established in [25].

Theorem 2.1. *Let (X, T) be a TDS and $f \in C(X, \mathbb{R})$. Then,*

$$P_{\text{pre}}(T, f) = \sup_{\mu \in \mathcal{M}(X, T)} \left\{ h_{\text{pre}, \mu}(T) + \int_X f \, d\mu \right\}. \quad (2.12)$$

We also have (see, e.g., [26]) the following proposition.

Proposition 2.2. *Let (X, T) be a TDS, $\alpha \in \mathcal{D}_X$ and $\mu \in \mathcal{M}(X, T)$. Then, $\mu \mapsto h_{\text{pre}, \mu}(T, \alpha)$ and $\mu \mapsto h_{\text{pre}, \mu}(T)$ are both affine functions on $\mathcal{M}(X, T)$. Moreover, if the ergodic decomposition of μ is*

$\mu = \int_{\mathcal{M}^e(X,T)} \theta d\lambda(\theta)$, then

$$h_{\text{pre},\mu}(T, \alpha) = \int_{\mathcal{M}^e(X,T)} h_{\text{pre},\theta}(T, \alpha) d\lambda(\theta), \quad h_{\text{pre},\mu}(T) = \int_{\mathcal{M}^e(X,T)} h_{\text{pre},\theta}(T) d\lambda(\theta). \quad (2.13)$$

2.4. Equilibrium States and Tangent Functionals to Pre-Image Pressure

Given $f \in C(X, \mathbb{R})$. A finite signed Borel measure μ on $(X, \mathcal{B}(X))$ is called a *tangent functional* to $P_{\text{pre}}(T, \bullet)$ at f if

$$P_{\text{pre}}(T, f + g) - P_{\text{pre}}(T, f) \geq \int g d\mu, \quad \forall g \in C(X, \mathbb{R}). \quad (2.14)$$

Let $\mathcal{T}_f(X, T)$ denote the collection of all tangent functionals to $P_{\text{pre}}(T, \bullet)$ at f . An application of the Hahn-Banach theorem gives $\mathcal{T}_f(X, T) \neq \emptyset$. It is easy to see that $\mu \in \mathcal{T}_f(X, T)$ if and only if

$$P_{\text{pre}}(T, f) - \int f d\mu = \inf \left\{ P_{\text{pre}}(T, g) - \int g d\mu : g \in C(X, \mathbb{R}) \right\}. \quad (2.15)$$

Also, we have $\mathcal{T}_f(X, T) \subseteq \mathcal{M}(X, T)$ (see [25] for details).

Theorem 2.3. *The following holds.*

(1) For $f \in C(X, \mathbb{R})$,

$$\mathcal{T}_f(X, T) = \overline{\bigcap_{n=1}^{\infty} \left\{ \mu \in \mathcal{M}(X, T) : h_{\text{pre},\mu}(T) + \int f d\mu > P_{\text{pre}}(T, f) - \frac{1}{n} \right\}}. \quad (2.16)$$

(2) If $f_1, f_2 \in C(X, \mathbb{R})$ and $\mu \in \mathcal{T}_{f_1}(X, T) \cap \mathcal{T}_{f_2}(X, T)$, then

$$P_{\text{pre}}(T, pf_1 + (1-p)f_2) = P_{\text{pre}}(T, f_1) + (1-p) \int (f_1 - f_2) d\mu, \quad \forall p \in [0, 1], \quad (2.17)$$

and $\mathcal{T}_{pf_1+(1-p)f_2}(X, T) \subseteq \mathcal{T}_{f_1}(X, T) \cap \mathcal{T}_{f_2}(X, T)$.

Proof. (1) Let $\mu \in I_f \equiv \overline{\bigcap_{n=1}^{\infty} \left\{ \mu \in \mathcal{M}(X, T) : h_{\text{pre},\mu}(T) + \int f d\mu > P_{\text{pre}}(T, f) - 1/n \right\}}$. By Theorem 2.1, there is $\mu_n \in \mathcal{M}(X, T)$ with $\mu_n \rightarrow \mu$ and $h_{\text{pre},\mu_n}(T) + \int f d\mu_n \rightarrow P_{\text{pre}}(T, f)$. Hence, for each $g \in C(X, \mathbb{R})$,

$$P_{\text{pre}}(T, f + g) - P_{\text{pre}}(T, f) \geq \lim_{n \rightarrow \infty} \left(h_{\text{pre},\mu_n}(T) + \int (f + g) d\mu_n - P_{\text{pre}}(T, f) \right) = \int g d\mu, \quad (2.18)$$

which follows that $\mu \in \mathcal{T}_f(X, T)$. Now suppose there is $\mu_0 \in \mathcal{T}_f(X, T) \setminus I_f$. Since I_f is convex, the standard separation theorem [27, page 417] follows that there exists $g \in C(X, \mathbb{R})$ with

$$\int g \, d\mu_0 > \sup \left\{ \int g \, d\mu : \mu \in I_f \right\}. \quad (2.19)$$

By Theorem 2.1, we can choose $\mu_n \in \mathcal{M}(X, T)$ such that

$$h_{\text{pre}, \mu_n}(T) + \int \left(f + \frac{g}{n} \right) d\mu_n > P_{\text{pre}} \left(T, f + \frac{g}{n} \right) - \frac{1}{n^2}. \quad (2.20)$$

Without loss of generality, we can assume $\mu_n \rightarrow \mu^*$. Then,

$$\begin{aligned} \int g \, d\mu_0 &= n \int \frac{g}{n} d\mu_0 \leq n \left(P_{\text{pre}} \left(T, f + \frac{g}{n} \right) - P_{\text{pre}}(T, f) \right) \\ &< n \left(h_{\text{pre}, \mu_n}(T) + \int \left(f + \frac{g}{n} \right) d\mu_n + \frac{1}{n^2} - h_{\text{pre}, \mu_n}(T) - \int f \, d\mu_n \right) \\ &= \int g \, d\mu_n + \frac{1}{n} \rightarrow \int g \, d\mu^*. \end{aligned} \quad (2.21)$$

However, $\mu^* \in I_f$ follows from the fact:

$$\begin{aligned} h_{\text{pre}, \mu_n}(T) + \int f \, d\mu_n &> P_{\text{pre}} \left(T, f + \frac{g}{n} \right) - \int \frac{g}{n} \, d\mu_n - \frac{1}{n^2} \quad (\text{by (2.20)}) \\ &> P_{\text{pre}}(T, f) - \frac{2 \cdot \|g\|}{n} - \frac{1}{n^2}, \quad (\text{by [25, Lemma 4.1(3)]}), \end{aligned} \quad (2.22)$$

which is a contradiction.

(2) If $0 < p < 1$, then

$$\begin{aligned} &pP_{\text{pre}}(T, f_1) + (1-p)P_{\text{pre}}(T, f_2) \\ &\geq P_{\text{pre}}(T, pf_1 + (1-p)f_2) \quad (\text{by [25, Lemma 4.1(3)]}) \\ &= P_{\text{pre}}(T, f_1 + (1-p)(f_2 - f_1)) \\ &\geq P_{\text{pre}}(T, f_1) + (1-p) \int (f_2 - f_1) \, d\mu \quad (\text{since } \mu \in \mathcal{T}_{f_1}(X, T)). \end{aligned} \quad (2.23)$$

Hence,

$$P_{\text{pre}}(T, f_2) - \int f_2 \, d\mu \geq P_{\text{pre}}(T, f_1) - \int f_1 \, d\mu. \quad (2.24)$$

By symmetry, $P_{\text{pre}}(T, f_2) - \int f_2 d\mu = P_{\text{pre}}(T, f_1) - \int f_1 d\mu$, which implies

$$P_{\text{pre}}(T, f_1) + (1-p) \int (f_2 - f_1) d\mu = p P_{\text{pre}}(T, f_1) + (1-p) P_{\text{pre}}(T, f_2) \geq P_{\text{pre}}(T, p f_1 + (1-p) f_2). \quad (2.25)$$

A member $\mu \in \mathcal{M}(X, T)$ is called an *equilibrium state* for $P_{\text{pre}}(T, \bullet)$ at f if

$$P_{\text{pre}}(T, f) = h_{\text{pre}, \mu}(T) + \int f d\mu. \quad (2.26)$$

Let $\mathcal{M}_f(X, T)$ denote the collection of all equilibrium states for $P_{\text{pre}}(T, \bullet)$ at f .

The set $\mathcal{T}_f(X, T)$ is convex and compact in the weak*-topology. The set $\mathcal{M}_f(X, T)$ is convex but it may be not closed in the weak*-topology. Note that $\mathcal{M}_f(X, T) \subseteq \mathcal{T}_f(X, T) \subseteq \mathcal{M}(X, T)$ and $\mathcal{M}_f(X, T)$ could be empty (see Example 5.1, [25]). We also have $\mathcal{M}_f(X, T) = \mathcal{T}_f(X, T)$ if and only if the metric pre-image entropy map $h_{\text{pre}, \bullet}(T)$ is upper semicontinuous at every element of $\mathcal{T}_f(X, T)$, Theorem 5.2 [25]. The extreme points of $\mathcal{M}_f(X, T)$ are precisely the ergodic members of $\mathcal{M}_f(X, T)$ and if $\mu \in \mathcal{M}_f(X, T)$, then almost every measure in the ergodic decomposition of μ is a member of $\mathcal{M}_f(X, T)$ (see Proposition 2.1, [26]). When the metric pre-image entropy map $h_{\text{pre}, \bullet}(T)$ is upper semicontinuous on $\mathcal{M}(X, T)$, then $\bigcup_{f \in C(X, \mathbb{R})} \mathcal{M}_f(X, T)$ is dense in $\mathcal{M}(X, T)$ in the norm topology, and given any finite collection of ergodic measures $\{\mu_1, \dots, \mu_n\}$, there is some $f \in C(X, \mathbb{R})$ such that $\{\mu_1, \dots, \mu_n\} \subseteq \mathcal{M}_f(X, T)$ [26, Theorem 4.2].

The following theorem shows when tangent functionals to pre-image pressure are not equilibrium states.

Theorem 2.4. *Let (X, T) be a TDS and $f \in C(X, \mathbb{R})$. The following statements are mutually equivalent:*

- (1) $\mu \in \mathcal{T}_f(X, T) \setminus \mathcal{M}_f(X, T)$;
- (2) $h_{\text{pre}, \mu}(T) + \int f d\mu < P_{\text{pre}}(T, f)$ and there exist $\{\mu_n\}_{n=1}^{\infty} \subseteq \mathcal{M}(X, T)$ with $\mu_n \rightarrow \mu$ and $h_{\text{pre}, \mu_n}(T) + \int f d\mu_n \rightarrow P_{\text{pre}}(T, f)$;
- (3) $\mu \in \mathcal{T}_f(X, T)$ and $h_{\text{pre}, \bullet}(T)$ is not upper semicontinuous at μ .

Proof. (1) \Rightarrow (2) Follows from the variational principle and Theorem 2.3(1).

(2) \Rightarrow (3) By Theorem 2.3(1), $\mu \in \mathcal{T}_f(X, T)$. If $h_{\text{pre}, \bullet}(T)$ is upper semicontinuous at μ , then

$$P_{\text{pre}}(T, f) = \lim_{n \rightarrow \infty} \left(h_{\text{pre}, \mu_n}(T) + \int f d\mu_n \right) \leq h_{\text{pre}, \mu}(T) + \int f d\mu. \quad (2.27)$$

Hence, $\mu \in \mathcal{M}_f(X, T)$ by the variational principle.

(3) \Rightarrow (1) If (3) holds, then there are $\mu_n \in \mathcal{M}(X, T)$ with $\mu_n \rightarrow \mu$ and $\lim_{n \rightarrow \infty} h_{\text{pre}, \mu_n}(T) > h_{\text{pre}, \mu}(T)$. Hence,

$$P_{\text{pre}}(T, f) \geq \lim_{n \rightarrow \infty} \left(h_{\text{pre}, \mu_n}(T) + \int f d\mu_n \right) > h_{\text{pre}, \mu}(T) + \int f d\mu. \quad (2.28)$$

Therefore, $\mu \in \mathcal{T}_f(X, T) \setminus \mathcal{M}_f(X, T)$. □

3. Gateaux Differentiability of the Pre-Image Pressure

In [26], we studied the uniqueness of the equilibrium state for the pre-image pressure. We showed that when the metric pre-image entropy map $h_{\text{pre},\bullet}(T)$ is upper semicontinuous on $\mathcal{M}(X, T)$, then the set of all functions with unique equilibrium state is dense in $C(X, \mathbb{R})$. Without the upper semicontinuity assumption, one can show that all functions with unique tangent functional are dense in $C(X, \mathbb{R})$ (can see [27, page 450] or [11, Appendix A.3.6]). In this section, we will show a continuous function with unique tangent functional to pre-image pressure if and only if it is Gateaux differentiable.

Given $f, g \in C(X, \mathbb{R})$. Since $P_{\text{pre}}(T, \bullet)$ is convex, the map $t \mapsto (P_{\text{pre}}(T, f + tg) - P_{\text{pre}}(T, f))/t$ is increasing and hence

$$\begin{aligned} d^+ P_{\text{pre}}(T, f)(g) &= \lim_{t \rightarrow 0^+} \frac{P_{\text{pre}}(T, f + tg) - P_{\text{pre}}(T, f)}{t}, \\ d^- P_{\text{pre}}(T, f)(g) &= \lim_{t \rightarrow 0^-} \frac{P_{\text{pre}}(T, f + tg) - P_{\text{pre}}(T, f)}{t} \end{aligned} \quad (3.1)$$

exist. Note that $d^+ P_{\text{pre}}(T, f)(g) = -d^- P_{\text{pre}}(T, f)(-g)$. The pre-image pressure function $P_{\text{pre}}(T, \bullet)$ is said to be *Gateaux differentiable at f* if, for all $g \in C(X, \mathbb{R})$,

$$\lim_{t \rightarrow 0} \frac{P_{\text{pre}}(T, f + tg) - P_{\text{pre}}(T, f)}{t} \quad (3.2)$$

exist. It is easy to check that $P_{\text{pre}}(T, \bullet)$ is Gateaux differentiable at f if and only if $g \mapsto d^+ P_{\text{pre}}(T, f)(g)$ is linear.

Lemma 3.1. *Let (X, T) be a TDS and $f, g \in C(X, \mathbb{R})$. Then,*

$$d^+ P_{\text{pre}}(T, f)(g) = \sup \left\{ \int g \, d\mu : \mu \in \mathcal{T}_f(X, T) \right\}. \quad (3.3)$$

Proof. If $\mu \in \mathcal{T}_f(X, T)$, then, for $g \in C(X, \mathbb{R})$,

$$\int g \, d\mu \leq \frac{P_{\text{pre}}(T, f + tg) - P_{\text{pre}}(T, f)}{t}, \quad \forall t > 0. \quad (3.4)$$

Hence,

$$\sup \left\{ \int g \, d\mu : \mu \in \mathcal{T}_f(X, T) \right\} \leq d^+ P_{\text{pre}}(T, f)(g). \quad (3.5)$$

Next, we prove the converse inequality. Set $a = d^+ P_{\text{pre}}(T, f)(g)$. Define a continuous linear functional $\gamma : \{tg : t \in \mathbb{R}\} \rightarrow \mathbb{R}$ by

$$\gamma(tg) = ta, \quad t \in \mathbb{R}. \quad (3.6)$$

The convexity of $P_{\text{pre}}(T, \bullet)$ implies

$$\gamma(tg) = t \cdot d^+ P_{\text{pre}}(T, f)(g) \leq P_{\text{pre}}(T, f + tg) - P_{\text{pre}}(T, f), \quad \forall t \in \mathbb{R}. \quad (3.7)$$

By the Hahn-Banach theorem, γ can be extended to a continuous linear functional on $C(X, \mathbb{R})$ such that

$$\gamma(h) \leq P_{\text{pre}}(T, f + h) - P_{\text{pre}}(T, f), \quad \forall h \in C(X, \mathbb{R}). \quad (3.8)$$

By the Riesz representation theorem, there is $\mu \in \mathcal{M}(X)$ with

$$\gamma(h) = \int h \, d\mu, \quad \forall g \in C(X, \mathbb{R}). \quad (3.9)$$

Combining (3.6), (3.8), and (3.9), we have $\mu \in \mathcal{T}_f(X, T)$, and

$$\int g \, d\mu = \gamma(g) = a = d^+ P_{\text{pre}}(T, f)(g). \quad (3.10)$$

The lemma is proved. \square

Theorem 3.2 (Uniqueness of tangent functional and Gateaux differentiability). *The following statements are mutually equivalent:*

- (1) the pre-image pressure function $P_{\text{pre}}(T, \bullet)$ is Gateaux differentiable at $f \in C(X, \mathbb{R})$;
- (2) the unique tangent functional to $P_{\text{pre}}(T, \bullet)$ at f is μ ;
- (3) for each $g \in C(X, \mathbb{R})$,

$$\lim_{t \rightarrow 0} \frac{P_{\text{pre}}(T, f + tg) - P_{\text{pre}}(T, f)}{t} = \int g \, d\mu. \quad (3.11)$$

Proof. (1) \Rightarrow (2) If the pre-image pressure function $P_{\text{pre}}(T, \bullet)$ is Gateaux differentiable at f , then the function $g \mapsto d^+ P_{\text{pre}}(T, f)(g)$ is linear. By Lemma 3.1,

$$\begin{aligned} \sup \left\{ \int g \, d\mu : \mu \in \mathcal{T}_f(X, T) \right\} &= d^+ P_{\text{pre}}(T, f)(g) \\ &= -d^+ P_{\text{pre}}(T, f)(-g) \\ &= -\sup \left\{ \int -g \, d\mu : \mu \in \mathcal{T}_f(X, T) \right\} \\ &= \inf \left\{ \int g \, d\mu : \mu \in \mathcal{T}_f(X, T) \right\} \end{aligned} \quad (3.12)$$

for each $g \in C(X, \mathbb{R})$. This implies there is a unique tangent functional to $P_{\text{pre}}(T, \bullet)$ at f .

(2) \Rightarrow (3) It directly follows from Lemma 3.1.

(3) \Rightarrow (1) It follows from the definition. \square

4. Fréchet Differentiability of the Pre-Image Pressure

In this section, we will study the Fréchet differentiability of pre-image pressure. The pre-image pressure function $P_{\text{pre}}(T, \bullet)$ is said to be *Fréchet differentiable at f* if there is $\gamma \in C(X, \mathbb{R})^*$ such that

$$\lim_{g \rightarrow 0} \frac{|P_{\text{pre}}(T, f + g) - P_{\text{pre}}(T, f) - \gamma(g)|}{\|g\|} = 0. \quad (4.1)$$

The pre-image pressure function $P_{\text{pre}}(T, \bullet)$ is said to be *Fréchet differentiable* if it is Fréchet differentiable at each $f \in C(X, \mathbb{R})$.

Note that if $P_{\text{pre}}(T, \bullet)$ is Fréchet differentiable at f , then it is Gateaux differentiable at f and $\gamma(g) = \int g \, d\mu$, where μ is the unique tangent functional to $P_{\text{pre}}(T, \bullet)$ at f .

Theorem 4.1. *The following conditions are mutually equivalent:*

- (1) $P_{\text{pre}}(T, \bullet)$ is Fréchet differentiable at f ;
- (2) $\mathcal{T}_f(X, T) = \{\mu_f\}$ and $\|\mu_n - \mu_f\| \rightarrow 0$ for each $\{\mu_n\} \subseteq \mathcal{M}(X, T)$ with $h_{\text{pre}, \mu_n}(T) + \int f \, d\mu_n \rightarrow P_{\text{pre}}(T, f)$;
- (3) $P_{\text{pre}}(T, \bullet)$ is locally affine at f ;
- (4) $\mathcal{T}_f(X, T) = \{\mu_f\}$ and

$$\limsup_{g \rightarrow 0} \{\|\mu - \mu_f\| : \mu \in \mathcal{T}_{f+g}(X, T)\} = 0. \quad (4.2)$$

Proof. (1) \Rightarrow (2) Suppose $P_{\text{pre}}(T, \bullet)$ is Fréchet differentiable at f . Then, f has a unique tangent functional μ_f to $P_{\text{pre}}(T, \bullet)$ at f . Let $\{\mu_n\}_{n=1}^{\infty} \subseteq \mathcal{M}(X, T)$ with $h_{\text{pre}, \mu_n}(T) + \int f \, d\mu_n \rightarrow P_{\text{pre}}(T, f)$. Given $\varepsilon > 0$, there are $N \in \mathbb{N}$ and $\delta > 0$ such that

$$P_{\text{pre}}(T, f) \leq h_{\text{pre}, \mu_n}(T) + \int f \, d\mu_n + \varepsilon\delta, \quad \forall n \geq N, \quad (4.3)$$

$$0 \leq P_{\text{pre}}(T, f + g) - P_{\text{pre}}(T, f) - \int g \, d\mu_f \leq \varepsilon \cdot \|g\| \quad \text{whenever } \|g\| < \delta. \quad (4.4)$$

Hence, if $n \geq N$ and $\|g\| < \delta$, then

$$\begin{aligned} \int g \, d\mu_n - \int g \, d\mu_f &= P_{\text{pre}}(T, f) + \int g \, d\mu_n - P_{\text{pre}}(T, f) - \int g \, d\mu_f \\ &\leq h_{\text{pre}, \mu_n}(T) + \int (f + g) \, d\mu_n + \varepsilon\delta - P_{\text{pre}}(T, f) - \int g \, d\mu_f \quad (\text{by 4.4}) \end{aligned}$$

$$\begin{aligned}
&\leq P_{\text{pre}}(T, f + g) - P_{\text{pre}}(T, f) - \int g \, d\mu_f + \epsilon\delta \quad (\text{by Theorem 2.1}) \\
&\leq \epsilon \cdot (\|g\| + \delta) \leq 2\epsilon\delta \quad (\text{by 4.4}).
\end{aligned} \tag{4.5}$$

Note that (4.5) is also true when $-g$ instead of g . So,

$$\left| \int g \, d\mu_n - \int g \, d\mu_f \right| \leq 2\epsilon\delta \quad \text{whenever } \|g\| < \delta, \quad n \geq N. \tag{4.6}$$

Therefore,

$$\begin{aligned}
\|\mu_n - \mu_f\| &= \sup \left\{ \left| \int g \, d\mu_n - \int g \, d\mu_f \right| : \|g\| < 1 \right\} \\
&= \frac{1}{\delta} \sup \left\{ \left| \int g \, d\mu_n - \int g \, d\mu_f \right| : \|g\| < \delta \right\} \\
&\leq 2\epsilon \quad (\text{by 4.6}).
\end{aligned} \tag{4.7}$$

By arbitrary of ϵ , $\|\mu_n - \mu_f\| \rightarrow 0$.

(2) \Rightarrow (3) By the variational principle of pre-image pressure, we can pick ergodic measures μ_n such that $h_{\text{pre}, \mu_n}(T) + \int f \, d\mu_n \rightarrow P_{\text{pre}}(T, f)$, Then $\|\mu_n - \mu_f\| \rightarrow 0$. Note that two distinct ergodic measures have norm-distance 2. So there is N such that $\mu_n = \mu_f$ for all $n \geq N$. Hence, $\mu_f \in \mathcal{M}^e(X, T)$ and

$$\epsilon := P_{\text{pre}}(T, f) - \sup \left\{ h_{\text{pre}, \mu}(T) + \int f \, d\mu : \mu \in \mathcal{M}^e(X, T), \mu \neq \mu_f \right\} > 0. \tag{4.8}$$

This implies for each $g \in C(X, \mathbb{R})$, with $\|f - g\| < \epsilon/2$, that

$$\begin{aligned}
&\sup \left\{ h_{\text{pre}, \mu}(T) + \int g \, d\mu : \mu \in \mathcal{M}^e(X, T), \mu \neq \mu_f \right\} \\
&\leq P_{\text{pre}}(T, f) - \epsilon + \|f - g\| \\
&\leq P_{\text{pre}}(T, g) - \epsilon + 2\|f - g\| \\
&< P_{\text{pre}}(T, g).
\end{aligned} \tag{4.9}$$

By the variational principle of pre-image pressure again, we have

$$P_{\text{pre}}(T, g) = h_{\text{pre}, \mu_f}(T) + \int g \, d\mu_f \quad \text{whenever } \|f - g\| < \frac{\epsilon}{2}. \tag{4.10}$$

Hence, $P_{\text{pre}}(T, \bullet)$ is affine on the neighborhood $B(f, \epsilon/2) = \{g \in C(X, \mathbb{R}) : \|f - g\| < \epsilon/2\}$ of f .

(3) \Rightarrow (4) Is obvious.

(4) \Rightarrow (1) Let $g \in C(X, \mathbb{R})$ and $\mu \in \mathcal{T}_{f+g}(X, T)$. By definition, we have

$$P_{\text{pre}}(T, f) - P_{\text{pre}}(T, f + g) \geq - \int g \, d\mu. \quad (4.11)$$

Hence,

$$\begin{aligned} 0 &\leq P_{\text{pre}}(T, f + g) - P_{\text{pre}}(T, f) - \int g \, d\mu_f \\ &\leq \int g \, d\mu - \int g \, d\mu_f \quad (\text{by (4.11)}) \\ &\leq \|g\| \cdot \|\mu - \mu_f\|. \end{aligned} \quad (4.12)$$

Therefore,

$$0 \leq \frac{P_{\text{pre}}(T, f + g) - P_{\text{pre}}(T, f) - \int g \, d\mu_f}{\|g\|} \leq \sup\{\|\mu - \mu_f\| : \mu \in \mathcal{T}_{f+g}(X, T)\} \rightarrow 0 \quad (4.13)$$

as $g \rightarrow 0$. That is $P_{\text{pre}}(T, \bullet)$ is Fréchet differentiable at f . \square

Corollary 4.2. *Let (X, T) be a TDS with finite pre-image entropy. Then, $P_{\text{pre}}(T, \bullet)$ is Fréchet differentiable if and only if T is uniquely ergodic.*

Proof. Using Theorem 4.1, $P_{\text{pre}}(T, \bullet)$ is locally affine whenever $P_{\text{pre}}(T, \bullet)$ is Fréchet differentiable. Hence, the map $f \in C(X, \mathbb{R}) \mapsto \mu_f \in \mathcal{M}(X, T)$ is locally constant, where $\mathcal{T}_f(X, T) = \{\mu_f\}$ for each $f \in C(X, \mathbb{R})$. Since $C(X, \mathbb{R})$ is connected, the map is constant. So $\mu_f = \mu_0$ for all $f \in C(X, \mathbb{R})$. If $\mu \in \mathcal{M}(X, T) \setminus \{\mu_0\}$, then we can choose $f \in C(X, \mathbb{R})$ such that $\int f \, d\mu > \int f \, d\mu_0$. Then for sufficiently large k , we have

$$P_{\text{pre}}(T, kf) \geq h_{\text{pre}, \mu}(T) + \int kf \, d\mu > h_{\text{pre}, \mu_0}(T) + \int kf \, d\mu_0 = P_{\text{pre}}(T, kf), \quad (4.14)$$

which is a contradiction. Therefore, $\mathcal{M}(X, T) = \{\mu_0\}$. \square

Remark 4.3. In the situation of Corollary 4.2, there is only one invariant measure, and the pre-image pressure is the expectation with respect to this measure, hence, it is linear.

Acknowledgments


The authors would like to thank the referee for many valuable and constructive comments and suggestions for improving this paper. The first and second authors are supported by the National Natural Science Foundation of China (no. 11161029) and the Department of

Education Research Foundation of Guangxi Province (nos. 200911MS298, 200103YB157); the third author is supported by the National Natural Science Foundation of China (no. 11126342) and the Natural Science Foundation of Guangxi Province (nos. 0897012, 2010-GXNSFA013109).

References

- [1] A. N. Kolmogorov, "A new metric invariant of transient dynamical systems and automorphisms of Lebesgue spaces," *Doklady Akademii Nauk SSSR*, vol. 119, pp. 861–864, 1958 (Russian).
- [2] R. L. Adler, A. G. Konheim, and M. H. McAndrew, "Topological entropy," *Transactions of the American Mathematical Society*, vol. 114, pp. 309–319, 1965.
- [3] T. N. T. Goodman, "Relating topological entropy and measure entropy," *The Bulletin of the London Mathematical Society*, vol. 3, pp. 176–180, 1971.
- [4] L. W. Goodwyn, "Topological entropy bounds measure-theoretic entropy," *Proceedings of the American Mathematical Society*, vol. 23, pp. 679–688, 1969.
- [5] D. Ruelle, "Statistical mechanics on a compact set with Z^n action satisfying expansiveness and specification," *Transactions of the American Mathematical Society*, vol. 187, pp. 237–251, 1973.
- [6] P. Walters, "A variational principle for the pressure of continuous transformations," *American Journal of Mathematics*, vol. 97, no. 4, pp. 937–971, 1975.
- [7] R. Bowen, *Equilibrium States and the Ergodic Theory of Anosov Diffeomorphisms*, vol. 470 of *Lecture Notes in Mathematics*, Springer, Berlin, Germany, 1975.
- [8] R. B. Israel, *Tangents to the Pressure as Invariant Equilibrium States in Statistical Mechanics of Lattice Systems*, Princeton University Press, Princeton, NJ, USA, 1978.
- [9] R. B. Israel, *Convexity in the Theory of Lattice Gases*, Princeton Series in Physics, Princeton University Press, Princeton, NJ, USA, 1979.
- [10] G. Keller, *Equilibrium States in Ergodic Theory*, vol. 42, Cambridge University Press, Cambridge, UK, 1998.
- [11] D. Ruelle, *Thermodynamic Formalism: The Mathematical Structures of Equilibrium Statistical Mechanics*, Cambridge University Press, Cambridge, UK, 2nd edition, 2004.
- [12] P. Walters, *An Introduction to Ergodic Theory*, vol. 79 of *Graduate Texts in Mathematics*, Springer, New York, NY, USA, 1982.
- [13] R. Bowen, "Hausdorff dimension of quasicircles," *Institut des Hautes Études Scientifiques*, no. 50, pp. 11–25, 1979.
- [14] D. Ruelle, "Repellers for real analytic maps," *Ergodic Theory and Dynamical Systems*, vol. 2, no. 1, pp. 99–107, 1982.
- [15] H. A. M. Daniëls and A. C. D. van Enter, "Differentiability properties of the pressure in lattice systems," *Communications in Mathematical Physics*, vol. 71, no. 1, pp. 65–76, 1980.
- [16] R. B. Israel and R. R. Phelps, "Some convexity questions arising in statistical mechanics," *Mathematica Scandinavica*, vol. 54, no. 1, pp. 133–156, 1984.
- [17] R. R. Phelps, "Generic Fréchet differentiability of the pressure in certain lattice systems," *Communications in Mathematical Physics*, vol. 91, no. 4, pp. 557–562, 1983.
- [18] P. Walters, "Differentiability properties of the pressure of a continuous transformation on a compact metric space," *Journal of the London Mathematical Society*, vol. 46, no. 3, pp. 471–481, 1992.
- [19] W.-C. Cheng and S. E. Newhouse, "Pre-image entropy," *Ergodic Theory and Dynamical Systems*, vol. 25, no. 4, pp. 1091–1113, 2005.
- [20] D. Fiebig, U.-R. Fiebig, and Z. H. Nitecki, "Entropy and preimage sets," *Ergodic Theory and Dynamical Systems*, vol. 23, no. 6, pp. 1785–1806, 2003.
- [21] M. Hurley, "On topological entropy of maps," *Ergodic Theory and Dynamical Systems*, vol. 15, no. 3, pp. 557–568, 1995.
- [22] Z. Nitecki and F. Przytycki, "Preimage entropy for mappings," *International Journal of Bifurcation and Chaos in Applied Sciences and Engineering*, vol. 9, no. 9, pp. 1815–1843, 1999.
- [23] K. Yan, F. Zeng, and B. Qin, "A variational principle for conditional pre-image entropy," *Journal of Computational Analysis and Applications*, vol. 13, no. 6, pp. 1180–1187, 2011.

- [24] F. Zeng, "Partial entropy, bundle-like entropy and topological entropy," in *Dynamical Systems*, Proceedings of the International Conference in Honor of Professor Liao Shantao, pp. 329–333, World Scientific, Singapore, 1999.
- [25] F. Zeng, K. Yan, and G. Zhang, "Pre-image pressure and invariant measures," *Ergodic Theory and Dynamical Systems*, vol. 27, no. 3, pp. 1037–1052, 2007.
- [26] K. Yan, F. Zeng, and G. Zhang, "Equilibrium states for pre-image pressure," *Nonlinear Dynamics and Systems Theory*, vol. 10, no. 4, pp. 409–416, 2010.
- [27] N. Dunford and J. T. Schwartz, *Linear Operators. I. General Theory*, vol. 7 of *Pure and Applied Mathematics*, Interscience Publishers, 1958.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

