

Research Article

Restricted p -Isometry Properties of Partially Sparse Signal Recovery

Haini Bi, Lingchen Kong, and Naihua Xiu

Department of Applied Mathematics, Beijing Jiaotong University, Beijing 100044, China

Correspondence should be addressed to Lingchen Kong; konglchen@126.com

Received 20 January 2013; Revised 3 March 2013; Accepted 5 March 2013

Academic Editor: Binggen Zhang

Copyright © 2013 Haini Bi et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

By generalizing the restricted p -isometry property to the partially sparse signal recovery problem, we give a sufficient condition for exactly recovering partially sparse signal via the partial l_p minimization (truncated l_p minimization) problem with $p \in (0, 1]$. Based on this, we establish a simpler sufficient condition which can show how the p -RIP bounds vary corresponding to different ps .

1. Introduction

The partially sparse signal recovery (PSSR) is the problem of recovering a partially sparse signal from a certain number of linear measurements when the part of the signal is known to be sparse, which was coined by Bandeira et al. [1, 2]. This type of problems has many applications in signal and image processing, derivative-free optimizations, and so on; see, for example, [1–4]. Clearly, PSSR includes *sparse signal recovery* (SSR) as a special case. The latter is the well-known NP-hard problem in the compressed sensing (CS), which is also called *cardinality minimization problem* (CMP, or l_0 -norm minimization problems); see, for example, [5–8]. In particular, Candès and Tao [8] introduced a restricted isometry property (RIP) of a sensing matrix which guarantees to recover a sparse solution of SSR by minimizing its convex relaxation (l_1 -norm minimization). However, there are some problems which cannot be reformulated as an SSR, but a PSSR. As we know, PSSR happens naturally in sparse Hessian recovery; see, for example, [2], where Bandeira et al. employed partially sparse recovery approach for building sparse quadratic interpolation models of functions with sparse Hessian. They have successfully applied the l_1 -norm minimization of PSSR in interpolation-based trust-region methods for derivative-free optimization. Vaswani and Lu [3] successfully applied modified CS (partially sparse recovery) in image reconstruction, where the sufficient RIP condition is weaker than the RIP for SSR. Moreover, Bandeira et al. [1]

considered the RIP and null space properties (NSP) for PSSR and extended recovery results under noisy measurements to the partially sparse case, where partial NSP is a necessary and sufficient condition for PSSR. In [4], Jacques also established the partial RIP condition for PSSR with noise via its convex relaxation problem.

Note that in the CS context, the SSR problem can also be relaxed to a l_p -norm minimization (truncated l_p -minimization) problem with $0 < p < 1$; see, for example, [9–19]. It is well known that Chartrand [20] firstly show that fewer measurements are required for exact reconstruction if we replace l_1 -norm with l_p -norm ($0 < p < 1$), and Chartrand and Staneva [10] established p -RIP conditions for exact SSR via l_p -minimization. In particular, the numerical experiments in magnetic resonance imaging (MRI) showed that this approach works very efficiently; see [9] for details. Wang et al. [19] studied the performance of l_p -minimization for strong recovery and *weak recovery* where we need to recover all the sparse vectors on one support with one sign pattern. Moreover, Saab et al. [16] provided a sufficient condition for SSR via l_p -minimization and provided a lower bound of the support size up to which l_p -minimization can recover all such sparse vectors, and Foucart and Lai [14] improved this bound by considering a generalized version of RIP condition. While SSR and l_p -minimization have been the focus point of some recent research, there are fewer research related to PSSR and the partially l_p -minimization. One may naturally wonder whether we can generalize the p -RIP conditions

introduced by [10] from the SSR to the PSSR case. This paper will deal with this issue. We will give a different p -RIP recovery condition for PSSR via its nonconvex relaxation. Furthermore, based on the recent work by Oymak et al. [21], we also extend our result to the matrix setting.

In the next section, we give the PSSR model and review some preliminaries on p -RIP conditions. In Section 3, we establish the exact partially p -RIP recovery conditions for PSSR via its nonconvex l_p -minimization. In Section 4, we give a sufficient condition for partially low-rank matrix recovery via the partially Schatten- p minimization problem.

2. Preliminaries

In this section, we will review some basic concepts and results on the p -RIP recovery conditions for SSR and introduce the p -RIP definition for PSSR. We begin with defining the mathematical model of the PSSR problem as follows:

$$\min \|x\|_0, \quad \text{s.t. } A_1 x + A_2 y = b, \quad (1)$$

where the l_0 -norm $\|x\|_0$ is defined as $\|x\|_0 := |\{i : x_i \neq 0\}|$ (which is not really a norm since it is not positive homogeneous). For any positive number s , we say x is s -sparse if $\|x\|_0 \leq s$. $A = [A_1, A_2] \in \mathbb{R}^{M \times N}$ is a sensing matrix with $A_1 \in \mathbb{R}^{M \times (N-r)}$, $A_2 \in \mathbb{R}^{M \times r}$, and $b \in \mathbb{R}^M$. It means that the unknown vector consists of two parts (x, y) , where $x \in \mathbb{R}^{N-r}$ is sparse and $y \in \mathbb{R}^r$ is possibly dense. When $A = A_1$, the previous problem reduces to the following l_0 -norm minimization problem (sparse signal recovery, SSR):

$$\min \|x\|_0, \quad \text{s.t. } Ax = b. \quad (2)$$

The previous PSSR problem (1) is an NP-hard problem, since its special case SSR (2) is well-known NP-hard problem in the compressed sensing (CS). As we mentioned in Section 1, one popular and powerful approach is to solve it via l_1 -norm minimization (its convex relaxation), where the l_0 -norm is replaced by the l_1 -norm in SSR (2). Moreover, we can also use a nonconvex approach for exact reconstruction with fewer measurements than the convex relaxation; see, for example, [9, 10]. That is the l_p -norm minimization problem with $0 < p < 1$, where we replace the l_0 -norm with the l_p -norm in (2) as follows:

$$\min \|x\|_p^p, \quad \text{s.t. } Ax = b. \quad (3)$$

Note that $\|\cdot\|_p$ is not a norm when $p \in (0, 1)$, but it is much close to l_0 -norm. Moreover, the numerical experiments in MRI showed that the approach via l_p -minimization works very efficiently; see [9] for details. In particular, Chartrand and Staneva [10] introduced the concept of restricted isometry constant via l_p -norm.

Definition 1 (p -RIC [10]). Given a matrix $A \in \mathbb{R}^{M \times N}$, where $M < N$, s is a positive number and $0 < p < 1$, then we say that δ_s is the restricted p -isometry constant (or p -RIC) of order s of the matrix A if δ_s is the smallest number, such that

$$(1 - \delta_s) \|x\|_2^p \leq \|Ax\|_p^p \leq (1 + \delta_s) \|x\|_2^p, \quad (4)$$

for all s -sparse vectors x .

In the same paper, Chartrand and Staneva gave the following sufficient condition for exact SSR via l_p -minimization.

Theorem 2 (see [10]). Let $A \in \mathbb{R}^{M \times N}$, $x \in \mathbb{R}^N$, and $k = \|x\|_0$ be the size of the support of x , $0 < p < 1$, $a_1 > 1$, and $a_2 = a_1^{2/(2-p)}$, rounded up, so that $a_2 k$ is an integer ($a_2 = \lceil a_1^{2/(2-p)} k \rceil / k$). If A satisfies

$$\delta_{a_2 k} + a_1 \delta_{(a_2+1)k} < a_1 - 1, \quad (5)$$

then x is the unique minimizer of problem (2).

Inspired by the previous analysis, it is natural to give the partially l_p -norm minimization problem for PSSR (1) as follows:

$$\min \|x\|_p^p, \quad \text{s.t. } A_1 x + A_2 y = b. \quad (6)$$

In order to establish the link between the PSSR (1) and its partially l_p -norm minimization problem, we need to give a partially p -RIC definition. Here we borrow the idea from Bandeira et al. [1]. Assume that A_2 is full column rank. For $A = [A_1, A_2]$ as mentioned above, let

$$B := I - A_2 (A_2^T A_2)^{-1} A_2^T, \quad (7)$$

which is the matrix of the orthogonal projection from \mathbb{R}^N to $\mathfrak{R}(A_2)^\perp$.

Definition 3 (Partially p -RIC). Let $A = [A_1, A_2] \in \mathbb{R}^{M \times N}$, where $A_1 \in \mathbb{R}^{M \times (N-r)}$, and $A_2 \in \mathbb{R}^{M \times r}$ is full column rank. We say that δ_{s-r}^r is the Partially Restricted Isometry Constant (Partially p -RIC) of order $s-r$ of the matrix A if δ_{s-r}^r is the p -RIC of order $s-r$ of the matrix BA_1 ; that is, δ_{s-r}^r is the smallest number, such that

$$(1 - \delta_{s-r}^r) \|x\|_2^p \leq \|BA_1 x\|_p^p \leq (1 + \delta_{s-r}^r) \|x\|_2^p, \quad (8)$$

for all $(s-r)$ -sparse vectors x , where B is given by (7).

3. Main Results

We will give our main results which state sufficient p -RIP recovery conditions on the exact PSSR via the nonconvex l_p -norm minimization. We begin with the following useful lemma.

Lemma 4. For $0 < p \leq 1$, let $c_1 = a_1^{1-p/2}$ and $c_2 = a_2^{1-p/2}$ with $a_1 > 0$ and $a_2 > 0$. If $(c_1 - 1)/c_2 > |a_1 - a_2|/a_2$, then $a_1 > 1$ and $a_2 > 1$.

Proof. In order to prove the lemma, we consider the following two cases.

Case 1 ($a_1 \geq a_2$). In this case, from the fact $(c_1 - 1)/c_2 > |a_1 - a_2|/a_2$, we have

$$\frac{a_1}{a_2} - \left(\frac{a_1}{a_2}\right)^{1-p/2} < 1 - \left(\frac{1}{a_2}\right)^{1-p/2}. \quad (9)$$

If $0 < a_2 \leq 1$, from the previous inequality we easily obtain

$$0 < \frac{a_1}{a_2} - \left(\frac{a_1}{a_2}\right)^{1-p/2} < 1 - \left(\frac{1}{a_2}\right)^{1-p/2} \leq 0, \quad (10)$$

which is a contradiction. Hence $a_1 \geq a_2 > 1$.

Case 2 ($a_1 < a_2$). Similarly, in this case, from $(c_1 - 1)/c_2 > |a_1 - a_2|/a_2$, we obtain

$$0 < 1 - \frac{a_1}{a_2} < \frac{c_1 - 1}{c_2}. \quad (11)$$

If $0 < a_1 \leq 1$, then

$$c_1 = a_1^{1-p/2} \leq 1. \quad (12)$$

Combining the previous inequalities we obtain

$$0 < 1 - \frac{a_1}{a_2} < \frac{c_1 - 1}{c_2} \leq 0, \quad (13)$$

which is a contradiction. Hence $1 < a_1 < a_2$.

Therefore, taking into account the previous two cases, we completed the proof. \square

We below propose a general recovery condition for PSSR via its p -norm minimization.

Theorem 5. Let $A = [A_1, A_2] \in \mathbb{R}^{M \times N}$ with $A_1 \in \mathbb{R}^{M \times (N-r)}$ and $A_2 \in \mathbb{R}^{M \times r}$. Suppose that A_2 is full column rank, and let $A_1 x + A_2 y = b$ with $\|x\|_0 = k$. For $0 < p \leq 1$, $a_1 > 0$, and $a_2 > 0$, let $c_1 = a_1^{1-p/2}$, $c_2 = a_2^{1-p/2}$ with $(c_1 - 1)/c_2 > |a_1 - a_2|/a_2$. If

$$\begin{aligned} & a_2 c_1 \delta_{(a_2+1)k} + (|a_1 - a_2| c_2 + a_2) \delta_{a_1 k} \\ & < a_2 c_1 - |a_1 - a_2| c_2 - a_2, \end{aligned} \quad (14)$$

then (x, y) is the unique minimizer of problem (6).

Proof. Note that (x, y) is a feasible solution to optimization problem (6). We remain to show that the solution set is a singleton $\{(x, y)\}$. This proof generally modifies that of [10], but under different assumptions. (Specifically, we use a different way to arrange the elements of T_0^C in the following.) Let (u, v) be an arbitrary solution to problem (6). We will show that $u = x$ and $y = v$. We will prove $u = x$ firstly. Taking $h = u - x$, we will show that $h = 0$. Let $\Phi = BA_1$. For $T \in \{1, \dots, N - r\}$, Φ_T denotes the matrix equaling Φ in those columns whose indices belong to T and otherwise zero. Similarly, we define the vector h_T . Let T_0 be the support of x . Then, the supports of x and $h_{T_0^C}$ are disjoint since $T_0 \cap T_0^C = \emptyset$. From direct calculation, we obtain

$$\begin{aligned} \|x\|_p^p & \geq \|u\|_p^p = \|x + h\|_p^p \\ & = \|x + h_{T_0} + h_{T_0^C}\|_p^p \\ & = \|x + h_{T_0}\|_p^p + \|h_{T_0^C}\|_p^p \\ & \geq \|x\|_p^p - \|h_{T_0}\|_p^p + \|h_{T_0^C}\|_p^p, \end{aligned} \quad (15)$$

where the first inequality holds because (u, v) solves (6), and the last one holds by the triangle inequality for $\|\cdot\|_p^p$. Then we have

$$\|h_{T_0^C}\|_p^p \leq \|h_{T_0}\|_p^p. \quad (16)$$

Now we arrange the elements of T_0^C in order of decreasing magnitude of $|h|$ and partition into $T_0^C = T_1 \cup T_2 \cup \dots \cup T_J$, where T_1 has $a_2 k$ elements and T_j ($j \geq 2$) each has $a_1 k$ elements (except possibly T_J). Set $T_{01} = T_0 \cup T_1$. Note that

$$BA_2 = \left[I - A_2 (A_2^T A_2)^{-1} A_2^T \right] A_2 = 0. \quad (17)$$

Direct calculations yield

$$\begin{aligned} 0 & = \|B(A_1 x + A_2 y - A_1 u - A_2 v)\|_p^p \\ & = \|BA_1 x - BA_1 u\|_p^p = \|\Phi x - \Phi u\|_p^p \\ & = \|\Phi h\|_p^p = \left\| \Phi h_{T_{01}} + \sum_{j \geq 2} \Phi h_{T_j} \right\|_p^p \\ & \geq \|\Phi h_{T_{01}}\|_p^p - \left\| \sum_{j \geq 2} \Phi h_{T_j} \right\|_p^p \\ & \geq \|\Phi h_{T_{01}}\|_p^p - \sum_{j \geq 2} \|\Phi h_{T_j}\|_p^p \\ & \geq (1 - \delta_{(a_2+1)k}) \|h_{T_{01}}\|_2^p - (1 + \delta_{a_1 k}) \\ & \quad \times \sum_{j \geq 2} \|h_{T_j}\|_2^p. \end{aligned} \quad (18)$$

Now we discuss the relation between l_2 -norm and l_p -norm. For each $t \in T_j$ and $s \in T_{j-1}$, it holds $|h_t| \leq |h_s|$. So, we have for $j = 2$,

$$\begin{aligned} |h_t|^p & \leq \frac{\|h_{T_1}\|_p^p}{a_2 k} \\ \implies |h_t|^2 & \leq \frac{\|h_{T_1}\|_p^2}{(a_2 k)^{2/p}} \\ \implies \frac{\|h_{T_2}\|_2^2}{a_1 k} & \leq \frac{\|h_{T_1}\|_p^2}{(a_2 k)^{2/p}} \\ \implies \|h_{T_2}\|_2^p & \leq \frac{a^{p/2}}{a_2 k^{1-p/2}} \|h_{T_1}\|_p^p. \end{aligned} \quad (19)$$

Similarly, we obtain that for $j \geq 3$,

$$\|h_{T_j}\|_2^p \leq \frac{1}{(a_1 k)^{1-p/2}} \|h_{T_{j-1}}\|_2^p. \quad (20)$$

Applying the Holder's inequality, we obtain

$$\begin{aligned} \|h_{T_0}\|_p^p &= \sum_{t \in T_0} |h_t|^p \cdot 1 \\ &\leq \left(\sum_{t \in T_0} |h_t|^2 \right)^{p/2} \left(\sum_{t \in T_0} 1 \right)^{1-p/2} \\ &= \|h_{T_0}\|_2^p \cdot k^{1-p/2}. \end{aligned} \quad (21)$$

Similarly, we have

$$\|h_{T_1}\|_p^p \leq \|h_{T_1}\|_2^p \cdot (a_2 k)^{1-p/2}. \quad (22)$$

Therefore,

$$\begin{aligned} &\sum_{j \geq 2} \|h_{T_j}\|_2^p \\ &\leq \frac{a_1^{p/2}}{a_2 k^{1-p/2}} \|h_{T_1}\|_p^p \\ &\quad + \frac{1}{(a_1 k)^{1-p/2}} \sum_{j \geq 2} \|h_{T_j}\|_p^p \quad (\text{By (19) and (20)}) \\ &= \frac{1}{(a_1 k)^{1-p/2}} \sum_{j \geq 1} \|h_{T_j}\|_p^p + \frac{(a_1 - a_2) a_1^{p/2}}{a_1 a_2 k^{1-p/2}} \|h_{T_1}\|_p^p \\ &= \frac{1}{(a_1 k)^{1-p/2}} \|h_{T_0^c}\|_p^p + \frac{(a_1 - a_2) a_1^{p/2}}{a_1 a_2 k^{1-p/2}} \|h_{T_1}\|_p^p \\ &\leq \frac{1}{(a_1 k)^{1-p/2}} \|h_{T_0}\|_p^p + \frac{(a_1 - a_2) a_1^{p/2}}{a_1 a_2 k^{1-p/2}} \|h_{T_1}\|_p^p \quad (\text{By (16)}) \\ &\leq \frac{k^{1-p/2}}{(a_1 k)^{1-p/2}} \|h_{T_0}\|_2^p \\ &\quad + \frac{(a_1 - a_2) (a_2 k)^{1-p/2}}{a_2 (a_1 k)^{1-p/2}} \|h_{T_1}\|_2^p \quad (\text{By (21) and (22)}) \\ &= \frac{1}{c_1} \|h_{T_0}\|_2^p + \frac{(a_1 - a_2) c_2}{a_2 c_1} \|h_{T_1}\|_2^p \\ &\leq \frac{1}{c_1} \|h_{T_0}\|_2^p + \frac{|a_1 - a_2| c_2}{a_2 c_1} \|h_{T_1}\|_2^p \\ &\leq \frac{1}{c_1} \|h_{T_{01}}\|_2^p + \frac{|a_1 - a_2| c_2}{a_2 c_1} \|h_{T_{01}}\|_2^p. \end{aligned} \quad (23)$$

Thus by (18) and (23), we have

$$\begin{aligned} 0 &\geq (1 - \delta_{(a_2+1)k}) \|h_{T_{01}}\|_2^p \\ &\quad - (1 + \delta_{a_1 k}) \sum_{j \geq 2} \|h_{T_j}\|_2^p \quad (\text{By (18)}) \\ &\geq (1 - \delta_{(a_2+1)k}) \|h_{T_{01}}\|_2^p - \frac{1 + \delta_{a_1 k}}{c_1} \|h_{T_{01}}\|_2^p \\ &\quad - \frac{|a_1 - a_2| c_2 (1 + \delta_{a_1 k})}{a_2 c_1} \|h_{T_{01}}\|_2^p \quad (\text{By (23)}) \\ &= \left[1 - \delta_{(a_2+1)k} - \frac{1 + \delta_{a_1 k}}{c_1} - \frac{|a_1 - a_2| c_2 (1 + \delta_{a_1 k})}{a_2 c_1} \right] \\ &\quad \times \|h_{T_{01}}\|_2^p. \end{aligned} \quad (24)$$

Clearly, the assumption ensures that the scalar factor is positive, and hence we obtain $h_{T_{01}} = 0$. That means $h_{T_0} = 0$. Using $\|h_{T_0^c}\|_p^p \leq \|h_{T_0}\|_p^p$, we obtain $h_{T_0^c} = 0$. Therefore, $h = 0$, which means $x = u$.

Now we remain to show that $y = v$. It is obvious that $A_1 x + A_2 y = A_1 u + A_2 v$. Since $x = u$, we have $A_2(y - v) = 0$. Then $y = v$ because A_2 is full column rank. \square

Theorem 5 states a different sufficient condition for the exactly PSSR via its nonconvex relaxation from the existing conditions for SSR.

Theorem 6. Let $A = [A_1, A_2] \in \mathbb{R}^{M \times N}$ with $A_1 \in \mathbb{R}^{M \times (N-r)}$ and $A_2 \in \mathbb{R}^{M \times r}$. Suppose that A_2 is full column rank, and let $A_1 x + A_2 y = b$ with $\|x\|_0 = k$. For $0 < p \leq 1$ and $a > 1$, if

$$\delta_{(a+1)k} < \frac{a^{1-p/2} - 1}{a^{1-p/2} + 1}, \quad (25)$$

then (x, y) is the unique minimizer of problem (6). Specifically, for all $0 < p \leq 1$, if

$$\delta_{(a+1)k} < \frac{\sqrt{a} - 1}{\sqrt{a} + 1}, \quad (26)$$

then (x, y) is the unique minimizer of problem (6).

Proof. Applying Theorem 5, here we only need to show that if (25) holds, we can find a_1 and a_2 , such that (14) holds. We consider the three cases in the following.

Case i ($a_1 \geq a_2 + 1$). In this case, we easily obtain $a_1 - a_2 \geq 1$ and $\delta_{a_1 k} \geq \delta_{(a_2+1)k}$. Therefore the following condition can guarantee the inequality (14):

$$a_2 c_1 \delta_{a_1 k} + [(a_1 - a_2) c_2 + a_2] \delta_{a_1 k} < a_2 (c_1 + c_2 - 1) - a_1 c_2. \quad (27)$$

Simplifying the previous inequality, we obtain

$$\begin{aligned} \delta_{a_1 k} &< \frac{a_2 c_1 - (a_1 - a_2) c_2 - a_2}{a_2 c_1 + (a_1 - a_2) c_2 + a_2} \\ &= \frac{a_1^{1-p/2} - (a_1 - a_2) a_2^{-p/2} - 1}{a_1^{1-p/2} + (a_1 - a_2) a_2^{-p/2} + 1}. \end{aligned} \quad (28)$$

In this case, employing $a_2^{-p/2} > 0$, we easily get that $a_1 = a_2 + 1$ gives the maximum value of the right of the inequality (the strongest result) which satisfies the condition (14). That is,

$$\delta_{(a_2+1)k} < \frac{(a_2 + 1)^{1-p/2} - a_2^{-p/2} - 1}{(a_2 + 1)^{1-p/2} + a_2^{-p/2} + 1}. \quad (29)$$

Case ii ($a_2 \leq a_1 < a_2 + 1$). In this case, we can get that $0 \leq a_1 - a_2 < 1$ and $\delta_{a_1 k} < \delta_{(a_2+1)k}$. Similarly, the following condition can guarantee the inequality (14):

$$\begin{aligned} a_2 c_1 \delta_{(a_2+1)k} + [(a_1 - a_2) c_2 + a_2] \delta_{(a_2+1)k} \\ < a_2 (c_1 + c_2 - 1) - a_1 c_2. \end{aligned} \quad (30)$$

Simplifying the previous inequality, we obtain

$$\delta_{(a_2+1)k} < \frac{a_1^{1-p/2} - (a_1 - a_2) a_2^{-p/2} - 1}{a_1^{1-p/2} + (a_1 - a_2) a_2^{-p/2} + 1}. \quad (31)$$

In this case, employing $a_2^{-p/2} > 0$, we get that $a_1 = a_2$ give the maximum value of the right of the inequality; that is,

$$\delta_{(a_2+1)k} < \frac{a_2^{1-p/2} - 1}{a_2^{1-p/2} + 1}. \quad (32)$$

Case iii ($a_1 < a_2$). In this case, it is clear that $0 < a_2 - a_1$, $a_1 < a_2 + 1$, and $\delta_{a_1 k} < \delta_{(a_2+1)k}$. So the following condition can guarantee the inequality (14):

$$\begin{aligned} a_2 c_1 \delta_{(a_2+1)k} + [(a_2 - a_1) c_2 + a_2] \delta_{(a_2+1)k} \\ < a_2 (c_1 - c_2 - 1) + a_1 c_2. \end{aligned} \quad (33)$$

Simplifying the previous inequality, we obtain

$$\delta_{(a_2+1)k} < \frac{a_1^{1-p/2} - (a_2 - a_1) a_2^{-p/2} - 1}{a_1^{1-p/2} + (a_2 - a_1) a_2^{-p/2} + 1}. \quad (34)$$

In this case, employing $a_2^{-p/2} > 0$, we chose $a_2 - a_1 \rightarrow 0$ to give the maximum value of the right of the inequality. That is,

$$\delta_{(a_2+1)k} < \frac{a_2^{1-p/2} - 1}{a_2^{1-p/2} + 1}. \quad (35)$$

It is easy to see that $((a_2 + 1)^{1-p/2} - a_2^{-p/2} - 1) / ((a_2 + 1)^{1-p/2} + a_2^{-p/2} + 1) < (a_2^{1-p/2} - 1) / (a_2^{1-p/2} + 1)$. In fact, $(a_2 + 1)^{1-p/2} + a_2^{-p/2} + 1 > a_2^{1-p/2} + 1$. On the other hand, $(a_2 + 1)^{1-p/2} - a_2^{-p/2} - 1 - (a_2^{1-p/2} - 1) = (a_2 + 1)[(a_2 + 1)^{-p/2} - a_2^{-p/2}] < 0$, which means $(a_2 + 1)^{1-p/2} - a_2^{-p/2} - 1 < a_2^{1-p/2} - 1$.




FIGURE 1: The upper bound of p -RIC of order $(a+1)k$ with particular values of p .

Therefore, combining the previous three cases, we obtain that one can choose $a_1 = a_2$ to get the weakest sufficient condition. It is easy to see that $a_1 = a_2$ satisfying the assumptions of Theorem 5.

After the previous discussion, using condition (25) and choosing $a_1 = a_2 = a$, we can derive condition (14).

Specifically, we consider the following function:

$$f(p) = \frac{a^{1-p/2} - 1}{a^{1-p/2} + 1} = 1 - \frac{2}{a^{1-p/2} + 1}. \quad (36)$$

Clearly,

$$f'(p) = -\frac{a^{1-p/2} \ln a}{(a^{1-p/2} + 1)^2} < 0, \quad (37)$$

and hence $f(p)$ is a decreasing function of p . Thus, for all $0 < p \leq 1$, condition

$$\delta_{(a+1)k} < \frac{\sqrt{a} - 1}{\sqrt{a} + 1} \quad (38)$$

can guarantee condition (14).

The proof is completed. \square

Applying Theorem 6, we understand how the p -RIP bounds related to p as in Figure 1. From Figure 1, it is easy to give a stronger result (i.e., weaker sufficient condition) for smaller p . Moreover, by taking different values of p with $p = 1/4, 1/2, 3/4$, we obtain some interesting p -RIP bounds as in Table 1.

4. Final Remark

In this paper, we studied the restricted p -isometry property to the partially sparse signal recovery problem and proposed a sufficient p -RIP condition for exactly recovering partially sparse signal via the partially l_p -minimization problem with $p \in (0, 1]$. It is worth generalizing the p -RIP condition

TABLE I: Bounds comparison on different values of p .

$p = \frac{1}{4}$	0.1756	0.2943	0.3807	0.4468	0.4991	0.5417
$p = \frac{1}{2}$	0.1509	0.2542	0.3307	0.3902	0.4380	0.4776
$p = \frac{3}{4}$	0.1260	0.2133	0.2788	0.3304	0.3726	0.4080
$p = 1$	0.1010	0.1716	0.2251	0.2679	0.3033	0.3333





from the vector case to the matrix case. Note that the well-known low-rank matrix recovery (LMR) problem has many applications and appeared in the literature of a diverse set of fields including matrix completion, quantum state tomography, face recognition, magnetic resonance imaging (MRI), computer vision, and system identification and control; see, for example, [21, 22] for more details and the reference therein. In particular, Oymak et al. [21] showed that several sufficient RIP recovery conditions for k sparse vector are also sufficient for recovery of matrices of rank up to $2k$ via Schatten p -norm minimization. According to our approach to extend the p -RIP bound from SSP to partially SSR, we can obtain some different restricted p -isometry properties for LMR problem by using the idea in [21].

Acknowledgments

The work was supported in part by the National Basic Research Program of China (2010CB732501), the National Natural Science Foundation of China (11171018), and the Fundamental Research Funds for the Central Universities (2011JBM128, 2013JBM095).

References

- [1] A. Bandeira, K. Scheinberg, and L. N. Vicente, "On partially sparse recovery," Department of Mathematics, University of Coimbra, 2011.
- [2] A. S. Bandeira, K. Scheinberg, and L. N. Vicente, "Computation of sparse low degree interpolating polynomials and their application to derivative-free optimization," *Mathematical Programming*, vol. 134, no. 1, pp. 223–257, 2012.
- [3] N. Vaswani and W. Lu, "Modified-CS: modifying compressive sensing for problems with partially known support," *IEEE Transactions on Signal Processing*, vol. 58, no. 9, pp. 4595–4607, 2010.
- [4] L. Jacques, "A short note on compressed sensing with partially known signal support," *Signal Processing*, vol. 90, no. 12, pp. 3308–3312, 2010.
- [5] D. L. Donoho, "Compressed sensing," *IEEE Transactions on Information Theory*, vol. 52, no. 4, pp. 1289–1306, 2006.
- [6] A. M. Bruckstein, D. L. Donoho, and M. Elad, "From sparse solutions of systems of equations to sparse modeling of signals and images," *SIAM Review*, vol. 51, no. 1, pp. 34–81, 2009.
- [7] E. J. Candés, J. Romberg, and T. Tao, "Robust uncertainty principles: exact signal reconstruction from highly incomplete frequency information," *IEEE Transactions on Information Theory*, vol. 52, no. 2, pp. 489–509, 2006.
- [8] E. J. Candés and T. Tao, "Decoding by linear programming," *IEEE Transactions on Information Theory*, vol. 51, no. 12, pp. 4203–4215, 2005.
- [9] R. Chartrand, "Fast algorithms for nonconvex compressive sensing: MRI reconstruction from very few data," in *Proceedings of the IEEE International Symposium on Biomedical Imaging (ISBI '09)*, pp. 262–265, 2009.
- [10] R. Chartrand and V. Staneva, "Restricted isometry properties and nonconvex compressive sensing," *Inverse Problems*, vol. 24, no. 3, pp. 1–14, 2008.
- [11] X. Chen, F. Xu, and Y. Ye, "Lower bound theory of nonzero entries in solutions of ℓ_2 - ℓ_p minimization," *SIAM Journal on Scientific Computing*, vol. 32, no. 5, pp. 2832–2852, 2010.
- [12] X. Chen and W. Zhou, "Convergence of reweighted ℓ_1 minimization algorithms and unique solution of truncated lp minimization," Tech. Rep., 2010.
- [13] M. E. Davies and R. Gribonval, "Restricted isometry constants where ℓ_p sparse recovery can fail for $0 < p \leq 1$," *IEEE Transactions on Information Theory*, vol. 55, no. 5, pp. 2203–2214, 2009.
- [14] S. Foucart and M.-J. Lai, "Sparsest solutions of underdetermined linear systems via ℓ_q -minimization for $0 < p \leq 1$," *Applied and Computational Harmonic Analysis*, vol. 26, no. 3, pp. 395–407, 2009.
- [15] D. Ge, X. Jiang, and Y. Ye, "A note on the complexity of ℓ_p minimization," *Mathematical Programming*, vol. 129, no. 2, pp. 285–299, 2011.
- [16] R. Saab, R. Chartrand, and O. Yilmaz, "Stable sparse approximations via nonconvex optimization," in *Proceedings of the IEEE International Conference on Acoustics, Speech and Signal Processing*, pp. 3885–3888, 2008.
- [17] Y. Shen and S. Li, "Restricted p -isometry property and its application for nonconvex compressive sensing," *Advances in Computational Mathematics*, vol. 37, no. 3, pp. 441–452, 2012.
- [18] R. Saab and O. Yilmaz, "Sparse recovery by non-convex optimization—instance optimality," *Applied and Computational Harmonic Analysis*, vol. 29, no. 1, pp. 30–48, 2010.
- [19] M. Wang, W. Xu, and A. Tang, "On the performance of sparse recovery via ℓ_p -minimization ($0 \leq p \leq 1$)," *IEEE Transactions on Information Theory*, vol. 57, no. 11, pp. 7255–7278, 2011.
- [20] R. Chartrand, "Exact reconstructions of sparse signals via nonconvex minimization," *IEEE Signal Processing Letters*, vol. 14, no. 10, pp. 707–710, 2007.
- [21] S. Oymak, K. Mohan, M. Fazel, and B. Hassibi, "A simplified approach to recovery conditions for low rank matrices," in *Proceedings of the IEEE International Symposium on Information Theory Proceedings (ISIT '11)*, pp. 2318–2322, 2011.
- [22] B. Recht, M. Fazel, and P. A. Parrilo, "Guaranteed minimum-rank solutions of linear matrix equations via nuclear norm minimization," *SIAM Review*, vol. 52, no. 3, pp. 471–501, 2010.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

