

Research Article

Fixed Point Iterations of a Pair of Hemirelatively Nonexpansive Mappings

Yan Hao^{1,2} and Sun Young Cho³

¹ School of Mathematics Physics and Information Science, Zhejiang Ocean University, Zhoushan 316004, China

² College of Mathematics, Physics and Information Engineering, Zhejiang Normal University, Jinhua 321004, China

³ Department of Mathematics, Gyeongsang National University, Jinju 660-701, South Korea

Correspondence should be addressed to Yan Hao, zjhaoyan@yahoo.cn

Received 27 September 2009; Accepted 22 March 2010

Academic Editor: Tomonari Suzuki

Copyright © 2010 Y. Hao and S. Y. Cho. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We introduce an iterative method for a pair of hemirelatively nonexpansive mappings. Strong convergence of the purposed iterative method is obtained in a Banach space.

1. Introduction and Preliminaries

Let E be a Banach space with the dual E^* . We denote by J the normalized duality mapping from E to 2^{E^*} defined by

$$Jx = \left\{ f^* \in E^* : \langle x, f^* \rangle = \|x\|^2 = \|f^*\|^2 \right\}, \quad (1.1)$$

where $\langle \cdot, \cdot \rangle$ denotes the generalized duality pairing. A Banach space E is said to be strictly convex if $\|(x + y)/2\| < 1$ for all $x, y \in E$ with $\|x\| = \|y\| = 1$ and $x \neq y$. It is said to be uniformly convex if $\lim_{n \rightarrow \infty} \|x_n - y_n\| = 0$ for any two sequences $\{x_n\}$ and $\{y_n\}$ in E such that $\|x_n\| = \|y_n\| = 1$ and $\lim_{n \rightarrow \infty} \|(x_n + y_n)/2\| = 1$. Let $U_E = \{x \in E : \|x\| = 1\}$ be the unit sphere of E . Then the Banach space E is said to be smooth provided that

$$\lim_{t \rightarrow 0} \frac{\|x + ty\| - \|x\|}{t} \quad (1.2)$$

exists for each $x, y \in U_E$. It is also said to be uniformly smooth if the limit (1.2) is attained uniformly for $x, y \in U_E$. It is well known that if E is uniformly smooth, then J is uniformly norm-to-norm continuous on each bounded subset of E . It is also well known that E is uniformly smooth if and only if E^* is uniformly convex.

Recall that a Banach space E has the Kadec-Klee property if for any sequences $\{x_n\} \subset E$ and $x \in E$ with $x_n \rightarrow x$ and $\|x_n\| \rightarrow \|x\|$, then $\|x_n - x\| \rightarrow 0$ as $n \rightarrow \infty$; for more details on Kadec-Klee property, the readers is referred to [1, 2] and the references therein. It is well known that if E is a uniformly convex Banach space, then E enjoys the Kadec-Klee property.

Let C be a nonempty closed and convex subset of a Banach space E and $T: C \rightarrow C$ a mapping. The mapping T is said to be closed if for any sequence $\{x_n\} \subset C$ such that $\lim_{n \rightarrow \infty} x_n = x_0$ and $\lim_{n \rightarrow \infty} Tx_n = y_0$, then $Tx_0 = y_0$. A point $x \in C$ is a fixed point of T provided $Tx = x$. In this paper, we use $F(T)$ to denote the fixed point set of T and use \rightarrow and \rightharpoonup to denote the strong convergence and weak convergence, respectively.

Recall that the mapping T is said to be nonexpansive if

$$\|Tx - Ty\| \leq \|x - y\|, \quad \forall x, y \in C. \quad (1.3)$$

It is well known that if C is a nonempty bounded closed and convex subset of a uniformly convex Banach space E , then every nonexpansive self-mapping T on C has a fixed point. Further, the fixed point set of T is closed and convex.

As we all know that if C is a nonempty closed convex subset of a Hilbert space H and $P_C: H \rightarrow C$ is the metric projection of H onto C , then P_C is nonexpansive. This fact actually characterizes Hilbert spaces and consequently, it is not available in more general Banach spaces. In this connection, Alber [3] recently introduced a generalized projection operator Π_C in a Banach space E which is an analogue of the metric projection in Hilbert spaces.

Next, we assume that E is a smooth Banach space. Consider the functional defined by

$$\phi(x, y) = \|x\|^2 - 2\langle x, Jy \rangle + \|y\|^2 \quad \text{for } x, y \in E. \quad (1.4)$$

Observe that, in a Hilbert space H , (1.4) is reduced to $\phi(x, y) = \|x - y\|^2$, $x, y \in H$. The generalized projection $\Pi_C: E \rightarrow C$ is a map that assigns to an arbitrary point $x \in E$, the minimum point of the functional $\phi(x, y)$, that is, $\Pi_C x = \bar{x}$, where \bar{x} is the solution to the minimization problem

$$\phi(\bar{x}, x) = \min_{y \in C} \phi(y, x). \quad (1.5)$$

Existence and uniqueness of the operator Π_C follow from the properties of the functional $\phi(x, y)$ and strict monotonicity of the mapping J (see, e.g., [1–4]). In Hilbert spaces, $\Pi_C = P_C$. It is obvious from the definition of function ϕ that

$$(\|y\| - \|x\|)^2 \leq \phi(y, x) \leq (\|y\| + \|x\|)^2, \quad \forall x, y \in E. \quad (1.6)$$

Remark 1.1. If E is a reflexive, strictly convex and smooth Banach space, then for $x, y \in E$, $\phi(x, y) = 0$ if and only if $x = y$. It is sufficient to show that if $\phi(x, y) = 0$ then $x = y$. From

(1.6), we have $\|x\| = \|y\|$. This implies that $\langle x, Jy \rangle = \|x\|^2 = \|Jy\|^2$. From the definition of J , we have $Jx = Jy$. Therefore, we have $x = y$; see [1, 2] for more details.

Let C be a nonempty closed convex subset of E and T a mapping from C into itself. A point p in C is said to be an asymptotic fixed point of T [5] if C contains a sequence $\{x_n\}$ which converges weakly to p such that $\lim_{n \rightarrow \infty} \|x_n - Tx_n\| = 0$. The set of asymptotic fixed points of T will be denoted by $\tilde{F}(T)$. A mapping T from C into itself is said to be relatively nonexpansive [3, 6, 7] if $\tilde{F}(T) = F(T) \neq \emptyset$ and $\phi(p, Tx) \leq \phi(p, x)$ for all $x \in C$ and $p \in F(T)$. The mapping T is said to be hemirelatively nonexpansive [8–12] if $F(T) \neq \emptyset$ and $\phi(p, Tx) \leq \phi(p, x)$ for all $x \in C$ and $p \in F(T)$. The asymptotic behavior of a relatively nonexpansive mappings was studied in [3, 6, 7].

Remark 1.2. The class of hemirelatively nonexpansive mappings is more general than the class of relatively nonexpansive mappings which requires the restriction: $F(T) = \tilde{F}(T)$. From Su et al. [11], we see that every hemirelatively nonexpansive mapping is relatively nonexpansive, but the inverse is not true. Hemirelatively nonexpansive mapping is also said to be quasi- ϕ -nonexpansive; see [13–17].

Recently, fixed point iterations of relatively nonexpansive mappings and hemirelatively nonexpansive mappings have been considered by many authors; see, for example [14–25] and the references therein. In 2005, Matsushita and Takahashi [8] considered fixed point problems of a single relatively nonexpansive mapping in a Banach space. To be more precise, they proved the following theorem.

Theorem MT. *Let E be a uniformly convex and uniformly smooth Banach space; let C be a nonempty closed convex subset of E ; let T be a relatively nonexpansive mapping from C into itself; let $\{\alpha_n\}$ be a sequence of real numbers such that $0 \leq \alpha_n < 1$ and $\limsup_{n \rightarrow \infty} \alpha_n < 1$. Suppose that $\{x_n\}$ is given by*

$$\begin{aligned} x_0 &= x \in C, \\ y_n &= J^{-1}(\alpha_n Jx_n + (1 - \alpha_n)JT x_n), \\ H_n &= \{z \in C : \phi(z, y_n) \leq \phi(z, x_n)\}, \\ W_n &= \{z \in C : \langle x_n - z, Jx - Jx_n \rangle \geq 0\}, \\ x_{n+1} &= P_{H_n \cap W_n} x_0, \quad n = 0, 1, 2, \dots, \end{aligned} \tag{1.7}$$

where J is the duality mapping on E . If $F(T)$ is nonempty, then $\{x_n\}$ converges strongly to $P_{F(T)}x$, where $P_{F(T)}$ is the generalized projection from C onto $F(T)$.

In 2007, Plubtieng and Ungchittrakool [9] further improved Theorem MT by considering a pair of relatively nonexpansive mappings. To be more precise, they proved the following theorem.

Theorem PU. Let E be a uniformly convex and uniformly smooth Banach space, and let C be a nonempty closed convex subset of E . Let S and T be two relatively nonexpansive mappings from C into itself with $F := F(T) \cap F(S)$ being nonempty. Let a sequence $\{x_n\}$ be defined by

$$\begin{aligned}
x_0 &= x \in C, \\
y_n &= J^{-1}(\alpha_n Jx_n + (1 - \alpha_n)Jz_n), \\
z_n &= J^{-1}(\beta_n^1 Jx_n + \beta_n^2 JT x_n + \beta_n^3 JS x_n), \\
H_n &= \{z \in C : \phi(z, y_n) \leq \phi(z, x_n)\}, \\
W_n &= \{z \in C : \langle x_n - z, Jx - Jx_n \rangle \geq 0\}, \\
x_{n+1} &= \Pi_{H_n \cap W_n} x, \quad \forall n \geq 0,
\end{aligned} \tag{1.8}$$

with the following restrictions:

- (1) $0 \leq \alpha_n < 1$ for each $n \geq 0$ and $\limsup_{n \rightarrow \infty} \alpha_n < 1$;
- (2) $0 \leq \beta_n^1, \beta_n^2, \beta_n^3 \leq 1$, $\beta_n^1 + \beta_n^2 + \beta_n^3 = 1$ for each $n \geq 0$, $\lim_{n \rightarrow \infty} \beta_n^1 = 0$ and $\liminf_{n \rightarrow \infty} \beta_n^2 \beta_n^3 > 0$.

Then the sequence $\{x_n\}$ converges strongly to $\Pi_F x$, where Π_F is the generalized projection from C onto F .

Very recently, Su et al. [11] improved Theorem PU partially by considering a pair of hemirelatively nonexpansive mappings. To be more precise, they obtained the following results.

Theorem SWX. Let E be a uniformly convex and uniformly smooth Banach space, and let C be a nonempty closed convex subset of E . Let S and T be two closed hemirelatively nonexpansive mappings from C into itself with $F := F(T) \cap F(S)$ being nonempty. Let a sequence $\{x_n\}$ be defined by

$$\begin{aligned}
x_0 &= x \in C, \\
y_n &= J^{-1}(\alpha_n Jx_n + (1 - \alpha_n)Jz_n), \\
z_n &= J^{-1}(\beta_n^1 Jx_n + \beta_n^2 JT x_n + \beta_n^3 JS x_n), \\
C_n &= \{z \in C_{n-1} \cap Q_{n-1} : \phi(z, y_n) \leq \phi(z, x_n)\}, \\
C_0 &= \{z \in C \cap Q_{n-1} : \phi(z, y_0) \leq \phi(z, x_0)\}, \\
Q_n &= \{z \in C_{n-1} \cap Q_{n-1} : \langle x_n - z, Jx - Jx_n \rangle \geq 0\}, \\
Q_0 &= C, \\
x_{n+1} &= \Pi_{C_n \cap Q_n} x_0, \quad \forall n \geq 0,
\end{aligned} \tag{1.9}$$

with the following restrictions:

- (1) $\liminf_{n \rightarrow \infty} \beta_n^1 \beta_n^2 > 0$;
- (2) $\liminf_{n \rightarrow \infty} \beta_n^1 \beta_n^3 > 0$;
- (3) $0 \leq \alpha_n \leq \alpha < 1$ for some $\alpha \in (0, 1)$.

Then the sequence $\{x_n\}$ converges strongly to $\Pi_F x$, where Π_F is the generalized projection from C onto F .

In this paper, motivated by Theorems MT, PU, and SWX, we consider the problem of finding a common fixed point of a pair of hemirelatively nonexpansive mappings by shrinking projection methods which were introduced by Takahashi et al. [26] in Hilbert spaces. Strong convergence theorems of common fixed points are established in a Banach space. The results presented in this paper mainly improve the corresponding results announced in Matsushita and Takahashi [8], Nakajo and Takahashi [27], and Su et al. [11].

In order to prove our main results, we need the following lemmas.

Lemma 1.3 (see [3]). *Let C be a nonempty closed convex subset of a smooth Banach space E and $x \in E$. Then, $x_0 = \Pi_C x$ if and only if*

$$\langle x_0 - y, Jx - Jx_0 \rangle \geq 0 \quad \forall y \in C. \quad (1.10)$$

Lemma 1.4 (see [3]). *Let E be a reflexive, strictly convex and smooth Banach space, C a nonempty closed convex subset of E , and $x \in E$. Then*

$$\phi(y, \Pi_C x) + \phi(\Pi_C x, x) \leq \phi(y, x) \quad \forall y \in C. \quad (1.11)$$

The following lemma can be deduced from Matsushita and Takahashi [8].

Lemma 1.5. *Let E be a strictly convex and smooth Banach space, C a nonempty closed convex subset of E and $T : C \rightarrow C$ a hemirelatively nonexpansive mapping. Then $F(T)$ is a closed convex subset of C .*

Lemma 1.6 (see [28]). *Let E be a uniformly convex Banach space and $B_r(0)$ a closed ball of E . Then there exists a continuous strictly increasing convex function $g : [0, \infty) \rightarrow [0, \infty)$ with $g(0) = 0$ such that*

$$\|\lambda x + \mu y + \gamma z\|^2 \leq \lambda \|x\|^2 + \mu \|y\|^2 + \gamma \|z\|^2 - \lambda \mu g(\|x - y\|) \quad (1.12)$$

for all $x, y, z \in B_r(0)$ and $\lambda, \mu, \gamma \in [0, 1]$ with $\lambda + \mu + \gamma = 1$.

2. Main Results

Theorem 2.1. *Let E be a uniformly smooth and strictly convex Banach space which enjoys the Kadec-Klee property and C a nonempty closed and convex subset of E . Let $T : C \rightarrow C$ and $S : C \rightarrow C$ be*

two closed and hemirelatively nonexpansive mappings such that $\mathcal{F} = F(T) \cap F(S)$ is nonempty. Let $\{x_n\}$ be a sequence generated in the following manner:

$$\begin{aligned}
x_0 &\in E \text{ chosen arbitrarily,} \\
C_1 &= C, \\
x_1 &= \Pi_{C_1} x_0, \\
z_n &= J^{-1}(\beta_{n,0}Jx_n + \beta_{n,1}JT x_n + \beta_{n,2}JSx_n), \\
y_n &= J^{-1}(\alpha_n Jx_n + (1 - \alpha_n)Jz_n), \\
C_{n+1} &= \{z \in C_n : \phi(z, y_n) \leq \phi(z, x_n)\}, \\
x_{n+1} &= \Pi_{C_{n+1}} x_0, \quad \forall n \geq 0,
\end{aligned} \tag{2.1}$$

where $\{\alpha_n\}$, $\{\beta_{n,0}\}$, $\{\beta_{n,1}\}$, and $\{\beta_{n,2}\}$ are real sequences in $[0, 1]$ satisfying the following restrictions:

- (a) $\limsup_{n \rightarrow \infty} \alpha_n < 1$;
- (b) $\beta_{n,0} + \beta_{n,1} + \beta_{n,2} = 1$;
- (c) $\liminf_{n \rightarrow \infty} \beta_{n,0}\beta_{n,1} > 0$ and $\liminf_{n \rightarrow \infty} \beta_{n,0}\beta_{n,2} > 0$.

Then $\{x_n\}$ converges strongly to $\Pi_{\mathcal{F}} x_0$, where $\Pi_{\mathcal{F}}$ is the generalized projection from E onto \mathcal{F} .

Proof. First, we show that C_n is closed and convex for each $n \geq 1$. It is obvious that $C_1 = C$ is closed and convex. Suppose that C_h is closed and convex for some h . For $z \in C_h$, we see that $\phi(z, y_h) \leq \phi(z, x_h)$ is equivalent to

$$2\langle z, Jx_h - Jy_h \rangle \leq \|x_h\|^2 - \|y_h\|^2. \tag{2.2}$$

It is easy to see that C_{h+1} is closed and convex. Then, for each $n \geq 1$, C_n is closed and convex. Now, we are in a position to show that $\mathcal{F} \subset C_n$ for each $n \geq 1$. Indeed, $\mathcal{F} \subset C_1 = C$ is obvious. Suppose that $\mathcal{F} \subset C_h$ for some h . Then, for all $w \in \mathcal{F} \subset C_h$, we have

$$\begin{aligned}
\phi(w, z_h) &= \phi\left(w, J^{-1}(\beta_{h,0}Jx_h + \beta_{h,1}JT x_h + \beta_{h,2}JSx_h)\right) \\
&= \|w\|^2 - 2\langle w, \beta_{h,0}Jx_h + \beta_{h,1}JT x_h + \beta_{h,2}JSx_h \rangle \\
&\quad + \|\beta_{h,0}Jx_h + \beta_{h,1}JT x_h + \beta_{h,2}JSx_h\|^2 \\
&\leq \|w\|^2 - 2\beta_{h,0}\langle w, Jx_h \rangle - 2\beta_{h,1}\langle w, JT x_h \rangle - 2\beta_{h,2}\langle w, JSx_h \rangle \\
&\quad + \beta_{h,0}\|x_h\|^2 + \beta_{h,1}\|Tx_h\|^2 + \beta_{h,2}\|Sx_h\|^2 \\
&= \beta_{h,0}\phi(w, x_h) + \beta_{h,1}\phi(w, Tx_h) + \beta_{h,2}\phi(w, Sx_h) \\
&\leq \beta_{h,0}\phi(w, x_h) + \beta_{h,1}\phi(w, x_h) + \beta_{h,2}\phi(w, x_h) \\
&= \phi(w, x_h).
\end{aligned} \tag{2.3}$$

It follows that

$$\begin{aligned}
\phi(w, y_h) &= \phi\left(w, J^{-1}(\alpha_h Jx_h + (1 - \alpha_h)Jz_h)\right) \\
&= \|w\|^2 - 2\langle w, \alpha_h Jx_h + (1 - \alpha_h)Jz_h \rangle + \|\alpha_h Jx_h + (1 - \alpha_h)Jz_h\|^2 \\
&\leq \|w\|^2 - 2\alpha_h \langle w, Jx_h \rangle - 2(1 - \alpha_h) \langle w, Jz_h \rangle + \alpha_h \|x_h\|^2 + (1 - \alpha_h) \|z_h\|^2 \\
&= \alpha_h \phi(w, x_h) + (1 - \alpha_h) \phi(w, z_h) \\
&\leq \alpha_h \phi(w, x_h) + (1 - \alpha_h) \phi(w, x_h) \\
&= \phi(w, x_h),
\end{aligned} \tag{2.4}$$

which shows that $w \in C_{h+1}$. This implies that $\mathcal{F} \subset C_n$ for each $n \geq 1$. On the other hand, we obtain from Lemma 1.4 that

$$\phi(x_n, x_0) = \phi(\Pi_{C_n} x_0, x_0) \leq \phi(w, x_0) - \phi(w, x_n) \leq \phi(w, x_0), \tag{2.5}$$

for each $w \in \mathcal{F} \subset C_n$ and for each $n \geq 1$. This shows that the sequence $\phi(x_n, x_0)$ is bounded. From (1.6), we see that the sequence $\{x_n\}$ is also bounded. Since the space is reflexive, we may, without loss of generality, assume that $x_n \rightharpoonup \bar{x}$. Note that C_n is closed and convex for each $n \geq 1$. It is easy to see that $\bar{x} \in C_n$ for each $n \geq 1$. Note that

$$\phi(x_n, x_0) \leq \phi(\bar{x}, x_0). \tag{2.6}$$

It follows that

$$\phi(\bar{x}, x_0) \leq \liminf_{n \rightarrow \infty} \phi(x_n, x_0) \leq \limsup_{n \rightarrow \infty} \phi(x_n, x_0) \leq \phi(\bar{x}, x_0). \tag{2.7}$$

This implies that

$$\lim_{n \rightarrow \infty} \phi(x_n, x_0) = \phi(\bar{x}, x_0). \tag{2.8}$$

Hence, we have $\|x_n\| \rightarrow \|\bar{x}\|$ as $n \rightarrow \infty$. In view of the Kadec-Klee property of E , we obtain that $x_n \rightarrow \bar{x}$ as $n \rightarrow \infty$.

Next, we show that $\bar{x} \in F(T)$. By the construction of C_n , we have that $C_{n+1} \subset C_n$ and $x_{n+1} = \Pi_{C_{n+1}} x_0 \in C_n$. It follows that

$$\begin{aligned}
\phi(x_{n+1}, x_n) &= \phi(x_{n+1}, \Pi_{C_n} x_0) \\
&\leq \phi(x_{n+1}, x_0) - \phi(\Pi_{C_n} x_0, x_0) \\
&= \phi(x_{n+1}, x_0) - \phi(x_n, x_0).
\end{aligned} \tag{2.9}$$

Letting $n \rightarrow \infty$ in (2.9), we obtain that $\phi(x_{n+1}, x_n) \rightarrow 0$. In view of $x_{n+1} \in C_{n+1}$, we arrive at $\phi(x_{n+1}, y_n) \leq \phi(x_{n+1}, x_n)$. It follows that

$$\lim_{n \rightarrow \infty} \phi(x_{n+1}, y_n) = 0. \quad (2.10)$$

From (1.6), we can obtain that

$$\|y_n\| \rightarrow \|\bar{x}\| \quad \text{as } n \rightarrow \infty. \quad (2.11)$$

It follows that

$$\|Jy_n\| \rightarrow \|J\bar{x}\| \quad \text{as } n \rightarrow \infty. \quad (2.12)$$

This implies that $\{Jy_n\}$ is bounded. Note that E is reflexive and E^* is also reflexive. We may assume that $Jy_n \rightharpoonup x^* \in E^*$. In view of the reflexivity of E , we see that $J(E) = E^*$. This shows that there exists an $x \in E$ such that $Jx = x^*$. It follows that

$$\begin{aligned} \phi(x_{n+1}, y_n) &= \|x_{n+1}\|^2 - 2\langle x_{n+1}, Jy_n \rangle + \|y_n\|^2 \\ &= \|x_{n+1}\|^2 - 2\langle x_{n+1}, Jy_n \rangle + \|Jy_n\|^2. \end{aligned} \quad (2.13)$$

Taking $\liminf_{n \rightarrow \infty}$, the both sides of equality above yield that

$$\begin{aligned} 0 &\geq \|\bar{x}\|^2 - 2\langle \bar{x}, x^* \rangle + \|x^*\|^2 \\ &= \|\bar{x}\|^2 - 2\langle \bar{x}, Jx \rangle + \|Jx\|^2 \\ &= \|\bar{x}\|^2 - 2\langle \bar{x}, Jx \rangle + \|x\|^2 \\ &= \phi(\bar{x}, x). \end{aligned} \quad (2.14)$$

That is, $\bar{x} = x$, which in turn implies that $x^* = J\bar{x}$. It follows that $Jy_n \rightharpoonup J\bar{x} \in E^*$. From (2.12) and since E^* enjoys the Kadec-Klee property, we obtain that

$$Jy_n - J\bar{x} \rightarrow 0 \quad \text{as } n \rightarrow \infty. \quad (2.15)$$

Note that $J^{-1} : E^* \rightarrow E$ is demicontinuous. It follows that $y_n \rightarrow \bar{x}$. From (2.11) and since E enjoys the Kadec-Klee property, we obtain that

$$y_n \rightarrow \bar{x} \quad \text{as } n \rightarrow \infty. \quad (2.16)$$

Note that

$$\|x_n - y_n\| \leq \|x_n - \bar{x}\| + \|\bar{x} - y_n\|. \quad (2.17)$$

It follows that

$$\lim_{n \rightarrow \infty} \|x_n - y_n\| = 0. \quad (2.18)$$

Since J is uniformly norm-to-norm continuous on any bounded sets, we have

$$\lim_{n \rightarrow \infty} \|Jx_n - Jy_n\| = 0. \quad (2.19)$$

On the other hand, we see from the definition of y_n that

$$\|Jy_n - Jx_n\| = (1 - \alpha_n)\|Jz_n - Jx_n\|. \quad (2.20)$$

In view of the assumption on $\{\alpha_n\}$ and (2.19), we see that

$$\lim_{n \rightarrow \infty} \|Jx_n - Jz_n\| = 0. \quad (2.21)$$

On the other hand, since $J : E \rightarrow E^*$ is demicontinuous, we have $Jx_n \rightarrow J\bar{x} \in E^*$. In view of

$$\| \|Jx_n\| - \|J\bar{x}\| \| = \| \|x_n\| - \|\bar{x}\| \| \leq \|x_n - \bar{x}\|, \quad (2.22)$$

we arrive at $\|Jx_n\| \rightarrow \|J\bar{x}\|$ as $n \rightarrow \infty$. By virtue of the Kadec-Klee property of E^* , we obtain that $\|Jx_n - J\bar{x}\| \rightarrow 0$ as $n \rightarrow \infty$. Note that

$$\|Jz_n - J\bar{x}\| \leq \|Jz_n - Jx_n\| + \|Jx_n - J\bar{x}\|. \quad (2.23)$$

In view of (2.21), we arrive at $\lim_{n \rightarrow \infty} \|Jz_n - J\bar{x}\| = 0$. Since $J^{-1} : E^* \rightarrow E$ is demicontinuous, we have $z_n \rightarrow \bar{x}$. Note that

$$\| \|z_n\| - \|x_n\| \| = \| \|Jz_n\| - \|J\bar{x}\| \| \leq \|Jz_n - J\bar{x}\|. \quad (2.24)$$

It follows that $\|z_n\| \rightarrow \|\bar{x}\|$ as $n \rightarrow \infty$. Since E enjoys the Kadec-Klee property, we obtain that $\lim_{n \rightarrow \infty} \|z_n - \bar{x}\| = 0$. Note that

$$\|z_n - x_n\| \leq \|z_n - \bar{x}\| + \|\bar{x} - x_n\|. \quad (2.25)$$

It follows that

$$\lim_{n \rightarrow \infty} \|z_n - x_n\| = 0. \quad (2.26)$$

Let $r = \max\{\sup_{n \geq 1}\{\|x_n\|\}, \sup_{n \geq 1}\{\|Tx_n\|\}, \sup_{n \geq 1}\{\|Sx_n\|\}\}$. Fixing $q \in \mathcal{F}$, we have from Lemma 1.6 that

$$\begin{aligned}
\phi(q, z_n) &= \phi\left(q, J^{-1}(\beta_{n,0}Jx_n + \beta_{n,1}JT x_n + \beta_{n,2}JSx_n)\right) \\
&= \|q\|^2 - 2\langle q, \beta_{n,0}Jx_n + \beta_{n,1}JT x_n + \beta_{n,2}JSx_n \rangle \\
&\quad + \|\beta_{n,0}Jx_n + \beta_{n,1}JT x_n + \beta_{n,2}JSx_n\|^2 \\
&\leq \|q\|^2 - 2\beta_{n,0}\langle q, Jx_n \rangle - 2\beta_{n,1}\langle q, JT x_n \rangle - 2\beta_{n,2}\langle q, JSx_n \rangle \\
&\quad + \beta_{n,0}\|Jx_n\|^2 + \beta_{n,1}\|JT x_n\|^2 + \beta_{n,2}\|JSx_n\|^2 - \beta_{n,0}\beta_{n,1}g(\|Jx_n - JT x_n\|) \\
&= \beta_{n,0}\phi(q, x_n) + \beta_{n,1}\phi(q, Tx_n) + \beta_{n,2}\phi(q, Sx_n) - \beta_{n,0}\beta_{n,1}g(\|Jx_n - JT x_n\|) \\
&\leq \beta_{n,0}\phi(q, x_n) + \beta_{n,1}\phi(q, x_n) + \beta_{n,2}\phi(q, x_n) - \beta_{n,0}\beta_{n,1}g(\|Jx_n - JT x_n\|) \\
&= \phi(q, x_n) - \beta_{n,0}\beta_{n,1}g(\|Jx_n - JT x_n\|).
\end{aligned} \tag{2.27}$$

It follows that

$$\beta_{n,0}\beta_{n,1}g(\|Jx_n - JT x_n\|) \leq \phi(q, x_n) - \phi(q, z_n). \tag{2.28}$$

On the other hand, we have

$$\begin{aligned}
\phi(q, x_n) - \phi(q, z_n) &= \|x_n\|^2 - \|z_n\|^2 - 2\langle q, Jx_n - Jz_n \rangle \\
&\leq \|x_n - z_n\|(\|x_n\| + \|z_n\|) + 2\|q\|\|Jx_n - Jz_n\|.
\end{aligned} \tag{2.29}$$

It follows from (2.21) and (2.26) that

$$\lim_{n \rightarrow \infty} (\phi(q, x_n) - \phi(q, z_n)) = 0. \tag{2.30}$$

In view of (2.28) and the assumption $\liminf_{n \rightarrow \infty} \beta_{n,0}\beta_{n,1} > 0$, we see that

$$\lim_{n \rightarrow \infty} g(\|Jx_n - JT x_n\|) = 0. \tag{2.31}$$

It follows from the property of g that

$$\lim_{n \rightarrow \infty} \|Jx_n - JT x_n\| = 0. \tag{2.32}$$

Note that

$$\lim_{n \rightarrow \infty} \|Jx_n - J\bar{x}\| = 0. \tag{2.33}$$

On the other hand, we have

$$\|JT x_n - J\bar{x}\| \leq \|JT x_n - Jx_n\| + \|Jx_n - J\bar{x}\|. \quad (2.34)$$

From (2.32) and (2.33), we arrive at

$$\lim_{n \rightarrow \infty} \|JT x_n - J\bar{x}\| = 0. \quad (2.35)$$

Note that $J^{-1} : E^* \rightarrow E$ is demicontinuous. It follows that $Tx_n \rightarrow \bar{x}$. On the other hand, we have

$$\|Tx_n\| - \|\bar{x}\| = \|JT x_n\| - \|J\bar{x}\| \leq \|JT x_n - J\bar{x}\|. \quad (2.36)$$

In view of (2.35), we obtain that $\|Tx_n\| \rightarrow \|\bar{x}\|$ as $n \rightarrow \infty$. Since E enjoys the Kadec-Klee property, we obtain that

$$\lim_{n \rightarrow \infty} \|Tx_n - \bar{x}\| = 0. \quad (2.37)$$

It follows from the closedness of T_1 that $T\bar{x} = \bar{x}$. By repeating (2.27)–(2.37), we can obtain that $\bar{x} \in F(S)$. This shows that $\bar{x} \in \mathcal{F}$.

Finally, we show that $\bar{x} = \Pi_{\mathcal{F}} x_0$. From $x_n = \Pi_{C_n} x_0$, we have

$$\langle x_n - w, Jx_0 - Jx_n \rangle \geq 0, \quad \forall w \in \mathcal{F} \subset C_n. \quad (2.38)$$

Taking the limit as $n \rightarrow \infty$ in (2.38), we obtain that

$$\langle \bar{x} - w, Jx_0 - J\bar{x} \rangle \geq 0, \quad \forall w \in \mathcal{F}, \quad (2.39)$$

and hence $\bar{x} = \Pi_{F(T)} x_0$ by Lemma 1.3. This completes the proof. \square

Remark 2.2. Theorem 2.1 improves Theorem SWX in the following aspects:

- (a) from the point of view on computation, we remove the set “ Q_n ” in Theorem SWX;
- (b) from the point of view on the framework of spaces, we extend Theorem SWX from a uniformly smooth and uniformly convex Banach space to a uniformly smooth and strictly convex Banach space which enjoys the Kadec-Klee property. Note that every uniformly convex Banach space enjoys the Kadec-Klee property.

If $\alpha_n = 0$ for each $n \geq 0$, then Theorem 2.1 is reduced to the following.

Corollary 2.3. *Let E be a uniformly smooth and strictly convex Banach space which enjoys the Kadec-Klee property and C a nonempty closed and convex subset of E . Let $T : C \rightarrow C$ and $S : C \rightarrow C$ be two closed and hemirelatively nonexpansive mappings such that $\mathcal{F} = F(T) \cap F(S)$ is nonempty. Let $\{x_n\}$ be a sequence generated in the following manner:*

$$\begin{aligned}
x_0 &\in E \text{ chosen arbitrarily,} \\
C_1 &= C, \\
x_1 &= \Pi_{C_1} x_0, \\
y_n &= J^{-1}(\beta_{n,0} Jx_n + \beta_{n,1} JT x_n + \beta_{n,2} JS x_n), \\
C_{n+1} &= \{z \in C_n : \phi(z, y_n) \leq \phi(z, x_n)\}, \\
x_{n+1} &= \Pi_{C_{n+1}} x_0, \quad \forall n \geq 0,
\end{aligned} \tag{2.40}$$

where $\{\beta_{n,0}\}$, $\{\beta_{n,1}\}$, and $\{\beta_{n,2}\}$ are real sequences in $[0, 1]$ satisfying the following restrictions:

- (a) $\beta_{n,0} + \beta_{n,1} + \beta_{n,2} = 1$;
- (b) $\liminf_{n \rightarrow \infty} \beta_{n,0} \beta_{n,1} > 0$ and $\liminf_{n \rightarrow \infty} \beta_{n,0} \beta_{n,2} > 0$.

Then $\{x_n\}$ converges strongly to $\Pi_{\mathcal{F}} x_0$, where $\Pi_{\mathcal{F}}$ is the generalized projection from E onto \mathcal{F} .

If $T = S$, then Corollary 2.3 is reduced to the following.

Corollary 2.4. *Let E be a uniformly smooth and strictly convex Banach space which enjoys the Kadec-Klee property and C a nonempty closed and convex subset of E . Let $T : C \rightarrow C$ be a closed and hemirelatively nonexpansive mapping with a nonempty fixed point set. Let $\{x_n\}$ be a sequence generated in the following manner:*

$$\begin{aligned}
x_0 &\in E \text{ chosen arbitrarily,} \\
C_1 &= C, \\
x_1 &= \Pi_{C_1} x_0, \\
y_n &= J^{-1}(\beta_n Jx_n + (1 - \beta_n) JT x_n), \\
C_{n+1} &= \{z \in C_n : \phi(z, y_n) \leq \phi(z, x_n)\}, \\
x_{n+1} &= \Pi_{C_{n+1}} x_0, \quad \forall n \geq 0,
\end{aligned} \tag{2.41}$$

where $\{\beta_n\}$ is a real sequence in $[0, 1]$ satisfying $\liminf_{n \rightarrow \infty} \beta_n(1 - \beta_n) > 0$. Then $\{x_n\}$ converges strongly to $\Pi_{\mathcal{F}} x_0$, where $\Pi_{\mathcal{F}}$ is the generalized projection from E onto \mathcal{F} .

Acknowledgment

This project is supported by the National Natural Science Foundation of China (no. 10901140).

References

- [1] I. Cioranescu, *Geometry of Banach Spaces, Duality Mappings and Nonlinear Problems*, vol. 62 of *Mathematics and Its Applications*, Kluwer Academic Publishers, Dordrecht, The Netherlands, 1990.
- [2] W. Takahashi, *Nonlinear Functional Analysis, Fixed Point Theory and Its Application*, Yokohama, Yokohama, Japan, 2000.
- [3] Ya. I. Alber, "Metric and generalized projection operators in Banach spaces: properties and applications," in *Theory and Applications of Nonlinear Operators of Accretive and Monotone Type*, A. G. Kartsatos, Ed., vol. 178 of *Lecture Notes in Pure and Applied Mathematics*, pp. 15–50, Marcel Dekker, New York, NY, USA, 1996.
- [4] Ya. I. Alber and S. Reich, "An iterative method for solving a class of nonlinear operator equations in Banach spaces," *Panamerican Mathematical Journal*, vol. 4, no. 2, pp. 39–54, 1994.
- [5] S. Reich, "A weak convergence theorem for the alternating method with Bregman distances," in *Theory and Applications of Nonlinear Operators of Accretive and Monotone Type*, A. G. Kartsatos, Ed., vol. 178 of *Lecture Notes in Pure and Applied Mathematics*, pp. 313–318, Marcel Dekker, New York, NY, USA, 1996.
- [6] D. Butnariu, S. Reich, and A. J. Zaslavski, "Asymptotic behavior of relatively nonexpansive operators in Banach spaces," *Journal of Applied Analysis*, vol. 7, no. 2, pp. 151–174, 2001.
- [7] D. Butnariu, S. Reich, and A. J. Zaslavski, "Weak convergence of orbits of nonlinear operators in reflexive Banach spaces," *Numerical Functional Analysis and Optimization*, vol. 24, no. 5-6, pp. 489–508, 2003.
- [8] S. Y. Matsushita and W. Takahashi, "A strong convergence theorem for relatively nonexpansive mappings in a Banach space," *Journal of Approximation Theory*, vol. 134, no. 2, pp. 257–266, 2005.
- [9] S. Plubtieng and K. Ungchittrakool, "Strong convergence theorems for a common fixed point of two relatively nonexpansive mappings in a Banach space," *Journal of Approximation Theory*, vol. 149, no. 2, pp. 103–115, 2007.
- [10] X. Qin and Y. Su, "Strong convergence theorems for relatively nonexpansive mappings in a Banach space," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 67, no. 6, pp. 1958–1965, 2007.
- [11] Y. Su, Z. Wang, and H. Xu, "Strong convergence theorems for a common fixed point of two hemi-relatively nonexpansive mappings," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 71, no. 11, pp. 5616–5628, 2009.
- [12] Y. Su, D. Wang, and M. Shang, "Strong convergence of monotone hybrid algorithm for hemi-relatively nonexpansive mappings," *Fixed Point Theory and Applications*, vol. 2008, Article ID 284613, 8 pages, 2008.
- [13] X. Qin, Y. J. Cho, S. M. Kang, and H. Zhou, "Convergence of a modified Halpern-type iteration algorithm for quasi- ϕ -nonexpansive mappings," *Applied Mathematics Letters*, vol. 22, no. 7, pp. 1051–1055, 2009.
- [14] X. Qin, Y. J. Cho, and S. M. Kang, "Convergence theorems of common elements for equilibrium problems and fixed point problems in Banach spaces," *Journal of Computational and Applied Mathematics*, vol. 225, no. 1, pp. 20–30, 2009.
- [15] X. Qin, S. Y. Cho, and S. M. Kang, "Strong convergence of shrinking projection methods for quasi- ϕ -nonexpansive mappings and equilibrium problems," *Journal of Computational and Applied Mathematics*, vol. 234, pp. 750–760, 2010.
- [16] X. Qin, Y. J. Cho, S. Y. Cho, and S. M. Kang, "Strong convergence theorems of common fixed points for a family of quasi- ϕ -nonexpansive mappings," *Fixed Point Theory and Applications*, vol. 2010, Article ID 754320, 11 pages, 2010.
- [17] H. Zhou, G. Gao, and B. Tan, "Convergence theorems of a modified hybrid algorithm for a family of quasi- ϕ -asymptotically nonexpansive mappings," *Journal of Applied Mathematics and Computing*, vol. 32, pp. 453–464, 2010.
- [18] Y. Kimura and W. Takahashi, "On a hybrid method for a family of relatively nonexpansive mappings in a Banach space," *Journal of Mathematical Analysis and Applications*, vol. 357, no. 2, pp. 356–363, 2009.

- [19] G. Lewicki and G. Marino, "On some algorithms in Banach spaces finding fixed points of nonlinear mappings," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 71, no. 9, pp. 3964–3972, 2009.
- [20] X. Qin, S. Y. Cho, and S. M. Kang, "On hybrid projection methods for asymptotically quasi- ϕ -nonexpansive mappings," *Applied Mathematics and Computation*, vol. 215, no. 11, pp. 3874–3883, 2010.
- [21] X. Qin and Y. Su, "Approximation of a zero point of accretive operator in Banach spaces," *Journal of Mathematical Analysis and Applications*, vol. 329, no. 1, pp. 415–424, 2007.
- [22] X. L. Qin, Y. J. Cho, S. M. Kang, and H. Y. Zhou, "Convergence of a hybrid projection algorithm in Banach spaces," *Acta Applicandae Mathematicae*, vol. 108, no. 2, pp. 299–313, 2009.
- [23] L. Wei, Y. J. Cho, and H. Zhou, "A strong convergence theorem for common fixed points of two relatively nonexpansive mappings and its applications," *Journal of Applied Mathematics and Computing*, vol. 29, no. 1-2, pp. 95–103, 2009.
- [24] K. Wattanawitton and P. Kumam, "Strong convergence theorems by a new hybrid projection algorithm for fixed point problems and equilibrium problems of two relatively quasi-nonexpansive mappings," *Nonlinear Analysis: Hybrid Systems*, vol. 3, no. 1, pp. 11–20, 2009.
- [25] H. Zegeye and N. Shahzad, "Strong convergence theorems for monotone mappings and relatively weak nonexpansive mappings," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 70, no. 7, pp. 2707–2716, 2009.
- [26] W. Takahashi, Y. Takeuchi, and R. Kubota, "Strong convergence theorems by hybrid methods for families of nonexpansive mappings in Hilbert spaces," *Journal of Mathematical Analysis and Applications*, vol. 341, no. 1, pp. 276–286, 2008.
- [27] K. Nakajo and W. Takahashi, "Strong convergence theorems for nonexpansive mappings and nonexpansive semigroups," *Journal of Mathematical Analysis and Applications*, vol. 279, no. 2, pp. 372–379, 2003.
- [28] Y. J. Cho, H. Zhou, and G. Guo, "Weak and strong convergence theorems for three-step iterations with errors for asymptotically nonexpansive mappings," *Computers & Mathematics with Applications*, vol. 47, no. 4-5, pp. 707–717, 2004.