

Research Article

Iterative Algorithms with Variable Coefficients for Multivalued Generalized Φ -Hemicontractive Mappings without Generalized Lipschitz Assumption

Ci-Shui Ge

Department of Mathematics and Physics, Anhui University of Architecture, Hefei, Anhui 230022, China

Correspondence should be addressed to Ci-Shui Ge, gecishui@sohu.com

Received 17 August 2010; Accepted 8 November 2010

Academic Editor: Tomonari Suzuki

Copyright © 2010 Ci-Shui Ge. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We introduce and study some new Ishikawa-type iterative algorithms with variable coefficients for multivalued generalized Φ -hemicontractive mappings. Several new fixed-point theorems for multivalued generalized Φ -hemicontractive mappings without generalized Lipschitz assumption are established in p -uniformly smooth real Banach spaces. A result for multivalued generalized Φ -hemicontractive mappings with bounded range is obtained in uniformly smooth real Banach spaces. As applications, several theorems for multivalued generalized Φ -hemicontractive mapping equations are given.

1. Introduction

Let X be a real Banach space and X^* the dual space of X . $\langle \cdot, \cdot \rangle$ denotes the generalized duality pairing between X and X^* . J is the normalized duality mapping from X to 2^{X^*} given by $J(x)$

$$J(x) := \{f \in X^* : \langle x, f \rangle = \|f\| \cdot \|x\|, \|f\| = \|x\|\}, \quad x \in X. \quad (1.1)$$

Let D be a nonempty convex subset of X and $CB(D)$ the family of all nonempty bounded closed subsets of D . $H(\cdot, \cdot)$ denotes the Hausdorff metric on $CB(D)$ defined by

$$H(A, B) := \max \left\{ \sup_{y \in B} \inf_{x \in A} \|x - y\|, \sup_{y \in A} \inf_{x \in B} \|x - y\| \right\}, \quad A, B \in CB(D). \quad (1.2)$$

We use $F(T)$ to denote the fixed-point set of T , that is, $F(T) := \{x : x \in Tx\}$. \mathbb{N} denotes the set of nonnegative integers.

Recall that a mapping $T : D \rightarrow D$ is called to be a generalized Lipschitz mapping [1], if there exists a constant $L > 0$ such that

$$\|Tx - Ty\| \leq L(1 + \|x - y\|), \quad \forall x, y \in D. \quad (1.3)$$

Similarly, a multivalued mapping $T : D \rightarrow CB(D)$ is said to be a generalized Lipschitz mapping, if there exists a constant $L > 0$ such that

$$H(Tx, Ty) \leq L(1 + \|x - y\|), \quad \forall x, y \in D. \quad (1.4)$$

A multivalued mapping $T : D \rightarrow 2^D$ is said to be a bounded mapping if for any bounded subset A of D ,

$$T(A) := \{x : x \in T(y), \exists y \in A\} \quad (1.5)$$

is a bounded subset of D .

Clearly, every mapping with bounded range is a generalized Lipschitz mapping [1, Example]. Furthermore, every generalized Lipschitz mapping is a bounded mapping. The following example shows that the class of generalized Lipschitz mappings is a proper subset of the class of bounded mappings.

Example 1.1. Take $D = (0, \infty)$ and define $T : D \rightarrow D$ by

$$Tx = \exp(x) + x \operatorname{sgn}(\sin x), \quad (1.6)$$

where $\operatorname{sgn}(\cdot)$ denotes sign function. Then, T is a bounded mapping but not a generalized Lipschitz mapping.

Definition 1.2 (see [2]). Let D be a nonempty subset of X . $T : D \rightarrow 2^D$ is said to be a multivalued Φ -hemiccontractive mapping if the fixed point set $F(T)$ of T is nonempty, and there exists a strictly increasing function $\Phi : [0, \infty) \rightarrow [0, \infty)$ with $\Phi(0) = 0$ such that for each $x \in D$ and $x^* \in F(T)$, there exists a $j(x - x^*) \in J(x - x^*)$ such that

$$\langle u - x^*, j(x - x^*) \rangle \leq \|x - x^*\|^2 - \Phi(\|x - x^*\|) \cdot \|x - x^*\|, \quad (1.7)$$

for all $u \in Tx$.

T is said to be a multivalued Φ -hemiaccretive mapping if $I - T$ is a multivalued Φ -hemiccontractive mapping.

Definition 1.3. Let D be a nonempty subset of X . $T : D \rightarrow 2^D$ is said to be a multivalued generalized Φ -hemiccontractive mapping if the fixed point set $F(T)$ of T is nonempty,

and there exists a strictly increasing function $\Phi : [0, \infty) \rightarrow [0, \infty)$ with $\Phi(0) = 0$ such that for each $x \in D$ and $x^* \in F(T)$, there exists a $j(x - x^*) \in J(x - x^*)$ such that

$$\langle u - x^*, j(x - x^*) \rangle \leq \|x - x^*\|^2 - \Phi(\|x - x^*\|), \quad (1.8)$$

for all $u \in Tx$.

T is said to be a multivalued generalized Φ -hemiccretive mapping if $I - T$ is a multivalued generalized Φ -hemiccontractive mapping.

The following example shows that the class of Φ -hemiccontractive mappings is a proper subset of the class of generalized Φ -hemiccontractive mappings.

Example 1.4. Let $X = \mathbb{R}^2$ with the Euclidean norm $\|\cdot\|$, where \mathbb{R} denotes the set of the real numbers. Define $T : X \rightarrow X$ by

$$Tx = \frac{\|x\|^2}{1 + \|x\|^2}x. \quad (1.9)$$

Thus, $F(T) = \{(0,0)\} \neq \emptyset$. It is easy to verify that T is a generalized Φ -hemiccontractive mapping with $\Phi(t) = t^2/(1+t^2)$. However, T is not Φ -hemiccontractive. Indeed, if there exists a strictly increasing function $\phi : [0, \infty) \rightarrow [0, \infty)$ with $\phi(0) = 0$ such that for each $x \in X$ and $x^* = (0,0) \in F(T)$,

$$\langle Tx - x^*, J(x - x^*) \rangle \leq \|x - x^*\|^2 - \phi(\|x - x^*\|) \cdot \|x - x^*\|, \quad (1.10)$$

then we get $\phi(t) \leq t/(1+t^2)$ for all $t \in (0, \infty)$. Thus, $\lim_{t \rightarrow \infty} \phi(t) = 0$. This is in contradiction with the hypotheses that $\phi(t)$ is strictly increasing and $\phi(0) = 0$.

In the last twenty years or so, numerous papers have been written on the existence and convergence of fixed points for nonlinear mappings, and strong and weak convergence theorems have been obtained by using some well-known iterative algorithms (see, e.g., [1–9] and the references therein).

For multivalued ϕ -hemiccontractive mappings, Hirano and Huang [2] obtained the following result.

Theorem HH (See [2, Theorem 1]). *Let E be a uniformly smooth Banach space and $T : E \rightarrow 2^E$ be a multivalued ϕ -hemiccontractive operator with bounded range. Suppose $\{a_n\}$, $\{b_n\}$, $\{c_n\}$ and $\{a'_n\}$, $\{b'_n\}$, $\{c'_n\}$ are real sequences in $[0, 1)$ satisfying the following conditions:*

- (i) $a_n + b_n + c_n = a'_n + b'_n + c'_n = 1$, for all $n \in \mathbb{N}$,
- (ii) $\lim_{n \rightarrow \infty} b_n = \lim_{n \rightarrow \infty} b'_n = \lim_{n \rightarrow \infty} c_n = 0$,
- (iii) $\sum_{n=1}^{\infty} b_n = \infty$,
- (iv) $c_n = o(b_n)$.

For arbitrary $x_1, u_1, v_1 \in E$, define the sequence $\{x_n\}_{n=1}^\infty$ by

$$\begin{aligned} x_{n+1} &= a_n x_n + b_n \eta_n + c_n u_n, & \exists \eta_n \in T y_n, n \in \mathbb{N}, \\ y_n &= a'_n x_n + b'_n \xi_n + c'_n v_n, & \exists \xi_n \in T x_n, n \in \mathbb{N}, \end{aligned} \quad (1.11)$$

where $\{u_n\}_{n=1}^\infty, \{v_n\}_{n=1}^\infty$ are arbitrary bounded sequences in E . Then, $\{x_n\}_{n=1}^\infty$ converges strongly to the unique fixed point of T .

Further, for general multivalued generalized Φ -hemicontractive mappings, C. E. Chidume and C. O. Chidume [1] gave the following interesting result.

Theorem CC (see [1, Theorem 3.8]). *Let E be a uniformly smooth real Banach space. Let $F(T) := \{x \in E : x \in Tx\} \neq \emptyset$. Suppose $T : E \rightarrow 2^E$ is a multivalued generalized Lipschitz and generalized Φ -hemicontractive mapping. Let $\{a_n\}, \{b_n\}$ and $\{c_n\}$ be real sequences in $[0, 1)$ satisfying the following conditions: (i) $a_n + b_n + c_n = 1$, (ii) $\sum (b_n + c_n) = \infty$, (iii) $\sum c_n < \infty$, and (iv) $\lim b_n = 0$. Let $\{x_n\}$ be generated iteratively from arbitrary $x_0 \in E$ by*

$$x_{n+1} = a_n x_n + b_n \eta_n + c_n u_n, \quad \exists \eta_n \in T x_n, n \geq 0, \quad (1.12)$$

where $\{u_n\}$ is an arbitrary bounded sequence in E . Then, there exists $\gamma_0 \in \mathbb{R}$ such that if $b_n + c_n \leq \gamma_0$ for all $n \geq 0$, the sequence $\{x_n\}$ converges strongly to the unique fixed point of T .

Remark 1.5. (1) Theorem CC [1, Theorem 3.8] is a multivalued version of Theorem 3.2 of [1]. Theorem 3.2 of [1] was obtained directly from Theorem 3.1 of [1]. However, it seems that there exists a gap in the proof of Theorem 3.1 in [1]. Indeed, the following inequality in the proof of Theorem 3.1 in [1].

$$a_0 \sum_{j=0}^n \alpha_j \leq \sum_{j=0}^n \left(\|x_j - x^*\|^2 - \|x_{j+1} - x^*\|^2 \right) + M \sum_{j=0}^n c_j < \infty \quad (*)$$

was obtained by using implicitly the following conditions:

$$\|x_j - x^*\| \leq 2\Phi^{-1}(a_0), \quad \|x_{j+1} - x^*\| > 2\Phi^{-1}(a_0), \quad j = 0, 1, \dots, n. \quad (1.13)$$

Thus, (*) is dubious in the remainder of [1, Theorem 3.1]. Hence, Theorem 3.1 of [1] is dubious, as is Theorem CC [1, Theorem 3.8].

(2) The real number γ_0 in Theorem CC is not easy to get.

It is our purpose in this paper to try to obtain some fixed-point theorems for multivalued generalized Φ -hemicontractive mappings without generalized Lipschitz assumption as in Theorem CC. Motivated and inspired by [1, 2, 5, 7], we introduce and study some new Ishikawa-type iterative algorithms with variable coefficients for multivalued generalized Φ -hemicontractive mappings. Our results improve essentially the corresponding results of [1] in the framework of p -uniformly smooth real Banach spaces and the corresponding results of [2] in uniformly smooth real Banach spaces.

2. Preliminaries

Let X be a real Banach space of dimension $\dim X \geq 2$. The modulus of smoothness of X is the function $\rho_X : [0, \infty) \rightarrow [0, \infty)$ defined by

$$\rho_X(\tau) := \sup \left\{ 2^{-1}(\|x + y\| + \|x - y\|) - 1 : \|x\| = 1, \|y\| \leq \tau \right\}, \quad \tau > 0. \quad (2.1)$$

The function $\rho_X(\tau)$ is convex, continuous, and increasing, and $\rho_X(0) = 0$.

The space X is called uniformly smooth if and only if

$$\lim_{\tau \rightarrow 0^+} \frac{\rho_X(\tau)}{\tau} = 0. \quad (2.2)$$

The space X is called p -uniformly smooth if and only if there exist a constant C_p and a real number $1 < p \leq 2$, such that

$$\rho_X(\tau) \leq C_p \tau^p. \quad (2.3)$$

Typical examples of uniformly smooth spaces are the Lebesgue L_p , the sequence ℓ_p , and Sobolev W_p^m spaces for $1 < p < \infty$. In particular, for $1 < p \leq 2$, these spaces are p -uniformly smooth and for $2 \leq p < \infty$, they are 2-uniformly smooth.

It is well known that if X is uniformly smooth, then the normalized duality mapping J is single-valued and uniformly continuous on any bounded subset of X .

Lemma 2.1 (see [3, 9]). *If X is a uniformly smooth Banach space, then for all $x, y \in X$ with $\|x\| \leq R, \|y\| \leq R$,*

$$\begin{aligned} \langle x - y, Jx - Jy \rangle &\leq 2L_F R^2 \rho_X \left(\frac{4\|x - y\|}{R} \right), \\ \|Jx - Jy\| &\leq 8Rh_X \left(\frac{16L_F \|x - y\|}{R} \right), \end{aligned} \quad (2.4)$$

where $h_X(\tau) := \rho_X(\tau)/\tau$, L_F is the Figiel's constant, $1 < L_F < 1.7$.

Lemma 2.2 (see [1]). *Let X be a real Banach space and J be the normalized duality mapping. Then, for any given $x, y \in X$, we have*

$$\|x + y\|^2 \leq \|x\|^2 + 2\langle y, j(x + y) \rangle, \quad \forall j(x + y) \in J(x + y). \quad (2.5)$$

Lemma 2.3 (see [8]). *Let $\{\alpha_n\}_{n \geq 1}$, $\{\beta_n\}_{n \geq 1}$ and $\{\gamma_n\}_{n \geq 1}$ be nonnegative sequences satisfying*

$$\alpha_{n+1} \leq (1 + \gamma_n)\alpha_n + \beta_n, \quad n \geq 1, \quad \sum_{n=1}^{\infty} \beta_n < \infty, \quad \sum_{n=1}^{\infty} \gamma_n < \infty. \quad (2.6)$$

Then, $\lim_{n \rightarrow \infty} \alpha_n$ exists. Moreover, if $\liminf_{n \rightarrow \infty} \alpha_n = 0$, then $\lim_{n \rightarrow \infty} \alpha_n = 0$.

Lemma 2.4 (see [4]). Let $f, g : \mathbb{N} \rightarrow [0, \infty)$ be sequences and suppose that

$$g(n) \leq 1, \quad \forall n \in \mathbb{N}, \quad g(n) \longrightarrow 0, \quad \text{as } n \longrightarrow \infty, \quad \sum_{n=1}^{\infty} g(n) = \infty. \quad (2.7)$$

Then,

$$\sum_{n=1}^{\infty} f(n) < \infty \implies f = o(g), \quad \text{as } n \longrightarrow \infty. \quad (2.8)$$

The converse is false.

3. Main Results and Their Proofs

Theorem 3.1. Let X be a p -uniformly smooth real Banach space and D a nonempty convex subset of X . Suppose $T : D \rightarrow 2^D$ is a multivalued generalized Φ -hemiccontractive and bounded mapping. For any given $x_0, u_0, v_0 \in D$, let $\{x_n\}$ be the sequence generated by the following Ishikawa-type iterative algorithm with variable coefficients:

$$\begin{aligned} y_n &= \hat{a}_n x_n + \hat{b}_n \xi_n + \hat{c}_n v_n, & \exists \xi_n \in Tx_n, \\ x_{n+1} &= \hat{\alpha}_n x_n + \hat{\beta}_n \eta_n + \hat{\gamma}_n u_n, & \exists \eta_n \in Ty_n, \end{aligned} \quad n \in \mathbb{N}, \quad (3.1)$$

where $\{u_n\}$ and $\{v_n\}$ are arbitrary bounded sequences in D ,

$$\begin{aligned} \hat{a}_n &= 1 - \hat{b}_n - \hat{c}_n, & \hat{b}_n &= \frac{b_n}{r_n^2}, & \hat{c}_n &= \frac{c_n}{r_n^2}, & r_n &= 2 + \|x_n\| + \|\xi_n\| + \|v_n\|, \\ \hat{\alpha}_n &= 1 - \hat{\beta}_n - \hat{\gamma}_n, & \hat{\beta}_n &= \frac{\beta_n}{R_n^2}, & \hat{\gamma}_n &= \frac{\gamma_n}{R_n^2}, & R_n &= r_n + \|\eta_n\| + \|u_n\|, \end{aligned} \quad (3.2)$$

$\{\beta_n\}$, $\{\gamma_n\}$, $\{b_n\}$ and $\{c_n\}$ are four sequences in $[0, 1]$ satisfying the following conditions:

$$\sum_{n=0}^{\infty} \beta_n = \infty, \quad \sum_{n=0}^{\infty} \beta_n^p < \infty, \quad \sum_{n=0}^{\infty} \gamma_n < \infty, \quad b_n \leq O(\beta_n), \quad c_n \leq O(\beta_n). \quad (3.3)$$

Then, $\{x_n\}$ converges strongly to the unique fixed point of T .

Proof. Since T is generalized Φ -hemiccontractive, then the fixed-point set $F(T)$ of T is nonempty and there exists a strictly increasing function $\Phi : [0, \infty) \rightarrow [0, \infty)$ with $\Phi(0) = 0$ such that for each $x \in D$ and $x^* \in F(T)$, the following inequality holds:

$$\langle \xi - x^*, J(x - x^*) \rangle \leq \|x - x^*\|^2 - \Phi(\|x - x^*\|), \quad \forall \xi \in Tx. \quad (3.4)$$

If $z \in F(T)$, that is, $z \in Tz$, then, by (3.4), we have

$$\|z - x^*\|^2 = \langle z - x^*, J(z - x^*) \rangle \leq \|z - x^*\|^2 - \Phi(\|z - x^*\|). \quad (3.5)$$

So, T has a unique fixed point, say x^* .

From (3.1) and (3.2), we have $\|x_n - x^*\| \leq r_n + \|x^*\|$, $\|y_n - x^*\| \leq r_n + \|x^*\|$, $\|\xi_n - x^*\| \leq r_n + \|x^*\|$, $\|\eta_n - x^*\| \leq R_n + \|x^*\|$ and $\|x_{n+1} - x^*\| \leq R_n + \|x^*\|$.

By Lemma 2.4 and (3.3), we know $\gamma_n = o(\beta_n)$. Since D is a convex subset of X and $T : D \rightarrow 2^D$, it follows from (3.1), (3.2), and (3.3) that

$$\begin{aligned}
\|x_{n+1} - y_n\| &= \|(x_{n+1} - x^*) - (y_n - x^*)\| \\
&= \left\| \left(1 - \widehat{\beta}_n - \widehat{\gamma}_n\right)(x_n - x^*) + \widehat{\beta}_n (\eta_n - x^*) + \widehat{\gamma}_n (u_n - x^*) \right. \\
&\quad \left. - \left(1 - \widehat{b}_n - \widehat{c}_n\right)(x_n - x^*) + \widehat{b}_n (\xi_n - x^*) + \widehat{c}_n (v_n - x^*) \right\| \\
&\leq \frac{O(\beta_n)}{r_n^2} (r_n + \|x^*\|) + \frac{O(\beta_n)}{R_n^2} (R_n + \|x^*\|) \\
&\leq \frac{O(\beta_n)}{r_n} \longrightarrow 0 \quad (n \longrightarrow \infty).
\end{aligned} \tag{3.6}$$

From (3.6) and $\|y_n - x^*\| \leq r_n + \|x^*\|$, we have $\|x_{n+1} - x^*\| \leq r_n + \|x^*\| + (O(\beta_n)/r_n)$.

Considering $1 < p \leq 2$ and $r_n \geq 2$, by Lemma 2.1, we have

$$\begin{aligned}
\|J(x_{n+1} - x^*) - J(y_n - x^*)\| &\leq 8 \left(r_n + \|x^*\| + \frac{O(\beta_n)}{r_n} \right) C_p \cdot \left(\frac{16L_F \|x_{n+1} - y_n\|}{r_n + \|x^*\| + O(\beta_n)/r_n} \right)^{p-1} \\
&\leq \left(r_n + \|x^*\| + \frac{O(\beta_n)}{r_n} \right)^{2-p} \frac{O(\beta_n^{p-1})}{r_n^{p-1}} \\
&\leq \left(r_n^2 + r_n \|x^*\| + O(\beta_n) \right)^{2-p} \frac{O(\beta_n^{p-1})}{r_n} \\
&\leq r_n \cdot O(\beta_n^{p-1}).
\end{aligned} \tag{3.7}$$

By (3.1), (3.2), (3.3) and Lemma 2.2, we have

$$\begin{aligned}
\|y_n - x^*\|^2 &\leq \left\| \widehat{a}_n (x_n - x^*) + \widehat{b}_n (\eta_n - x^*) + \widehat{c}_n (v_n - x^*) \right\|^2 \\
&\leq \widehat{a}_n^2 \|x_n - x^*\|^2 + 2 \left\langle \widehat{b}_n (\xi_n - x^*) + \widehat{c}_n (v_n - x^*), J(y_n - x^*) \right\rangle \\
&\leq \|x_n - x^*\|^2 + O(\beta_n).
\end{aligned} \tag{3.8}$$

From (3.1), (3.2), (3.7), and (3.8) and Lemma 2.2, it can be concluded that

$$\begin{aligned}
\|x_{n+1} - x^*\|^2 &= \left\| \widehat{\alpha}_n(x_n - x^*) + \widehat{\beta}_n(\eta_n - x^*) + \widehat{\gamma}_n(u_n - x^*) \right\|^2 \\
&\leq \widehat{\alpha}_n^2 \|x_n - x^*\|^2 + 2\widehat{\beta}_n \langle \eta_n - x^*, J(x_{n+1} - x^*) - J(y_n - x^*) \rangle \\
&\quad + 2\widehat{\beta}_n \langle \eta_n - x^*, J(y_n - x^*) \rangle + 2\widehat{\gamma}_n \langle u_n - x^*, J(x_{n+1} - x^*) \rangle \\
&\leq \widehat{\alpha}_n^2 \|x_n - x^*\|^2 + 2\widehat{\beta}_n \|\eta_n - x^*\| \cdot \|J(x_{n+1} - x^*) - J(y_n - x^*)\| \\
&\quad + 2\widehat{\beta}_n \left(\|y_n - x^*\|^2 - \Phi(\|y_n - x^*\|) \right) + 2\widehat{\gamma}_n \|u_n - x^*\| \cdot \|J(x_{n+1} - x^*)\| \\
&\leq \left(1 - \widehat{\beta}_n - \widehat{\gamma}_n \right)^2 \|x_n - x^*\|^2 + 2\widehat{\beta}_n (R_n + \|x^*\|) \cdot r_n \cdot O(\beta_n^{p-1}) \\
&\quad + 2\widehat{\beta}_n \left(\|x_n - x^*\|^2 + O(\beta_n) \right) - 2\widehat{\beta}_n \Phi(\|y_n - x^*\|) + 2\widehat{\gamma}_n \cdot (R_n + \|x^*\|)^2 \\
&\leq \|x_n - x^*\|^2 + \left(\widehat{\beta}_n + \widehat{\gamma}_n \right)^2 \|x_n - x^*\|^2 + O(\beta_n^p) + O(\beta_n^2) + O(\gamma_n) \\
&\quad - 2\widehat{\beta}_n \Phi(\|y_n - x^*\|) \\
&\leq \|x_n - x^*\|^2 + O(\beta_n^2) \|x_n - x^*\|^2 + O(\beta_n^p) + O(\gamma_n) - 2\widehat{\beta}_n \Phi(\|y_n - x^*\|).
\end{aligned} \tag{3.9}$$

From (3.3) and (3.9), we have

$$\|x_{n+1} - x^*\|^2 \leq \left(1 + O(\beta_n^2) \right) \|x_n - x^*\|^2 + O(\beta_n^p) + O(\gamma_n). \tag{3.10}$$

Thus, by (3.3), (3.10) and Lemma 2.3, we have $\{\|x_n - x^*\|\}$ bounded. It implies the sequences $\{x_n\}$ and $\{y_n\}$ are bounded. Since T is a bounded mapping, we have $T\{x_n\}$ and $T\{y_n\}$ bounded. Since $\eta_n \in Ty_n$ and $\xi_n \in Tx_n$, $\{R_n\}$ is bounded. Let its bound be $R > 0$. From (3.9), there exists a number $M > 0$ such that

$$\|x_{n+1} - x^*\|^2 \leq \left(1 + M\beta_n^2 \right) \|x_n - x^*\|^2 + M(\beta_n^p + \gamma_n) - \frac{2\beta_n}{R^2} \Phi(\|y_n - x^*\|). \tag{3.11}$$

Next, we will show

$$\liminf_{n \rightarrow \infty} \Phi(\|y_n - x^*\|) = 0. \tag{3.12}$$

If it is not true, then there exist a $n_0 \in \mathbb{N}$ and a positive constant m_0 such that for any positive integer $n \geq n_0$

$$\Phi(\|y_n - x^*\|) \geq m_0. \tag{3.13}$$

In view of (3.11) and (3.13), for any positive integer $n \geq n_0$, we have

$$\|x_{n+1} - x^*\|^2 \leq \left(1 + M\beta_n^2\right)\|x_n - x^*\|^2 + M\left(\beta_n^p + \gamma_n\right) - \frac{2m_0\beta_n}{R^2}. \quad (3.14)$$

Taking $n = n_0, n_0 + 1, \dots, k$ in (3.14) above, we have

$$\begin{aligned} \sum_{n=n_0}^k \|x_{n+1} - x^*\|^2 &\leq \sum_{n=n_0}^k \|x_n - x^*\|^2 + \sum_{n=n_0}^k M\beta_n^2(R + \|x^*\|)^2 \\ &\quad + \sum_{n=n_0}^k M\left(\beta_n^p + \gamma_n\right) - \sum_{n=n_0}^k \frac{2m_0\beta_n}{R^2}. \end{aligned} \quad (3.15)$$

So,

$$\frac{2m_0}{R^2} \sum_{n=n_0}^k \beta_n \leq M(R + \|x^*\|)^2 \sum_{n=n_0}^k \beta_n^2 + M\left(\sum_{n=n_0}^k \beta_n^p + \sum_{n=n_0}^k \gamma_n\right). \quad (3.16)$$

This leads to a contradiction as $k \rightarrow \infty$. Hence, $\liminf_{n \rightarrow \infty} \Phi(\|y_n - x^*\|) = 0$.

By the definition of Φ and (3.12), there exists a subsequence $\{y_{n_i}\}$ of $\{y_n\}$ such that $\{y_{n_i}\} \rightarrow x^*$ as $i \rightarrow \infty$. Thus, by (3.6), we have $\liminf_{n \rightarrow \infty} \|x_n - x^*\| = 0$. Further, Using Lemma 2.3 and (3.11), we obtain $\lim_{n \rightarrow \infty} \|x_n - x^*\| = 0$. It means that $\{x_n\}$ converges strongly to the unique fixed point of T . The proof is finished. \square

From Theorem 3.1, we can obtain the following theorems.

Theorem 3.2. *Let X be a p -uniformly smooth Banach space, D be a nonempty convex subset of X , and $T : D \rightarrow 2^D$ a multivalued generalized Φ -hemiccontractive and bounded mapping. For any given $x_0, u_0 \in D$, let $\{x_n\}$ be the sequence generated by the following Mann-type iterative algorithm with variable coefficients:*

$$x_{n+1} = \hat{\alpha}_n x_n + \hat{\beta}_n \eta_n + \hat{\gamma}_n u_n, \quad \exists \eta_n \in Tx_n, \quad n \in \mathbb{N}, \quad (3.17)$$

where $\{u_n\}$ is an arbitrary bounded sequence in D ,

$$\hat{\alpha}_n = 1 - \hat{\beta}_n - \hat{\gamma}_n, \quad \hat{\beta}_n = \frac{\beta_n}{R_n^2}, \quad \hat{\gamma}_n = \frac{\gamma_n}{R_n^2}, \quad R_n = 2 + \|x_n\| + \|\eta_n\| + \|u_n\|, \quad (3.18)$$

$\{\beta_n\}$ and $\{\gamma_n\}$ are sequences in $[0, 1]$ satisfying the following conditions:

$$\sum_{n=0}^{\infty} \beta_n = \infty, \quad \sum_{n=0}^{\infty} \beta_n^p < \infty, \quad \sum_{n=0}^{\infty} \gamma_n < \infty. \quad (3.19)$$

Then, $\{x_n\}$ converges strongly to the unique fixed point of T .

Remark 3.3. Theorems 3.1 and 3.2 improve Theorem CC [1, Theorem 3.8] in p -uniformly smooth real Banach spaces since the class of multivalued generalized Lipschitz mappings is a proper subset of the class of bounded mappings and the number γ_0 in Theorem CC [1, Theorem 3.8] is dropped off.

In uniformly smooth real Banach spaces, we have the following theorem.

Theorem 3.4. *Let X be a uniformly smooth real Banach space and D a nonempty convex subset of X . Suppose $T : D \rightarrow 2^D$ is a multivalued generalized Φ -hemicontractive mapping with bounded range. For any given $x_0, u_0, v_0 \in D$, let $\{x_n\}$ be the sequence generated by the following Ishikawa-type iterative algorithm with variable coefficients:*

$$\begin{aligned} y_n &= \widehat{a}_n x_n + \widehat{b}_n \xi_n + \widehat{c}_n v_n, & \exists \xi_n \in Tx_n, \\ x_{n+1} &= \widehat{\alpha}_n x_n + \widehat{\beta}_n \eta_n + \widehat{\gamma}_n u_n, & \exists \eta_n \in Ty_n, \end{aligned} \quad n \in \mathbb{N}, \quad (3.20)$$

where $\{u_n\}$ and $\{v_n\}$ are arbitrary bounded sequences in D ,

$$\begin{aligned} \widehat{a}_n &= 1 - \widehat{b}_n - \widehat{c}_n, & \widehat{b}_n &= \frac{b_n}{r_n^2}, & \widehat{c}_n &= \frac{c_n}{r_n^2}, & r_n &= 2 + \|x_n\| + \|\xi_n\| + \|v_n\|, \\ \widehat{\alpha}_n &= 1 - \widehat{\beta}_n - \widehat{\gamma}_n, & \widehat{\beta}_n &= \frac{\beta_n}{R_n^2}, & \widehat{\gamma}_n &= \frac{\gamma_n}{R_n^2}, & R_n &= r_n + \|\eta_n\| + \|u_n\|, \end{aligned} \quad (3.21)$$

$\{\beta_n\}$, $\{\gamma_n\}$, $\{b_n\}$ and $\{c_n\}$ are four sequences in $[0, 1]$ satisfying the following conditions:

$$\sum_{n=0}^{\infty} \beta_n = \infty, \quad \sum_{n=0}^{\infty} \beta_n^2 < \infty, \quad \sum_{n=0}^{\infty} \gamma_n < \infty, \quad b_n \leq O(\beta_n), \quad c_n \leq O(\beta_n). \quad (3.22)$$

Then, $\{x_n\}$ converges strongly to the unique fixed point of T .

Proof. From Theorem 3.1, T has a unique fixed point, say x^* . Let $\{x_n\}$, $\{y_n\}$ be the sequences generated by the algorithm (3.20). Since T has a bounded range, we set

$$\begin{aligned} d &:= \sup\{\|\xi - \eta\| : x, y \in D, \xi \in Tx, \eta \in Ty\} + \sup\{\|u_n - x^*\|, n \in \mathbb{N}\} \\ &\quad + \sup\{\|v_n - x^*\|, n \in \mathbb{N}\}. \end{aligned} \quad (3.23)$$

Obviously, $d < \infty$. Next, we will prove that for $n \geq 0$, $\|x_n - x^*\| \leq d + \|x_0 - x^*\|$. In fact, for $n = 0$, the above inequality holds. Assume the inequality is true for $n = k$. Then, for $n = k + 1$, there exists a $\eta_k \in Ty_k$ such that

$$\begin{aligned} \|x_{k+1} - x^*\| &\leq \widehat{\alpha}_k \|x_k - x^*\| + \widehat{\beta}_k \|\eta_k - x^*\| + \widehat{\gamma}_k \|u_k - x^*\| \\ &\leq \widehat{\alpha}_k (d + \|x_0 - x^*\|) + \widehat{\beta}_k d + \widehat{\gamma}_k d \\ &\leq d + \|x_0 - x^*\|. \end{aligned} \quad (3.24)$$

By induction, we have the sequence $\{x_n\}$ bounded. Similarly, we have the sequence $\{y_n\}$ also bounded.

From the proof of Theorem 3.1, we have $\|x_{n+1} - y_n\| \rightarrow 0$ as $n \rightarrow \infty$. Since X is a real uniformly smooth Banach space, so that the normalized duality mapping J is single valued and uniformly continuous on any bounded subset of X , thus

$$d_n := \|J(x_{n+1} - x^*) - J(y_n - x^*)\| \rightarrow 0 \quad (3.25)$$

as $n \rightarrow \infty$.

Next, following the reasoning in the proof of Theorem 3.1, we deduce the conclusion of Theorem 3.4. \square

Remark 3.5. In view of Example 1.4, the class of Φ -hemicontractive mappings is a proper subset of the class of generalized Φ -hemicontractive mappings. Hence, Theorem 3.4 improves essentially the result of [2, Theorem 2].

As applications, we give the following theorems.

Theorem 3.6. *Let X be a p -uniformly smooth Banach space $T : X \rightarrow 2^X$, a multivalued generalized Φ -hemiaccretive and bounded mapping. For any given $f \in X$, define $S : X \rightarrow 2^X$ by $Sx := x - Tx + f$ for all $x \in X$. For any given $x_0, u_0, v_0 \in X$, let $\{x_n\}$ be the Ishikawa-type iterative sequence with variable coefficients, defined by*

$$\begin{aligned} y_n &= \hat{a}_n x_n + \hat{b}_n \xi_n + \hat{c}_n v_n, & \exists \xi_n \in Sx_n, \\ x_{n+1} &= \hat{\alpha}_n x_n + \hat{\beta}_n \eta_n + \hat{\gamma}_n u_n, & \exists \eta_n \in Sy_n, \end{aligned} \quad n \in \mathbb{N}, \quad (3.26)$$

where $\{u_n\}, \{v_n\}$ are bounded sequences in X ,

$$\begin{aligned} \hat{a}_n &= 1 - \hat{b}_n - \hat{c}_n, & \hat{b}_n &= \frac{b_n}{r_n^2}, & \hat{c}_n &= \frac{c_n}{r_n^2}, & r_n &= 2 + \|x_n\| + \|\xi_n\| + \|v_n\|, \\ \hat{\alpha}_n &= 1 - \hat{\beta}_n - \hat{\gamma}_n, & \hat{\beta}_n &= \frac{\beta_n}{R_n^2}, & \hat{\gamma}_n &= \frac{\gamma_n}{R_n^2}, & R_n &= r_n + \|\eta_n\| + \|u_n\|, \end{aligned} \quad (3.27)$$

$\{\beta_n\}, \{\gamma_n\}, \{b_n\}$, and $\{c_n\}$ are four sequences in $[0, 1]$ satisfying the following conditions:

$$\sum_{n=0}^{\infty} \beta_n = \infty, \quad \sum_{n=0}^{\infty} \beta_n^p < \infty, \quad \sum_{n=0}^{\infty} \gamma_n < \infty, \quad b_n \leq O(\beta_n), \quad c_n \leq O(\beta_n). \quad (3.28)$$

Then, $\{x_n\}$ converges strongly to the unique solution of the generalized Φ -hemiaccretive mapping equation $f \in Tx$.

Theorem 3.7. Let X be a uniformly smooth Banach space and $T : X \rightarrow 2^X$ a generalized Φ -hemiccretive with bounded range. For any given $f \in X$, define $S : X \rightarrow 2^X$ by $Sx := x - Tx + f$ for all $x \in X$. For any given $x_0, u_0, v_0 \in X$, let $\{x_n\}$ be the Ishikawa-type iterative sequence with variable coefficients, defined by

$$\begin{aligned} y_n &= \widehat{a}_n x_n + \widehat{b}_n \xi_n + \widehat{c}_n v_n, & \exists \xi_n \in Sx_n, \\ x_{n+1} &= \widehat{\alpha}_n x_n + \widehat{\beta}_n \eta_n + \widehat{\gamma}_n u_n, & \exists \eta_n \in Sy_n, \end{aligned} \quad n = 0, 1, 2, \dots, \quad (3.29)$$

where $\{u_n\}, \{v_n\}$ are bounded sequences in X ,

$$\begin{aligned} \widehat{a}_n &= 1 - \widehat{b}_n - \widehat{c}_n, & \widehat{b}_n &= \frac{b_n}{r_n^2}, & \widehat{c}_n &= \frac{c_n}{r_n^2}, & r_n &= 2 + \|x_n\| + \|\xi_n\| + \|v_n\|, \\ \widehat{\alpha}_n &= 1 - \widehat{\beta}_n - \widehat{\gamma}_n, & \widehat{\beta}_n &= \frac{\beta_n}{R_n^2}, & \widehat{\gamma}_n &= \frac{\gamma_n}{R_n^2}, & R_n &= r_n + \|\eta_n\| + \|u_n\|, \end{aligned} \quad (3.30)$$

$\{\beta_n\}, \{\gamma_n\}, \{b_n\}$ and $\{c_n\}$ are four sequences in $[0, 1]$ satisfying the following conditions:

$$\sum_{n=0}^{\infty} \beta_n = \infty, \quad \sum_{n=0}^{\infty} \beta_n^2 < \infty, \quad \sum_{n=0}^{\infty} \gamma_n < \infty, \quad b_n \leq O(\beta_n), \quad c_n \leq O(\beta_n). \quad (3.31)$$

Then, $\{x_n\}$ converges strongly to the unique solution of the generalized Φ -hemiccretive mapping equation $f \in Tx$.

Remark 3.8. (1) Theorem 3.6 improves some recent results, for example, [1, Theorem 3.7] and [2, Theorem 2] in p -uniformly smooth real Banach spaces since the multivalued generalized Φ -hemiccretive mapping within the equation has no generalized Lipschitz assumption.

(2) In view of Example 1.4, the class of Φ -hemiccontractive mappings is a proper subset of the class of generalized Φ -hemiccontractive mappings. Hence, Theorem 3.7 improves essentially the result of [2, Theorem 2] in uniformly smooth real Banach spaces.

Acknowledgment

The work was partially supported by the Specialized Research Fund 2010 for the Doctoral Program of Anhui University of Architecture and the Natural Science Foundation of Anhui Educational Committee.

References

- [1] C. E. Chidume and C. O. Chidume, "Convergence theorem for zeros of generalized Lipschitz generalized ϕ -quasi-accretive operators," *Proceedings of the American Mathematical Society*, vol. 134, no. 1, pp. 243–251, 2006.
- [2] N. Hirano and Z. Huang, "Convergence theorems for multivalued Φ -hemiccontractive operators and Φ -strongly accretive operators," *Computers & Mathematics with Applications*, vol. 46, no. 10-11, pp. 1461–1471, 2003.

- [3] Ya. I. Alber, "On the stability of iterative approximations to fixed points of nonexpansive mappings," *Journal of Mathematical Analysis and Applications*, vol. 328, no. 2, pp. 958–971, 2007.
- [4] C. E. Chidume and C. Moore, "The solution by iteration of nonlinear equations in uniformly smooth Banach spaces," *Journal of Mathematical Analysis and Applications*, vol. 215, no. 1, pp. 132–146, 1997.
- [5] C.-S. Ge and J. Liang, "Convergence theorems of new Ishikawa iterative procedures with errors for multi-valued Φ -hemicontractive mappings," *Communications in Mathematical Analysis*, vol. 7, no. 1, pp. 12–20, 2009.
- [6] C.-S. Ge, J. Liang, and T.-J. Xiao, "Iterative algorithms with variable coefficients for asymptotically strict pseudocontractions," *Fixed Point Theory and Applications*, vol. 2010, Article ID 948529, 8 pages, 2010.
- [7] N.-J. Huang, C.-J. Gao, and X.-P. Huang, "New iteration procedures with errors for multivalued φ -strongly pseudocontractive and φ -strongly accretive mappings," *Computers & Mathematics with Applications*, vol. 43, no. 10-11, pp. 1381–1390, 2002.
- [8] M. O. Osilike and S. C. Aniagbosor, "Weak and strong convergence theorems for fixed points of asymptotically nonexpansive mappings," *Mathematical and Computer Modelling*, vol. 32, no. 10, pp. 1181–1191, 2000.
- [9] L.-C. Zeng, T. Tanaka, and J.-C. Yao, "Iterative construction of fixed points of nonself-mappings in Banach spaces," *Journal of Computational and Applied Mathematics*, vol. 206, no. 2, pp. 814–825, 2007.