

## GHIZZETTI'S THEOREM FOR PIECEWISE CONTINUOUS SOLUTIONS

MANUEL PINTO

Departamento de Matemáticas  
 Facultad de Ciencias, Universidad de Chile  
 Casilla 653, Santiago - Chile

(Received March 30, 1992)

**ABSTRACT.** We obtain that certain second order differential equations have discontinuous solutions which behaves asymptotically as straight lines.

**KEY WORDS AND PHRASES.** Fixed impulse times, discontinuous solutions.

**1991 AMS SUBJECT CLASSIFICATION CODES.** 34A37, 34G20.

### 1. INTRODUCTION.

Many applied problems in physics, biology, economics, etc., are submitted to noncontinuous perturbations. Biological systems such as heart beats, blood flows, pulse frequency modulated systems and models for biological neural nets exhibit an impulsive behavior. This justify the increasing interest on the differential equations with impulse action.

We wish to study the second order impulsive differential equation

$$y''(t) = f(t, y(t), y'(t)), \quad t \neq t_i, t \geq a > 0 \quad (P_1)$$

$$\Delta y(t) = g_1(t, y(t), y'(t)), \quad t = t_i \quad (P_2)$$

$$\Delta y'(t) = g_2(t, y(t), y'(t)), \quad t = t_i \quad (P_3)$$

$$y(t_0) = y_0, a \leq t_0 < t_1 \quad (P_4)$$

where

$$\Delta y(t_i) = y(t_i^+) - y(t_i)$$

and

$$\nu = \{t_i\}_{i=1}^{\infty} \subset I = [a, \infty), t_i < t_{i+1} \rightarrow \infty \text{ as } i \rightarrow \infty.$$

We assume the following basic hypotheses

(F)  $f: I \times C^2 \rightarrow C$  a continuous function such that

$$|f(t, y_1, y_2)| \leq \lambda_1(t) |y_1| + \lambda_2(t) |y_2|. \quad (1.1)$$

(G)  $g_i: \nu \times C \rightarrow C (i = 1, 2)$  are two continuous functions such that

$$|g_i(t, y_1, y_2)| \leq \gamma_i(t) (|y_1| + |y_2|) \quad i = 1, 2. \quad (1.2)$$

(I)  $\lambda(s) = s\lambda_1(s) + \lambda_2(s) \in L_1(I), \gamma(t_k) = t_k\gamma_2(t_k) + \gamma_1(t_k) \in \ell_1.$

We will demonstrate that for every  $t_0 \geq a > 0$ , any solution  $y$  of problem (P) is defined on all of  $[t_0, \infty)$  and as  $t \rightarrow \infty$ , it satisfies

$$y(t) = \delta_1 + 0(\Lambda(t)) + (\delta_2 + 0(\Lambda(t)))t \quad (1.3)$$

and

$$y'(t) = \delta_2 + 0(\Lambda(t)), \tag{1.4}$$

where  $\delta_i (i = 1, 2)$  are constants and

$$\Lambda(t) = \int_t^\infty \lambda(s)ds + \sum_{t_k \in (t, \infty)} \gamma(t_k). \tag{1.5}$$

Furthermore, under the stronger condition

$$(I') \quad s^2 \lambda_1(s), s \lambda_2(s) \in L_1(I); t_k^2 \gamma_2(t_k), t_k \gamma_1(t_k) \in \ell_1.$$

the relation

$$y(t) = \delta_1 + \delta_2 t + 0(\Lambda_1(t)), t \rightarrow \infty \tag{1.6}$$

holds, where

$$\Lambda_1(t) = \int_t^\infty s \lambda(s)ds + \sum_{t_k \in (t, \infty)} t_k \gamma(t_k). \tag{1.7}$$

In this way, we extend the classical Ghizzetti's Theorem [2] for ordinary differential equations (see [5] to [9]) to problem (P).

Let  $I_\nu = I - \nu$  and  $C_\nu^+(I) = \{u \in C(I_\nu) / u(t_i^-) = u(t_i)$  and  $u(t_i^+) \text{ exist } i = 1, 2, \dots\}$ . As usual,  $u(t_i^-)$  and  $u(t_i^+)$  denote the left and right limit of  $u(t)$  as  $t \rightarrow t_i$ .

Next, we establish without proof (see [1], [3]) a Gronwall's inequality.

**LEMMA 1.** Let be  $0 \leq u, \lambda \in C_\nu^+([t_0, \infty)), \gamma: \nu \rightarrow [0, \infty)$  and  $c \geq 0$ . The inequality

$$u(t) \leq c + \int_{t_0}^t \lambda(s)u(s)ds + \sum_{(t_0, t)} \gamma(t_k)u(t_k), t \geq t_0$$

implies

$$u(t) \leq c \exp \left[ \int_{t_0}^t \lambda(s)ds + \sum_{(t_0, t)} \gamma(t_k) \right],$$

where

$$\sum_{(t_0, t)} \gamma(t_k) = \sum_{t_k \in (t_0, t)} \gamma(t_k).$$

**2. MAIN RESULTS.**

We will prove the following results:

**THEOREM 1.** Under conditions (F), (G), and (I), for every  $t_0 \geq a > 0$  any solution  $y$  of problem (P) is defined on all of  $[t_0, \infty)$  and it satisfies formulae (1.3) and (1.4).

**THEOREM 2.** Under conditions (F), (G), and (I), there exist nonoscillatory solutions  $y$  of problem (P).

**THEOREM 3.** Under conditions (F), (G), and (I'), the conclusion of Theorem 1 is true and formula (1.6) holds. Moreover, there exists solutions  $y$  of problem (P) satisfying (1.6) with  $\delta_i \neq 0 (i = 1, 2)$ .

**PROOF OF THEOREM 1.** Let

$$y = A(t) + B(t)t \tag{2.1}$$

under the condition

$$A'(t) + B'(t)t = 0 \text{ for } t \neq t_i. \tag{2.2}$$

Then

$$y'(t) = B(t) \tag{2.3}$$

and hence

$$B'(t) = \tilde{f}(t), \text{ for } t \neq t_i, \tag{2.4}$$

where  $\tilde{f}(t) = \tilde{f}(t, y(t), y'(t))$ . Solving equations (2.1)-(2.4) we get for  $t \neq t_i$ .

$$\begin{aligned} A(t) &= A(t_0) - \int_{t_0}^t s \tilde{f}(s)ds \\ B(t) &= B(t_0) + \int_{t_0}^t \tilde{f}(s)ds. \end{aligned} \tag{2.5}$$

For the impulse effect, from

$$\Delta y = \Delta A = \Delta(Bt) = \Delta A + t\Delta B$$

we get

$$\Delta A + t\Delta B = \tilde{g}_1$$

where

$$\Delta B = \tilde{g}_2,$$

So

$$\tilde{g}_i(t) = g_i(t, y(t), y'(t)).$$

$$\begin{cases} \Delta A = \tilde{g}_1(t) - t\tilde{g}_2(t), & t = t_i \\ \Delta B = \tilde{g}_2(t) & t = t_i \end{cases} \quad (2.6)$$

Next, we consider the impulsive integral system (2.5)-(2.6). It is equivalent to the system

$$A(t) = A(t_0) - \int_{t_0}^t s\tilde{f}(s)ds + \sum_{t_k \in (t_0, t)} [g_1(t_k, y(t_k), y'(t_k)) - t_k g_2(t_k, y(t_k), y'(t_k))] \quad (2.7)$$

$$B(t) = B(t_0) + \int_{t_0}^t \tilde{f}(s)ds + \sum_{t_k \in (t_0, t)} g_2(t_k, y(t_k), y'(t_k)) \quad (2.8)$$

From (2.1), (2.3) and the hypotheses with  $v(s) = s^{-1} |A(s)| + |B(s)|$  we get

$$\begin{aligned} |f(s, y(s), y'(s))| &\leq \lambda_1(s) |y(s)| + \lambda_2(s) |y'(s)| \\ &\leq s\lambda_1(s) |s^{-1}A(s) + B(s)| + \lambda_2(s) |B(s)| \leq (s\lambda_1(s) + \lambda_2(s))v(s) \end{aligned}$$

and

$$\begin{aligned} |g_1(t_k, y(t_k), y'(t_k))| &\leq \gamma_1(t_k) \{ |A(t_k) + t_k B(t_k)| + |B(t_k)| \} \leq t_k \gamma_1(t_k) v(t_k), \\ |g_2(t_k, y(t_k), y'(t_k))| &\leq t_k \gamma_2(t_k) v(t_k). \end{aligned} \quad (2.9)$$

Now, from (2.7) and (2.9) we obtain for  $t \geq t_0$

$$\begin{aligned} \frac{|A(t)|}{t} &\leq \frac{|A(t_0)|}{t} + \int_{t_0}^t \frac{s}{t} |\tilde{f}(s)| ds + \sum_{t_0 < t_k < t} \frac{1}{t} |\tilde{g}_1(t_k) - t_k \tilde{g}_2(t_k)| \\ &\leq t_0^{-1} |A(t_0)| + \int_{t_0}^t (s\lambda_1(s) + \lambda_2(s))v(s)ds + \sum_{t_0 < t_k < t} [\gamma_1(t_k) + t_k \gamma_2(t_k)]v(t_k) \end{aligned} \quad (2.10)$$

Similarly, from (2.8) and (2.9) we get

$$|B(t)| \leq |B(t_0)| + \int_{t_0}^t (s\lambda_1(s) + \lambda_2(s))v(s)ds + \sum_{(t_0, t)} t_k \gamma_2(t_k) v(t_k). \quad (2.11)$$

Adding (2.10) to (2.11) we get

$$v(t) \leq v(t_0) + \int_{t_0}^t 2\lambda(s)v(s)ds + \sum_{(t_0, t)} 2\gamma(t_k)v(t_k)$$

where  $\lambda(s) = s\lambda_1(s) + \lambda_2(s)$ ,  $\gamma(s) = s\gamma_2(s) + \gamma_1(s)$ . From Lemma 1, we obtain

$$v(t) \leq v(t_0) \prod_{(t_0, t)} (1 + 2\gamma(t_k)) \cdot \exp(2 \int_{t_0}^t \lambda(s)ds) (t \geq t_0).$$

Since  $\lambda \in L_1, \gamma \in \ell_1(v)$  we get that  $v$  is bounded and hence both  $t^{-1}A(t), B(t)$  are also bounded. From (2.9) we get that  $f(s, y(s), y'(s)) \in L_1([a, \infty))$  and  $g_1(t_k, y(t_k), y'(t_k)), g_2(t_k, y(t_k), y'(t_k)) \in \ell_1$ . Then, from (2.8),  $B$  converges, i.e.,

$$B(t) = \delta_2 + o(1) \text{ as } t \rightarrow \infty$$

and by (2.8)

$$B(t) = \delta_2 + o(\Lambda(t))$$

where

$$\Lambda(t) = \int_t^\infty \lambda(s)ds + \sum_{(t, \infty)} \gamma(t_k).$$

On the other hand, the functions of  $t$

$$\int_{t_0}^t \frac{(t-s)}{t} |\tilde{f}(s)| ds \text{ and } \sum_{(t_0, t)} \frac{(t-t_k)}{t} |\tilde{g}_i(t_k)| \ (i = 1, 2)$$

are increasing and bounded since  $\tilde{f} \in L_1; \tilde{g}_i(t_k) \in \ell_1 \ (i = 1, 2)$ . Then they converge as  $t \rightarrow \infty$ . Therefore the functions

$$\int_{t_0}^t \frac{(t-s)}{t} \tilde{f}(s) ds \text{ and } \sum_{(t_0, t)} \frac{(t-t_k)}{t} \tilde{g}_i(t_k) \ (i = 1, 2)$$

converge as  $t \rightarrow \infty$ . So, by (2.7) and (2.8), we get

$$\begin{aligned} \frac{A(t)}{t} + B(t) &= \frac{A(t_0)}{t} + B(t_0) + \int_{t_0}^t \frac{(t-s)}{t} \tilde{f}(s) ds \\ &+ \sum_{(t_0, t)} \tilde{g}_1(t_k) + \sum_{(t_0, t)} \frac{(t-t_k)}{t} \tilde{g}_2(t_k) \end{aligned}$$

Then  $t^{-1}A(t) + B(t)$  converges at  $t \rightarrow \infty$  and hence  $t^{-1}A(t)$  converges also, i.e.,

$$A(t) = (\delta_1 + o(1))t.$$

**PROOF OF THEOREMS 2 AND 3.** From the proof of Theorem 1, we get that  $t^{-1}y(t) = t^{-1}A(t) + B(t)$  and  $y' = B$  converge as  $t \rightarrow \infty$ . Furthermore

$$\lim_{t \rightarrow \infty} \frac{y(t)}{t} = \lim_{t \rightarrow \infty} y'(t) \tag{2.12}$$

Let  $y'(t_0) = y_1 \neq 0$ . Then there exists  $T$  large enough so that

$$\left| \int_T^\infty \tilde{f}(s) ds + \sum_{(T, \infty)} \tilde{g}_2(t_k) \right| < |y_1| = |B(t_0)|$$

Thus (2.8) implies that  $\lim_{t \rightarrow \infty} B(t) = \lim_{t \rightarrow \infty} y'(t) \neq 0$  and Theorem 2 follows from (2.12).

Finally, Theorem 3 follows at once from

$$\begin{aligned} t(y'(t) - \delta_2) &= t \left( \int_t^\infty \tilde{f}(s) ds + \sum_{(t, \infty)} \tilde{g}_2(t_k) \right) \\ &\leq \int_t^\infty s \tilde{f}(s) ds + \sum_{(t, \infty)} t_k \tilde{g}_2(t_k) \rightarrow 0 \text{ as } t \rightarrow \infty \end{aligned}$$

because of condition (I') implies  $s\tilde{f}(s) \in L_1$  and  $t_k \tilde{g}_2(t_k) \in \ell_1$ . In this case, by (2.7)  $A$  itself converges as  $t \rightarrow \infty$  and in the same way as it was proved previously we can demonstrate that there exist  $A$  so that  $A(\infty) \neq 0$ , i.e., in formula (1.6),  $\delta_1$  can be taken nonzero in this case.

**REFERENCES**

1. BAINOV, D.; LAKSHMIKANTHAM, V. & SIMEONOV, P., *Theory of Impulsive Differential System*, World Scientific, 1989.
2. GHIZZETTI, A., Un teorema sul comportamento asintotico degli integrali delle equazioni differenziali lineari omogenee, *Rend. Mat. e Appl.* **8** (5) (1949), 28-42.
3. PINTO, M., Impulsive integral inequalities of Bihari-Type, *Libertas Math* **XII** (1992), 57-70.
4. PINTO, M., Second order impulsive differential equations, submitted.
5. KONG, Q., Asymptotic behavior of a class of nonlinear differential equations of  $n$ -th order, *Proc. Amer. Math. Soc.* **103** (1988), 831-838.
6. KREITH, K., Extremal solutions for a class of nonlinear differential equations, *Proc. Amer. Math. Soc.* **79** (1980), 415-421.
7. TONG, J., The asymptotic behavior of a class of nonlinear differential equations of second order, *Proc. Amer. Math. Soc.* **84** (1982), 235-236.
8. PINTO, M., Continuation, nonoscillation and asymptotic formulae of solutions of second order nonlinear differential equations, *Nonlinear Analysis T.M.A.* **16** (1991), 981-995.
9. PINTO, M., Integral inequalities of Bihari-Type and applications, *Funck. Ekvac.* **33** (1990), 387-403.