

## Research Article

# Domination Conditions for Families of Quasilinearly Subharmonic Functions

**Juhani Riihenta**

*Department of Mathematical Sciences, University of Oulu, P.O. Box 3000, 90014 Oulun Yliopisto, Finland*

Correspondence should be addressed to Juhani Riihenta, juhani.riihenta@gmail.com

Received 16 February 2011; Accepted 12 April 2011

Academic Editor: Teodor Bulboacă

Copyright © 2011 Juhani Riihenta. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Domar has given a condition that ensures the existence of the largest subharmonic minorant of a given function. Later Rippon pointed out that a modification of Domar's argument gives in fact a better result. Using our previous, rather general and flexible, modification of Domar's original argument, we extend their results both to the subharmonic and quasilinearly subharmonic settings.

## 1. Introduction

### 1.1. Results of Domar and Rippon

Suppose that  $D$  is a domain of  $\mathbb{R}^n$ ,  $n \geq 2$ . Let  $F : D \rightarrow [0, +\infty]$  be an upper semicontinuous function. Let  $\mathcal{F}$  be a family of subharmonic functions  $u : D \rightarrow [0, +\infty)$  which satisfy

$$u(x) \leq F(x), \quad (1.1)$$

for all  $x \in D$ . Write

$$w(x) = \sup_{u \in \mathcal{F}} u(x), \quad x \in D, \quad (1.2)$$

and let  $w^* : D \rightarrow [0, +\infty]$  be the upper semicontinuous regularization of  $w$ , that is,

$$w^*(x) := \limsup_{y \rightarrow x} w(y). \quad (1.3)$$

Domar gave the following result.

**Theorem A.** *If for some  $\varepsilon > 0$ ,*

$$\int_D [\log^+ F(x)]^{n-1+\varepsilon} dm_n(x) < +\infty, \quad (1.4)$$

*then  $w$  is locally bounded above in  $D$ , and thus  $w^*$  is subharmonic in  $D$ .*

See [1, Theorems 1 and 2, pages 430 and 431]. As Domar points out, the original case of subharmonic functions in the result of Theorem 1 is due to Sjöberg [2] and Brelot [3] (cf. also [4]). Observe, however, that Domar also sketches a new proof for Theorem 1 which uses elementary methods and applies to more general functions.

Rippon [5, Theorem 1, page 128] generalized Domar's result in the following form.

**Theorem B.** *Let  $\varphi : [0, +\infty] \rightarrow [0, +\infty]$  be an increasing function such that*

$$\int_1^{+\infty} \frac{dt}{[\varphi(t)]^{1/(n-1)}} < +\infty. \quad (1.5)$$

*If*

$$\int_D \varphi(\log^+ F(x)) dm_n(x) < +\infty, \quad (1.6)$$

*then  $w$  is locally bounded above in  $D$ , and thus  $w^*$  is subharmonic in  $D$ .*

As pointed out by Domar [1, pages 436–440] and by Rippon [5, page 129], the above results are for many particular cases sharp.

As Domar points out, in [1, page 430], the result of his Theorem A holds in fact for more general functions, that is, for functions which by good reasons might be—and indeed already have been—called quasilinearly subharmonic functions. See Section 1.2 below for the definition of this function class. In addition, Domar has given a related result for an even more general function class  $K(A, \alpha)$ , where the above conditions (1.4) and (1.6) are replaced by a certain integrability condition on the decreasing rearrangement of  $\log F$ , see [6, Theorem 1, page 485]. Observe, however, that in the case  $\alpha = n$  Domar's class  $K(A, n)$  equals the class of nonnegative quasilinearly subharmonic functions: if  $u \in K(A, n)$ , then  $u$  is  $\nu_n A^{n+1}$ -quasilinearly subharmonic. Here (and below)  $\nu_n$  is the Lebesgue measure of the unit ball  $B^n(0, 1)$  in  $\mathbb{R}^n$ . Conversely, if  $u \geq 0$  is  $C$ -quasilinearly subharmonic, then  $u \in K(C, n)$ .

Below we give a general and at the same time flexible result which includes both Domar's and Rippon's results, Theorems A and B above. See Theorem 2.1, Corollary 2.4, and Remark 2.5 below. For previous preliminary, more or less standard results, see also [7, Theorem 2(d), page 15], [8, Theorem 2, page 71], and [9, Theorem 2.2(vi), page 55] (see Remark 1.2(v)).

*Notation.* Our notation is rather standard, see, for example, [7, 9]. For the convenience of the reader we, however, recall the following.  $m_n$  is the Lebesgue measure in the Euclidean space

$\mathbb{R}^n$ ,  $n \geq 2$ .  $D$  is always a domain in  $\mathbb{R}^n$ . Constants will be denoted by  $C$  and  $K$ . They are always nonnegative and may vary from line to line.

## 1.2. Subharmonic Functions and Generalizations

We recall that an upper semicontinuous function  $u : D \rightarrow [-\infty, +\infty)$  is *subharmonic* if for all closed balls  $\overline{B^n(x, r)} \subset D$ ,

$$u(x) \leq \frac{1}{v_n r^n} \int_{B^n(x, r)} u(y) dm_n(y). \quad (1.7)$$

The function  $u \equiv -\infty$  is considered subharmonic.

We say that a function  $u : D \rightarrow [-\infty, +\infty)$  is *nearly subharmonic*, if  $u$  is Lebesgue measurable,  $u^+ \in \mathcal{L}_{\text{loc}}^1(D)$ , and for all  $\overline{B^n(x, r)} \subset D$ ,

$$u(x) \leq \frac{1}{v_n r^n} \int_{B^n(x, r)} u(y) dm_n(y). \quad (1.8)$$

Observe that in the standard definition of nearly subharmonic functions one uses the slightly stronger assumption that  $u \in \mathcal{L}_{\text{loc}}^1(D)$ , see, for example, [7, page 14]. However, our above, slightly more general definition seems to be more practical, see, for example [9, Propositions 2.1(iii) and 2.2(vi)-(vii), pages 54 and 55], and also Remark 1.2(i)-(vi) below. The following lemma emphasizes this fact still more.

**Lemma 1.1** (see [9, Lemma, page 52]). *Let  $u : D \rightarrow [-\infty, +\infty)$  be Lebesgue measurable. Then  $u$  is nearly subharmonic (in the sense defined above) if and only if there exists a function  $u^*$ , subharmonic in  $D$  such that  $u^* \geq u$  and  $u^* = u$  almost everywhere in  $D$ . Here  $u^*$  is the upper semicontinuous regularization of  $u$ :*

$$u^*(x) = \limsup_{x' \rightarrow x} u(x'). \quad (1.9)$$

The proof follows at once from [7, proof of Theorem 1, pages 14 and 15], (and referring also to [9, Propositions 2.1(iii) and 2.2(vii), pages 54 and 55]).

We say that a Lebesgue measurable function  $u : D \rightarrow [-\infty, +\infty)$  is *K-quasilinearly subharmonic*, if  $u^+ \in \mathcal{L}_{\text{loc}}^1(D)$  and if there is a constant  $K = K(n, u, D) \geq 1$  such that for all  $\overline{B^n(x, r)} \subset D$ ,

$$u_M(x) \leq \frac{K}{v_n r^n} \int_{B^n(x, r)} u_M(y) dm_n(y), \quad (1.10)$$

for all  $M \geq 0$ , where  $u_M := \max\{u, -M\} + M$ . A function  $u : D \rightarrow [-\infty, +\infty)$  is *quasilinearly subharmonic*, if  $u$  is  $K$ -quasilinearly subharmonic for some  $K \geq 1$ .

A Lebesgue measurable function  $u : D \rightarrow [-\infty, +\infty)$  is  $K$ -quasilinearly subharmonic n.s. (in the narrow sense), if  $u^+ \in \mathcal{L}_{\text{loc}}^1(D)$  and if there is a constant  $K = K(n, u, D) \geq 1$  such that for all  $\overline{B^n(x, r)} \subset D$ ,

$$u(x) \leq \frac{K}{v_n r^n} \int_{B^n(x, r)} u(y) dm_n(y). \quad (1.11)$$

A function  $u : D \rightarrow [-\infty, +\infty)$  is quasilinearly subharmonic n.s., if  $u$  is  $K$ -quasilinearly subharmonic n.s. for some  $K \geq 1$ .

As already pointed out, Domar [1, 6] considered nonnegative quasilinearly subharmonic functions. Later on, quasilinearly subharmonic functions (perhaps with a different terminology, and sometimes in certain special cases, or just the corresponding generalized mean value inequality (1.10) or (1.11)) have been considered in many papers, see, for example, [8–13] and the references therein.

We recall here only that this function class includes, among others, subharmonic functions, and, more generally, quasisubharmonic and nearly subharmonic functions (see, e.g., [7, pages 14 and 26]), also functions satisfying certain natural growth conditions, especially certain eigenfunctions, and polyharmonic functions. Also, the class of Harnack functions is included, thus, among others, nonnegative harmonic functions as well as nonnegative solutions of some elliptic equations. In particular, the partial differential equations associated with quasiregular mappings belong to this family of elliptic equations.

*Remark 1.2.* For the sake of convenience of the reader we recall the following, see [9, Propositions 2.1 and 2.2, pages 54 and 55].

- (i) A  $K$ -quasilinearly subharmonic function n.s. is  $K$ -quasilinearly subharmonic, but not necessarily conversely.
- (ii) A nonnegative Lebesgue measurable function is  $K$ -quasilinearly subharmonic if and only if it is  $K$ -quasilinearly subharmonic n.s.
- (iii) A Lebesgue measurable function is 1-quasilinearly subharmonic if and only if it is 1-quasilinearly subharmonic n.s. and if and only if it is nearly subharmonic (in the sense defined above).
- (iv) If  $u : D \rightarrow [-\infty, +\infty)$  is  $K_1$ -quasilinearly subharmonic and  $v : D \rightarrow [-\infty, +\infty)$  is  $K_2$ -quasilinearly subharmonic, then  $\max\{u, v\}$  is  $\max\{K_1, K_2\}$ -quasilinearly subharmonic in  $D$ . Especially,  $u^+ := \max\{u, 0\}$  is  $K_1$ -quasilinearly subharmonic in  $D$ .
- (v) Let  $\mathcal{F}$  be a family of  $K$ -quasilinearly subharmonic (resp.,  $K$ -quasilinearly subharmonic n.s.) functions in  $D$  and let  $w := \sup_{u \in \mathcal{F}} u$ . If  $w$  is Lebesgue measurable and  $w^+ \in \mathcal{L}_{\text{loc}}^1(D)$ , then  $w$  is  $K$ -quasilinearly subharmonic (resp.,  $K$ -quasilinearly subharmonic n.s.) in  $D$ .
- (vi) If  $u : D \rightarrow [-\infty, +\infty)$  is quasilinearly subharmonic n.s., then either  $u \equiv -\infty$  or  $u$  is finite almost everywhere in  $D$ , and  $u \in \mathcal{L}_{\text{loc}}^1(D)$ .

## 2. The Result

**Theorem 2.1.** Let  $K \geq 1$ . Let  $\varphi : [0, +\infty] \rightarrow [0, +\infty]$  and let  $\psi : [0, +\infty] \rightarrow [0, +\infty]$  be increasing functions for which there are  $s_0, s_1 \in \mathbb{N}$ ,  $s_0 < s_1$ , such that

- (i) the inverse functions  $\varphi^{-1}$  and  $\psi^{-1}$  are defined on  $[\min\{\varphi(s_1 - s_0), \psi(s_1 - s_0)\}, +\infty]$ ,  
(ii)  $2K(\psi^{-1} \circ \varphi)(s - s_0) \leq (\psi^{-1} \circ \varphi)(s)$  for all  $s \geq s_1$ ,  
(iii) the function

$$[s_1 + 1, +\infty] \ni s \mapsto \frac{(\psi^{-1} \circ \varphi)(s + 1)}{(\psi^{-1} \circ \varphi)(s)} \in \mathbb{R} \quad (2.1)$$

is bounded,

- (iv) the following integral is convergent:

$$\int_{s_1}^{+\infty} \frac{ds}{\varphi(s - s_0)^{1/(n-1)}} < +\infty. \quad (2.2)$$

Let  $\mathcal{F}_K$  be a family of  $K$ -quasilinearly subharmonic functions  $u : D \rightarrow [-\infty, +\infty)$  such that

$$u(x) \leq F_K(x), \quad (2.3)$$

for all  $x \in D$ , where  $F_K : D \rightarrow [0, +\infty]$  is a Lebesgue measurable function. If for each compact set  $E \subset D$ ,

$$\int_E \varphi(F_K(x)) dm_n(x) < +\infty, \quad (2.4)$$

then the family  $\mathcal{F}_K$  is locally (uniformly) bounded in  $D$ . Moreover, the function  $w^* : D \rightarrow [0, +\infty)$  is a  $K$ -quasilinearly subharmonic function. Here

$$w^*(x) := \limsup_{y \rightarrow x} w(y), \quad (2.5)$$

where

$$w(x) := \sup_{u \in \mathcal{F}_K} u^+(x). \quad (2.6)$$

The proof of the theorem will be based on the following lemma, which has its origin in [1, Lemma 1, pages 431 and 432], see also [14, Proposition 2, pages 257–259]. Observe that we have applied our rather general and flexible lemma already before (unlike previously, now we allow also the value  $+\infty$  for our “test functions”  $\varphi$  and  $\psi$ ; this does not, however, cause any changes in the proof of our lemma, see [15, pages 5–8]) when considering quasilinearly subharmonicity of separately quasilinearly subharmonic functions. As a matter of fact, this lemma enabled us to slightly improve Armitage’s and Gardiner’s almost sharp condition, see [14, Theorem 1, page 256], which ensures a separately subharmonic function to be subharmonic. See [15, Corollary 4.5, page 13], and [12, 13, Corollary 3.3.3, page 2622].

**Lemma 2.2** (see [15, Lemma 3.2, page 5 and Remark 3.3, page 8]). Let  $K, \varphi, \psi$  and  $s_0, s_1 \in \mathbb{N}$  be as in Theorem 2.1. Let  $u : D \rightarrow [0, +\infty)$  be a  $K$ -quasilinearly subharmonic function. Let  $\tilde{s}_1 \in \mathbb{N}$ ,  $\tilde{s}_1 \geq s_3$ , be arbitrary, where  $s_3 := \max\{s_1 + 3, (\psi^{-1} \circ \varphi)(s_1 + 3)\}$ . Then for each  $x \in D$  and  $r > 0$  such that  $\overline{B^n(x, r)} \subset D$  either

$$u(x) \leq (\psi^{-1} \circ \varphi)(\tilde{s}_1 + 1) \quad (2.7)$$

or

$$\Phi(u(x)) \leq \frac{C}{r^n} \int_{B^n(x, r)} \psi(u(y)) dm_n(y), \quad (2.8)$$

where  $C = C(n, K, s_0)$  and  $\Phi : [0, +\infty) \rightarrow [0, +\infty)$ ,

$$\Phi(t) := \begin{cases} \left[ \int_{(\varphi^{-1} \circ \psi)(t)-2}^{+\infty} \frac{ds}{\varphi(s-s_0)^{1/(n-1)}} \right]^{1-n}, & \text{when } t \geq s_3, \\ \frac{t}{s_3} \Phi(s_3), & \text{when } 0 \leq t < s_3. \end{cases} \quad (2.9)$$

*Proof of Theorem 2.1.* Let  $E$  be an arbitrary compact subset of  $D$ . Write  $\rho_0 := \text{dist}(E, \partial D)$ . Clearly  $\rho_0 > 0$ . Write

$$E_1 := \bigcup_{x \in E} \overline{B^n\left(x, \frac{\rho_0}{2}\right)}. \quad (2.10)$$

Then  $E_1$  is compact, and  $E \subset E_1 \subset D$ . Take  $u \in \mathcal{F}_K^+$  arbitrarily, where

$$\mathcal{F}_K^+ := \{u^+ : u \in \mathcal{F}_K\}. \quad (2.11)$$

Let  $\tilde{s}_1 = s_1 + 2$ , say. Take  $x \in E$  arbitrarily and suppose that  $u(x) > \tilde{s}_3$ , where  $\tilde{s}_3 := \max\{\tilde{s}_1 + 3, (\psi^{-1} \circ \varphi)(\tilde{s}_1 + 3)\}$ , say. Using our lemma and the assumption, we get

$$\begin{aligned} \left( \int_{(\varphi^{-1} \circ \psi)(u(x))-2}^{+\infty} \frac{ds}{\varphi(s-s_0)^{1/(n-1)}} \right)^{1-n} &\leq \frac{C}{(\rho_0/2)^n} \int_{B^n(x, \rho_0/2)} \psi(u(y)) dm_n(y) \\ &\leq \frac{C}{(\rho_0/2)^n} \int_{E_1} \psi(F_K(y)) dm_n(y) < +\infty. \end{aligned} \quad (2.12)$$

Since

$$\int_{s_1}^{+\infty} \frac{ds}{\varphi(s-s_0)^{1/(n-1)}} < +\infty, \quad (2.13)$$

and  $1 - n < 0$ , the set of values

$$(\varphi^{-1} \circ \varphi)(u(x)) - 2, \quad x \in E, u \in \mathcal{F}_K^+, \tag{2.14}$$

is bounded. Thus also the set of values

$$u(x), \quad x \in E, u \in \mathcal{F}_K^+, \tag{2.15}$$

is bounded.

To show that  $w^*$  is  $K$ -quasilinearly subharmonic in  $D$ , proceed as follows. Take  $x \in D$  and  $r > 0$  such that  $\overline{B^n(x, r)} \subset D$ . For each  $u \in \mathcal{F}_K^+$  we have then

$$u(x) \leq \frac{K}{v_n r^n} \int_{B^n(x, r)} u(y) dm_n(y). \tag{2.16}$$

Since

$$u(x) \leq \sup_{u \in \mathcal{F}_K^+} u(x) = w(x) \leq w^*(x), \tag{2.17}$$

we have

$$w(x) \leq \frac{K}{v_n r^n} \int_{B^n(x, r)} w^*(y) dm_n(y). \tag{2.18}$$

Then just take the upper semicontinuous regularizations on both sides of (2.18) and use Fatou's lemma on the right-hand side (this is of course possible, since  $w^*$  is locally bounded in  $D$ ), say

$$\begin{aligned} \limsup_{y \rightarrow x} w(y) &\leq \limsup_{y \rightarrow x} \frac{K}{v_n r^n} \int_{B^n(y, r)} w^*(z) dm_n(z) \\ &\leq \limsup_{y \rightarrow x} \frac{K}{v_n r^n} \int w^*(z) \chi_{\overline{B^n(y, r)}}(z) dm_n(z) \\ &\leq \frac{K}{v_n r^n} \int w^*(z) \left( \limsup_{y \rightarrow x} \chi_{\overline{B^n(y, r)}}(z) \right) dm_n(z). \end{aligned} \tag{2.19}$$

Since for all  $z \in D$ ,

$$\limsup_{y \rightarrow x} \chi_{\overline{B^n(y, r)}}(z) \leq \chi_{\overline{B^n(x, r)}}(z), \tag{2.20}$$

we get the desired inequality

$$\omega^*(x) \leq \frac{K}{\nu_n r^n} \int_{B^n(x,r)} \omega^*(y) dm_n(y). \quad (2.21)$$

□

*Remark 2.3.* If  $\omega$  is Lebesgue measurable, it follows that already  $\omega$  is  $K$ -quasilinearly subharmonic.

**Corollary 2.4.** Let  $\varphi : [0, +\infty] \rightarrow [0, +\infty]$  be a strictly increasing function such that for some  $s_0, s_1 \in \mathbb{N}$ ,  $s_0 < s_1$ ,

$$\int_{s_1}^{+\infty} \frac{ds}{[\varphi(s - s_0)]^{1/(n-1)}} < +\infty. \quad (2.22)$$

Let  $\mathcal{F}_K$  be a family of  $K$ -quasilinearly subharmonic functions  $u : D \rightarrow [-\infty, +\infty)$  such that

$$u(x) \leq F_K(x), \quad (2.23)$$

for all  $x \in D$ , where  $F_K : D \rightarrow [0, +\infty]$  is a Lebesgue measurable function.

Let  $p > 0$  be arbitrary. If for each compact set  $E \subset D$ ,

$$\int_E \varphi(\log^+[F(x)]^p) dm_n(x) < +\infty, \quad (2.24)$$

then the family  $\mathcal{F}_K$  is locally (uniformly) bounded in  $D$ . Moreover, the function  $\omega^* : D \rightarrow [0, +\infty)$  is a  $K$ -quasilinearly subharmonic function. Here

$$\omega^*(x) := \limsup_{y \rightarrow x} \omega(y), \quad (2.25)$$

where

$$\omega(x) := \sup_{u \in \mathcal{F}_K} u^+(x). \quad (2.26)$$

The case  $p = 1$  and  $K = 1$  gives Domar's and Rippon's results, Theorems A and B above. For the proof, take  $p > 0$  arbitrarily, choose  $\psi(t) = (\varphi \circ \log^+)(t^p)$ , and just check that the conditions (i)–(iv) indeed hold.


*Remark 2.5.* As already pointed out, Theorem 2.1 is indeed flexible. To get another simple, but still slightly more general corollary, just choose, say,  $\psi(t) = (\varphi \circ \log^+)(\phi(t))$ , where  $\phi : [0, +\infty] \rightarrow [0, +\infty]$  is any strictly increasing function which satisfies the following two conditions:

- (a)  $\phi^{-1}$  satisfies the  $\Delta_2$ -condition,
- (b)  $2K\phi^{-1}(e^{s-s_0}) \leq \phi^{-1}(e^s)$  for all  $s \geq s_1$ .



## References

- [1] Y. Domar, "On the existence of a largest subharmonic minorant of a given function," *Arkiv för Matematik*, vol. 3, no. 39, pp. 429–440, 1957.
- [2] N. Sjöberg, "Sur les minorantes sousharmoniques d'une fonction donnée," in *Neuvième Congrès des mathématiciens Scandinaves*, pp. 309–319, Helsingfors, 1938.
- [3] M. Brelot, "Minorantes sous-harmoniques, extrémales et capacités," *Journal de Mathématiques Pures et Appliquées. Neuvième Série*, vol. 24, pp. 1–32, 1945.
- [4] J. W. Green, "Approximately subharmonic functions," *Proceedings of the American Mathematical Society*, vol. 3, pp. 829–833, 1952.
- [5] P. J. Rippon, "Some remarks on largest subharmonic minorants," *Mathematica Scandinavica*, vol. 49, no. 1, pp. 128–132, 1981.
- [6] Y. Domar, "Uniform boundedness in families related to subharmonic functions," *Journal of the London Mathematical Society. Second Series*, vol. 38, no. 3, pp. 485–491, 1988.
- [7] M. Hervé, *Analytic and Plurisubharmonic Functions in Finite and Infinite Dimensional Spaces*, Lecture Notes in Mathematics, Vol. 198, Springer, Berlin, Germany, 1971.
- [8] J. Riihenta, "On a theorem of Avanissian-Arsove," *Expositiones Mathematicae*, vol. 7, no. 1, pp. 69–72, 1989.
- [9] J. Riihenta, "Subharmonic functions, generalizations and separately subharmonic functions," in *XIV Conference on Analytic Functions*, vol. 2 of *Scientific Bulletin of Chełm, Section of Mathematics and Computer Science*, pp. 49–76, Chełm, Poland, 2007.
- [10] M. Pavlović, "On subharmonic behaviour and oscillation of functions on balls in  $\mathbb{R}^n$ ," *Institut Mathématique. Publications. Nouvelle Série*, vol. 55(69), pp. 18–22, 1994.
- [11] M. Pavlović and J. Riihenta, "Classes of quasi-nearly subharmonic functions," *Potential Analysis*, vol. 29, no. 1, pp. 89–104, 2008.
- [12] J. Riihenta, "Subharmonic functions, generalizations, weighted boundary behavior, and separately subharmonic functions: A survey," in *5th World Congress of Nonlinear Analysts (WCNA '08)*, Orlando, Fla, USA, 2008.
- [13] J. Riihenta, "Subharmonic functions, generalizations, weighted boundary behavior, and separately subharmonic functions: A survey," *Nonlinear Analysis, Series A: Theory, Methods & Applications*, vol. 71, no. 12, pp. e2613–e26267, 2009.
- [14] D. H. Armitage and S. J. Gardiner, "Conditions for separately subharmonic functions to be subharmonic," *Potential Analysis*, vol. 2, no. 3, pp. 255–261, 1993.
- [15] J. Riihenta, "Quasi-nearly subharmonicity and separately quasi-nearly subharmonic functions," *Journal of Inequalities and Applications*, vol. 2008, Article ID 149712, 15 pages, 2008.



# Hindawi

Submit your manuscripts at  
<http://www.hindawi.com>

