

# BOUNDARY VALUE PROBLEM WITH INTEGRAL CONDITIONS FOR A LINEAR THIRD-ORDER EQUATION

M. DENCHE AND A. MEMOU

*Received 6 March 2003 and in revised form 29 July 2003*

We prove the existence and uniqueness of a strong solution for a linear third-order equation with integral boundary conditions. The proof uses energy inequalities and the density of the range of the generated operator.

## 1. Introduction

In the rectangle  $\Omega = [0, 1] \times [0, T]$ , we consider the equation

$$\mathcal{L}u = \frac{\partial^3 u}{\partial t^3} + \frac{\partial}{\partial x} \left( a(x, t) \frac{\partial u}{\partial x} \right) = f(x, t), \quad (1.1a)$$

with the initial conditions

$$u(x, 0) = 0, \quad \frac{\partial u}{\partial t}(x, 0) = 0, \quad x \in (0, 1), \quad (1.1b)$$

the final condition

$$\frac{\partial^2 u}{\partial t^2}(x, T) = 0, \quad x \in (0, 1), \quad (1.1c)$$

the Dirichlet condition

$$u(0, t) = 0, \quad \forall t \in (0, T), \quad (1.1d)$$

and the integral condition

$$\int_0^1 u(x,t) dx = 0, \quad \forall t \in (0,T). \quad (1.1e)$$

In addition, we assume that the function  $a(x,t)$  is bounded with

$$0 < a_0 \leq a(x,t) \leq a_1, \quad (1.2)$$

and has bounded partial derivatives such that

$$c'_k \leq \frac{\partial^k a}{\partial t^k}(x,t) \leq c_k, \quad \forall x \in (0,1), t \in (0,T), k = \overline{1,3}, \text{ with } c'_1 \geq 0, \quad (1.3)$$

$$\left| \frac{\partial a}{\partial x}(x,t) \right| \leq b_1, \quad \text{for } (x,t) \in \Omega.$$

Various problems arising in heat conduction [4, 6, 14, 15], chemical engineering [9], underground water flow [13], thermoelasticity [21], and plasmaphysics [19] can be reduced to the nonlocal problems with integral boundary conditions. This type of boundary value problems has been investigated in [1, 2, 3, 5, 6, 7, 9, 14, 15, 16, 20, 23] for parabolic equations, in [18, 22] for hyperbolic equations, and in [10, 11, 12] for mixed-type equations. The basic tool in [4, 10, 11, 12, 16, 23] is the energy inequality method which, of course, requires appropriate multipliers and functional spaces. In this paper, we extend this method to the study of a linear third-order partial differential equation. This type of problems is encountered in the study of thermal conductivity [17] and microscale heat transfer [8].

## 2. Preliminaries

In this paper, we prove the existence and uniqueness of a strong solution of problem (1.1). For this, we consider the solution of problem (1.1) as a solution of the operator equation  $Lu = \mathcal{F}$ , where  $L$  is the operator with domain of definition  $D(L)$  consisting of functions  $u \in E$  such that  $\sqrt{1-x}(\partial^{k+1}u/\partial t^k \partial x)(x,t) \in L^2(\Omega)$ ,  $k = \overline{0,3}$  and  $u$  satisfies conditions (1.1d) and (1.1e). The operator  $L$  is considered from  $E$  to  $F$ , where  $E$  is the Banach space of the functions  $u$ ,  $u \in L^2(\Omega)$ , with the finite norm

$$\|u\|_E^2 = \int_{\Omega} \frac{(1-x)^2}{2} \left\{ \left| \frac{\partial^3 u}{\partial t^3} \right|^2 + \left| \frac{\partial^2 u}{\partial x^2} \right|^2 \right\} dx dt \quad (2.1)$$

$$+ \int_{\Omega} \left( \frac{(1-x)^2}{2} \left| \frac{\partial u}{\partial x} \right|^2 + |u|^2 \right) dx dt,$$

and  $F$  is the Hilbert space of the functions  $\mathcal{F} = (f, 0, 0, 0)$ ,  $f \in L^2(\Omega)$ , with the finite norm

$$\|\mathcal{F}\|_F^2 = \int_{\Omega} (1-x)^2 |f|^2 dx dt. \tag{2.2}$$

Then we establish an energy inequality

$$\|u\|_E \leq k \|Lu\|_F, \quad \forall u \in D(L), \tag{2.3}$$

and we show that the operator  $L$  has the closure  $\bar{L}$ .

*Definition 2.1.* A solution of the operator equation  $\bar{L}u = \mathcal{F}$  is called a strong solution of problem (1.1).

Inequality (2.3) can be extended to  $u \in D(\bar{L})$ , that is,

$$\|u\|_E \leq k \|\bar{L}u\|_F, \quad \forall u \in D(\bar{L}). \tag{2.4}$$

From this inequality, we obtain the uniqueness of a strong solution, if it exists, and the equality of the sets  $R(\bar{L})$  and  $\overline{R(L)}$ . Thus, to prove the existence of a strong solution of problem (1.1) for any  $\mathcal{F} \in F$ , it remains to prove that the set  $R(L)$  is dense in  $F$ .

### 3. An energy inequality and its applications

**THEOREM 3.1.** *For any function  $u \in D(L)$ , there exists the a priori estimate*

$$\|u\|_E \leq k \|Lu\|_F, \tag{3.1}$$

where

$$k^2 = \frac{17 \exp(ct) [5 + 4(b_1)^2 / (c'_3 - 3cc_2 + 3c^2c'_1 - c^3a_1 - b_1^2)] + 1}{\min(1, a_0^2, c'_3 - 3cc_2 + 3c^2c'_1 - c^3a_1 - b_1^2)}, \tag{3.2}$$

with the constant  $c$  satisfying

$$\begin{aligned} \sup_{(x,t) \in \Omega} \left( \frac{1}{a} \frac{\partial a}{\partial t} \right) &\leq c < \inf_{(x,t) \in \Omega} \left( \frac{1}{a} \frac{\partial a}{\partial t} + 1 \right), \\ c'_3 - 3cc_2 + 3c^2c'_1 - c^3a_1 - (b_1)^2 &> 0, \\ c_2 - 2cc'_1 + c^2a_1 - c'_1 + ca_1 &< 0. \end{aligned} \tag{3.3}$$

*Proof.* Let

$$Mu = (1-x)^2 \frac{\partial^3 u}{\partial t^3} + 2(1-x) J_x \frac{\partial^3 u}{\partial t^3}, \quad (3.4)$$

where

$$J_x u = \int_0^x u(\zeta, t) d\zeta. \quad (3.5)$$

We consider the quadratic form

$$\Phi(u, u) = \operatorname{Re} \int_{\Omega} \exp(-ct) \varepsilon u \overline{Mu} dx dt, \quad (3.6)$$

with the constant  $c$  satisfying (3.3), obtained by multiplying (1.1a) by  $\exp(-ct) \overline{Mu}$ , integrating over  $\Omega$ , and taking the real part. Substituting the expression of  $Mu$  in (3.6), we obtain

$$\begin{aligned} & \operatorname{Re} \int_{\Omega} \exp(-ct) \varepsilon u \overline{Mu} dx dt \\ &= \operatorname{Re} \int_{\Omega} \exp(-ct) (1-x)^2 \left| \frac{\partial^3 u}{\partial t^3} \right|^2 dx dt \\ &+ 2 \operatorname{Re} \int_{\Omega} \exp(-ct) (1-x) \frac{\partial^3 u}{\partial t^3} J_x \frac{\partial^3 \overline{u}}{\partial t^3} dx dt \\ &+ \operatorname{Re} \int_{\Omega} \exp(-ct) \frac{\partial}{\partial x} \left( a(x, t) \frac{\partial u}{\partial x} \right) \overline{Mu} dx dt. \end{aligned} \quad (3.7)$$

Integrating the last two terms on the right-hand side by parts with respect to  $x$  in (3.7) and using the Dirichlet condition (1.1d), we obtain

$$2 \operatorname{Re} \int_0^1 (1-x) \exp(-ct) \frac{\partial^3 u}{\partial t^3} J_x \frac{\partial^3 \overline{u}}{\partial t^3} dx = \int_0^1 \exp(-ct) \left| J_x \frac{\partial^3 u}{\partial t^3} \right|^2 dx, \quad (3.8)$$

$$\begin{aligned} & \operatorname{Re} \int_{\Omega} \exp(-ct) \frac{\partial}{\partial x} \left( a \frac{\partial u}{\partial x} \right) \overline{Mu} dx dt \\ &= - \operatorname{Re} \int_{\Omega} \exp(-ct) (1-x)^2 a \frac{\partial u}{\partial x} \frac{\partial^4 \overline{u}}{\partial t^3 \partial x} dx dt \\ &- 2 \operatorname{Re} \int_{\Omega} \exp(-ct) \frac{\partial a}{\partial x} u J_x \frac{\partial^3 \overline{u}}{\partial t^3} dx dt \\ &- 2 \operatorname{Re} \int_{\Omega} \exp(-ct) a u \frac{\partial^3 \overline{u}}{\partial t^3} dx dt. \end{aligned} \quad (3.9)$$

Integrating each term by parts in (3.9) with respect to  $t$  and using the initial and final conditions (1.1b) and (1.1c), we get

$$\begin{aligned}
 & \operatorname{Re} \int_{\Omega} \exp(-ct) \frac{\partial}{\partial x} \left( a \frac{\partial u}{\partial x} \right) \overline{Mu} \, dx \, dt \\
 &= -2 \operatorname{Re} \int_{\Omega} \exp(-ct) \frac{\partial a}{\partial x} u J_x \frac{\partial^3 \bar{u}}{\partial t^3} \, dx \, dt \\
 &+ \int_{\Omega} \exp(-ct) \left( \frac{\partial^3 a}{\partial t^3} - 3c \frac{\partial^2 a}{\partial t^2} + 3c^2 \frac{\partial a}{\partial t} - c^3 a \right) \\
 &\quad \times \left[ \frac{(1-x)^2}{2} \left| \frac{\partial u}{\partial x} \right|^2 + |u|^2 \right] \, dx \, dt \\
 &- 3 \int_{\Omega} \exp(-ct) \left( \frac{\partial a}{\partial t} - ca \right) \left[ \frac{(1-x)^2}{2} \left| \frac{\partial^2 u}{\partial t \partial x} \right|^2 + \left| \frac{\partial u}{\partial t} \right|^2 \right] \, dx \, dt \\
 &+ \int_0^1 \exp(-ct) a \left[ \frac{(1-x)^2}{2} \left| \frac{\partial^2 u}{\partial t \partial x} \right|^2 + \left| \frac{\partial u}{\partial t} \right|^2 \right] \, dx \Big|_{T=t} \\
 &- \int_0^1 \exp(-ct) \left( \frac{\partial^2 a}{\partial t^2} - 2c \frac{\partial a}{\partial t} + c^2 a \right) \left[ \frac{(1-x)^2}{2} \left| \frac{\partial u}{\partial x} \right|^2 + |u|^2 \right] \, dx \Big|_{t=T} \\
 &+ \operatorname{Re} \int_0^1 \exp(-ct) \left( \frac{\partial a}{\partial t} - ca \right) \left\{ (1-x)^2 \frac{\partial^2 \bar{u}}{\partial t \partial x} \frac{\partial u}{\partial x} + 2u \frac{\partial \bar{u}}{\partial t} \right\} \Big|_{T=t} \, dx.
 \end{aligned} \tag{3.10}$$

Substituting (3.8) and (3.10) in (3.7) and using conditions (1.2), (1.3), and (3.3), we obtain

$$\begin{aligned}
 & \int_{\Omega} \exp(-ct) (1-x)^2 \left| \frac{\partial^3 u}{\partial t^3} \right|^2 \, dx \, dt \\
 &\quad + \int_{\Omega} \exp(-ct) \{ c'_3 - 3cc_2 + 3c^2 c'_1 - c^3 a_1 - b_1^2 \} \\
 &\quad \times \left[ \frac{(1-x)^2}{2} \left| \frac{\partial u}{\partial x} \right|^2 + |u|^2 \right] \, dx \, dt \\
 &\leq \operatorname{Re} \int_{\Omega} \exp(-ct) \epsilon u M \bar{u} \, dx \, dt.
 \end{aligned} \tag{3.11}$$

Again, substituting the expression of  $Mu$  in (3.11) and using elementary inequality, we get

$$\begin{aligned}
& \int_{\Omega} \exp(-ct) \frac{(1-x)^2}{2} \left| \frac{\partial^3 u}{\partial t^3} \right|^2 dx dt \\
& \quad + \int_{\Omega} \exp(-ct) \{ c'_3 - 3cc_2 + 3c^2c'_1 - c^3a_1 - b_1^2 \} \\
& \quad \quad \times \left[ \frac{(1-x)^2}{2} \left| \frac{\partial u}{\partial x} \right|^2 + |u|^2 \right] dx dt \\
& \leq 17 \int_{\Omega} \exp(-ct) (1-x)^2 |f|^2 dx dt.
\end{aligned} \tag{3.12}$$

By virtue of (1.1a), we have

$$\begin{aligned}
& \int_{\Omega} a_0 \left| \frac{\partial^2 u}{\partial x^2} \right|^2 \frac{(1-x)^2}{2} dx dt \\
& \leq \int_{\Omega} (1-x)^2 |f|^2 dx dt + \int_{\Omega} 2(1-x)^2 \left| \frac{\partial^3 u}{\partial t^3} \right|^2 dx dt \\
& \quad + 4 \int_{\Omega} b_1^2 \left\{ \frac{(1-x)^2}{2} \left| \frac{\partial u}{\partial x} \right|^2 + |u|^2 \right\} dx dt.
\end{aligned} \tag{3.13}$$

This last inequality combined with (3.12) yields

$$\begin{aligned}
& \int_{\Omega} \frac{(1-x)^2}{2} \left| \frac{\partial^3 u}{\partial t^3} \right|^2 dx dt \\
& \quad + \int_{\Omega} (c'_3 - 3cc_2 + 3c^2c'_1 - c^3a_1 - b_1^2) \left\{ \frac{(1-x)^2}{2} \left| \frac{\partial u}{\partial x} \right|^2 + |u|^2 \right\} dx dt \\
& \quad + \int_{\Omega} a_0^2 \frac{(1-x)^2}{2} \left| \frac{\partial^2 u}{\partial x^2} \right|^2 dx dt \\
& \leq \left\{ 17 \exp(cT) \left[ 5 + \frac{4b_1^2}{c'_3 - 3cc_2 + 3c^2c'_1 - c^3a_1 - b_1^2} \right] + 1 \right\} \\
& \quad \times \int_{\Omega} (1-x)^2 |f|^2 dx dt.
\end{aligned} \tag{3.14}$$

Thus, this inequality implies

$$\begin{aligned}
& \int_{\Omega} \frac{(1-x)^2}{2} \left\{ \left| \frac{\partial^3 u}{\partial t^3} \right|^2 + \left| \frac{\partial^2 u}{\partial x^2} \right|^2 \right\} dx dt + \int_{\Omega} \frac{(1-x)^2}{2} \left| \frac{\partial u}{\partial x} \right|^2 + |u|^2 dx dt \\
& \leq k^2 \int_{\Omega} (1-x)^2 |f|^2 dx dt,
\end{aligned} \tag{3.15}$$

where

$$k^2 = \frac{17 \exp(cT) [5 + 4b_1^2 / (c'_3 - 3cc_2 + 3c^2c'_1 - c^3a_1 - b_1^2)] + 1}{\min(1, a_0^2, c'_3 - 3cc_2 + 3c^2c'_1 - c^3a_1 - b_1^2)}. \tag{3.16}$$

Then,

$$\|u\|_E \leq k \|Lu\|_F, \quad \forall u \in D(L). \tag{3.17}$$

Thus, we obtain the desired inequality. □

LEMMA 3.2. *The operator L from E to F admits a closure.*

*Proof.* Suppose that  $(u_n) \in D(L)$  is a sequence such that

$$u_n \longrightarrow 0 \text{ in } E, \quad Lu_n \longrightarrow \mathcal{F} \text{ in } F. \tag{3.18}$$

We need to show that  $\mathcal{F} = 0$ . We introduce the operator

$$\mathcal{E}_0 v = -(1-x)^2 \frac{\partial^3 v}{\partial t^3} + \frac{\partial}{\partial x} \left\{ a(x,t) \frac{\partial}{\partial x} [(1-x)^2 v] \right\}, \tag{3.19}$$

with domain  $D(\mathcal{E}_0)$  consisting of functions  $v \in W_2^{2,3}(\Omega)$  satisfying

$$v|_{t=0} = 0, \quad \frac{\partial v}{\partial t} \Big|_{t=0} = 0, \quad \frac{\partial^2 v}{\partial t^2} \Big|_{t=0} = 0, \quad v|_{x=0} = 0, \quad \frac{\partial v}{\partial x} \Big|_{x=0} = 0. \tag{3.20}$$

We note that  $D(\mathcal{E}_0)$  is dense in the Hilbert space obtained by completing  $L^2(\Omega)$  with respect to the norm

$$\int_{\Omega} (1-x)^2 |v|^2 dx dt = \|v\|^2. \tag{3.21}$$

Since

$$\begin{aligned} \int_{\Omega} (1-x)^2 f \bar{v} dx dt &= \lim_{n \rightarrow +\infty} \int_{\Omega} (1-x)^2 \mathcal{E} u_n \bar{v} dx dt \\ &= \lim_{n \rightarrow +\infty} \int_{\Omega} u_n \mathcal{E}_0 \bar{v} dx dt = 0, \end{aligned} \tag{3.22}$$

for any function  $v \in D(\mathcal{E}_0)$ , it follows that  $f = 0$ . □

**Theorem 3.1** is valid for a strong solution, then we have the inequality

$$\|u\|_E \leq k \|\bar{L}u\|_F, \quad \forall u \in D(\bar{L}). \tag{3.23}$$

Hence we obtain the following corollary.

**COROLLARY 3.3.** *A strong solution of problem (1.1) is unique if it exists, and depends continuously on  $\mathcal{F}$ .*

**COROLLARY 3.4.** *The range  $R(\bar{L})$  of the operator  $\bar{L}$  is closed in  $F$ , and  $R(\bar{L}) = \overline{R(\bar{L})}$ .*

#### 4. Solvability of problem (1.1)

To prove the solvability of problem (1.1), it is sufficient to show that  $R(L)$  is dense in  $F$ . The proof is based on the following lemma.

**LEMMA 4.1.** *Suppose that  $a(x, t)$  and its derivatives  $\partial^4 a / \partial t^3 \partial x$  and  $\partial^2 a / \partial t \partial x$  are bounded. Let  $D_0(L) = \{u \in D(L) : u(x, 0) = 0, (\partial u / \partial t)(x, 0) = 0, (\partial^2 u / \partial t^2)(x, T) = 0\}$ . If, for  $u \in D_0(L)$  and for some functions  $w \in L^2(\Omega)$ ,*

$$\int_{\Omega} (1-x) \varepsilon u \bar{w} \, dx \, dt = 0, \tag{4.1}$$

then  $w = 0$ .

*Proof.* Equality (4.1) can be written as follows:

$$\int_{\Omega} (1-x) \bar{w} \frac{\partial^3 u}{\partial t^3} \, dx \, dt = - \int_{\Omega} \frac{\partial}{\partial x} \left( a(1-x) \frac{\partial u}{\partial x} \right) \left\{ \bar{w} - \int_0^x \frac{\bar{w}}{1-\zeta} \, d\zeta \right\} \, dx \, dt. \tag{4.2}$$

For a given  $w(x, t)$ , we introduce the function  $v(x, t)$  such that

$$v(x, t) = w(x, t) - \int_0^x \frac{w(\zeta, t)}{1-\zeta} \, d\zeta. \tag{4.3}$$

From (4.3), we conclude that  $\int_0^1 v(x, t) \, dx = 0$ , and thus, we have

$$\int_{\Omega} \frac{\partial^3 u}{\partial t^3} N \bar{v} \, dx \, dt = - \int_{\Omega} A(t) u \bar{v} \, dx \, dt, \tag{4.4}$$

where  $A(t)u = (\partial / \partial x)(a(1-x)(\partial u / \partial x))$  and  $Nv = (1-x)v + Jv$ .



Following [23], we introduce the smoothing operators

$$J_\epsilon^{-1} = \left( I - \epsilon \left( \frac{\partial^3}{\partial t^3} \right) \right)^{-1}, \quad (J_\epsilon^{-1})^* = \left( I + \epsilon \left( \frac{\partial^3}{\partial t^3} \right) \right)^{-1}, \quad (4.5)$$

with respect to  $t$ , which provide the solutions of the respective problems

$$\begin{aligned} g_\epsilon - \epsilon \frac{\partial^3 g_\epsilon}{\partial t^3} = g, & \quad g_\epsilon(0) = 0, & \quad \frac{\partial g_\epsilon}{\partial t}(0) = 0, & \quad \frac{\partial^2 g_\epsilon}{\partial t^2}(T) = 0, \\ g_\epsilon^* + \epsilon \frac{\partial^3 g_\epsilon^*}{\partial t^3} = g, & \quad g_\epsilon^*(0) = 0, & \quad \frac{\partial g_\epsilon^*}{\partial t}(T) = 0, & \quad \frac{\partial^2 g_\epsilon^*}{\partial t^2}(T) = 0. \end{aligned} \quad (4.6)$$

We also have the following properties: for any  $g \in L^2(0, T)$ , the functions  $J_\epsilon^{-1}(g)$ ,  $(J_\epsilon^{-1})^*g \in W_2^3(0, T)$ . If  $g \in D(L)$ , then  $J_\epsilon^{-1}(g) \in D(L)$  and we have

$$\begin{aligned} \lim \| (J_\epsilon^{-1})^*g - g \|_{L^2[0, T]} &= 0 \quad \text{for } \epsilon \rightarrow 0, \\ \lim \| (J_\epsilon^{-1})g - g \|_{L^2[0, T]} &= 0 \quad \text{for } \epsilon \rightarrow 0. \end{aligned} \quad (4.7)$$

Substituting the function  $u$  in (4.4) by the smoothing function  $u_\epsilon$  and using the relation

$$A(t)u_\epsilon = J_\epsilon^{-1}Au - \epsilon J_\epsilon^{-1}\beta_\epsilon(t)u_\epsilon, \quad (4.8)$$

where

$$\beta_\epsilon(t)u_\epsilon = 3 \frac{\partial^2 A(t)}{\partial t^2} \frac{\partial u_\epsilon}{\partial t} + 3 \frac{\partial A(t)}{\partial t} \frac{\partial^2 u_\epsilon}{\partial t^2} + \frac{\partial^3 A(t)}{\partial t^3} u_\epsilon, \quad (4.9)$$

we obtain

$$-\int_\Omega uN \frac{\partial^3 \overline{v_\epsilon^*}}{\partial t^3} dx dt = \int_\Omega A(t)u\overline{v_\epsilon^*} dx dt - \epsilon \int_\Omega \beta_\epsilon(t)u_\epsilon\overline{v_\epsilon^*} dx dt. \quad (4.10)$$

Passing to the limit, the equality in the relation (4.10) remains true for all functions  $u \in L^2(\Omega)$  such that  $(1-x)(\partial u/\partial x)$ ,  $(\partial/\partial x)((1-x)(\partial u/\partial x)) \in L^2(\Omega)$ , and satisfying condition (1.1d).

The operator  $A(t)$  has a continuous inverse in  $L^2(0,1)$  defined by

$$A^{-1}(t)g = - \int_0^x \frac{1}{1-\zeta} \frac{1}{a(\zeta,t)} \int_0^\zeta g(\eta,t) d\eta d\zeta + C(t) \int_0^x \frac{1}{1-\zeta} \frac{1}{a(\zeta,t)} d\zeta, \quad (4.11)$$

where

$$C(t) = \frac{\int_0^1 (d\zeta/a(\zeta,t)) \int_0^\zeta g(\eta,t) d\eta}{\int_0^1 (d\zeta/a(\zeta,t))}. \quad (4.12)$$

Then, we have  $\int_0^1 A^{-1}(t)g dx = 0$ , hence the function  $u_\varepsilon = (J_\varepsilon)^{-1}u$  can be represented in the form

$$u_\varepsilon = (J_\varepsilon)^{-1}A^{-1}(t)A(t)u. \quad (4.13)$$

Then

$$\begin{aligned} B_\varepsilon(t)g &= \frac{\partial^4 a}{\partial t^3 \partial x} J_\varepsilon^{-1} \left[ \frac{1}{a(x,t)} \left( \int_0^x g(\eta,t) d\eta - C(t) \right) \right] \\ &+ \frac{\partial^3 a}{\partial t^3} J_\varepsilon^{-1} \left[ \frac{g}{a} - \frac{a_x}{a^2(x,t)} \left( \int_0^x g(\eta,t) d\eta - C(t) \right) \right] \\ &+ 3 \frac{\partial}{\partial t} \frac{\partial^2 a}{\partial t^2 \partial x} \frac{\partial}{\partial t} J_\varepsilon^{-1} \frac{1}{a(x,t)} \left( \int_0^x g(\eta,t) d\eta - C(t) \right) \\ &+ \frac{\partial a}{\partial t} \frac{\partial}{\partial t} J_\varepsilon^{-1} \frac{g}{a} - \frac{a_x}{a^2(x,t)} \left( \int_0^x g(\eta,t) d\eta - C(t) \right). \end{aligned} \quad (4.14)$$

The adjoint of  $B_\varepsilon(t)$  has the form

$$\begin{aligned} B_\varepsilon^*(t) &= \frac{1}{a} (J_\varepsilon^{-1})^* \left[ \frac{\partial^3 a}{\partial t^3} \bar{h} \right] + \frac{3}{a} (J_\varepsilon^{-1})^* \frac{\partial}{\partial t} \left( \frac{\partial a}{\partial t} \frac{\partial \bar{h}}{\partial t} \right) \\ &+ (G_\varepsilon h)(x) - \frac{\int_0^x (1/a(\eta,t)) d\eta}{\int_0^1 (1/a(x,t)) dx} (G_\varepsilon h)(1), \end{aligned} \quad (4.15)$$

where

$$\begin{aligned}
 (G_\varepsilon h)(x) = & \int_0^x \left( -\frac{3}{a(\zeta,t)} (J_\varepsilon^{-1})^* \frac{\partial}{\partial t} \left( \frac{\partial^2}{\partial t \partial \zeta} \frac{\partial h}{\partial t} \right) \right. \\
 & + 3 \frac{\partial a}{\partial \zeta} \frac{1}{a^2(\zeta,t)} (J_\varepsilon^{-1})^* \frac{\partial}{\partial t} \left( \frac{\partial a}{\partial t} \frac{\partial h}{\partial t} \right) \\
 & \left. - \frac{1}{a(\zeta,t)} (J_\varepsilon^{-1})^* \left( \frac{\partial^4 a}{\partial t^3 \partial \zeta} h \right) + \frac{\partial a}{\partial \zeta} \frac{1}{a^2(\zeta,t)} (J_\varepsilon^{-1})^* \left( \frac{\partial^3 a}{\partial t^3} h \right) \right) d\zeta.
 \end{aligned} \tag{4.16}$$

Consequently, equality (4.10) becomes

$$- \int_\Omega u N \frac{\partial^3 \overline{v_\varepsilon^*}}{\partial t^3} dx dt = \int_\Omega A(t) u \overline{h_\varepsilon} dx dt, \tag{4.17}$$

where  $h_\varepsilon = v_\varepsilon^* - \varepsilon B_\varepsilon^* v_\varepsilon^*$ .

The left-hand side of (4.17) is a continuous linear functional of  $u$ . Hence the function  $h_\varepsilon$  has the derivatives  $(1-x)(\partial h_\varepsilon / \partial x)$ ,  $(\partial / \partial x)((1-x)(\partial h_\varepsilon / \partial x)) \in L^2(\Omega)$  and the following conditions are satisfied:  $h_\varepsilon|_{x=0} = 0$ ,  $h_\varepsilon|_{x=1} = 0$ , and  $(1-x)(\partial h_\varepsilon / \partial x)|_{x=1} = 0$ .

From the equality

$$\begin{aligned}
 (1-x) \frac{\partial h_\varepsilon}{\partial x} = & \left[ I - \varepsilon \frac{1}{a} (J_\varepsilon^{-1})^* \frac{\partial^3 a}{\partial t^3} \right] (1-x) \frac{\partial v_\varepsilon^*}{\partial x} \\
 & - 3\varepsilon \frac{1}{a} (J_\varepsilon^{-1})^* \frac{\partial}{\partial t} \left( \frac{\partial a}{\partial t} \frac{\partial}{\partial t} (1-x) \frac{\partial v_\varepsilon^*}{\partial x} \right),
 \end{aligned} \tag{4.18}$$

and since the operator  $(J_\varepsilon^{-1})^*$  is bounded in  $L^2(\Omega)$ , for sufficiently small  $\varepsilon$ , we have  $\|\varepsilon(1/a)(J_\varepsilon^{-1})^*(\partial^3 a / \partial t^3)\| < 1$ . Hence the operator  $I - \varepsilon(1/a)(J_\varepsilon^{-1})^*(\partial^3 a / \partial t^3)$  has a bounded inverse in  $L^2(\Omega)$ . We conclude that  $(1-x)(\partial v_\varepsilon^* / \partial x) \in L^2(\Omega)$ .

Similarly, we conclude that  $(\partial / \partial x)((1-x)(\partial v_\varepsilon^* / \partial x))$  exists and belongs to  $L^2(\Omega)$ , and the following conditions are satisfied:

$$v_\varepsilon^*|_{x=0} = 0, \quad v_\varepsilon^*|_{x=1} = 0, \quad (1-x) \frac{\partial v_\varepsilon^*}{\partial x} \Big|_{x=1} = 0. \tag{4.19}$$

Substituting  $u = \int_0^t \int_\zeta^\eta \exp(c\tau) v_\varepsilon^*(\tau) d\tau d\zeta d\eta$  in (4.4), where the constant  $c$  satisfies (3.3), we obtain

$$\int_\Omega \exp(ct) v_\varepsilon^* N \overline{v} dx dt = - \int_\Omega A(t) u \overline{v} dx dt. \tag{4.20}$$

Using the properties of smoothing operators, we have

$$\int_{\Omega} \exp(ct) v_{\varepsilon}^* N \bar{v} dx dt = - \int_{\Omega} A(t) u \bar{v}_{\varepsilon}^* dx dt - \varepsilon \int_{\Omega} A(t) u \frac{\partial^3 \bar{v}_{\varepsilon}^*}{\partial t^3} dx dt, \tag{4.21}$$

and from

$$\begin{aligned} \varepsilon \operatorname{Re} \int_{\Omega} A(t) u \frac{\partial^3 \bar{v}_{\varepsilon}^*}{\partial t^3} dx dt &= \varepsilon \int_{\Omega} (1-x) a \frac{\partial u}{\partial x} \frac{\partial}{\partial x} \frac{\partial^3 \bar{v}_{\varepsilon}^*}{\partial t^3} dx dt \\ &= -\varepsilon \operatorname{Re} \int_{\Omega} (1-x) \frac{\partial a}{\partial t} \frac{\partial u}{\partial x} \frac{\partial^2}{\partial t^2} \frac{\partial \bar{v}_{\varepsilon}^*}{\partial x} dx dt \\ &\quad + \varepsilon \operatorname{Re} \int_{\Omega} (1-x) \frac{\partial a}{\partial t} \frac{\partial^2 u}{\partial t \partial x} \frac{\partial}{\partial t} \frac{\partial \bar{v}_{\varepsilon}^*}{\partial x} dx dt \tag{4.22} \\ &\quad + \varepsilon \int_{\Omega} a \exp(-ct) (1-x) \left| \frac{\partial \bar{v}_{\varepsilon}^*}{\partial x} \right|^2 dx dt \\ &\quad + \varepsilon \operatorname{Re} \int_{\Omega} (1-x) \frac{\partial a}{\partial t} \frac{\partial^2 u}{\partial t \partial x} \frac{\partial \bar{v}_{\varepsilon}^*}{\partial x} dx dt, \end{aligned}$$

we have

$$\begin{aligned} \varepsilon \operatorname{Re} \int_{\Omega} A(t) u \frac{\partial^3 \bar{v}_{\varepsilon}^*}{\partial t^3} dx dt &\geq \varepsilon \int_{\Omega} a \exp(+ct) (1-x) \left| \frac{\partial \bar{v}_{\varepsilon}^*}{\partial x} \right|^2 dx dt \\ &\quad - \varepsilon \int_{\Omega} (1-x) \frac{1}{4a} \left( \frac{\partial a}{\partial t} \right)^2 \exp(-ct) \left| \frac{\partial^3 u}{\partial t^2 \partial x} \right|^2 dx dt \\ &\quad - \varepsilon \int_{\Omega} a \exp(+ct) (1-x) \left| \frac{\partial \bar{v}_{\varepsilon}^*}{\partial x} \right|^2 dx dt \tag{4.23} \\ &\quad - \varepsilon \int_{\Omega} \frac{1-x}{2} \left( \frac{\partial a}{\partial t} \right)^2 \exp(-ct) \left| \frac{\partial u}{\partial x} \right|^2 dx dt \\ &\quad - \varepsilon \int_{\Omega} \exp(+ct) \frac{1-x}{2} \left| \frac{\partial^3 \bar{v}_{\varepsilon}^*}{\partial t^2 \partial x} \right|^2 dx dt \\ &\quad - \varepsilon \int_{\Omega} \exp(+ct) \frac{1}{2} \left| \frac{\partial^2 \bar{v}_{\varepsilon}^*}{\partial t \partial x} \right|^2 dx dt \\ &\quad - \varepsilon \int_{\Omega} \frac{1-x}{2} \left( \frac{\partial a}{\partial t} \right)^2 \exp(-ct) \left| \frac{\partial^2 u}{\partial x \partial t} \right|^2 dx dt. \end{aligned}$$

Integrating the first term on the right-hand side by parts in (4.21), we obtain

$$\begin{aligned}
 & \operatorname{Re} \int_{\Omega} A(t) u \bar{v}_{\varepsilon}^* dx dt \\
 & \geq -\frac{3}{2} \int_{\Omega} (1-x) \exp(-ct) \left( \frac{\partial a}{\partial t} - ca \right) \left| \frac{\partial^2 \bar{u}}{\partial t \partial x} \right|^2 dx dt \\
 & \quad + \frac{1}{2} \int_0^1 (1-x) \exp(-ct) \left( a - \left| \frac{\partial a}{\partial t} - ca \right| \right) \left| \frac{\partial^2 \bar{u}}{\partial t \partial x} \right|^2 dx \Big|_{t=T} \\
 & \quad - \frac{1}{2} \int_0^1 (1-x) \exp(-ct) \left\{ \frac{\partial^2 a}{\partial t^2} - 2c \frac{\partial a}{\partial t} + c^2 a + \left| \frac{\partial a}{\partial t} - ca \right| \right\} \left| \frac{\partial u}{\partial x} \right|^2 \Big|_{t=T} dx \\
 & \quad + \frac{1}{2} \int_{\Omega} (1-x) \exp(-ct) \left\{ \frac{\partial^3 a}{\partial t^3} - 3c \frac{\partial^2 a}{\partial t^2} + 3c^2 \frac{\partial a}{\partial t} - c^3 a \right\} \left| \frac{\partial u}{\partial x} \right|^2 dx dt.
 \end{aligned} \tag{4.24}$$

Combining (4.23) and (4.24), we get

$$\begin{aligned}
 & \operatorname{Re} \int_{\Omega} \exp(ct) v_{\varepsilon}^* N \bar{v} dx dt \\
 & \leq \frac{3}{2} \int_{\Omega} (1-x) \exp(-ct) (c_1 - ca_0) \left| \frac{\partial^2 \bar{u}}{\partial t \partial x} \right|^2 dx dt \\
 & \quad - \frac{1}{2} \int_0^1 (1-x) \exp(-ct) \{ a_0 - c'_1 - ca_1 \} \left| \frac{\partial^2 u}{\partial t \partial x} \right|^2 dx \Big|_{t=T} \\
 & \quad + \frac{1}{2} \int_0^1 (1-x) \exp(-ct) \{ c_2 - 2c'_1 c - c^2 a_1 - c'_1 + ca_1 \} \left| \frac{\partial u}{\partial x} \right|^2 \Big|_{t=T} dx \\
 & \quad - \frac{1}{2} \int_{\Omega} (1-x) \exp(-ct) \{ c'_3 - 3c_2 c + 3c^2 c'_1 - c^3 a_1 \} \left| \frac{\partial u}{\partial x} \right|^2 dx dt \\
 & \quad + \varepsilon \left( \int_{\Omega} (1-x) \exp(-ct) \frac{c_1^2}{4a_0} \left| \frac{\partial^3 \bar{u}}{\partial t^2 \partial x} \right|^2 dx dt \right. \\
 & \quad \quad + \int_{\Omega} (1-x) \exp(-ct) \frac{c_1^2}{2} \left| \frac{\partial u}{\partial x} \right|^2 dx dt \\
 & \quad \quad + \int_{\Omega} \frac{1-x}{2} \exp(ct) \left| \frac{\partial^3 v_{\varepsilon}^*}{\partial t^2 \partial x} \right|^2 dx dt \\
 & \quad \quad + \int_{\Omega} (1-x) \exp(-ct) \frac{c_1^2}{2} \left| \frac{\partial^2 \bar{u}}{\partial t \partial x} \right|^2 dx dt \\
 & \quad \quad \left. + \int_{\Omega} \frac{1-x}{2} \exp(ct) \left| \frac{\partial^2 v_{\varepsilon}^*}{\partial t \partial x} \right|^2 dx dt \right).
 \end{aligned} \tag{4.25}$$

Using conditions (3.3) and inequalities (4.23) and (4.24), we obtain

$$\operatorname{Re} \int_{\Omega} \exp(ct) v N \bar{v} \, dx \, dt \leq 0, \quad \text{as } \varepsilon \rightarrow 0. \quad (4.26)$$

Since  $\operatorname{Re} \int_{\Omega} \exp(ct) v J_x \bar{v} \, dx \, dt = 0$ , then  $v = 0$  a.e.

Finally, from the equality  $(1-x)v + J_x v = (1-x)w$ , we conclude  $w = 0$ .  $\square$

**THEOREM 4.2.** *The range  $R(\bar{L})$  of  $\bar{L}$  coincides with  $F$ .*

*Proof.* Since  $F$  is Hilbert space, then  $R(\bar{L}) = F$  if and only if the relation

$$\int_{\Omega} (1-x)^2 \mathcal{E} u \bar{f} \, dx \, dt = 0, \quad (4.27)$$

for arbitrary  $u \in D_0(L)$  and  $\mathcal{F} \in F$ , implies that  $f = 0$ .

Taking  $u \in D_0(L)$  in (4.27) and using Lemma 4.1, we obtain that  $w = (1-x)f = 0$ , then  $f = 0$ .  $\square$

## References

- [1] G. W. Batten Jr., *Second-order correct boundary conditions for the numerical solution of the mixed boundary problem for parabolic equations*, Math. Comp. **17** (1963), 405–413.
- [2] S. A. Beilin, *Existence of solutions for one-dimensional wave equations with nonlocal conditions*, Electron. J. Differential Equations **2001** (2001), no. 76, 1–8.
- [3] N. E. Benouar and N. I. Yurchuk, *Mixed problem with an integral condition for parabolic equations with the Bessel operator*, Differ. Equ. **27** (1991), no. 12, 1482–1487.
- [4] A. Bouziani and N. E. Benouar, *Mixed problem with integral conditions for a third order parabolic equation*, Kobe J. Math. **15** (1998), no. 1, 47–58.
- [5] B. Cahlon, D. M. Kulkarni, and P. Shi, *Stepwise stability for the heat equation with a nonlocal constraint*, SIAM J. Numer. Anal. **32** (1995), no. 2, 571–593.
- [6] J. R. Cannon, *The solution of the heat equation subject to the specification of energy*, Quart. Appl. Math. **21** (1963), 155–160.
- [7] ———, *The One-Dimensional Heat Equation*, Encyclopedia of Mathematics and Its Applications, vol. 23, Addison-Wesley Publishing, Massachusetts, 1984.
- [8] A. U. Çoskun, *Investigation of microscale heat transfer effects in nanoscale electronic devices by spectral methods*, Abstracts of the International Conference on Mathematical Modeling and Scientific Computing (Konya, Turkey), 2001.
- [9] Y. S. Choi and K. Y. Chan, *A parabolic equation with nonlocal boundary conditions arising from electrochemistry*, Nonlinear Anal. **18** (1992), no. 4, 317–331.

- [10] M. Denche and A. L. Marhoune, *High-order mixed-type differential equations with weighted integral boundary conditions*, Electron. J. Differential Equations **2000** (2000), no. 60, 1–10.
- [11] ———, *A three-point boundary value problem with an integral condition for parabolic equations with the Bessel operator*, Appl. Math. Lett. **13** (2000), no. 6, 85–89.
- [12] ———, *Mixed problem with nonlocal boundary conditions for a third-order partial differential equation of mixed type*, Int. J. Math. Math. Sci. **26** (2001), no. 7, 417–426.
- [13] R. E. Ewing and T. Lin, *A class of parameter estimation techniques for fluid flow in porous media*, Adv. in Water Res. **14** (1991), no. 2, 89–97.
- [14] N. I. Ionkin, *The solution of a certain boundary value problem of the theory of heat conduction with a nonclassical boundary condition*, Differ. Uravn. **13** (1977), no. 2, 294–304 (Russian).
- [15] N. I. Kamynin, *A boundary value problem in the theory of heat conduction with non-classical boundary conditions*, U.S.S.R. Comput. Math. and Math. Phys. **4** (1964), 33–59.
- [16] A. V. Kartynnik, *A three-point mixed problem with an integral condition with respect to the space variable for second-order parabolic equations*, Differ. Equ. **26** (1990), no. 9, 1160–1166.
- [17] V. I. Korzyuk and V. B. Kiselev, *A boundary value problem for a third-order linear nonclassical equation generated by the heat equation*, Vestsi Nats. Akad. Navuk Belarusi Ser. Fiz.-Mat. Navuk **2001** (2001), no. 3, 5–11 (Russian).
- [18] L. S. Pulkina, *A non-local problem with integral conditions for hyperbolic equations*, Electron. J. Differential Equations **1999** (1999), no. 45, 1–6.
- [19] A. A. Samarski, *Some problems in the modern theory of differential equations*, Differ. Uravn. **16** (1980), 1925–1935 (Russian).
- [20] P. Shi, *Weak solution to an evolution problem with a nonlocal constraint*, SIAM J. Math. Anal. **24** (1993), no. 1, 46–58.
- [21] P. Shi and M. Shillor, *On design of contact patterns in one-dimensional thermoelasticity*, Theoretical Aspects of Industrial Design (Wright-Patterson Air Force Base, Ohio, 1990), SIAM, Pennsylvania, 1992, pp. 76–82.
- [22] V. F. Volkodavov and V. E. Zhukov, *Two problems for the string vibration equation with integral conditions and special matching conditions on the characteristic*, Differ. Equ. **34** (1998), no. 4, 501–505.
- [23] N. I. Yurchuk, *Mixed problem with an integral condition for certain parabolic equations*, Differ. Equ. **22** (1986), 1457–1463.

M. Denche: Laboratoire Equations Différentielles, Département de Mathématiques, Faculté des Sciences, Université Mentouri, 25000 Constantine, Algeria  
*E-mail address:* [denech@wissal.dz](mailto:denech@wissal.dz)

A. Memou: Laboratoire Equations Différentielles, Département de Mathématiques, Faculté des Sciences, Université Mentouri, 25000 Constantine, Algeria