

Research Article

Some New Delay Integral Inequalities in Two Independent Variables on Time Scales

Bin Zheng, Yaoming Zhang, and Qinghua Feng

School of Science, Shandong University of Technology, Zibo, Shandong, 255049, China

Correspondence should be addressed to Bin Zheng, zhengbin2601@126.com

Received 9 August 2011; Accepted 10 October 2011

Academic Editor: C. Conca

Copyright © 2011 Bin Zheng et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Some new Gronwall-Bellman type delay integral inequalities in two independent variables on time scales are established, which can be used as a handy tool in the research of boundedness of solutions of delay dynamic equations on time scales. Some of the established results are 2D extensions of several known results in the literature, while some results unify existing continuous and discrete analysis.

1. Introduction

In the research of solutions of certain differential and difference equations, if the solutions are unknown, then it is necessary to study their qualitative and quantitative properties such as boundedness, uniqueness, and continuous dependence on initial data. The Gronwall-Bellman inequality [1, 2] and its various generalizations, which provide explicit bounds, play a fundamental role in the research of this domain. During the past decades, much effort has been done for developing such inequalities (e.g., see [3–15] and the references therein). On the other hand, Hilger [16] initiated the theory of time scales as a theory capable to contain both difference and differential calculus in a consistent way. Since then many authors have expounded on various aspects of the theory of dynamic equations on time scales (e.g., see [17–19] and the references therein). In these investigations, integral inequalities on time scales have been paid much attention by many authors, which play a fundamental role in the research of quantitative as well as qualitative properties of solutions of certain dynamic equations on time scales. A lot of integral inequalities on time scales have been established (e.g., see [20–26]), which have been designed to unify continuous and discrete analysis. But to our best knowledge, the Gronwall-Bellman-type delay integral inequalities on time scales have been paid little attention in the literature so far. Recent results in this direction include the work of Li [27] and that of Ma and Pečarić [28]. Furthermore, nobody has studied

the Gronwall-Bellman-type delay integral inequalities in two independent variables on time scales.

The aim of this paper is to establish some new Gronwall-Bellman-type delay integral inequalities in two independent variables on time scales, which provide new bounds for the unknown functions concerned. Some of our results are 2D extensions of many known inequalities in the literature, while some results unify existing continuous and discrete analysis. For illustrating the validity of the established results, we will present some applications of them.

First we will give some preliminaries on time scales and some universal symbols for further use.

Throughout this paper, \mathbb{R} denotes the set of real numbers and $\mathbb{R}_+ = [0, \infty)$, while \mathbb{Z} denotes the set of integers. For two given sets G, H , we denote the set of maps from G to H by (G, H) .

A time scale is an arbitrary nonempty closed subset of the real numbers. In this paper, \mathbb{T} denotes an arbitrary time scale. On \mathbb{T} we define the forward and backward jump operators $\sigma \in (\mathbb{T}, \mathbb{T})$ and $\rho \in (\mathbb{T}, \mathbb{T})$ such that $\sigma(t) = \inf\{s \in \mathbb{T}, s > t\}$, $\rho(t) = \sup\{s \in \mathbb{T}, s < t\}$.

Definition 1.1. The graininess $\mu \in (\mathbb{T}, \mathbb{R}_+)$ is defined by $\mu(t) = \sigma(t) - t$.

Remark 1.2. Obviously, $\mu(t) = 0$ if $\mathbb{T} = \mathbb{R}$ while $\mu(t) = 1$ if $\mathbb{T} = \mathbb{Z}$.

Definition 1.3. A point $t \in \mathbb{T}$ is said to be left-dense if $\rho(t) = t$ and $t \neq \inf \mathbb{T}$, right-dense if $\sigma(t) = t$ and $t \neq \sup \mathbb{T}$, left-scattered if $\rho(t) < t$, and right-scattered if $\sigma(t) > t$.

Definition 1.4. The set \mathbb{T}^κ is defined to be \mathbb{T} if \mathbb{T} does not have a left-scattered maximum; otherwise it is \mathbb{T} without the left-scattered maximum.

Definition 1.5. A function $f \in (\mathbb{T}, \mathbb{R})$ is called rd-continuous if it is continuous at right-dense points and if the left-sided limits exist at left-dense points, while f is called regressive if $1 + \mu(t)f(t) \neq 0$. C_{rd} denotes the set of rd-continuous functions, while \mathfrak{R} denotes the set of all regressive and rd-continuous functions, and $\mathfrak{R}^+ = \{f \mid f \in \mathfrak{R}, 1 + \mu(t)f(t) > 0, \forall t \in \mathbb{T}\}$.

Definition 1.6. For some $t \in \mathbb{T}^\kappa$ and a function $f \in (\mathbb{T}, \mathbb{R})$, the *delta derivative* of f at t is denoted by $f^\Delta(t)$ (provided it exists) with the property such that for every $\varepsilon > 0$ there exists a neighborhood \mathfrak{U} of t satisfying

$$\left| f(\sigma(t)) - f(s) - f^\Delta(t)(\sigma(t) - s) \right| \leq \varepsilon |\sigma(t) - s| \quad \forall s \in \mathfrak{U}. \quad (1.1)$$

Similarly, for some $y \in \mathbb{T}^\kappa$ and a function $f \in (\mathbb{T} \times \mathbb{T}, \mathbb{R})$, the *partial delta* of f with respect to y is denoted by $(f(x, y))_y^\Delta$ or $f_y^\Delta(x, y)$ and satisfies

$$\left| f(x, \sigma(y)) - f(x, s) - f_y^\Delta(x, y)(\sigma(y) - s) \right| \leq \varepsilon |\sigma(y) - s| \quad \forall \varepsilon > 0, \quad (1.2)$$

where $s \in \mathfrak{U}$ and \mathfrak{U} is a neighborhood of y .

Remark 1.7. If $\mathbb{T} = \mathbb{R}$, then $f^\Delta(t)$ becomes the usual derivative $f'(t)$, while $f^\Delta(t) = f(t+1) - f(t)$ if $\mathbb{T} = \mathbb{Z}$, which represents the forward difference.

For more details about the calculus of time scales, see [29]. In the rest of this paper, for the convenience of notation, we always assume that $\mathbb{T}_0 = [x_0, \infty) \cap \mathbb{T}$, $\tilde{\mathbb{T}}_0 = [y_0, \infty) \cap \mathbb{T}$, where $x_0, y_0 \in \mathbb{T}^\kappa$ and furthermore assume $\mathbb{T}_0 \subseteq \mathbb{T}^\kappa, \tilde{\mathbb{T}}_0 \subseteq \mathbb{T}^\kappa$.

2. Main Results

We will give some lemmas for further use.

Lemma 2.1. *Suppose $X \in \mathbb{T}_0$ is a fixed number, and $u(X, y), a(X, y), b(X, y) \in C_{\text{rd}}, m(X, y) \in \mathfrak{R}_+$ with respect to $y, m(X, y) \geq 0$, then*

$$u(X, y) \leq a(X, y) + b(X, y) \int_{y_0}^y m(X, t) u(X, t) \Delta t, \quad y \in \tilde{\mathbb{T}}_0, \quad (2.1)$$

implies

$$u(X, y) \leq a(X, y) + b(X, y) \int_{y_0}^y e_{\bar{m}}(y, \sigma(t)) a(X, t) m(X, t) \Delta t, \quad y \in \tilde{\mathbb{T}}_0, \quad (2.2)$$

where $\bar{m}(X, y) = m(X, y)b(X, y)$: and $e_{\bar{m}}(y, y_0)$ is the unique solution of the following equation

$$z_y^\Delta(X, y) = m(X, y)z(X, y), z(X, y_0) = 1. \quad (2.3)$$

The proof of Lemma 2.1 is similar to [26, Theorem 5.6].

Lemma 2.2. *Under the conditions of Lemma 2.1, and furthermore assuming $a(x, y)$ is nondecreasing in y for every fixed $x, b(x, y) \equiv 1$, then one has*

$$u(X, y) \leq a(X, y)e_m(y, y_0). \quad (2.4)$$

Proof. Since $a(x, y)$ is nondecreasing in y for every fixed x , then from Lemma 2.1 we have

$$u(X, y) \leq a(X, y) + \int_{y_0}^y e_m(y, \sigma(t)) a(X, t) m(X, t) \Delta t \leq a(X, y) \left[1 + \int_{y_0}^y e_m(y, \sigma(t)) m(X, t) \Delta t \right]. \quad (2.5)$$

On the other hand, from [29, Theorems 2.39 and 2.36 (i)] we have $1 + \int_{y_0}^y e_m(y, \sigma(t)) m(X, t) \Delta t = e_m(y, y_0)$. Then collecting the above information, we can obtain the desired inequality. \square

Lemma 2.3 (see [11]). *Assume that $a \geq 0, p \geq q \geq 0$, and $p \neq 0$; then, for any $K > 0$*

$$a^{q/p} \leq \frac{q}{p} K^{(q-p)/p} a + \frac{p-q}{p} K^{q/p}. \quad (2.6)$$

Lemma 2.4. Let $h : \mathbb{T} \times \mathbb{R} \rightarrow \mathbb{R}$ be continuous and nondecreasing in the second variable, and assume X is a fixed number in \mathbb{T} . Suppose $v(X, y)$ and $w(X, y)$ satisfy the dynamics inequalities:

$$v_y^\Delta \leq h(y, v), \quad w_y^\Delta \geq h(y, w). \quad (2.7)$$

Then $v(X, y_0) \leq w(X, y_0)$ for some $y_0 \in \mathbb{T}$ implies $v(X, y) \leq w(X, y)$ for all $y \in \mathbb{T}$.

The proof of Lemma 2.4 is similar to [26, Theorem 5.7].

Theorem 2.5. Suppose $u, a, b, f \in C_{rd}(\mathbb{T}_0 \times \tilde{\mathbb{T}}_0, \mathbb{R}_+)$, and $a(x, y), b(x, y)$ are nondecreasing. p is a constant, and $p \geq 1$. $\tau_1 \in (\mathbb{T}_0, \mathbb{T})$, $\tau_1(x) \leq x, -\infty < \alpha = \inf\{\tau_1(x), x \in \mathbb{T}_0\} \leq x_0$. $\tau_2 \in (\tilde{\mathbb{T}}_0, \mathbb{T})$, $\tau_2(y) \leq y, -\infty < \beta = \inf\{\tau_2(y), y \in \tilde{\mathbb{T}}_0\} \leq y_0$. $\phi \in C_{rd}([\alpha, x_0] \times [\beta, y_0]) \cap \mathbb{T}^2, \mathbb{R}_+$. If for $(x, y) \in \mathbb{T}_0 \times \tilde{\mathbb{T}}_0$, $u(x, y)$ satisfies the following inequality:

$$u^p(x, y) \leq a(x, y) + b(x, y) \int_{y_0}^y \int_{x_0}^x [f(s, t)u(\tau_1(s), \tau_2(t))] \Delta s \Delta t, \quad (2.8)$$

with the initial condition

$$\begin{aligned} u(x, y) &= \phi(x, y), \quad \text{if } x \in [\alpha, x_0] \cap \mathbb{T} \text{ or } y \in [\beta, y_0] \cap \mathbb{T}, \\ \phi(\tau_1(x), \tau_2(y)) &\leq a^{1/p}(x, y), \quad \forall (x, y) \in \mathbb{T}_0 \times \tilde{\mathbb{T}}_0, \text{ if } \tau_1(x) \leq x_0 \text{ or } \tau_2(y) \leq y_0, \end{aligned} \quad (2.9)$$

then

$$u(x, y) \leq \left[H_1(x, y) + b(x, y) \int_{y_0}^y e_{\bar{H}_2}(y, \sigma(t)) H_2(x, t) H_1(x, t) \Delta t \right]^{1/p}, \quad (x, y) \in \mathbb{T}_0 \times \tilde{\mathbb{T}}_0, \quad (2.10)$$

where

$$\begin{aligned} H_1(x, y) &= a(x, y) + b(x, y) \int_{y_0}^y \int_{x_0}^x f(s, t) \frac{p-1}{p} K^{1/p} \Delta s \Delta t, \quad \forall K > 0, \\ H_2(x, y) &= \int_{x_0}^x f(s, y) \frac{1}{p} K^{(1-p)/p} \Delta s, \\ \bar{H}_2(x, y) &= b(x, y) H_2(x, y). \end{aligned} \quad (2.11)$$

Proof. Fix $X \in \mathbb{T}_0$, and $x \in [x_0, X] \cap \mathbb{T}$, $y \in \tilde{\mathbb{T}}_0$. Let

$$v(x, y) = a(x, y) + b(x, y) \int_{y_0}^y \int_{x_0}^x [f(s, t)u(\tau_1(s), \tau_2(t))] \Delta s \Delta t. \quad (2.12)$$

Then

$$u(x, y) \leq v^{1/p}(x, y) \leq v^{1/p}(X, y), \quad \forall x \in [x_0, X] \cap \mathbb{T}, y \in \tilde{\mathbb{T}}_0. \quad (2.13)$$

If $\tau_1(x) \geq x_0$ and $\tau_2(y) \geq y_0$, then $\tau_1(x) \in [x_0, X] \cap \mathbb{T}$, $\tau_2(y) \in \tilde{\mathbb{T}}_0$, and

$$u(\tau_1(x), \tau_2(y)) \leq v^{1/p}(\tau_1(x), \tau_2(y)) \leq v^{1/p}(x, y). \quad (2.14)$$

If $\tau_1(x) \leq x_0$ or $\tau_2(y) \leq y_0$, then from (2.9) we have

$$u(\tau_1(x), \tau_2(y)) = \phi(\tau_1(x), \tau_2(y)) \leq a^{1/p}(x, y) \leq v^{1/p}(x, y). \quad (2.15)$$

From (2.14) and (2.15) we always have

$$u(\tau_1(x), \tau_2(y)) \leq v^{1/p}(x, y), \quad x \in [x_0, X] \cap \mathbb{T}, y \in \tilde{\mathbb{T}}_0. \quad (2.16)$$

Moreover

$$\begin{aligned} v(X, y) &= a(X, y) + b(X, y) \int_{y_0}^y \int_{x_0}^X [f(s, t)u(\tau_1(s), \tau_2(t))] \Delta s \Delta t \\ &\leq a(X, y) + b(X, y) \int_{y_0}^y \int_{x_0}^X f(s, t)v^{1/p}(s, t) \Delta s \Delta t. \end{aligned} \quad (2.17)$$

From Lemma 2.3, we have

$$v^{1/p}(s, t) \leq \frac{1}{p}K^{(1-p)/p}v(s, t) + \frac{p-1}{p}K^{1/p}, \quad \forall K > 0. \quad (2.18)$$

So

$$\begin{aligned} v(X, y) &\leq a(X, y) + b(X, y) \int_{y_0}^y \int_{x_0}^X f(s, t) \left[\frac{1}{p}K^{(1-p)/p}v(s, t) + \frac{p-1}{p}K^{1/p} \right] \Delta s \Delta t \\ &\leq a(X, y) + b(X, y) \int_{y_0}^y \int_{x_0}^X f(s, t) \frac{p-1}{p}K^{1/p} \Delta s \Delta t \\ &\quad + b(X, y) \int_{y_0}^y \left[\int_{x_0}^X f(s, t) \frac{1}{p}K^{(1-p)/p} \Delta s \right] v(X, t) \Delta t \\ &= H_1(X, y) + b(X, y) \int_{y_0}^y H_2(X, t)v(X, t) \Delta t. \end{aligned} \quad (2.19)$$

Then applying Lemma 2.1 to (2.19), we obtain

$$v(X, y) \leq H_1(X, y) + b(X, y) \int_{y_0}^y e_{\overline{H_2}}(y, \sigma(t)) H_2(X, t) H_1(X, t) \Delta t. \quad (2.20)$$

So

$$u(x, y) \leq v^{1/p}(X, y) \leq \left[H_1(X, y) + b(X, y) \int_{y_0}^y e_{\overline{H_2}}(y, \sigma(t)) H_2(X, t) H_1(X, t) \Delta t \right]^{1/p}, \quad (2.21)$$

$$x \in [x_0, X] \cap \mathbb{T}, \quad y \in \tilde{\mathbb{T}}_0.$$

Setting $x = X$ in (2.21), it follows that

$$u(X, y) \leq \left[H_1(X, y) + b(X, y) \int_{y_0}^y e_{\overline{H_2}}(y, \sigma(t)) H_2(X, t) H_1(X, t) \Delta t \right]^{1/p}. \quad (2.22)$$

Replacing X with x in (2.22), we obtain the desired inequality. \square

Remark 2.6. Theorem 2.5 is the 2D extension of [27, Theorem 1]. For its special case $\mathbb{T} = \mathbb{R}$, the established bound for $u(x, y)$ in (2.10) is a new bound compared with the result in [12, Theorem 2.2].

Remark 2.7. Assume $b(x, y) \equiv 1$ in Theorem 2.5. If we apply Lemma 2.2 instead of Lemma 2.1 to (2.19) in the proof of Theorem 2.5, then we obtain another bound for $u(x, y)$ as follows:

$$u(x, y) \leq [H_1(x, y) e_{H_2}(y, y_0)]^{1/p}, \quad (x, y) \in \mathbb{T}_0 \times \tilde{\mathbb{T}}_0. \quad (2.23)$$

Now we will establish a more general inequality than that in Theorem 2.5.

Theorem 2.8. Suppose $u, a, b, f, \phi, \tau_1, \tau_2, \alpha, \beta$ are the same as in Theorem 2.5, and $g \in C_{rd}(\mathbb{T}_0 \times \tilde{\mathbb{T}}_0, \mathbb{R}_+)$. p, q, r are constants, and $p \geq q \geq 0$, $p \geq r \geq 0$, $p \neq 0$. If for $(x, y) \in \mathbb{T}_0 \times \tilde{\mathbb{T}}_0$, $u(x, y)$ satisfies the following inequality:

$$u^p(x, y) \leq a(x, y) + b(x, y) \int_{y_0}^y \int_{x_0}^x [f(s, t) u^q(\tau_1(s), \tau_2(t)) + g(s, t) u^r(\tau_1(s), \tau_2(t))] \Delta s \Delta t, \quad (2.24)$$

with the initial condition (2.9), then

$$u(x, y) \leq \left[\tilde{H}_1(x, y) + b(x, y) \int_{y_0}^y e_{\overline{\tilde{H}_2}}(y, \sigma(t)) \tilde{H}_2(x, t) \tilde{H}_1(x, t) \Delta t \right]^{1/p}, \quad (x, y) \in \mathbb{T}_0 \times \tilde{\mathbb{T}}_0, \quad (2.25)$$

where

$$\begin{aligned}\widetilde{H}_1(x, y) &= a(x, y) + b(x, y) \int_{y_0}^y \int_{x_0}^x \left[f(s, t) \frac{p-q}{p} K^{q/p} + g(s, t) \frac{p-r}{p} K^{r/p} \right] \Delta s \Delta t, \\ \widetilde{H}_2(x, y) &= \int_{x_0}^x \left[f(s, y) \frac{q}{p} K^{(q-p)/p} + g(s, y) \frac{r}{p} K^{(r-p)/p} \right] \Delta s, \\ \overline{\widetilde{H}}_2(x, y) &= b(x, y) \widetilde{H}_2(x, y).\end{aligned}\tag{2.26}$$

Proof. Fix $X \in \mathbb{T}_0$, and $x \in [x_0, X] \cap \mathbb{T}$, $y \in \widetilde{\mathbb{T}}_0$. Let

$$v(x, y) = a(x, y) + b(x, y) \int_{y_0}^y \int_{x_0}^x [f(s, t) u^q(\tau_1(s), \tau_2(t)) + g(s, t) u^r(\tau_1(s), \tau_2(t))] \Delta s \Delta t.\tag{2.27}$$

Then

$$u(x, y) \leq v^{1/p}(x, y) \leq v^{1/p}(X, y), \quad \forall x \in [x_0, X] \cap \mathbb{T}, y \in \widetilde{\mathbb{T}}_0.\tag{2.28}$$

Similar to (2.14)–(2.16), we obtain

$$u(\tau_1(x), \tau_2(y)) \leq v^{1/p}(x, y), \quad x \in [x_0, X] \cap \mathbb{T}, y \in \widetilde{\mathbb{T}}_0.\tag{2.29}$$

So

$$\begin{aligned}v(X, y) &= a(X, y) + b(X, y) \int_{y_0}^y \int_{x_0}^X [f(s, t) u^q(\tau_1(s), \tau_2(t)) + g(s, t) u^r(\tau_1(s), \tau_2(t))] \Delta s \Delta t \\ &\leq a(X, y) + b(X, y) \int_{y_0}^y \int_{x_0}^X [f(s, t) v^{q/p}(s, t) + g(s, t) v^{r/p}(s, t)] \Delta s \Delta t.\end{aligned}\tag{2.30}$$

From Lemma 2.3, we have

$$\begin{aligned}v^{q/p}(s, t) &\leq \frac{q}{p} K^{(q-p)/p} v(s, t) + \frac{p-q}{p} K^{q/p}, \quad \forall K > 0, \\ v^{r/p}(s, t) &\leq \frac{r}{p} K^{(r-p)/p} v(s, t) + \frac{p-r}{p} K^{r/p}, \quad \forall K > 0.\end{aligned}\tag{2.31}$$

Combining (2.30) and (2.31) we get that

$$\begin{aligned}
v(X, y) &\leq a(X, y) + b(X, y) \int_{y_0}^y \int_{x_0}^X \left[f(s, t) \left(\frac{q}{p} K^{(q-p)/p} v(s, t) + \frac{p-q}{p} K^{q/p} \right) \right. \\
&\quad \left. + g(s, t) \left(\frac{r}{p} K^{(r-p)/p} v(s, t) + \frac{p-r}{p} K^{r/p} \right) \right] \Delta s \Delta t \\
&\leq a(X, y) + b(X, y) \int_{y_0}^y \int_{x_0}^X \left[f(s, t) \frac{p-q}{p} K^{q/p} + g(s, t) \frac{p-r}{p} K^{r/p} \right] \Delta s \Delta t \quad (2.32) \\
&\quad + b(X, y) \int_{y_0}^y \left\{ \int_{x_0}^X \left[f(s, t) \frac{q}{p} K^{(q-p)/p} + g(s, t) \frac{r}{p} K^{(r-p)/p} \right] \Delta s \right\} v(X, t) \Delta t \\
&= \widetilde{H}_1(X, y) + b(X, y) \int_{y_0}^y \widetilde{H}_2(X, t) v(X, t) \Delta t.
\end{aligned}$$

Applying Lemma 2.1 to (2.32) yields

$$v(X, y) \leq \widetilde{H}_1(X, y) + b(X, y) \int_{y_0}^y e_{\widetilde{H}_2}^{\overline{\overline{}}}(y, \sigma(t)) \widetilde{H}_2(X, t) \widetilde{H}_1(X, t) \Delta t. \quad (2.33)$$

Then

$$\begin{aligned}
u(x, y) \leq v^{1/p}(X, y) &\leq \left[\widetilde{H}_1(X, y) + b(X, y) \int_{y_0}^y e_{\widetilde{H}_2}^{\overline{\overline{}}}(y, \sigma(t)) \widetilde{H}_2(X, t) \widetilde{H}_1(X, t) \Delta t \right]^{1/p}, \quad (2.34) \\
&x \in [x_0, X] \cap \mathbb{T}, \quad y \in \widetilde{\mathbb{T}}_0.
\end{aligned}$$

Setting $x = X$ in (2.34) yields

$$u(X, y) \leq \left[\widetilde{H}_1(X, y) + b(X, y) \int_{y_0}^y e_{\widetilde{H}_2}^{\overline{\overline{}}}(y, \sigma(t)) \widetilde{H}_2(X, t) \widetilde{H}_1(X, t) \Delta t \right]^{1/p}. \quad (2.35)$$

Considering $X \in \mathbb{T}_0$ is arbitrary and replacing X with x in (2.35), we obtain the desired inequality. \square

Remark 2.9. Assume $b(x, y) \equiv 1$ in Theorem 2.8. If we apply Lemma 2.2 instead of Lemma 2.1 to (2.32) in the proof of Theorem 2.8, then we obtain another bound for $u(x, y)$ as follows:

$$u(x, y) \leq \left[\widetilde{H}_1(x, y) e_{\widetilde{H}_2}^{\overline{\overline{}}}(y, y_0) \right]^{1/p}, \quad (x, y) \in \mathbb{T}_0 \times \widetilde{\mathbb{T}}_0. \quad (2.36)$$

Remark 2.10. Theorem 2.8 is the 2D extension of [27, Theorem 3].

Theorem 2.11. Suppose $u, f, \alpha, \beta, \phi, \tau_1, \tau_2$ are the same as in Theorem 2.5, and $C > 0$ is a constant. If for $(x, y) \in (\mathbb{T}_0 \times \tilde{\mathbb{T}}_0)$, $u(x, y)$ satisfies the following inequality:

$$u^2(x, y) \leq C + \int_{y_0}^y \int_{x_0}^x f(s, t) [u(\tau_1(s), \tau_2(t)) + u(\tau_1(s), \sigma(\tau_2(t)))] \Delta s \Delta t \quad (2.37)$$

with the initial condition

$$\begin{aligned} u(x, y) &= \phi(x, y), \quad \text{if } x \in [\alpha, x_0] \cap \mathbb{T} \text{ or } y \in [\beta, y_0] \cap \mathbb{T}, \\ \phi(\tau_1(x), \tau_2(y)) &\leq C^{1/2}, \quad \forall (x, y) \in \mathbb{T}_0 \times \tilde{\mathbb{T}}_0, \text{ if } \tau_1(x) \leq x_0 \text{ or } \tau_2(y) \leq y_0, \end{aligned} \quad (2.38)$$

then

$$u(x, y) \leq \sqrt{C} + \int_{y_0}^y \int_{x_0}^x f(s, t) \Delta s \Delta t, \quad (x, y) \in (\mathbb{T}_0 \times \tilde{\mathbb{T}}_0). \quad (2.39)$$

Proof. Let the right side of (2.37) be $v^2(x, y)$. Then

$$u(x, y) \leq v(x, y), \quad \forall (x, y) \in (\mathbb{T}_0 \times \tilde{\mathbb{T}}_0). \quad (2.40)$$

For $(x, y) \in (\mathbb{T}_0 \times \tilde{\mathbb{T}}_0)$, if $\tau_1(x) \geq x_0$ and $\tau_2(y) \geq y_0$, then $\tau_1(x) \in \mathbb{T}_0$ and $\tau_2(y) \in \tilde{\mathbb{T}}_0$, and from (2.40) we have

$$u(\tau_1(x), \tau_2(y)) \leq v(\tau_1(x), \tau_2(y)) \leq v(x, y). \quad (2.41)$$

If $\tau_1(x) \leq x_0$ or $\tau_2(y) \leq y_0$, from (2.38) we have

$$u(\tau_1(x), \tau_2(y)) = \phi(\tau_1(x), \tau_2(y)) \leq a^{1/2}(\tau_1(x), \tau_2(y)) \leq a^{1/2}(x, y) \leq v(x, y). \quad (2.42)$$

So from (2.41) and (2.42), we always have

$$u(\tau_1(x), \tau_2(y)) \leq v(x, y), \quad \forall (x, y) \in (\mathbb{T}_0 \times \tilde{\mathbb{T}}_0). \quad (2.43)$$

Similarly, when $\tau_1(x) \geq x_0$ and $\sigma(\tau_2(y)) \geq y_0$, then $\tau_1(x) \in \mathbb{T}_0$ and $\sigma(\tau_2(y)) \in \tilde{\mathbb{T}}_0$, and from (2.40) we have

$$u(\tau_1(x), \sigma(\tau_2(y))) \leq v(\tau_1(x), \sigma(\tau_2(y))) \leq v(x, \sigma(y)). \quad (2.44)$$

When $\tau_1(x) \leq x_0$ or $\sigma(\tau_2(y)) \leq y_0$, considering $\sigma(\tau_2(y)) \geq \tau_2(y) \geq \beta$, from (2.38) it follows that

$$u(\tau_1(x), \sigma(\tau_2(y))) = \phi(\tau_1(x), \sigma(\tau_2(y))) \leq C^{1/2} \leq v(x, y) \leq v(x, \sigma(y)). \quad (2.45)$$

Combining (2.44) and (2.45), we always have

$$u(\tau_1(x), \sigma(\tau_2(y))) \leq v(x, \sigma(y)), \quad \forall (x, y) \in (\mathbb{T}_0 \times \tilde{\mathbb{T}}_0). \quad (2.46)$$

By (2.43) and (2.46), we obtain

$$v^2(x, y) \leq C + \int_{y_0}^y \int_{x_0}^x f(s, t) [v(s, t) + v(s, \sigma(t))] \Delta s \Delta t, \quad x \in \mathbb{T}_0, y \in \tilde{\mathbb{T}}_0. \quad (2.47)$$

Let the right side of (2.47) be $z^2(x, y)$. Then

$$v(x, y) \leq z(x, y), \quad \forall (x, y) \in (\mathbb{T}_0 \times \tilde{\mathbb{T}}_0), \quad (2.48)$$

$$\begin{aligned} (z^2(x, y))_y^\Delta &= \int_{x_0}^x f(s, y) [v(s, y) + v(s, \sigma(y))] \Delta s \\ &\leq \left(\int_{x_0}^x f(s, y) \Delta s \right) [v(x, y) + v(x, \sigma(y))] \\ &\leq \left(\int_{x_0}^x f(s, y) \Delta s \right) [z(x, y) + z(x, \sigma(y))]. \end{aligned} \quad (2.49)$$

Considering $z(x, y) + z(x, \sigma(y)) \geq z(x_0, y_0) = C > 0$, and $(z^2(x, y))_y^\Delta = [z(x, y) + z(x, \sigma(y))] (z(x, y))_y^\Delta$, from (2.49) it follows that

$$(z(x, y))_y^\Delta \leq \int_{x_0}^x f(s, y) \Delta s. \quad (2.50)$$

An integration of (2.50) with respect to y from y_0 to y yields $z(x, y) - z(x, y_0) \leq \int_{y_0}^y \int_{x_0}^x f(s, t) \Delta s \Delta t$.

Considering $z(x, y_0) = \sqrt{C}$, it follows that

$$z(x, y) \leq \sqrt{C} + \int_{y_0}^y \int_{x_0}^x f(s, t) \Delta s \Delta t. \quad (2.51)$$

Then combining (2.40), (2.48), and (2.51), we obtain

$$u(x, y) \leq v(x, y) \leq z(x, y) \leq \sqrt{C} + \int_{y_0}^y \int_{x_0}^x f(s, t) \Delta s \Delta t, \quad (2.52)$$

and the proof is complete. \square

Remark 2.12. If we take $\mathbb{T} = \mathbb{R}$, then Theorem 2.11 becomes the extension of the known Ou-Iang's inequality [13] to the 2D case.

The following theorem provides a more general result than Theorem 2.11.

Theorem 2.13. Suppose p is a positive integer, and $p \geq 2$. Under the conditions of Theorem 2.11, if $u(x, y)$ satisfies

$$u^p(x, y) \leq C + \int_{y_0}^y \int_{x_0}^x f(s, t) \sum_{l=0}^{p-1} \left\{ u^l[(\tau_1(s), \tau_2(t))] u^{p-1-l}[(\tau_1(s), \sigma(\tau_2(t)))] \right\} \Delta s \Delta t, \quad (2.53)$$

$$(x, y) \in (\mathbb{T}_0 \times \tilde{\mathbb{T}}_0),$$

with the initial condition

$$u(x, y) = \phi(x, y), \quad \text{if } x \in [\alpha, x_0] \cap \mathbb{T} \text{ or } y \in [\beta, y_0] \cap \mathbb{T}, \quad (2.54)$$

$$\phi(\tau_1(x), \tau_2(y)) \leq C^{1/p}, \quad \forall (x, y) \in \mathbb{T}_0 \times \tilde{\mathbb{T}}_0, \text{ if } \tau_1(x) \leq x_0 \text{ or } \tau_2(y) \leq y_0,$$

then

$$u(x, y) \leq C^{1/p} + \int_{y_0}^y \int_{x_0}^x f(s, t) \Delta s \Delta t, \quad x \in \mathbb{T}_0, y \in \tilde{\mathbb{T}}_0. \quad (2.55)$$

The proof of Theorem 2.13 is similar to Theorem 2.11. As long as we notice a *delta d* differentiable function $z(x, y)$, the following formula [26, Equation (6.2)] holds:

$$(z^p(x, y))_y^\Delta = (z(x, y))_y^\Delta \sum_{l=0}^{p-1} \left[z^l(x, y) z^{p-1-l}(x, \sigma(y)) \right]. \quad (2.56)$$

Then following a similar manner as in Theorem 2.11, we can deduce the desired result.

Theorem 2.14. Suppose $u, f, \tau_1, \tau_2, \phi, \alpha, \beta$ are the same as in Theorem 2.5, $\omega \in C(\mathbb{R}_+, \mathbb{R}_+)$ is nondecreasing, and p, C are constants with $p \geq 1, C > 0$. Furthermore, define a bijective function G such that $[G(z(x, y))]_y^\Delta = (z(x, y))_y^\Delta / \omega(z^{1/p}(x, y))$. If for $(x, y) \in \mathbb{T}_0 \times \tilde{\mathbb{T}}_0$, $u(x, y)$ satisfies the following inequality:

$$u^p(x, y) \leq C + \int_{y_0}^y \int_{x_0}^x [f(s, t) \omega(u(\tau_1(s), \tau_2(t)))] \Delta s \Delta t, \quad (2.57)$$

with the initial condition

$$u(x, y) = \phi(x, y), \quad \text{if } x \in [\alpha, x_0] \cap \mathbb{T} \text{ or } y \in [\beta, y_0] \cap \mathbb{T}, \quad (2.58)$$

$$\phi(\tau_1(x), \tau_2(y)) \leq C^{1/p}, \quad \forall (x, y) \in \mathbb{T}_0 \times \tilde{\mathbb{T}}_0, \text{ if } \tau_1(x) \leq x_0 \text{ or } \tau_2(y) \leq y_0,$$

then

$$u(x, y) \leq \left\{ G^{-1} \left[G(C) + \int_{y_0}^y \eta_1(x, t) \Delta t \right] \right\}^{1/p}, \quad (x, y) \in (\mathbb{T}_0 \times \tilde{\mathbb{T}}_0), \quad (2.59)$$

where $\eta_1(x, y) = \int_{x_0}^x f(s, y) \Delta s$.

Proof. Fix $X \in \mathbb{T}_0$, and $x \in [x_0, X] \cap \mathbb{T}$, $y \in \tilde{\mathbb{T}}_0$. Let

$$v(x, y) = C + \int_{y_0}^y \int_{x_0}^x [f(s, t)] \Delta s \Delta t, \quad x \in [x_0, X] \cap \mathbb{T}, \quad y \in \tilde{\mathbb{T}}_0. \quad (2.60)$$

Then

$$u(x, y) \leq v^{1/p}(x, y) \leq v^{1/p}(X, y), \quad \forall x \in [x_0, X] \cap \mathbb{T}, \quad y \in \tilde{\mathbb{T}}_0. \quad (2.61)$$

Similar to (2.14)–(2.16), we obtain

$$u(\tau_1(x), \tau_2(y)) \leq v^{1/p}(x, y), \quad x \in [x_0, X] \cap \mathbb{T}, \quad y \in \tilde{\mathbb{T}}_0. \quad (2.62)$$

Moreover,

$$\begin{aligned} v_y^\Delta(X, y) &= \int_{x_0}^X [f(s, y) \omega(u(\tau_1(s), \tau_2(y)))] \Delta s \\ &\leq \int_{x_0}^X [f(s, y) \omega(v^{1/p}(s, y))] \Delta s \\ &\leq \left(\int_{x_0}^X f(s, y) \Delta s \right) \omega(v^{1/p}(X, y)) = \eta_1(X, y) \omega(v^{1/p}(X, y)). \end{aligned} \quad (2.63)$$

Let $\bar{v}(X, y)$ be the solution of the following problem:

$$\bar{v}_y^\Delta(X, y) = \eta_1(X, y) \omega(\bar{v}^{1/p}(X, y)), \quad \bar{v}(X, y_0) = C. \quad (2.64)$$

Considering $v(X, y_0) = C$ and ω is nondecreasing and continuous, then from (2.63), (2.64), and Lemma 2.4, we have

$$v(X, y) \leq \bar{v}(X, y), \quad y \in \tilde{\mathbb{T}}_0. \quad (2.65)$$

On the other hand, from the definition of G we have $(G(\bar{v}(X, y)))_y^\Delta = \bar{v}_y^\Delta(X, y) / \omega(\bar{v}(X, y)) = \eta_1(X, y)$. Then an integration with respect to y from y_0 to y yields

$$G(\bar{v}(X, y)) - G(\bar{v}(X, y_0)) = \int_{y_0}^y \eta_1(X, t) \Delta t, \quad (2.66)$$

that is,

$$\bar{v}(X, y) \leq G^{-1} \left[G(C) + \int_{y_0}^y \eta_1(X, t) \Delta t \right], \quad y \in \tilde{\mathbb{T}}_0. \quad (2.67)$$

Combining (2.61), (2.65), and (2.67), we have

$$u(x, y) \leq \left\{ G^{-1} \left[G(C) + \int_{y_0}^y \eta_1(X, t) \Delta t \right] \right\}^{1/p}, \quad x \in [x_0, X] \cap \mathbb{T}, \quad y \in \tilde{\mathbb{T}}_0. \quad (2.68)$$

Setting $x = X$ in (2.68), we get the desired result. \square

Remark 2.15. If we take $\mathbb{T} = \mathbb{R}$, then Theorem 2.14 reduces to [14, Theorem 2.1], while Theorem 2.14 reduces to [15, Theorem 2.1] if we take $\mathbb{T} = \mathbb{Z}$.

Theorem 2.16. Suppose $u, f, \tau_1, \tau_2, \phi, \alpha, \beta$ are the same as in Theorem 2.5, and furthermore, u is delta differential on $\tilde{\mathbb{T}}_0$ with respect to y , $g \in C_{rd}(\mathbb{T}_0 \times \tilde{\mathbb{T}}_0, \mathbb{R}_+)$. $\omega \in C(\mathbb{R}_+, \mathbb{R}_+)$ is nondecreasing, and ω is submultiplicative, that is, $\omega(\alpha\beta) \leq \omega(\alpha)\omega(\beta)$, for all $\alpha, \beta \in \mathbb{R}_+$. $C > 0$ is a constant. \tilde{G} is a bijective function such that $[\tilde{G}(z(x, y))]_y^\Delta = (z(x, y))_y^\Delta / \omega(z(x, y))$. If for $(x, y) \in \mathbb{T}_0 \times \tilde{\mathbb{T}}_0$, $u(x, y)$ satisfies the following inequality:

$$u(x, y) \leq C + \int_{y_0}^y \int_{x_0}^x [f(s, t)\omega(u(\tau_1(s), \tau_2(t))) + g(s, t)u(\tau_1(s), \tau_2(t))] \Delta s \Delta t, \quad (2.69)$$

with the initial condition

$$\begin{aligned} u(x, y) &= \phi(x, y), \quad \text{if } x \in [\alpha, x_0] \cap \mathbb{T} \text{ or } y \in [\beta, y_0] \cap \mathbb{T}, \\ \phi(\tau_1(x), \tau_2(y)) &\leq C, \quad \forall (x, y) \in (\mathbb{T}_0 \times \tilde{\mathbb{T}}_0), \text{ if } \tau_1(x) \leq x_0 \text{ or } \tau_2(y) \leq y_0, \end{aligned} \quad (2.70)$$

then

$$u(x, y) \leq \tilde{G}^{-1} \left[\tilde{G}(C) + \int_{y_0}^y \eta_2(x, t) \Delta t \right] e_{B_1}(y, y_0), \quad (x, y) \in (\mathbb{T}_0 \times \tilde{\mathbb{T}}_0), \quad (2.71)$$

where $B_1(x, y) = \int_{x_0}^x g(s, y) \Delta s$, $\eta_2(x, y) = \omega(e_{B_1}(y, y_0)) \int_{x_0}^x f(s, y) \Delta s$ and $e_{B_1}(y, y_0)$ is the unique solution of the following equation:

$$z_y^\Delta(x, y) = B_1(x, y)z(x, y), \quad z(x, y_0) = 1. \quad (2.72)$$

Proof. Fix $X \in \mathbb{T}_0$, and $x \in [x_0, X] \cap \mathbb{T}$, $y \in \tilde{\mathbb{T}}_0$. Let

$$v(x, y) = C + \int_{y_0}^y \int_{x_0}^x [f(s, t)\omega(u(\tau_1(s), \tau_2(t))) + g(s, t)u(\tau_1(s), \tau_2(t))] \Delta s \Delta t. \quad (2.73)$$

Then

$$u(x, y) \leq v(x, y) \leq v(X, y), \quad \forall x \in [x_0, X] \cap \mathbb{T}, \quad y \in \tilde{\mathbb{T}}_0. \quad (2.74)$$

Similar to (2.14)–(2.16), we can obtain

$$u(\tau_1(x), \tau_2(y)) \leq v(x, y), \quad x \in [x_0, X] \cap \mathbb{T}, \quad y \in \tilde{\mathbb{T}}_0. \quad (2.75)$$

Furthermore we have

$$\begin{aligned} v(X, y) &= C + \int_{y_0}^y \int_{x_0}^X [f(s, t)\omega(u(\tau_1(s), \tau_2(t))) + g(s, t)u(\tau_1(s), \tau_2(t))] \Delta s \Delta t \\ &\leq C + \int_{y_0}^y \int_{x_0}^X [f(s, t)\omega(v(s, t)) + g(s, t)v(s, t)] \Delta s \Delta t \\ &\leq C + \int_{y_0}^y \int_{x_0}^X f(s, t)\omega(v(s, t)) \Delta s \Delta t + \int_{y_0}^y \left(\int_{x_0}^X g(s, t) \Delta s \right) v(X, t) \Delta t, \quad y \in \tilde{\mathbb{T}}_0. \end{aligned} \quad (2.76)$$

Let $B_2(X, y) = C + \int_{y_0}^y \int_{x_0}^X f(s, t)\omega(v(s, t)) \Delta s \Delta t$. Then from (2.76) it follows that

$$v(X, y) \leq B_2(X, y) + \int_{y_0}^y B_1(X, t)v(X, t) \Delta t, \quad y \in \tilde{\mathbb{T}}_0. \quad (2.77)$$

Considering $B_2(X, y)$ is nondecreasing in y , by applying Lemma 2.2 to (2.77), we obtain

$$v(X, y) \leq B_2(X, y)e_{B_1}(y, y_0), \quad y \in \tilde{\mathbb{T}}_0. \quad (2.78)$$

On the other hand,

$$\begin{aligned} [B_2(X, y)]_y^\Delta &= \int_{x_0}^X [f(s, y)\omega(v(s, y))] \Delta s \leq \left[\int_{x_0}^X f(s, y) \Delta s \right] \omega(v(X, y)) \\ &\leq \left[\int_{x_0}^X f(s, y) \Delta s \right] \omega[B_2(X, y)e_{B_1}(y, y_0)] \\ &\leq \left[\int_{x_0}^X f(s, y) \Delta s \right] \omega(B_2(X, y))\omega(e_{B_1}(y, y_0)) \\ &= \omega(B_2(X, y))\eta_2(X, y). \end{aligned} \quad (2.79)$$

Let $\bar{v}(X, y)$ be the solution of the following equation:

$$\bar{v}_y^\Delta(X, y) = \eta_2(X, y)\omega(\bar{v}(X, y)), \quad \bar{v}(X, y_0) = C. \quad (2.80)$$

Considering $B_2(X, y_0) = C$ and ω is nondecreasing and continuous, then from (2.79), (2.80), and Lemma 2.4, we have

$$B_2(X, y) \leq \bar{v}(X, y), \quad y \in \tilde{\mathbb{T}}_0. \quad (2.81)$$

From the definition of \tilde{G} and (2.80), we have $(\tilde{G}(\bar{v}(X, y)))_y^\Delta = \bar{v}_y^\Delta(X, y) / \omega(\bar{v}(X, y)) = \eta_2(X, y)$. Then similar to (2.66) and (2.67), we obtain

$$B_2(X, y) \leq \bar{v}(X, y) \leq \tilde{G}^{-1} \left[\tilde{G}(C) + \int_{y_0}^y \eta_2(X, t) \Delta t \right], \quad y \in \tilde{\mathbb{T}}_0. \quad (2.82)$$

Combining (2.74), (2.78), and (2.82), we have

$$u(x, y) \leq \tilde{G}^{-1} \left[\tilde{G}(C) + \int_{y_0}^y \eta_2(X, t) \Delta t \right] e_{B_1}(y, y_0), \quad x \in [x_0, X] \cap \mathbb{T}, \quad y \in \tilde{\mathbb{T}}_0. \quad (2.83)$$

Setting $x = X$ in (2.83), we obtain

$$u(X, y) \leq \tilde{G}^{-1} \left[\tilde{G}(C) + \int_{y_0}^y \eta_2(X, t) \Delta t \right] e_{B_1}(y, y_0), \quad y \in \tilde{\mathbb{T}}_0. \quad (2.84)$$

Replacing X with x in (2.84) yields the desired inequality (2.71). \square

Theorem 2.17. *Under the conditions of Theorem 2.16, if p, C are constants with $p > 0, C > 0$, and for $(x, y) \in \mathbb{T}_0 \times \tilde{\mathbb{T}}_0$, $u(x, y)$ satisfies the following inequality:*

$$u^p(x, y) \leq C + \int_{y_0}^y \int_{x_0}^x [f(s, t) \omega(u(\tau_1(s), \tau_2(t))) + g(s, t) u^p(\tau_1(s), \tau_2(t))] \Delta s \Delta t, \quad (2.85)$$

with the initial condition (2.58), then

$$u(x, y) \leq \left\{ G^{-1} \left[G(C) + \int_{y_0}^y \eta_3(x, t) \Delta t \right] e_{J_1}(y, y_0) \right\}^{1/p}, \quad (x, y) \in \mathbb{T}_0 \times \tilde{\mathbb{T}}_0, \quad (2.86)$$

where G is defined as in Theorem 2.14, $J_1(x, y) = \int_{x_0}^x g(s, y) \Delta s$, $\eta_3(x, y) = \omega((e_{J_1}(y, y_0))^{1/p}) \int_{x_0}^x f(s, y) \Delta s$, and $e_{J_1}(y, y_0)$ is the unique solution of the following equation:

$$z_y^\Delta(x, y) = J_1(x, y) z(x, y), \quad z(x, y_0) = 1. \quad (2.87)$$

The proof of Theorem 2.17 is similar to that of Theorem 2.16, and we omit it here.

3. Some Simple Applications

In this section, we will present some examples to illustrate the validity of our results in deriving explicit bounds of solutions of certain delay dynamic equations on time scales.

Example 3.1. Consider the following delay dynamic integral equation:

$$u^p(x, y) = C + \int_{y_0}^y \int_{x_0}^x M[s, t, u(\tau_1(s), \tau_2(t))] \Delta s \Delta t, \quad (x, y) \in \mathbb{T}_0 \times \tilde{\mathbb{T}}_0, \quad (3.1)$$

with the initial condition

$$\begin{aligned} u(x, y) &= \phi(x, y), \quad \text{if } x \in [\alpha, x_0] \cap \mathbb{T} \text{ or } y \in [\beta, y_0] \cap \mathbb{T}, \\ \phi(\tau_1(x), \tau_2(y)) &\leq |C|^{1/p}, \quad \forall (x, y) \in (\mathbb{T}_0, \tilde{\mathbb{T}}_0), \text{ if } \tau_1(x) \leq x_0 \text{ or } \tau_2(y) \leq y_0, \end{aligned} \quad (3.2)$$

where $u \in C_{\text{rd}}(\mathbb{T}_0 \times \tilde{\mathbb{T}}_0, \mathbb{R})$, $\phi, \alpha, \beta, \tau_1, \tau_2$ are the same as in Theorem 2.8, and $M \in C_{\text{rd}}(\mathbb{T}_0 \times \tilde{\mathbb{T}}_0 \times \mathbb{R}, \mathbb{R})$. Furthermore, assume $|M(s, t, u)| \leq f(s, t)|u|^q + g(s, t)|u|^r$, where $f, g \in C_{\text{rd}}(\mathbb{T}_0 \times \tilde{\mathbb{T}}_0, \mathbb{R}_+)$, and p, q, r are the same as in Theorem 2.8.

From (3.1) we have

$$\begin{aligned} |u^p(x, y)| &\leq |C| + \int_{y_0}^y \int_{x_0}^x |M[s, t, u(\tau_1(s), \tau_2(t))]| \Delta s \Delta t \\ &\leq |C| + \int_{y_0}^y \int_{x_0}^x [f(s, t)|u(\tau_1(s), \tau_2(t))|^q + g(s, t)|u(\tau_1(s), \tau_2(t))|^r] \Delta s \Delta t. \end{aligned} \quad (3.3)$$

Then according to Theorem 2.8, we can obtain the following estimate:

$$|u(x, y)| \leq \left[\tilde{H}_1(x, y) + \int_{y_0}^y e_{\tilde{H}_2}(y, \sigma(t)) \tilde{H}_2(x, t) \tilde{H}_1(x, t) \Delta t \right]^{1/p}, \quad (x, y) \in \mathbb{T}_0 \times \tilde{\mathbb{T}}_0, \quad (3.4)$$

where

$$\begin{aligned} \tilde{H}_1(x, y) &= |C| + \int_{y_0}^y \int_{x_0}^x \left[f(s, t) \frac{p-q}{p} K^{q/p} + g(s, t) \frac{p-r}{p} K^{r/p} \right] \Delta s \Delta t, \quad \forall K > 0, \\ \tilde{H}_2(x, y) &= \int_{x_0}^x \left[f(s, y) \frac{q}{p} K^{(q-p)/p} + g(s, y) \frac{r}{p} K^{(r-p)/p} \right] \Delta s, \quad \forall K > 0. \end{aligned} \quad (3.5)$$

Example 3.2. Considering the following delay dynamic integral equation:

$$u^3(x, y) = C + \int_{y_0}^y \int_{x_0}^x N[s, t, u(\tau_1(s), \tau_2(t)), u(\tau_1(s), \sigma(\tau_2(t)))] \Delta s \Delta t, \quad (x, y) \in \mathbb{T}_0 \times \tilde{\mathbb{T}}_0, \quad (3.6)$$

with the initial condition

$$\begin{aligned} u(x, y) &= \phi(x, y), \quad \text{if } x \in [\alpha, x_0] \cap \mathbb{T} \text{ or } y \in [\beta, y_0] \cap \mathbb{T}; \\ \phi(\tau_1(x), \tau_2(y)) &\leq |C|^{1/3}, \quad \forall x \in \mathbb{T}_0, y \in \tilde{\mathbb{T}}_0, \text{ if } \tau_1(x) \leq x_0 \text{ or } \tau_2(y) \leq y_0, \end{aligned} \quad (3.7)$$

where $u \in C_{\text{rd}}(\mathbb{T}_0 \times \tilde{\mathbb{T}}_0, \mathbb{R})$, and $N \in C_{\text{rd}}(\mathbb{T}_0 \times \tilde{\mathbb{T}}_0 \times \mathbb{R}^2, \mathbb{R})$.

Assume $|N(x, y, u, v)| \leq f(x, y)(|u|^2 + |v|^2)$, where $f \in C_{rd}(\mathbb{T}_0 \times \tilde{\mathbb{T}}_0, \mathbb{R}_+)$, then from (3.6) we have

$$\begin{aligned} |u^3(x, y)| &\leq |C| + \int_{y_0}^y \int_{x_0}^x |N[s, t, u(\tau_1(s), \tau_2(t)), u(\tau_1(s), \sigma(\tau_2(t)))]| \Delta s \Delta t \\ &\leq |C| + \int_{y_0}^y \int_{x_0}^x f(s, t) \left[|u(\tau_1(s), \tau_2(t))|^2 + |u(\tau_1(s), \sigma(\tau_2(t)))|^2 \right] \Delta s \Delta t \\ &\leq |C| + \int_{y_0}^y \int_{x_0}^x f(s, t) \left[|u(\tau_1(s), \tau_2(t))|^2 + |u(\tau_1(s), \sigma(\tau_2(t)))|^2 \right. \\ &\quad \left. + |u(\tau_1(s), \tau_2(t))||u(\tau_1(s), \sigma(\tau_2(t)))| \right] \Delta s \Delta t. \end{aligned} \quad (3.8)$$

According to Theorem 2.13 ($p = 3$), we can reach the following estimate:

$$|u(x, y)| \leq |C|^{1/3} + \int_{y_0}^y \int_{x_0}^x f(s, t) \Delta s \Delta t, \quad (x, y) \in \mathbb{T}_0 \times \tilde{\mathbb{T}}_0. \quad (3.9)$$

4. Conclusions

In this paper, we established some new Gronwall-Bellman-type delay integral inequalities in two independent variables on time scales. As one can see, the presented results provide a handy tool for deriving bounds for solutions of certain delay dynamic equations on time scales. Furthermore, the process of constructing Theorems 2.5, 2.8, 2.14, 2.16 and 2.17 can be applied to the situation with n independent variables.

Acknowledgments

This work is supported by the Natural Science Foundation of Shandong Province (ZR2010AZ003) (China). The authors thank the referees very much for their careful comments and valuable suggestions on this paper.

References

- [1] T. H. Gronwall, "Note on the derivatives with respect to a parameter of the solutions of a system of differential equations," *Annals of Mathematics*, vol. 20, no. 4, pp. 292–296, 1919.
- [2] R. Bellman, "The stability of solutions of linear differential equations," *Duke Mathematical Journal*, vol. 10, pp. 643–647, 1943.
- [3] B. G. Pachpatte, *Inequalities for Differential and Integral Equations*, vol. 197 of *Mathematics in Science and Engineering*, Academic Press Inc., San Diego, Calif, USA, 1998.
- [4] W.-S. Cheung, "Some new nonlinear inequalities and applications to boundary value problems," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 64, no. 9, pp. 2112–2128, 2006.
- [5] O. Lipovan, "Integral inequalities for retarded Volterra equations," *Journal of Mathematical Analysis and Applications*, vol. 322, no. 1, pp. 349–358, 2006.
- [6] E. Yang, "On some nonlinear integral and discrete inequalities related to Ou-Iang's inequality," *Acta Mathematica Sinica*, vol. 14, no. 3, pp. 353–360, 1998.
- [7] Q.-H. Ma, "Estimates on some power nonlinear Volterra-Fredholm type discrete inequalities and their applications," *Journal of Computational and Applied Mathematics*, vol. 233, no. 9, pp. 2170–2180, 2010.

- [8] R. P. Agarwal, S. Deng, and W. Zhang, "Generalization of a retarded Gronwall-like inequality and its applications," *Applied Mathematics and Computation*, vol. 165, no. 3, pp. 599–612, 2005.
- [9] B. G. Pachpatte, "Inequalities applicable in the theory of finite difference equations," *Journal of Mathematical Analysis and Applications*, vol. 222, no. 2, pp. 438–459, 1998.
- [10] R. A. C. Ferreira and D. F. M. Torres, "Generalized retarded integral inequalities," *Applied Mathematics Letters*, vol. 22, no. 6, pp. 876–881, 2009.
- [11] F. Jiang and F. Meng, "Explicit bounds on some new nonlinear integral inequalities with delay," *Journal of Computational and Applied Mathematics*, vol. 205, no. 1, pp. 479–486, 2007.
- [12] H. Zhang and F. Meng, "Integral inequalities in two independent variables for retarded Volterra equations," *Applied Mathematics and Computation*, vol. 199, no. 1, pp. 90–98, 2008.
- [13] Ou Yang-Liang, "The boundedness of solutions of linear differential equations $y'' + A(t)y = 0$," *Advances in Mathematics*, vol. 3, pp. 409–415, 1957.
- [14] W. S. Cheung, "Some new nonlinear inequalities and applications to boundary value problems," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 64, no. 9, pp. 2112–2128, 2006.
- [15] W. S. Cheung and J. Ren, "Discrete non-linear inequalities and applications to boundary value problems," *Journal of Mathematical Analysis and Applications*, vol. 319, no. 2, pp. 708–724, 2006.
- [16] S. Hilger, "Analysis on measure chains—a unified approach to continuous and discrete calculus," *Results in Mathematics*, vol. 18, no. 1-2, pp. 18–56, 1990.
- [17] M. Bohner, L. Erbe, and A. Peterson, "Oscillation for nonlinear second order dynamic equations on a time scale," *Journal of Mathematical Analysis and Applications*, vol. 301, no. 2, pp. 491–507, 2005.
- [18] Y. Xing, M. Han, and G. Zheng, "Initial value problem for first-order integro-differential equation of Volterra type on time scales," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 60, no. 3, pp. 429–442, 2005.
- [19] R. Agarwal, M. Bohner, D. O'Regan, and A. Peterson, "Dynamic equations on time scales: a survey," *Journal of Computational and Applied Mathematics*, vol. 141, no. 1-2, pp. 1–26, 2002.
- [20] F.-H. Wong, C.-C. Yeh, and W.-C. Lian, "An extension of Jensen's inequality on time scales," *Advances in Dynamical Systems and Applications*, vol. 1, no. 1, pp. 113–120, 2006.
- [21] X.-L. Cheng, "Improvement of some Ostrowski-Grüss type inequalities," *Computers & Mathematics with Applications*, vol. 42, no. 1-2, pp. 109–114, 2001.
- [22] M. Bohner and T. Matthews, "The Grüss inequality on time scales," *Communications in Mathematical Analysis*, vol. 3, no. 1, pp. 1–8, 2007.
- [23] Q.-A. Ngô, "Some mean value theorems for integrals on time scales," *Applied Mathematics and Computation*, vol. 213, no. 2, pp. 322–328, 2009.
- [24] W. Liu and Q.-A. Ngô, "Some Iyengar-type inequalities on time scales for functions whose second derivatives are bounded," *Applied Mathematics and Computation*, vol. 216, no. 11, pp. 3244–3251, 2010.
- [25] H. M. Srivastava, K.-L. Tseng, S.-J. Tseng, and J.-C. Lo, "Some generalizations of Maroni's inequality on time scales," *Mathematical Inequalities & Applications*, vol. 14, no. 2, pp. 469–480, 2011.
- [26] R. Agarwal, M. Bohner, and A. Peterson, "Inequalities on time scales: a survey," *Mathematical Inequalities & Applications*, vol. 4, no. 4, pp. 535–557, 2001.
- [27] W. N. Li, "Some delay integral inequalities on time scales," *Computers & Mathematics with Applications*, vol. 59, no. 6, pp. 1929–1936, 2010.
- [28] Q.-H. Ma and J. Pečarić, "The bounds on the solutions of certain two-dimensional delay dynamic systems on time scales," *Computers & Mathematics with Applications*, vol. 61, no. 8, pp. 2158–2163, 2011.
- [29] M. Bohner and A. Peterson, *Dynamic Equations on Time Scales: An Introduction with Application*, Birkhäuser Boston, Boston, Mass, USA, 2001.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

