

Research Article

Eigenvalue Comparisons for Second-Order Linear Equations with Boundary Value Conditions on Time Scales

Chao Zhang and Shurong Sun

School of Mathematical Sciences, University of Jinan, Jinan, Shandong 250022, China

Correspondence should be addressed to Chao Zhang, zmaths@163.com

Received 29 January 2012; Revised 22 March 2012; Accepted 22 March 2012

Academic Editor: Kai Diethelm

Copyright © 2012 C. Zhang and S. Sun. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

This paper studies the eigenvalue comparisons for second-order linear equations with boundary conditions on time scales. Using results from matrix algebras, the existence and comparison results concerning eigenvalues are obtained.

1. Introduction

In this paper, we consider the eigenvalue problems for the following second-order linear equations:

$$\left(r(t)y^\Delta(t)\right)^\Delta + \lambda^{(1)}p(t)y^\sigma(t) = 0, \quad t \in [\rho(a), \rho(b)]_{\mathbb{T}}, \quad (1.1)$$

$$\left(r(t)y^\Delta(t)\right)^\Delta + \lambda^{(2)}q(t)y^\sigma(t) = 0, \quad t \in [\rho(a), \rho(b)]_{\mathbb{T}}, \quad (1.2)$$

with the boundary conditions

$$y(\rho(a)) - \tau y(a) = y(\sigma(b)) - \delta y(b) = 0, \quad (1.3)$$

where $\lambda^{(1)}$ and $\lambda^{(2)}$ are parameters, $\sigma(t)$ and $\rho(t)$ are the forward and backward jump operators, y^Δ is the delta derivative, $y^\sigma(t) := y(\sigma(t))$, and $[\rho(a), \rho(b)]_{\mathbb{T}}$ is a finite isolated time scale; the discrete interval is given by

$$[\rho(a), \rho(b)]_{\mathbb{T}} := \{\rho(a), a, \sigma(a), \sigma^2(a), \dots, \rho(b)\}. \quad (1.4)$$

We assume throughout this paper that

- (H₁) r^Δ , p , and q are real-valued functions on $[\rho(a), \rho(b)]_{\mathbb{T}}$, $p \geq 0$ ($\neq 0$), $q \geq 0$ ($\neq 0$) on $[\rho(a), \rho(b)]_{\mathbb{T}}$ and $r > 0$ on $[\rho(a), b]_{\mathbb{T}}$;
 (H₂) $\tau, \delta \in [0, 1)$.

First we briefly recall some existing results of eigenvalues comparisons for differential and difference equations. In 1973, Travis [1] considered the eigenvalue problem for boundary value problems of higher-order differential equations. He employed the theory of u_0 -positive linear operator on a Banach space with a cone of nonnegative elements to obtain comparison results for the smallest eigenvalues. A representative set of references for these works would be Davis et al. [2], Diaz and Peterson [3], Hankerson and Henderson [4], Hankerson and Peterson [5–7], Henderson and Prasad [8], and Kaufmann [9]. However, in all the above papers, the comparison results are for the smallest eigenvalues only. The main purpose of this paper is to establish the comparison theorems for all the eigenvalues of (1.1) with (1.3) and (1.2) with (1.3).

Like the eigenvalue comparison for the boundary value problems of linear equations, this type of comparison of eigenvalues in matrix algebra is known as Weyl's inequality [10, Corollary 6.5.]: If A, B are Hermitian matrices, that is, $A = A^*$, where A^* is the conjugate transpose of A and $A - B$ is positive semidefinite, then $\lambda_i^{(A)} \geq \lambda_i^{(B)}$, where $\lambda_i^{(A)}$ and $\lambda_i^{(B)}$ are all eigenvalues of A and B . Associated with this conclusion is spectral order of operators. The spectral order has proved to be useful for solving several open problems of spectral theory and has been studied in the context of von Neumann algebras, matrix algebras, and so forth in [10–15]. Recently, Hamhalter [15] studied the spectral order in a more general setting of Jordan operator algebras, which is a generalization of the result due to Kato [13]. And as a preparatory material, he extended Olson's characterization of the spectral order to JBW algebras [14]. Since the boundary value problems (1.1), (1.3) and (1.2), (1.3) can be rewritten into matrix equations, we employ some results from matrix algebras to establish the comparison theorems for the eigenvalues of (1.1), (1.3) and (1.2), (1.3).

This paper is organized as follows. Section 2 introduces some basic concepts and a fundamental theory about time scales, which will be used in Section 3. By some results from matrix algebras and time scales, the existence and comparison theorems of eigenvalues of boundary value problems (1.1), (1.3) and (1.2), (1.3) are obtained, which will be given in Section 3.

2. Preliminaries

In this section, some basic concepts and some fundamental results on time scales are introduced.

Let $\mathbb{T} \subset \mathbf{R}$ be a nonempty closed subset. Define the forward and backward jump operators $\sigma, \rho : \mathbb{T} \rightarrow \mathbb{T}$ by

$$\sigma(t) = \inf\{s \in \mathbb{T} : s > t\}, \quad \rho(t) = \sup\{s \in \mathbb{T} : s < t\}, \quad (2.1)$$

where $\inf \emptyset = \sup \mathbb{T}$, $\sup \emptyset = \inf \mathbb{T}$. We put $\mathbb{T}^k = \mathbb{T}$ if \mathbb{T} is unbounded above and $\mathbb{T}^k = \mathbb{T} \setminus (\rho(\max \mathbb{T}), \max \mathbb{T}]$ otherwise. The graininess functions $\nu, \mu : \mathbb{T} \rightarrow [0, \infty)$ are defined by

$$\mu(t) = \sigma(t) - t, \quad \nu(t) = t - \rho(t). \quad (2.2)$$

Let f be a function defined on \mathbb{T} . f is said to be (delta) differentiable at $t \in \mathbb{T}^k$ provided there exists a constant a such that for any $\varepsilon > 0$, there is a neighborhood U of t (i.e., $U = (t - \delta, t + \delta) \cap \mathbb{T}$ for some $\delta > 0$) with

$$|f(\sigma(t)) - f(s) - a(\sigma(t) - s)| \leq \varepsilon |\sigma(t) - s|, \quad \forall s \in U. \quad (2.3)$$

In this case, denote $f^\Delta(t) := a$. If f is (delta) differentiable for every $t \in \mathbb{T}^k$, then f is said to be (delta) differentiable on \mathbb{T} . If f is differentiable at $t \in \mathbb{T}^k$, then

$$f^\Delta(t) = \begin{cases} \lim_{\substack{s \rightarrow t \\ s \in \mathbb{T}}} \frac{f(t) - f(s)}{t - s} & \text{if } \mu(t) = 0, \\ \frac{f(\sigma(t)) - f(t)}{\mu(t)} & \text{if } \mu(t) > 0. \end{cases} \quad (2.4)$$

For convenience, we introduce the following results ([16, Chapter 1], [17, Chapter 1], and [18, Lemma 1]), which are useful in this paper.

Lemma 2.1. Let $f, g : \mathbb{T} \rightarrow \mathbf{R}$ and $t \in \mathbb{T}^k$.

(i) If f and g are differentiable at t , then fg is differentiable at t and

$$(fg)^\Delta(t) = f^\sigma(t)g^\Delta(t) + f^\Delta(t)g(t) = f^\Delta(t)g^\sigma(t) + f(t)g^\Delta(t). \quad (2.5)$$

(ii) If f and g are differentiable at t , and $f(t)f^\sigma(t) \neq 0$, then $f^{-1}g$ is differentiable at t and

$$(gf^{-1})^\Delta(t) = (g^\Delta(t)f(t) - g(t)f^\Delta(t))(f^\sigma(t)f(t))^{-1}. \quad (2.6)$$

3. Eigenvalue Comparisons

In the following, we will write $X \geq Y$ if X and Y are symmetric $n \times n$ matrices and $X - Y$ is positive semidefinite. A matrix is said to be positive if every component of the matrix is positive. We denote $\rho(a) = \sigma^{-1}(a)$, $a = \sigma^0(a)$, $\rho(b) = \sigma^{n-2}(a)$, $b = \sigma^{n-1}(a)$, $\mu_i = \sigma^{i+1}(a) - \sigma^i(a)$, and $r^{\sigma^i}(a) = r(\sigma^i(a))$, $i = -1, 0, 1, 2, \dots, n-1$.

It follows from Lemma 2.1(ii), (2.4), and (1.4) that the boundary value problem (1.1), (1.3) can be written in the form

$$(-D + \lambda^{(1)}P)y = 0, \quad (3.1)$$

where

$$D := \begin{pmatrix} \mathcal{A} + \mathcal{B} & -\mathcal{B} & 0 & \cdots & 0 & 0 & 0 \\ -\mathcal{B} & \mathcal{B} + \mathcal{C} & -\mathcal{C} & \cdots & 0 & 0 & 0 \\ 0 & -\mathcal{C} & \mathcal{C} + \frac{r^{\sigma^2}(a)}{\mu_2} & \cdots & 0 & 0 & 0 \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ 0 & 0 & 0 & \cdots & \frac{r^{\sigma^{n-4}}(a)}{\mu_{n-4}} + \mathfrak{D} & -\mathfrak{D} & 0 \\ 0 & 0 & 0 & \cdots & -\mathfrak{D} & \mathfrak{D} + \mathcal{E} & -\mathcal{E} \\ 0 & 0 & 0 & \cdots & 0 & -\mathcal{E} & \mathcal{E} + \frac{(1-\delta)r^{\sigma^{n-1}}(a)}{\mu_{n-1}} \end{pmatrix} \quad (3.2)$$

$$P = \text{diag}(\mu_{-1}p(\sigma^{-1}(a)), \mu_0p(\sigma^0(a)), \dots, \mu_{n-3}p(\sigma^{n-3}(a)), \mu_{n-2}p(\sigma^{n-2}(a))),$$

where \mathcal{A} donates $(1 - \tau)r^{\sigma^{-1}}(a)/\mu_{-1}$, \mathcal{B} donates $r^{\sigma^0}(a)/\mu_0$, \mathcal{C} donates $r^{\sigma}(a)/\mu_1$, \mathfrak{D} donates $r^{\sigma^{n-3}}(a)/\mu_{n-3}$, and \mathcal{E} donates $r^{\sigma^{n-2}}(a)/\mu_{n-2}$.

$$y = (y(\sigma^0(a)), y(\sigma(a)), \dots, y(\sigma^{n-2}(a)), y(\sigma^{n-1}(a)))^T. \quad (3.3)$$

And the problem (1.2), (1.3) is equivalent to the equation

$$(-D + \lambda^{(2)}Q)y = 0, \quad (3.4)$$

where

$$Q = \text{diag}(\mu_{-1}q(\sigma^{-1}(a)), \mu_0q(\sigma^0(a)), \dots, \mu_{n-3}q(\sigma^{n-3}(a)), \mu_{n-2}q(\sigma^{n-2}(a))). \quad (3.5)$$

Since the solutions of (1.1), (1.3) can be written into the form of vectors, then the nontrivial solution corresponding to λ is called an eigenvector.

Let e_i be the i th column of the identity matrix I of order n and

$$D_1 := \begin{pmatrix} \mathcal{A} + \mathcal{B} & -\mathcal{B} & 0 & \cdots & 0 & 0 & 0 \\ -\mathcal{B} & \mathcal{B} + \mathcal{C} & -\mathcal{C} & \cdots & 0 & 0 & 0 \\ 0 & -\mathcal{C} & \mathcal{C} + \frac{r^{\sigma^2}(a)}{\mu_2} & \cdots & 0 & 0 & 0 \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ 0 & 0 & 0 & \cdots & \frac{r^{\sigma^{n-4}}(a)}{\mu_{n-4}} + \mathfrak{D} & -\mathfrak{D} & 0 \\ 0 & 0 & 0 & \cdots & -\mathfrak{D} & \mathfrak{D} + \mathfrak{E} & -\mathfrak{E} \\ 0 & 0 & 0 & \cdots & 0 & -\mathfrak{E} & \mathfrak{E} \end{pmatrix} \quad (3.6)$$

Define $P_i = I + e_{i-1}e_i^T$. It is easily seen that

$$D = D_1 + e_n(1 - \delta) \frac{r^{\sigma^{n-1}}(a)}{\mu_{n-1}} e_n^T, \quad (3.7)$$

$$P_2 P_3 \cdots P_n D_1 P_n^T \cdots P_3^T P_2^T = \text{diag} \left((1 - \tau) \frac{r^{\sigma^{-1}}(a)}{\mu_{-1}}, \frac{r^{\sigma^0}(a)}{\mu_0}, \dots, \frac{r^{\sigma^{n-3}}(a)}{\mu_{n-3}}, \frac{r^{\sigma^{n-2}}(a)}{\mu_{n-2}} \right). \quad (3.8)$$

It follows from (H₁), (H₂), and (3.8) that

$$D_1 = P_n^{-1} \cdots P_3^{-1} P_2^{-1} \text{diag} \left((1 - \tau) \frac{r^{\sigma^{-1}}(a)}{\mu_{-1}}, \frac{r^{\sigma^0}(a)}{\mu_0}, \dots, \frac{r^{\sigma^{n-3}}(a)}{\mu_{n-3}}, \frac{r^{\sigma^{n-2}}(a)}{\mu_{n-2}} \right) P_2^{-T} P_3^{-T} \cdots P_n^{-T}, \quad (3.9)$$

$$D_1^{-1} = P_n^T \cdots P_3^T P_2^T \text{diag} \left(\frac{\mu_{-1}}{(1 - \tau)r^{\sigma^{-1}}(a)}, \frac{\mu_0}{r^{\sigma^0}(a)}, \dots, \frac{\mu_{n-3}}{r^{\sigma^{n-3}}(a)}, \frac{\mu_{n-2}}{r^{\sigma^{n-2}}(a)} \right) P_2 P_3 \cdots P_n. \quad (3.10)$$

For any $x = (x_1, x_2, \dots, x_n)^T$, we have

$$\begin{aligned} x^* D x &= x^* D_1 x + (1 - \delta) \frac{r^{\sigma^{n-1}}(a)}{\mu_{n-1}} x^* e_n e_n^T x \\ &= \left(P_2^{-T} P_3^{-T} \cdots P_n^{-T} x \right)^* \text{diag} \left((1 - \tau) \frac{r^{\sigma^{-1}}(a)}{\mu_{-1}}, \frac{r^{\sigma^0}(a)}{\mu_0}, \dots, \frac{r^{\sigma^{n-3}}(a)}{\mu_{n-3}}, \frac{r^{\sigma^{n-2}}(a)}{\mu_{n-2}} \right) \\ &\quad \times \left(P_2^{-T} P_3^{-T} \cdots P_n^{-T} x \right) + (1 - \delta) \frac{r^{\sigma^{n-1}}(a)}{\mu_{n-1}} \left| e_n^T x \right|^2 \geq 0. \end{aligned} \quad (3.11)$$

Moreover, $x^* D x = 0$ implies $x = 0$. Hence, the matrix D is positive definite.

Lemma 3.1. *If $\lambda^{(1)}$ is an eigenvalue of the boundary value problem (1.1), (1.3) and y is a corresponding eigenvector, then*

- (i) $y^*Py > 0$,
- (ii) $\lambda^{(1)}$ is real and positive.

*If $\rho \neq \lambda^{(1)}$ is an eigenvalue of the boundary value problem (1.1), (1.3) and x is a corresponding eigenvector, then $x^*Py = 0$.*

Proof. (i) It follows from (H_1) and (3.2) that $y^*Py \geq 0$. Assume the contrary that $y^*Py = 0$, we have $y^*Dy = \lambda^{(1)}y^*Py = 0$. Since D is positive definite, then $y = 0$, which is a contradiction.

(ii) We can write

$$\lambda^{(1)}y^*Py = y^*(\lambda^{(1)}Py) = y^*Dy = (Dy)^*y = (\lambda^{(1)}Py)^*y = \overline{\lambda^{(1)}}y^*P^*y = \overline{\lambda^{(1)}}y^*Py, \quad (3.12)$$

which implies $\lambda^{(1)} = \overline{\lambda^{(1)}}$, that is, λ is real. Since D is positive definite and $y^*Py > 0$, we have $\lambda^{(1)} = y^*Dy/y^*Py > 0$.

If $\rho Px = Dx$ and $\rho \neq \lambda^{(1)}$, then

$$(\lambda^{(1)} - \rho)x^*Py = \lambda^{(1)}x^*Py - \rho x^*Py = x^*(\lambda^{(1)}Py) - (\rho Px)^*y = x^*Dy - (Dx)^*y = 0. \quad (3.13)$$

Hence, $x^*Py = 0$. This completes the proof. \square

Lemma 3.2. *If $\lambda^{(1)}$ is an eigenvalue of the boundary value problem (1.1), (1.3), then $1/\lambda^{(1)}$ is an eigenvalue of $D^{-1/2}PD^{-1/2}$. If α is a positive eigenvalue of $D^{-1/2}PD^{-1/2}$, then $1/\alpha$ is an eigenvalue of (1.1), (1.3), respectively.*

Proof. If $\lambda^{(1)}$ is an eigenvalue of the boundary value problem (1.1), (1.3) and y is a corresponding eigenvector, then $\lambda^{(1)} > 0$ and $\lambda^{(1)}Py = Dy$. Therefore,

$$\begin{aligned} \lambda^{(1)}Py &= D^{1/2}D^{1/2}y, \\ D^{-1/2}PD^{-1/2}(D^{1/2}y) &= \frac{1}{\lambda^{(1)}}(D^{1/2}y). \end{aligned} \quad (3.14)$$

With a similar argument, one can get that if α is a positive eigenvalue of $D^{-1/2}PD^{-1/2}$, then $1/\alpha$ is an eigenvalue of (1.1), (1.3). This completes proof. \square

Lemma 3.3. *For any $1 \leq i, j \leq n$, define $\gamma = \min\{i, j\}$. We have*

- (i) $e_i^T D_1^{-1} e_j = \mu_{-1}/(1-\tau)r^{\sigma^{-1}}(a) + \sum_{k=0}^{\gamma-2} (\mu_k/r^{\sigma^k}(a))$;
- (ii) $e_i^T D^{-1} e_j = ((\mu_{-1}/(1-\tau)r^{\sigma^{-1}}(a) + \sum_{k=0}^{\gamma-2} (\mu_k/r^{\sigma^k}(a)))(\mu_{n-1}/(1-\delta)r^{\sigma^{n-1}}(a) + (\mu_{-1}/(1-\tau)r^{\sigma^{-1}}(a) + \sum_{k=0}^{n-2} (\mu_k/r^{\sigma^k}(a))) - ((\mu_{-1}/(1-\tau)r^{\sigma^{-1}}(a) + \sum_{k=0}^{i-2} (\mu_k/r^{\sigma^k}(a)))(\mu_{-1}/(1-\tau)r^{\sigma^{-1}}(a) + \sum_{k=0}^{j-2} (\mu_k/r^{\sigma^k}(a)))/(\mu_{n-1}/(1-\delta)r^{\sigma^{n-1}}(a) + (\mu_{-1}/(1-\tau)r^{\sigma^{-1}}(a) + \sum_{k=0}^{n-2} (\mu_k/r^{\sigma^k}(a)))$.

Proof. It is easy to see that $P_i e_j = e_j$ if $i \neq j$, while $P_i e_j = e_{j-1} + e_j$ if $i = j$. Hence,

$$P_2 P_3 \cdots P_n e_j = e_1 + e_2 + \cdots + e_j. \quad (3.15)$$

(i) It is seen from (3.10) and (3.15) that

$$\begin{aligned} e_i^T D_1^{-1} e_j &= (P_2 P_3 \cdots P_n e_i)^T \text{diag} \left(\frac{\mu_{-1}}{(1-\tau)r^{\sigma^{-1}}(a)}, \frac{\mu_0}{r^{\sigma^0}(a)}, \dots, \frac{\mu_{n-3}}{r^{\sigma^{n-3}}(a)}, \frac{\mu_{n-2}}{r^{\sigma^{n-2}}(a)} \right) (P_2 P_3 \cdots P_n e_j) \\ &= (e_1 + e_2 + \cdots + e_i)^T \text{diag} \left(\frac{\mu_{-1}}{(1-\tau)r^{\sigma^{-1}}(a)}, \frac{\mu_0}{r^{\sigma^0}(a)}, \dots, \frac{\mu_{n-3}}{r^{\sigma^{n-3}}(a)}, \frac{\mu_{n-2}}{r^{\sigma^{n-2}}(a)} \right) \\ &\quad \times (e_1 + e_2 + \cdots + e_j) \\ &= \frac{\mu_{-1}}{(1-\tau)r^{\sigma^{-1}}(a)} + \sum_{k=0}^{n-2} \frac{\mu_k}{r^{\sigma^k}(a)}. \end{aligned} \quad (3.16)$$

(ii) It follows from (3.7) and the Sherman-Morrison updating formula [19] that

$$D^{-1} = D_1^{-1} - \frac{D_1^{-1} e_n e_n^T D_1^{-1}}{(\mu_{n-1}/(1-\delta)r^{\sigma^{n-1}}(a)) + e_n^T D_1^{-1} e_n}, \quad (3.17)$$

leading to

$$e_i^T D^{-1} e_j = e_i^T D_1^{-1} e_j - \frac{e_i^T D_1^{-1} e_n e_n^T D_1^{-1} e_j}{(\mu_{n-1}/(1-\delta)r^{\sigma^{n-1}}(a)) + e_n^T D_1^{-1} e_n}, \quad (3.18)$$

which, together with (i), further implies the result (ii). This completes the proof. \square

Theorem 3.4. (i) If $\lambda^{(1)}$ is an eigenvalue of the boundary value problem (1.1), (1.3) and $y \neq 0$ is a corresponding eigenvector, then $y(a) \neq 0$ and $y(b) \neq 0$.

(ii) If $\lambda_1^{(1)} > 0$ is the smallest eigenvalue of the boundary value problem (1.1), (1.3), then there exists a positive eigenvector $y > 0$ corresponding to $\lambda_1^{(1)}$.

Proof. (i) Assume the contrary that either $y(a) = 0$ or $y(b) = 0$. By the boundary condition (1.3), we can easily deduce a contradiction $y(t) \equiv 0$.

(ii) It follows from $D^{-1} P y = (1/\lambda_1^{(1)}) y$ that $1/\lambda_1^{(1)}$ is the maximum eigenvalue of $D^{-1} P$ and the y is an eigenvector corresponding to $1/\lambda_1^{(1)}$. By Lemma 3.3(ii), we have that all the elements of D^{-1} are positive, then D^{-1} is a positive matrix. Since $p(t) \geq 0$ for all $t \in [\rho(a), \rho(b)]_{\mathbb{T}}$, hence, the following discussions are divided into two cases.

Case 1. If $p(t) > 0$ for all $t \in [\rho(a), \rho(b)]_{\mathbb{T}}$, then we obtain that the matrix $D^{-1}P$ is positive and therefore, the result follows from the Perron-Forbenius theorem [20].

Case 2. Let $p(t) = 0$ for some $t \in [\rho(a), \rho(b)]_{\mathbb{T}}$. Without loss of generality, we assume that $p(t) = 0$ for all $t \in [\rho(a), \sigma^{m-2}(a)]_{\mathbb{T}}$ and $p(t) > 0$ for all $t \in [\sigma^{m-1}(a), \rho(b)]_{\mathbb{T}}$; we can write $D^{-1}P$ as follows:

$$D^{-1}P = \begin{pmatrix} 0 & V \\ 0 & Z \end{pmatrix}, \quad (3.19)$$

where V is an $m \times (n - m)$ matrix and Z is an $(n - m) \times (n - m)$ matrix. Both V and Z are positive matrices. $1/\lambda_1^{(1)}$ is also the maximum eigenvalue of Z . Applying the Perron-Forbenius theorem to the positive matrix Z , there exists a positive vector $y_Z > 0$ such that $Zy_Z = (1/\lambda_1^{(1)})y_Z$. Let $y_V = \lambda_1^{(1)}Vy_Z$ and $y = (y_V^T, y_Z^T)^T$. Obviously, we have

$$D^{-1}Py = \frac{1}{\lambda_1^{(1)}}y, \quad \text{where } y > 0. \quad (3.20)$$

This completes the proof. \square

Lemma 3.5. *If $\lambda^{(1)}$ is an eigenvalue of the boundary value problem (1.1), (1.3), then the dimension of the null space of $(-D + \lambda^{(1)}P)$ is 1.*

Proof. Let $x \neq 0$ and $y \neq 0$ be any two eigenvectors of the boundary value problem (1.1), (1.3) corresponding to $\lambda^{(1)}$ and define $z = x(a)y - y(a)x$. Obviously, we have

$$(-D + \lambda^{(1)}P)z = x(a)(-D + \lambda^{(1)}P)y - y(a)(-D + \lambda^{(1)}P)x = 0, \quad (3.21)$$

which, together with $z(a) = 0$, indicates that $z = 0$, that is, $x(a)y = y(a)x$. Therefore, x and y are linearly dependent. So the dimension of the null space of $(-D + \lambda^{(1)}P)$ is 1. This completes the proof. \square

Lemma 3.6. *Let $N \geq 1$ be the number of positive elements in the set $\{p(t) \mid t \in [\rho(a), \rho(b)]_{\mathbb{T}}\}$. Then there are N distinct eigenvalues $\lambda_i^{(1)}$ ($i = 1, 2, \dots, N$) of the boundary value problem (1.1), (1.3) and $\alpha_i = 1/\lambda_i^{(1)}$ ($i = 1, 2, \dots, N$) are the only positive eigenvalues of $D^{-1/2}PD^{-1/2}$.*

Proof. Suppose that $\alpha_1 \geq \alpha_2 \geq \dots \geq \alpha_n \geq 0$ are all eigenvalues of $D^{-1/2}PD^{-1/2}$. Since $D^{-1/2}PD^{-1/2}$ is real and symmetric that there exists an orthogonal matrix C such that

$$C^T D^{-1/2} P D^{-1/2} C = \text{diag}(\alpha_1 \alpha_2 \cdots \alpha_n), \quad (3.22)$$

therefore, we have that

$$\text{rank}(P) = \text{rank}(C^T D^{-1/2} P D^{-1/2} C) = \text{rank}(\text{diag}(\alpha_1 \alpha_2 \cdots \alpha_n)) \quad (3.23)$$

indicating that the number of positive α_i is the same as that of positive number in P which is equal to N .

Suppose that $\alpha_{i_0} = \alpha_{i_0+1} > 0$ for some i_0 where $1 \leq i_0 \leq N - 1$. Observe that $C^T D^{-1/2} P D^{-1/2} C e_i = \alpha_i e_i$ in view of (3.22), which further implies that

$$D(D^{-1/2} C e_i) = \frac{1}{\alpha_i} P(D^{-1/2} C e_i) \quad i = i_0, \quad i_0 + 1. \quad (3.24)$$

Thus, we have two independent vectors in the null space of $(-D + \lambda^{(1)} P)$ for $\lambda^{(1)} = 1/\alpha_{i_0}$, which contradicts Lemma 3.5. Thus, from Lemma 3.2, we see that $\{\lambda_i^{(1)} = 1/\alpha_i \mid i = 1, 2, \dots, N\}$ gives the complete set of eigenvalues of the boundary value problem (1.1), (1.3). This completes the proof. \square

Theorem 3.7. *Let j be the number of positive elements in the set $\{p(t) \mid t \in [\rho(a), \rho(b)]_{\mathbb{T}}\}$ and k the number of positive elements in the set $\{q(t) \mid t \in [\rho(a), \rho(b)]_{\mathbb{T}}\}$. Let $\{\lambda_1^{(1)} < \lambda_2^{(1)} < \dots < \lambda_j^{(1)}\}$ be the set of all eigenvalues of the boundary value problem (1.1), (1.3) and $\{\lambda_1^{(2)} < \lambda_2^{(2)} < \dots < \lambda_k^{(2)}\}$ the set of all eigenvalues of the boundary value problem (1.2), (1.3). If $p(t) \geq q(t)$ for all $t \in [\rho(a), \rho(b)]_{\mathbb{T}}$, then $\lambda_i^{(1)} \leq \lambda_i^{(2)}$ for $1 \leq i \leq k$.*

Proof. It follows from Lemma 3.6 that

$$\begin{aligned} \alpha_1 = \frac{1}{\lambda_1^{(1)}} > \dots > \alpha_j = \frac{1}{\lambda_j^{(1)}} > 0, \quad \alpha_{j+1} = \dots = \alpha_n = 0, \\ \beta_1 = \frac{1}{\lambda_1^{(2)}} > \dots > \beta_k = \frac{1}{\lambda_k^{(2)}} > 0, \quad \beta_{k+1} = \dots = \alpha_n = 0 \end{aligned} \quad (3.25)$$

are the eigenvalues of $D^{-1/2} P D^{-1/2}$ and $D^{-1/2} Q D^{-1/2}$, respectively. If $p(t) \geq q(t)$ for all $t \in [\rho(a), \rho(b)]_{\mathbb{T}}$, then $P \geq Q$, implying

$$D^{-1/2} P D^{-1/2} \geq D^{-1/2} Q D^{-1/2}. \quad (3.26)$$

By Weyl's inequality and (3.26), we have

$$\alpha_i \geq \beta_i \quad 1 \leq i \leq n. \quad (3.27)$$

Finally, it is easily seen from (3.25) and (3.27) that

$$\frac{1}{\lambda_i^{(1)}} \geq \frac{1}{\lambda_i^{(2)}} \quad 1 \leq i \leq k, \quad (3.28)$$

implying that $\lambda_i^{(1)} \leq \lambda_i^{(2)}$ for $1 \leq i \leq k$. This completes the proof. \square



Acknowledgments

Many thanks are due to Kai Diethelm (the editor) and the anonymous reviewer(s) for helpful comments and suggestions. This paper was supported by the NNSF of China (Grants

nos. 11071143 and 11101241), the NNSF of Shandong Province (Grants nos. ZR2009AL003, ZR2010AL016, and ZR2011AL007), the Scientific Research and Development Project of Shandong Provincial Education Department (J11LA01), and the NSF of University of Jinan (XKY0918).

References

- [1] C. C. Travis, "Comparison of eigenvalues for linear differential equations of order $2n$," *Transactions of the American Mathematical Society*, vol. 177, pp. 363–374, 1973.
- [2] J. M. Davis, P. W. Eloe, and J. Henderson, "Comparison of eigenvalues for discrete Lidstone boundary value problems," *Dynamic Systems and Applications*, vol. 8, no. 3-4, pp. 381–388, 1999.
- [3] J. Diaz and A. Peterson, "Comparison theorems for a right disfocal eigenvalue problem," in *Inequalities and Applications*, vol. 3 of *World Scientific Series in Applicable Analysis*, pp. 149–177, World Scientific Publishing, River Edge, NJ, USA, 1994.
- [4] D. Hankerson and J. Henderson, "Comparison of eigenvalues for n -point boundary value problems for difference equations," in *Differential Equations (Colorado Springs, CO, 1989)*, vol. 127 of *Lecture Notes in Pure and Applied Mathematics*, pp. 203–208, Dekker, New York, NY, USA, 1991.
- [5] D. Hankerson and A. Peterson, "Comparison of eigenvalues for focal point problems for n th order difference equations," *Differential and Integral Equations*, vol. 3, no. 2, pp. 363–380, 1990.
- [6] D. Hankerson and A. Peterson, "A positivity result applied to difference equations," *Journal of Approximation Theory*, vol. 59, no. 1, pp. 76–86, 1989.
- [7] D. Hankerson and A. Peterson, "Comparison theorems for eigenvalue problems for n th order differential equations," *Proceedings of the American Mathematical Society*, vol. 104, no. 4, pp. 1204–1211, 1988.
- [8] J. Henderson and K. R. Prasad, "Comparison of eigenvalues for Lidstone boundary value problems on a measure chain," *Computers & Mathematics with Applications*, vol. 38, no. 11-12, pp. 55–62, 1999.
- [9] E. R. Kaufmann, "Comparison of eigenvalues for eigenvalue problems of a right disfocal operator," *Panamerican Mathematical Journal*, vol. 4, no. 4, pp. 103–124, 1994.
- [10] T. Ando, "Majorization, doubly stochastic matrices, and comparison of eigenvalues," *Linear Algebra and its Applications*, vol. 118, pp. 163–248, 1989.
- [11] C. A. Akemann and N. Weaver, "Minimal upper bounds of commuting operators," *Proceedings of the American Mathematical Society*, vol. 124, no. 11, pp. 3469–3476, 1996.
- [12] W. Arveson, "On groups of automorphisms of operator algebras," *Journal of Functional Analysis*, vol. 15, pp. 217–243, 1974.
- [13] T. Kato, "Spectral order and a matrix limit theorem," *Linear and Multilinear Algebra*, vol. 8, no. 1, pp. 15–19, 1979/80.
- [14] M. P. Olson, "The selfadjoint operators of a von Neumann algebra form a conditionally complete lattice," *Proceedings of the American Mathematical Society*, vol. 28, pp. 537–544, 1971.
- [15] J. Hamhalter, "Spectral order of operators and range projections," *Journal of Mathematical Analysis and Applications*, vol. 331, no. 2, pp. 1122–1134, 2007.
- [16] M. Bohner and A. Peterson, *Dynamic Equations on Time Scales*, Birkhäuser, Boston, Mass, USA, 2001, An introduction with Applications.
- [17] V. Lakshmikantham, S. Sivasundaram, and B. Kaymakcalan, *Dynamic Systems on Measure Chains*, vol. 370 of *Mathematics and its Applications*, Kluwer Academic Publishers, Dordrecht, The Netherlands, 1996.
- [18] R. P. Agarwal and M. Bohner, "Basic calculus on time scales and some of its applications," *Results in Mathematics*, vol. 35, no. 1-2, pp. 3–22, 1999.
- [19] J. Sherman and W. J. Morrison, "Adjustment of an inverse matrix corresponding to a change in one element of a given matrix," *Annals of Mathematical Statistics*, vol. 21, pp. 124–127, 1950.
- [20] R. S. Varga, *Matrix Iterative Analysis*, Prentice-Hall, Englewood Cliffs, NJ, USA, 1962.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

