

Research Article

A New Singular Impulsive Delay Differential Inequality and Its Application

Zhixia Ma¹ and Xiaohu Wang²

¹ College of Computer Science and Technology, Southwest University for Nationalities, Chengdu 610041, China

² Yangtze Center of Mathematics, Sichuan University, Chengdu 610064, China

Correspondence should be addressed to Xiaohu Wang, xiaohuwang111@163.com

Received 10 January 2009; Accepted 5 March 2009

Recommended by Wing-Sum Cheung

A new singular impulsive delay differential inequality is established. Using this inequality, the invariant and attracting sets for impulsive neutral neural networks with delays are obtained. Our results can extend and improve earlier publications.

Copyright © 2009 Z. Ma and X. Wang. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

It is well known that inequality technique is an important tool for investigating dynamical behavior of differential equation. The significance of differential and integral inequalities in the qualitative investigation of various classes of functional equations has been fully illustrated during the last 40 years [1–3]. Various inequalities have been established such as the delay integral inequality in [4], the differential inequalities in [5, 6], the impulsive differential inequalities in [7–10], Halanay inequalities in [11–13], and generalized Halanay inequalities in [14–17]. By using the technique of inequality, the invariant and attracting sets for differential systems have been studied by many authors [9, 18–21].

However, the inequalities mentioned above are ineffective for studying the invariant and attracting sets of impulsive nonautonomous neutral neural networks with time-varying delays. On the basis of this, this article is devoted to the discussion of this problem.

Motivated by the above discussions, in this paper, a new singular impulsive delay differential inequality is established. Applying this equality and using the methods in [10, 22], some sufficient conditions ensuring the invariant set and the global attracting set for a class of neutral neural networks system with impulsive effects are obtained.

2. Preliminaries

Throughout the paper, E_n means n -dimensional unit matrix, \mathbb{R} the set of real numbers, \mathbb{N} the set of positive integers, and $\mathcal{N} \triangleq \{1, 2, \dots, n\}$. $A \geq B$ ($A > B$) means that each pair of corresponding elements of A and B satisfies the inequality " \geq " (" $>$ "). Especially, A is called a nonnegative matrix if $A \geq 0$.

$C(X, Y)$ denotes the space of continuous mappings from the topological space X to the topological space Y . In particular, let $C \triangleq C([- \tau, 0], \mathbb{R}^n)$, where $\tau > 0$ is a constant.

$PC([a, b], \mathbb{R}^n)$ denotes the space of piecewise continuous functions $\varphi(s) : [a, b] \rightarrow \mathbb{R}^n$ with at most countable discontinuous points and at this points $\varphi(s)$ are right continuous.

Especially, let $PC \triangleq PC([- \tau, 0], \mathbb{R}^n)$. Furthermore, put $PC([a, b], \mathbb{R}^n) = \bigcup_{c \in [a, b]} PC([a, c], \mathbb{R}^n)$.

$PC^1([a, b], \mathbb{R}^n) = \{\varphi(s) : [a, b] \rightarrow \mathbb{R}^n \mid \varphi(s), \dot{\varphi}(s) \in PC([a, b], \mathbb{R}^n)\}$, where $\dot{\varphi}(s)$ denotes the derivative of $\varphi(s)$. In particular, let $PC^1 \triangleq PC^1([a, b], \mathbb{R}^n)$.

$\mathcal{H} = \{h(t) : \mathbb{R} \rightarrow \mathbb{R} \mid h(t) \text{ is a positive integrable function and satisfies } \sup_{a \leq t < b} \int_{t-\tau}^t h(s) ds = \sigma < \infty \text{ and } \lim_{t \rightarrow \infty} \int_a^t h(s) ds = \infty\}$.

For $x \in \mathbb{R}^n$, $A \in \mathbb{R}^{n \times n}$, $\varphi \in C$ or $\varphi \in PC$, we define $[x]^+ = (|x_1|, \dots, |x_n|)^T$, $[A]^+ = (|a_{ij}|)_{n \times n}$, $[\varphi(t)]_\tau = ([\varphi_1(t)]_\tau, \dots, [\varphi_n(t)]_\tau)^T$, $[\varphi(t)]_\tau^+ = [[\varphi(t)]^+]_\tau$, $[\varphi_i(t)]_\tau = \sup_{-\tau \leq \theta < 0} \{\varphi_i(t + \theta)\}$. And we introduce the following norm, respectively,

$$\|x\| = \max_{1 \leq i \leq n} |x_i|, \quad \|A\| = \max_{1 \leq i \leq n} \sum_{j=1}^n |a_{ij}|, \quad \|\varphi\|_\tau = \sup_{-\tau \leq s \leq 0} \|\varphi(s)\|. \quad (2.1)$$

For any $\varphi \in PC^1$, we define the following norm:

$$\|\varphi\|_{1\tau} = \max \{ \|\varphi\|_\tau, \|\varphi'\|_\tau \}. \quad (2.2)$$

For an M -matrix D defined in [23], we denote

$$\Omega_M(D) \triangleq \{z \in \mathbb{R}^n \mid Dz > 0, z > 0\}. \quad (2.3)$$

It is a cone without conical surface in \mathbb{R}^n . We call it an " M -cone".

3. Singular Impulsive Delay Differential Inequality

For convenience, we introduce the following conditions.

(C₁) Let the r -dimensional diagonal matrix $K = \text{diag}\{k_1, \dots, k_r\}$ satisfy

$$k_i > 0, \quad i \in S \subset \mathcal{N}^* \triangleq \{1, \dots, r\}, \quad k_i = 0, \quad i \in S^* \triangleq \mathcal{N}^* - S. \quad (3.1)$$

(C₂) Let $U = -(P + Q)$ be an M -matrix, where $Q = (q_{ij})_{r \times r} \geq 0$ and $P = (p_{ij})_{r \times r}$ satisfies

$$p_{ij} \geq 0, \quad i \neq j, \quad p_{ij} = 0, \quad i \neq j, \quad i \in \mathcal{N}^*, \quad j \in S^*. \quad (3.2)$$

Theorem 3.1. *Assume the conditions (C₁) and (C₂) hold. Let $L = (L_1, \dots, L_r)$ and $u(t) = (u_1(t), \dots, u_r(t))^T$ be a solution of the following singular delay differential inequality with the initial conditions $u(t) \in PC([a - \tau, a], \mathbb{R}^r)$:*

$$KD^+u(t) \leq h(t)[Pu(t) + Q[u(t)]_\tau + L], \quad t \in [a, b), \tag{3.3}$$

where $\tau > 0$, $a < b \leq +\infty$, and $u_i(t) \in C([a, b), \mathbb{R})$, $i \in S$, $u_i(t) \in PC([a, b), \mathbb{R})$, $i \in S^*$, $h(t) \in \mathcal{L}$. Then

$$u(t) \leq dze^{-\lambda \int_a^t h(s)ds} - (P + Q)^{-1}L, \quad t \in [a, b), \tag{3.4}$$

provided that the initial conditions satisfy

$$u(t) \leq dze^{-\lambda \int_a^t h(s)ds} - (P + Q)^{-1}L, \quad t \in [a - \tau, a], \tag{3.5}$$

where $d \geq 0$, $z = (z_1, \dots, z_r)^T \in \Omega_M(U)$ and the positive number λ satisfies the following inequality:

$$[\lambda K + P + Qe^{\lambda\sigma}]z < 0, \quad t \in [a, b). \tag{3.6}$$

Proof. By the conditions (C₂) and the definition of M -matrix, there is a constant vector $z = (z_1, \dots, z_r)^T$ such that $(P + Q)z < 0$, $-(P + Q)^{-1}$ exists and $-(P + Q)^{-1} \geq 0$.

By using continuity, we obtain that there must exist a positive constant λ satisfying the inequality (3.6), that is,

$$\sum_{j=1}^r [p_{ij} + q_{ij}e^{\lambda\sigma}]z_j < -\lambda k_i z_i, \quad i \in \mathcal{N}^*. \tag{3.7}$$

Denote by

$$v(t) = (v_1(t), \dots, v_r(t))^T = u(t) + (P + Q)^{-1}L, \quad t \in [a - \tau, b). \tag{3.8}$$

It follows from (3.3) and (3.5) that

$$\begin{aligned} KD^+v(t) &\leq h(t)[Pu(t) + Q[u(t)]_\tau + L] \\ &\leq h(t)[Pv(t) + Q[v(t)]_\tau], \quad t \in [a, b), \\ v(t) &\leq dze^{-\lambda \int_a^t h(s)ds}, \quad t \in [a - \tau, a]. \end{aligned} \tag{3.9}$$

In the following, we will prove that for any positive constant ε ,

$$v_i(t) \leq (d + \varepsilon)z_i e^{-\lambda \int_a^t h(s)ds} \triangleq \omega_i(t), \quad t \in [a, b), \quad i \in \mathcal{N}^*. \tag{3.10}$$

Let

$$\begin{aligned}\hat{\rho} &= \{i \in \mathcal{N}^* \mid v_i(t) > w_i(t) \text{ for some } t \in [a, b)\}, \\ \theta_i &= \inf \{t \in [a, b) \mid v_i(t) > w_i(t), i \in \hat{\rho}\}.\end{aligned}\quad (3.11)$$

If inequality (3.10) is not true, then $\hat{\rho}$ is a nonempty set and there must exist some integer $m \in \hat{\rho}$ such that $\theta_m = \min_{i \in \hat{\rho}} \{\theta_i\} \in [a, b)$.

If $m \in S$, by $v_m(t) \in C([a, b), \mathbb{R})$ and the inequality (3.5), we can get

$$\theta_m > a, \quad v_m(\theta_m) = w_m(\theta_m), \quad D^+ v_m(\theta_m) \geq \dot{w}_m(\theta_m), \quad (3.12)$$

$$v_i(t) \leq w_i(t), \quad t \in [a - \tau, \theta_m), i \in \mathcal{N}^*, \quad v_i(\theta_m) \leq w_i(\theta_m), \quad i \in S. \quad (3.13)$$

By using (C₂), (3.3), (3.7), (3.12), (3.13), and $[v_i(t)]_\tau = \sup_{-\tau \leq \theta < 0} \{u_i(t + \theta)\}$, $i \in \mathcal{N}^*$, we obtain that

$$\begin{aligned}k_m D^+ v_m(\theta_m) &\leq h(\theta_m) \sum_{j=1}^r [p_{mj} v_j(\theta_m) + q_{mj} [v_j(\theta_m)]_\tau] \\ &= h(\theta_m) \left[p_{mm} v_m(\theta_m) + \sum_{j \neq m, j \in S} p_{mj} v_j(\theta_m) \right. \\ &\quad \left. + \sum_{j \in S^*} p_{mj} v_j(\theta_m) + \sum_{j=1}^r q_{mj} [v_j(\theta_m)]_\tau \right] \\ &\leq h(\theta_m) \left[\sum_{j \in S} p_{mj} (d + \varepsilon) z_j e^{-\lambda \int_a^{\theta_m} h(s) ds} + \sum_{j=1}^r q_{mj} (d + \varepsilon) z_j e^{-\lambda \int_a^{\theta_m - \tau} h(s) ds} \right] \\ &\leq (d + \varepsilon) h(\theta_m) \sum_{j=1}^r [p_{mj} + q_{mj} e^{\lambda \sigma}] z_j e^{-\lambda \int_a^{\theta_m} h(s) ds} \\ &< -(d + \varepsilon) \lambda k_m z_m h(\theta_m) e^{-\lambda \int_a^{\theta_m} h(s) ds}.\end{aligned}\quad (3.14)$$

Since $m \in S$, we have $k_m > 0$ by (H₁). Then (3.14) becomes

$$D^+ u_m(\theta_m) < -(d + \varepsilon) \lambda z_m h(\theta_m) e^{-\lambda \int_a^{\theta_m} r(s) ds} = \dot{w}_m(\theta_m), \quad (3.15)$$

which contradicts the second inequality in (3.12).

If $m \in S^*$, then $k_m = 0$ by (C₁) and $v_m(t) \in PC([a, b), \mathbb{R})$. From the inequality (3.5), we can get

$$\begin{aligned}\theta_m &> a, \quad v_m(\theta_m) \geq w_m(\theta_m), \quad v_i(\theta_m) \leq w_i(\theta_m), \quad i \in S, \\ v_i(t) &\leq w_i(t), \quad t \in [a - \tau, \theta_m), i \in \mathcal{N}^*.\end{aligned}\quad (3.16)$$

By using (C₂), (3.3), (3.7), (3.16), and $[v_i(t)]_\tau = \sup_{-\tau \leq \theta < 0} \{v_i(t + \theta)\}$, $i \in \mathcal{N}^*$, we obtain that

$$\begin{aligned}
 0 &\leq \sum_{j=1}^r p_{mj} v_j(\theta_m) + \sum_{j=1}^r q_{mj} [v_j(\theta_m)]_\tau \\
 &= \sum_{j \in S} p_{mj} v_j(\theta_m) + p_{mm} v_m(\theta_m) + \sum_{j \neq m, j \in S^*} p_{mj} v_j(\theta_m) + \sum_{j=1}^r q_{mj} [v_j(\theta_m)]_\tau \\
 &\leq (d + \varepsilon) \left[\sum_{j \in S} p_{mj} z_j + p_{mm} z_m + \sum_{j=1}^r q_{mj} z_j e^{\lambda \int_{\theta_m - \tau}^{\theta_m} h(s) ds} \right] e^{-\lambda \int_a^{\theta_m} h(s) ds} \\
 &\leq (d + \varepsilon) \sum_{j=1}^r [p_{mj} z_j + q_{mj} z_j e^{\lambda \sigma}] e^{-\lambda \int_a^{\theta_m} h(s) ds} \\
 &< -(d + \varepsilon) k_m z_m h(\theta_m) e^{-\lambda \int_a^{\theta_m} h(s) ds} \\
 &= 0.
 \end{aligned} \tag{3.17}$$

This is a contradiction. Thus the inequality (3.10) holds. Therefore, letting $\varepsilon \rightarrow 0$ in (3.10), we have

$$v(t) = u(t) + (P + Q)^{-1} L \leq d z e^{-\lambda \int_a^t h(s) ds}, \quad t \in [a, b]. \tag{3.18}$$

The proof is complete. \square

Remark 3.2. In order to overcome the difficulty that $u(t)$ in (3.3) may be discontinuous, we introduce the notation $[u_i(t)]_\tau = \sup_{-\tau \leq s < 0} \{u_i(t + s)\}$ which is different from the notation $[u_i(t)]_\tau = \sup_{-\tau \leq s \leq 0} \{u_i(t + s)\}$ in [7]. However, when $u_i(t)$ is continuous in t , we have

$$[u_i(t)]_\tau = \sup_{-\tau \leq s < 0} \{u_i(t + s)\} = \sup_{-\tau \leq s \leq 0} \{u_i(t + s)\}, \quad i \in \mathcal{N}^*. \tag{3.19}$$

So we can get [7, Lemma 1] when we choose $K = E_r$, $S = \mathcal{N}^*$, $h(t) \equiv 1$ in Theorem 3.1.

Remark 3.3. Suppose that $L = (L_1, \dots, L_r)^T = 0$ and $h(t) \equiv 1$ in Theorem 3.1, then we can get [10, Theorem 3.1].

4. Applications

The singular impulsive delay differential inequality obtained in Section 3 can be widely applied to study the dynamics of impulsive neutral differential equations. To illustrate the theory, we consider the following nonautonomous impulsive neutral neural networks with delays

$$\begin{aligned}
 \dot{x}(t) &= -D(t)x(t) + A(t)F(x(t)) + B(t)G(x(t - \tau(t))) \\
 &\quad + C(t)H(\dot{x}(t - r(t))) + J(t), \quad t \neq t_k, \\
 x(t) &= I_k(t, x(t^-)), \quad t = t_k,
 \end{aligned} \tag{4.1}$$

where $x = (x_1, \dots, x_n)^T \in \mathbb{R}^n$ is the neural state vector; $D(t) = \text{diag}\{d_1(t), \dots, d_n(t)\} > 0$, $A(t) = (a_{ij}(t))_{n \times n}$, $B(t) = (b_{ij}(t))_{n \times n}$, $C(t) = (c_{ij}(t))_{n \times n}$ are the interconnection matrices representing the weighting coefficients of the neurons; $F(x) = (f_1(x_1), \dots, f_n(x_n))^T$, $G(x) = (g_1(x_1), \dots, g_n(x_n))^T$, $H(x) = (h_1(x_1), \dots, h_n(x_n))^T$ are activation functions; $\tau(t) = (\tau_{ij}(t))_{n \times n}$, $r(t) = (r_{ij}(t))_{n \times n}$ are transmission delays; $J(t) = (J_1(t), \dots, J_n(t))^T$ denotes the external inputs at time t . $I_k(t, y) = (I_{1k}(t, y), \dots, I_{nk}(t, y))^T$ represents impulsive perturbations; the fixed moments of time t_k satisfy $t_k < t_{k+1}$, $\lim_{k \rightarrow +\infty} t_k = +\infty$, $k \in \mathbb{N}$.

The initial condition for (4.1) is given by

$$x(t_0 + s) = \varphi(s) \in PC^1, \quad t_0 \in \mathbb{R}, \quad -\tau \leq s \leq 0. \quad (4.2)$$

We always assume that for any $\varphi \in PC^1$, (4.1) has at least one solution through (t_0, φ) , denoted by $x(t, t_0, \varphi)$ or $x_t(t_0, \varphi)$ (simply $x(t)$ or x_t if no confusion should occur).

Definition 4.1. The set $S \subset PC^1$ is called a positive invariant set of (4.1), if for any initial value $\varphi \in S$, we have the solution $x_t(t_0, \varphi) \in S$ for $t \geq t_0$.

Definition 4.2. The set $S \subset PC^1$ is called a global attracting set of (4.1), if for any initial value $\varphi \in PC^1$, the solution $x_t(t_0, \varphi)$ converges to S as $t \rightarrow +\infty$. That is,

$$\text{dist}(x_t, S) \rightarrow 0 \quad \text{as } t \rightarrow +\infty, \quad (4.3)$$

where $\text{dist}(\phi, S) = \inf_{\psi \in S} \text{dist}(\phi, \psi)$, $\text{dist}(\phi, \psi) = \sup_{s \in [-\tau, 0]} |\phi(s) - \psi(s)|$, for $\phi \in PC^1$.

Throughout this section, we suppose the following.

(H₁) $D(t) \in PC(\mathbb{R}, \mathbb{R}^n)$, $A(t), B(t), C(t), \tau(t), r(t)$ are continuous. Moreover, $0 \leq \tau_{ij}(t) \leq \tau$ and $0 < r_{ij}(t) \leq \tau$ ($i, j \in \mathcal{N}$).

(H₂) There exist nonnegative matrices $\tilde{D}_1 = \text{diag}\{\hat{d}_{11}, \dots, \hat{d}_{1n}\}$, $\tilde{D}_2 = \text{diag}\{\hat{d}_{21}, \dots, \hat{d}_{2n}\}$, $\hat{J} = (\hat{J}_1, \dots, \hat{J}_n)^T$, $h(s) \in \mathcal{A}$ and a constant $\delta > 0$ such that

$$\tilde{D}_1 h(t) \leq D(t) \leq \tilde{D}_2 h(t), \quad 0 < h(t) \leq \frac{1}{\delta}, \quad [J(t)]^+ \leq \hat{J} h(t). \quad (4.4)$$

(H₃) There exist nonnegative matrices $\tilde{A} = (\tilde{a}_{ij})_{n \times n}$, $\tilde{B} = (\tilde{b}_{ij})_{n \times n}$, $\tilde{C} = (\tilde{c}_{ij})_{n \times n}$ such that

$$[A(t)]^+ \leq \tilde{A} h(t), \quad [B(t)]^+ \leq \tilde{B} h(t), \quad [C(t)]^+ \leq \tilde{C} h(t). \quad (4.5)$$

(H₄) There exist nonnegative matrices $\tilde{F} = \text{diag}\{\alpha_1, \dots, \alpha_n\}$, $\tilde{G} = \text{diag}\{\beta_1, \dots, \beta_n\}$, $\tilde{H} = \text{diag}\{\gamma_1, \dots, \gamma_n\}$ such that for all $u \in \mathbb{R}^n$ the activating functions $F(\cdot), G(\cdot)$ and $H(\cdot)$ satisfy

$$[F(u)]^+ \leq \tilde{F}[u]^+, \quad [G(u)]^+ \leq \tilde{G}[u]^+, \quad [H(u)]^+ \leq \tilde{H}[u]^+. \quad (4.6)$$

(H₅) There exists nonnegative matrix $\widehat{I}_k = (\widehat{I}_{ij}^k)_{n \times n}$, such that for all $u \in \mathbb{R}^n$, $i \in \mathcal{N}$ and $k \in \mathbb{N}$

$$[I_k(t, u)]^+ \leq \widehat{I}_k[u]^+. \tag{4.7}$$

(H₆) Denote by

$$\begin{aligned} \widetilde{U} &= \widetilde{A}\widetilde{F}, & \widetilde{V} &= \widetilde{B}\widetilde{G}, & \widetilde{W} &= \widetilde{C}\widetilde{H}, \\ \overline{K} &= \begin{pmatrix} E_n & 0 \\ 0 & 0 \end{pmatrix} \triangleq \text{diag}\{\widehat{k}_1, \dots, \widehat{k}_{2n}\}, & \overline{L} &= (\widehat{J}, \widehat{J})^T, \\ \overline{P} &= \begin{pmatrix} -\widetilde{D}_1 + \widetilde{U} & 0 \\ \widetilde{D}_2 + \widetilde{U} & -\delta E_n \end{pmatrix} \triangleq (\widehat{p}_{ij}(t))_{2n \times 2n}, & \overline{Q} &= \begin{pmatrix} \widetilde{V} & \widetilde{W} \\ \widetilde{V} & \widetilde{W} \end{pmatrix} \triangleq (\widehat{q}_{ij}(t))_{2n \times 2n} \end{aligned} \tag{4.8}$$

and let $\overline{D} = -(\overline{P} + \overline{Q})$ be an M -matrix, and $\varrho = -(\overline{P} + \overline{Q})^{-1}\overline{L} = (\varrho_1, \varrho_2)^T \geq 0$, $\varrho_1, \varrho_2 \in \mathbb{R}^n$.

(H₇) There exists a constant ν such that

$$\ln \eta_k \leq \nu \int_{t_{k-1}}^{t_k} h(s) ds, \quad k \in \mathbb{N}, \quad \mu = \sum_{k=1}^{\infty} \ln \mu_k < \infty, \tag{4.9}$$

where $\nu < \lambda$, and the scalar $\lambda > 0$ is determined by the inequality

$$[\lambda \overline{K} + \overline{P} + \overline{Q}e^{\lambda \sigma}]z^* < 0, \tag{4.10}$$

where $z^* = (z_1, \dots, z_{2n})^T \in \Omega_M(\overline{D})$, and

$$\eta_k, \mu_k \geq 1, \quad \eta_k z_x^* \geq \widehat{I}_k z_x^*, \quad \mu_k \varrho_1 \geq \widehat{I}_k \varrho_1, \quad k \in \mathbb{N}, \quad z_x^* = (z_1, \dots, z_n)^T. \tag{4.11}$$

Theorem 4.3. Assume that (H₁)–(H₇) hold. Then $S = \{\phi \in PC^1 \mid [\phi]_{\tau}^+ \leq e^{\mu} \varrho_1\}$ is a global attracting set of (4.1).

Proof. Denote $\dot{x}(t) = y(t)$. Let $\text{sgn}(\cdot)$ be the sign function. For $x = (x_1, \dots, x_n)^T$, define $\text{Sgn}(x) = \text{diag}\{\text{sgn}(x_1), \dots, \text{sgn}(x_n)\}$.

Calculating the upper right derivative $D^+[x(t)]^+$ along system (4.1). From (4.1), (H₂) and (H₃) we have

$$\begin{aligned} D^+[x(t)]^+ &\leq h(t)\{(-D_1 + \widetilde{U})[x(t)]^+ + \widetilde{V}G[x(t)]_{\tau}^+ \\ &\quad + \widetilde{W}[y(t)]_{\tau}^+ + \widehat{J}\}, \quad t \in [t_{k-1}, t_k), \quad t \geq \sigma, \quad k \in \mathbb{N}. \end{aligned} \tag{4.12}$$

On the other hand, from (4.1) and (H_2) – (H_4) , we have

$$\begin{aligned} 0 &\leq h(t) \left\{ -\frac{1}{h(t)} [y(t)]^+ + (\tilde{D}_2 + \tilde{U}) [x(t)]^+ + \tilde{V} [x(t)]_\tau^+ + \tilde{W} [y(t)]_\tau^+ + \hat{J} \right\} \\ &\leq h(t) \{ -\delta [y(t)]^+ + (\tilde{D}_2 + \tilde{U}) [x(t)]^+ + \tilde{V} [x(t)]_\tau^+ + \tilde{W} [y(t)]_\tau^+ + \hat{J} \}, \quad t \in [t_{k-1}, t_k], \quad k \in \mathbb{N}. \end{aligned} \quad (4.13)$$

Let

$$u(t) = (x(t), y(t))^T \in \mathbb{R}^{2n}, \quad (4.14)$$

then from (4.12)–(4.14) and (H_6) , we have

$$\overline{KD}^+ [u(t)]^+ \leq h(t) [\overline{P}[u(t)]^+ + \overline{Q}[u(t)]_\tau^+ + \overline{L}], \quad t \in [t_{k-1}, t_k], \quad k \in \mathbb{N}. \quad (4.15)$$

By the conditions (H_6) and the definition of M -matrix, we may choose a vector $z^* = (z_1, \dots, z_{2n})^T \in \Omega_M(\overline{D})$ such that

$$\overline{D}z^* = -(\overline{P} + \overline{Q})z^* > 0. \quad (4.16)$$

By using continuity, we obtain that there must be a positive constant λ satisfying the inequality (4.10). Let $z_x^* = (z_1, \dots, z_n)^T$ and $z_y^* = (z_{n+1}, \dots, z_{2n})^T$, then $z^* = (z_x^*, z_y^*)^T$. Since $z^* > 0$, denote

$$d = \frac{1}{\min_{1 \leq i \leq 2n} \{z_i\}}, \quad (4.17)$$

then $dz^* \geq e_{2n} = (1, \dots, 1)^T \in \mathbb{R}^{2n}$. From the property of M -cone, we have, $dz^* \in \Omega_M(\overline{D})$.

For the initial conditions $x(t_0 + s) = \varphi(s)$, $s \in [-\tau, 0]$, where $\varphi \in PC^1$ and $t_0 \in \mathbb{R}$ (no loss of generality, we assume $t_0 \leq t_1$), and $t \in [t_0 - \tau, t_0]$, we can get

$$\begin{aligned} [x(t)]^+ &\leq \|\varphi(t)\|_\tau e^{-\lambda \int_{t_0}^t h(s) ds} dz_x^*, \\ [y(t)]^+ &\leq \|\varphi'(t)\|_\tau e^{-\lambda \int_{t_0}^t h(s) ds} dz_y^*. \end{aligned} \quad (4.18)$$

Then (4.18) yield

$$[u(t)]^+ \leq dz^* \|\varphi\|_{1\tau} e^{-\lambda \int_{t_0}^t h(s) ds} + \varrho, \quad t_0 - \tau \leq t \leq t_0. \quad (4.19)$$

Let $\mathcal{N}^* = \{1, \dots, 2n\}$, $S = \{1, \dots, n\} = \mathcal{N}$ and $S^* = \{n+1, \dots, 2n\} = \mathcal{N}^* - S$. Thus, all conditions of Theorem 3.1 are satisfied. By Theorem 3.1, we have

$$[u(t)]^+ \leq dz^* \|\varphi\|_{1\tau} e^{-\lambda \int_{t_0}^t h(s) ds} + \varrho, \quad t_0 \leq t < t_1. \quad (4.20)$$

Suppose that for all $m = 1, \dots, k$, the inequalities

$$[u(t)]^+ \leq \prod_{j=0}^{m-1} \eta_j dz_x^* \|\varphi\|_{1\tau} e^{-\lambda \int_{t_0}^t h(s) ds} + \prod_{j=0}^{m-1} \mu_j Q, \quad t_{m-1} \leq t < t_m, \quad t \geq t_0, \quad (4.21)$$

hold, where $\eta_0 = \mu_0 = 1$.

From (4.21), (H₅), and (H₇), we can get

$$\begin{aligned} [x(t_k)]^+ &\leq \widehat{I}_k [x(t_k^-)]^+ \\ &\leq \prod_{j=0}^k \eta_j dz_x^* \|\varphi(t)\|_{1\tau} e^{-\lambda \int_{t_0}^{t_k} h(s) ds} + \prod_{j=0}^k \mu_j Q_1. \end{aligned} \quad (4.22)$$

Since $\varrho = -(\overline{P} + \overline{Q})^{-1} \overline{L}$, we have

$$(\widetilde{D}_2 + \widetilde{U} + \widetilde{V})Q_1 + \widetilde{W}Q_2 + \widehat{J} = \delta Q_2. \quad (4.23)$$

On the other hand, it follows from (H₇) that

$$(\widetilde{D}_2 + \widetilde{U})z_x^* + (\widetilde{V}z_x^* + \widetilde{W}z_y^*)e^{\lambda\sigma} < \delta z_y^*. \quad (4.24)$$

Then from (4.21)–(4.24), we have

$$[y(t_k)]^+ \leq \prod_{j=0}^k \eta_j dz_y^* \|\varphi\|_{1\tau} e^{-\lambda \int_{t_0}^{t_k} h(s) ds} + \prod_{j=0}^k \mu_j Q_2, \quad (4.25)$$

which together with (4.22) yields that

$$[u(t_k)]^+ \leq \prod_{j=0}^k \eta_j dz_x^* \|\varphi\|_{1\tau} e^{-\lambda \int_{t_0}^{t_k} h(s) ds} + \prod_{j=0}^k \mu_j Q. \quad (4.26)$$

Then, it follows from (4.21) and (4.26) that

$$\begin{aligned} [u(t)]^+ &\leq \prod_{j=0}^k \eta_j dz_x^* \|\varphi\|_{1\tau} e^{-\lambda \int_{t_0}^t h(s) ds} + \prod_{j=0}^k \mu_j Q \\ &= \prod_{j=0}^k \eta_j dz_x^* \|\varphi\|_{1\tau} e^{-\lambda \int_{t_0}^{t_k} h(s) ds} e^{-\lambda \int_{t_k}^t h(s) ds} + \prod_{j=0}^k \mu_j Q, \quad \forall t \in [t_k - \tau, t_k]. \end{aligned} \quad (4.27)$$

Using Theorem 3.1 again, we have

$$\begin{aligned} [u(t)]^+ &\leq \prod_{j=0}^k \eta_j dz^* \|\varphi\|_{1\tau} e^{-\lambda \int_{t_0}^{t_k} h(s) ds} e^{-\lambda \int_{t_k}^t h(s) ds} + \prod_{j=0}^k \mu_j Q \\ &= \prod_{j=0}^k \eta_j dz^* \|\varphi\|_{1\tau} e^{-\lambda \int_{t_0}^t h(s) ds} + \prod_{j=0}^k \mu_j Q, \quad t_k \leq t < t_{k+1}. \end{aligned} \quad (4.28)$$

By mathematical induction, we can conclude that

$$[u(t)]^+ \leq \prod_{j=0}^k \eta_j dz^* \|\varphi\|_{1\tau} e^{-\lambda \int_{t_0}^t h(s) ds} + \prod_{j=0}^k \mu_j Q, \quad t_k \leq t < t_{k+1}, \quad k \in \mathbb{N}. \quad (4.29)$$

Noticing that $\eta_k \leq e^{\nu \int_{t_{k-1}}^{t_k} h(s) ds}$, by (H₇), we can use (4.29) to conclude that

$$\begin{aligned} [u(t)]^+ &\leq dz^* \|\varphi\|_{1\tau} e^{\nu \int_{t_0}^t h(s) ds} e^{-\lambda \int_{t_0}^t h(s) ds} + e^\mu Q \\ &= dz^* \|\varphi\|_{1\tau} e^{-(\lambda-\nu) \int_{t_0}^t h(s) ds} + e^\mu Q, \quad t_{k-1} \leq t < t_k, \quad k \in \mathbb{N}. \end{aligned} \quad (4.30)$$

This implies that the conclusion of the theorem holds. \square

By using Theorem 4.3 with $d = 0$, we can obtain a positive invariant set of (4.1), and the proof is similar to that of Theorem 4.3.

Theorem 4.4. Assume that (H₁)–(H₇) with $\widehat{I}_k = E_n$ hold. Then $S = \{\phi \in PC^1 \mid [\phi]_\tau^+ \leq Q_1\}$ is a positive invariant set and also a global attracting set of (4.1).

Remark 4.5. Suppose that $\widehat{c}_{ij} \equiv 0$, $i, j \in \mathcal{N}$ in (H₅), and $h(t) \equiv 1$, then we can get Theorems 1 and 2 in [9].

Remark 4.6. If $I_k(t, x(t^-)) = x \in \mathbb{R}^n$ then (4.1) becomes the nonautonomous neutral neural networks without impulses, we can get Theorem 4.1 in [22].

5. Illustrative Example

The following illustrative example will demonstrate the effectiveness of our results.

Example 5.1. Consider nonlinear impulsive neutral neural networks:

$$\begin{aligned} \dot{x}_1(t) &= -(7 + \cos^2 t)x_1(t) + \sin t \tan(x_1(t - \tau_{11}(t))) - \frac{1}{4} \cos t |\dot{x}_2(t - r_{12}(t))| - 1.5 \cos t, \\ \dot{x}_2(t) &= -(6 + \sin^2 t)x_2(t) - 2 \cos t |x_2(t - \tau_{22}(t))| + \frac{1}{4} \sin t \tan(x_1(t - r_{21}(t))) + 2.5 \sin t, \end{aligned} \quad t \neq k, \quad (5.1)$$

with

$$\begin{aligned} x_1(t) &= I_1(t, x(t^-)), \\ x_2(t) &= I_2(t, x(t^-)), \end{aligned} \quad t = k, \quad k \in \mathbb{N}, \tag{5.2}$$

where $\tau_{ij}(t) = (1/4)|\cos((i + j)t)| \leq 1/4 \stackrel{\Delta}{=} \tau$, $r_{ij}(t) = 1/4 - (1/8)|\sin((i + j)t)|$, $i, j = 1, 2$, $I_k(t, x) = (a_{1k}(t)x_1 + b_{1k}(t)x_2, a_{2k}(t)x_1 + b_{2k}(t)x_2)^T$, $k \in \mathbb{N}$.

The parameters of conditions (H_3) – (H_9) are as follows:

$$\begin{aligned} h(t) &= \delta = 1, & \tilde{D}_1 &= \text{diag}\{7, 6\}, & \tilde{D}_2 &= \text{diag}\{8, 7\}, & \sigma &= \frac{1}{4}, & \hat{J} &= (1.5, 2.5)^T, \\ \tilde{U} &= \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}, & \tilde{V} &= \begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix}, & \tilde{W} &= \frac{1}{4} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \\ \bar{K} &= \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, & \bar{P} &= \begin{pmatrix} -\tilde{D}_1 + \tilde{U} & 0 \\ \tilde{D}_2 + \tilde{U} & -\delta E_2 \end{pmatrix} = \begin{pmatrix} -7 & 0 & 0 & 0 \\ 0 & -6 & 0 & 0 \\ 8 & 0 & -1 & 0 \\ 0 & 7 & 0 & -1 \end{pmatrix}, \\ \bar{Q} &= \begin{pmatrix} \tilde{V} & \tilde{W} \\ \tilde{V} & \tilde{W} \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & \frac{1}{4} \\ 0 & 2 & \frac{1}{4} & 0 \\ 1 & 0 & 0 & \frac{1}{4} \\ 0 & 2 & \frac{1}{4} & 0 \end{pmatrix}, \\ \bar{D} &= -(\bar{P} + \bar{Q}) = \begin{pmatrix} 6 & 0 & 0 & -\frac{1}{4} \\ 0 & 4 & -\frac{1}{4} & 0 \\ -9 & 0 & 1 & -\frac{1}{4} \\ 0 & -9 & -\frac{1}{4} & 1 \end{pmatrix}. \end{aligned} \tag{5.3}$$

It is easy to prove that \bar{D} is an M -matrix and

$$\Omega_M(\bar{D}) = \{(z_1, z_2, z_3, z_4)^T > 0 \mid 9z_2 + \frac{1}{4}z_3 < z_4 < 24z_1, 9z_1 + \frac{1}{4}z_4 < z_3 < 16z_2\}. \tag{5.4}$$

Let $z^* = (1, 1, 15, 20)^T$, then $z^* \in \Omega_M(\bar{D})$ and $z_x^* = (1, 1)^T$. Let $\lambda = 0.1$ which satisfies the inequality

$$[\lambda \bar{K} + \bar{P} + \bar{Q} e^{\lambda \sigma}] z^* < 0. \tag{5.5}$$

Now, we discuss the asymptotical behavior of the system (5.1) as follows.

(i) If $a_{1k}(t) = b_{2k}(t) = 0$, $b_{1k}(t) = (1/2)e^{1/5^{2k}}(1 + \sin t)$, $a_{2k}(t) = (1/2)e^{1/5^{2k}}(1 - \cos t)$, then

$$\hat{I}_k = e^{1/5^{2k}} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}. \quad (5.6)$$

Thus $\eta_k = \mu_k = e^{1/5^{2k}} \geq 1$, $\ln \eta_k = e^{1/5^{2k}} \leq 0.04$, $\nu = 0.04 < \lambda$, and $\mu = 1/24$. Clearly, all conditions of Theorem 4.3 are satisfied, by Theorem 4.3, $S = \{\phi \in PC^1 \mid [\phi]_\tau^+ \leq e^{1/24} \varrho_1 \approx e^{1/24} (1.196, 1.746)^T\}$ is a global attracting set of (5.1).

(ii) If $a_{1k}(t) = \cos t$, $b_{2k}(t) = \sin t$, $b_{1k}(t) = a_{2k}(t) = 0$, then $\hat{I}_k = E_2$. By Theorem 4.4, $S = \{\phi \in PC^1 \mid [\phi]_\tau^+ \leq \varrho_1 \approx (1.196, 1.746)^T\}$ is a positive invariant set of (5.1).

Acknowledgments

This work was supported by the National Natural Science Foundation of China under Grant no. 10671133 and the Scientific Research Fund of Sichuan Provincial Education Department (08ZA044).

References

- [1] W. Walter, *Differential and Integral Inequalities*, Springer, New York, NY, USA, 1970.
- [2] V. Lakshmikantham and S. Leela, *Differential and Integral Inequalities: Theory and Applications. Vol. I: Ordinary Differential Equations*, vol. 55 of *Mathematics in Science and Engineering*, Academic Press, New York, NY, USA, 1969.
- [3] V. Lakshmikantham and S. Leela, *Differential and Integral Inequalities: Theory and Applications. Vol. II: Functional, Partial, Abstract, and Complex Differential Equations*, vol. 55 of *Mathematics in Science and Engineering*, Academic Press, New York, NY, USA, 1969.
- [4] D. Y. Xu, "Integro-differential equations and delay integral inequalities," *Tohoku Mathematical Journal*, vol. 44, no. 3, pp. 365–378, 1992.
- [5] L. Wang and D. Y. Xu, "Global exponential stability of Hopfield reaction-diffusion neural networks with time-varying delays," *Science in China. Series F*, vol. 46, no. 6, pp. 466–474, 2003.
- [6] Y. M. Huang, D. Y. Xu, and Z. G. Yang, "Dissipativity and periodic attractor for non-autonomous neural networks with time-varying delays," *Neurocomputing*, vol. 70, no. 16–18, pp. 2953–2958, 2007.
- [7] D. Y. Xu and Z. Yang, "Impulsive delay differential inequality and stability of neural networks," *Journal of Mathematical Analysis and Applications*, vol. 305, no. 1, pp. 107–120, 2005.
- [8] D. Y. Xu, W. Zhu, and S. Long, "Global exponential stability of impulsive integro-differential equation," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 64, no. 12, pp. 2805–2816, 2006.
- [9] D. Y. Xu and Z. Yang, "Attracting and invariant sets for a class of impulsive functional differential equations," *Journal of Mathematical Analysis and Applications*, vol. 329, no. 2, pp. 1036–1044, 2007.
- [10] D. Y. Xu, Z. Yang, and Z. Yang, "Exponential stability of nonlinear impulsive neutral differential equations with delays," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 67, no. 5, pp. 1426–1439, 2007.
- [11] R. D. Driver, *Ordinary and Delay Differential Equations*, vol. 20 of *Applied Mathematical Sciences*, Springer, New York, NY, USA, 1977.
- [12] K. Gopalsamy, *Stability and Oscillations in Delay Differential Equations of Population Dynamics*, vol. 74 of *Mathematics and Its Applications*, Kluwer Academic Publishers, Dordrecht, The Netherlands, 1992.
- [13] A. Halanay, *Differential Equations: Stability, Oscillations, Time Lags*, Academic Press, New York, NY, USA, 1966.
- [14] T. Amemiya, "Delay-independent stabilization of linear systems," *International Journal of Control*, vol. 37, no. 5, pp. 1071–1079, 1983.

- [15] E. Liz and S. Trofimchuk, "Existence and stability of almost periodic solutions for quasilinear delay systems and the Halanay inequality," *Journal of Mathematical Analysis and Applications*, vol. 248, no. 2, pp. 625–644, 2000.
- [16] A. Ivanov, E. Liz, and S. Trofimchuk, "Halanay inequality, Yorke 3/2 stability criterion, and differential equations with maxima," *Tohoku Mathematical Journal*, vol. 54, no. 2, pp. 277–295, 2002.
- [17] H. Tian, "The exponential asymptotic stability of singularly perturbed delay differential equations with a bounded lag," *Journal of Mathematical Analysis and Applications*, vol. 270, no. 1, pp. 143–149, 2002.
- [18] K. N. Lu, D. Y. Xu, and Z. C. Yang, "Global attraction and stability for Cohen-Grossberg neural networks with delays," *Neural Networks*, vol. 19, no. 10, pp. 1538–1549, 2006.
- [19] D. Y. Xu, S. Li, X. Zhou, and Z. Pu, "Invariant set and stable region of a class of partial differential equations with time delays," *Nonlinear Analysis: Real World Applications*, vol. 2, no. 2, pp. 161–169, 2001.
- [20] D. Y. Xu and H.-Y. Zhao, "Invariant and attracting sets of Hopfield neural networks with delay," *International Journal of Systems Science*, vol. 32, no. 7, pp. 863–866, 2001.
- [21] H. Zhao, "Invariant set and attractor of nonautonomous functional differential systems," *Journal of Mathematical Analysis and Applications*, vol. 282, no. 2, pp. 437–443, 2003.
- [22] Q. Guo, X. Wang, and Z. Ma, "Dissipativity of non-autonomous neutral neural networks with time-varying delays," *Far East Journal of Mathematical Sciences*, vol. 29, no. 1, pp. 89–100, 2008.
- [23] A. Berman and R. J. Plemmons, *Nonnegative Matrices in the Mathematical Sciences*, Computer Science and Applied Mathematic, Academic Press, New York, NY, USA, 1979.