

## Research Article

# Infinitely Many Periodic Solutions for Variable Exponent Systems

Xiaoli Guo,<sup>1</sup> Mingxin Lu,<sup>2</sup> and Qihu Zhang<sup>1</sup>

<sup>1</sup> Department of Mathematics and Information Science, Zhengzhou University of Light Industry, Zhengzhou, Henan 450002, China

<sup>2</sup> Department of Information Management, Nanjing University, Nanjing 210093, China

Correspondence should be addressed to Qihu Zhang, zhangqh1999@yahoo.com.cn

Received 4 December 2008; Accepted 14 April 2009

Recommended by Marta Garcia-Huidobro

We mainly consider the system  $-\Delta_{p(x)}u = f(v) + h(u)$  in  $\mathbb{R}$ ,  $-\Delta_{q(x)}v = g(u) + \omega(v)$  in  $\mathbb{R}$ , where  $1 < p(x), q(x) \in C^1(\mathbb{R})$  are periodic functions, and  $-\Delta_{p(x)}u = -(|u|^{p(x)-2}u)'$  is called  $p(x)$ -Laplacian. We give the existence of infinitely many periodic solutions under some conditions.

Copyright © 2009 Xiaoli Guo et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## 1. Introduction

The study of differential equations and variational problems with variable exponent growth conditions has been a new and interesting topic. Many results have been obtained on this kind of problems, for example [1–18]. On the applied background, we refer to [1, 3, 11, 18]. In this paper, we mainly consider the existence of infinitely many periodic solutions for the system

$$(P) \begin{cases} -\Delta_{p(x)}u = f(v) + h(u) & \text{in } \mathbb{R}, \\ -\Delta_{q(x)}v = g(u) + \omega(v) & \text{in } \mathbb{R}, \end{cases} \quad (1.1)$$

where  $p(x), q(x) \in C^1(\mathbb{R})$  are functions. The operator  $-\Delta_{p(x)}u = -(|u|^{p(x)-2}u)'$  is called one-dimensional  $p(x)$ -Laplacian. Especially, if  $p(x) \equiv p$  (a constant) and  $q(x) \equiv q$  (a constant), then (P) is the well-known constant exponent system.

$(u, v)$  is called a solution of (P), if  $u, v \in C^1(\mathbb{R})$ ,  $|u|^{p(x)-2}u'$  and  $|v|^{q(x)-2}v'$  are absolute continuous and satisfy (P) almost every where.

In [19], the authors consider the existence of positive weak solutions for the following constant exponent problems:

$$(I) \begin{cases} -\Delta_p u = \lambda f(v), & \text{in } \Omega, \\ -\Delta_p v = \lambda g(u), & \text{in } \Omega, \\ u = v = 0, & \text{on } \partial\Omega. \end{cases} \quad (1.2)$$

The first eigenfunction is used to construct the subsolution of constant exponent problems successfully. Under the condition that  $\lambda$  is large enough and

$$\lim_{u \rightarrow +\infty} \frac{f[M(g(u))^{1/(p-1)}]}{u^{p-1}} = 0, \quad \text{for every } M > 0, \quad (1.3)$$

the authors give the existence of positive solutions for problem (I).

In [20], the author considers the existence and nonexistence of positive weak solution to the following constant exponent elliptic system:

$$(II) \begin{cases} -\Delta_p u = \lambda u^\alpha v^\gamma, & \text{in } \Omega, \\ -\Delta_q v = \lambda u^\delta v^\beta, & \text{in } \Omega, \\ u = v = 0, & \text{on } \partial\Omega. \end{cases} \quad (1.4)$$

The first eigenfunction is used to construct the subsolution of constant exponent problems successfully.

Because of the nonhomogeneity of  $p(x)$ -Laplacian,  $p(x)$ -Laplacian problems are more complicated than those of  $p$ -Laplacian. Maybe the first eigenvalue and the first eigenfunction of  $p(x)$ -Laplacian do not exist (see [6]). Even if the first eigenfunction of  $p(x)$ -Laplacian exists, because of the nonhomogeneity of  $p(x)$ -Laplacian, the first eigenfunction cannot be used to construct the subsolution of  $p(x)$ -Laplacian problems.

There are many papers on the existence of periodic solutions for  $p$ -Laplacian elliptic systems, for example [21–24]. The results on the periodic solutions for variable exponent systems are rare. Through a new method of constructing sub-supersolution, this paper gives the existence of infinitely many periodic solutions for problem (P).

## 2. Main Results and Proofs

At first, we give an existence of positive solutions for variable exponent systems on bounded domain via sub-super-solution method. The result itself has dependent value.

Denote  $\Omega_R = (-R, R)$ . Let us consider the existence of positive solutions of the following:

$$(P_1) \begin{cases} -\Delta_{p(x)}u = f(v) + h(u), & \text{in } \Omega_R, \\ -\Delta_{q(x)}v = g(u) + \omega(v), & \text{in } \Omega_R, \\ u = v = 0, & \text{on } \partial\Omega_R. \end{cases} \tag{2.1}$$

Write  $z^+ = \sup_{x \in \mathbb{R}} z(x)$ ,  $z^- = \inf_{x \in \mathbb{R}} z(x)$ , for any  $z \in C(\mathbb{R})$ . Assume that

(H<sub>1</sub>)  $p(x), q(x) \in C^1(\mathbb{R})$  satisfy

$$1 < p^- \leq p^+ < \infty, \quad \sup |p'(x)| < \infty, \quad 1 < q^- \leq q^+ < \infty, \quad \sup |q'(x)| < \infty. \tag{2.2}$$

(H<sub>2</sub>)  $f, g, h, \omega : [0, +\infty) \rightarrow \mathbb{R}$  are  $C^1$ , monotone functions such that

$$\lim_{t \rightarrow +\infty} f(t) = \lim_{t \rightarrow +\infty} g(t) = \lim_{t \rightarrow +\infty} h(t) = \lim_{t \rightarrow +\infty} \omega(t) = +\infty. \tag{2.3}$$

(H<sub>3</sub>) For any positive constant  $M$ , there are  $\lim_{t \rightarrow +\infty} f[M(g(t))^{1/(q^- - 1)}] / t^{p^- - 1} = 0$ .

(H<sub>4</sub>)  $\lim_{t \rightarrow +\infty} h(t) / t^{p^- - 1} = \lim_{t \rightarrow +\infty} \omega(t) / t^{q^- - 1} = 0$ .

(H<sub>5</sub>)  $f, g, h$ , and  $\omega$  are odd functions such that  $f(0) = g(0) = h(0) = \omega(0) = 0$ ,  $p(x)$  and  $q(x)$  are even, and  $T$  is a periodic of  $p$  and  $q$ , namely,  $p(x) = p(x + T)$ ,  $q(x) = q(x + T)$ , for all  $x \in \mathbb{R}$ .

**Note.** In [14], the present author discussed the existence of solutions of  $(P_1)$ , under the conditions that  $(P_1)$  is radial,  $p(x) = q(x)$ , and  $h = \omega \equiv 0$ . Because of the nonhomogeneity of variable exponent problems, variable exponent problems are more complicated than constant exponent problems, and many results and methods for constant exponent problems are invalid for variable exponent problems. In many cases, the radial symmetric conditions are effective to deal with variable exponent problems. There are many results about the radial variable exponent problems (see [4, 14, 16]), but the following Theorem 2.1 does not assume any symmetric conditions.

We will establish.

**Theorem 2.1.** *If  $(H_1)$ – $(H_4)$  hold, then  $(P_1)$  possesses a positive solution, when  $R$  is sufficiently large.*

*Proof.* If we can construct a positive subsolution  $(\phi_1, \phi_2)$  and supersolution  $(z_1, z_2)$  of  $(P_1)$ , namely,

$$\begin{aligned} -\Delta_{p(x)}\phi_1 &\leq f(\phi_2) + h(\phi_1), & -\Delta_{q(x)}\phi_2 &\leq g(\phi_1) + \omega(\phi_2), & \text{for a.e. } x \in \mathbb{R}, \\ -\Delta_{p(x)}z_1 &\geq f(z_2) + h(z_1), & -\Delta_{q(x)}z_2 &\geq g(z_1) + \omega(z_2), & \text{for a.e. } x \in \mathbb{R}, \end{aligned} \tag{2.4}$$

which satisfy  $\phi_1 \leq z_1$  and  $\phi_2 \leq z_2$ , then  $(P_1)$  possesses a positive solution (see [5]).

Step 1. We will construct a subsolution of  $(P_1)$ .

Denfine

$$\phi_1(x) = \begin{cases} e^{-k_2(x-R)} - 1, & R - a < x \leq R, \\ e^{ak_2} - 1 + \int_x^{R-a} (k_2 e^{ak_2})^{(p(R-a)-1)/(p(r)-1)} \left[ \sin\left(\varepsilon_2(r - (R-a)) + \frac{\pi}{2}\right) \right]^{1/(p(r)-1)} dr, & R - a - \frac{\pi}{2\varepsilon_2} < x \leq R - a, \\ e^{ak_2} - 1 + \int_{R-a-\pi/2\varepsilon_2}^{R-a} (k_2 e^{ak_2})^{(p(R-a)-1)/(p(r)-1)} \left[ \sin\left(\varepsilon_2(r - (R-a)) + \frac{\pi}{2}\right) \right]^{1/(p(r)-1)} dr, & -R + a + \frac{\pi}{2\varepsilon_1} < x \leq R - a - \frac{\pi}{2\varepsilon_2}, \\ e^{ak_1} - 1 + \int_{-R+a}^x (k_1 e^{ak_1})^{(p(-R+a)-1)/(p(r)-1)} \left[ \sin\left(\frac{\pi}{2} - \varepsilon_1(r - (-R+a))\right) \right]^{1/(p(r)-1)} dr, & -R + a \leq x \leq -R + a + \frac{\pi}{2\varepsilon_1}, \\ e^{k_1(x+R)} - 1, & -R \leq x < -R + a. \end{cases}$$

$$\phi_2(x) = \begin{cases} e^{-k_4(x-R)} - 1, & R - a < x \leq R, \\ e^{ak_4} - 1 + \int_x^{R-a} (k_4 e^{ak_4})^{(q(R-a)-1)/(q(r)-1)} \left[ \sin\left(\varepsilon_4(r - (R-a)) + \frac{\pi}{2}\right) \right]^{1/(q(r)-1)} dr, & R - a - \frac{\pi}{2\varepsilon_4} < x \leq R - a, \\ e^{ak_4} - 1 + \int_{R-a-\pi/2\varepsilon_4}^{R-a} (k_4 e^{ak_4})^{(q(R-a)-1)/(q(r)-1)} \left[ \sin\left(\varepsilon_4(r - (R-a)) + \frac{\pi}{2}\right) \right]^{1/(q(r)-1)} dr, & -R + a + \frac{\pi}{2\varepsilon_3} < x \leq R - a - \frac{\pi}{2\varepsilon_4}, \\ e^{ak_3} - 1 + \int_{-R+a}^x (k_3 e^{ak_3})^{(q(-R+a)-1)/(q(r)-1)} \left[ \sin\left(\frac{\pi}{2} - \varepsilon_3(r - (-R+a))\right) \right]^{1/(q(r)-1)} dr, & -R + a \leq x \leq -R + a + \frac{\pi}{2\varepsilon_3}, \\ e^{k_3(x+R)} - 1, & -R \leq x < -R + a, \end{cases} \quad (2.5)$$

where

$$a = \min \left\{ \frac{\inf p(x) - 1}{4(\sup |p'(x)| + 1)}, \frac{\inf q(x) - 1}{4(\sup |q'(x)| + 1)} \right\}, \quad b = \min \{ f(0), g(0), h(0), \omega(0), -1 \},$$

$$\varepsilon_i = k_i^{-p^+} e^{-ak_i p^+}, \quad i = 1, 2, \quad \varepsilon_i = k_i^{-q^+} e^{-ak_i q^+}, \quad i = 3, 4; \quad R > \frac{\pi}{\varepsilon_i}, \quad i = 1, 2, 3, 4, \quad (2.6)$$

$k_1$  and  $k_2$  satisfy

$$\begin{aligned}
 & e^{ak_2} - 1 + \int_{R-a-\pi/2\varepsilon_2}^{R-a} (k_2 e^{ak_2})^{(p(R-a)-1)/(p(r)-1)} \left[ \sin \left( \varepsilon_2(r - (R-a)) + \frac{\pi}{2} \right) \right]^{1/(p(r)-1)} dr \\
 & = e^{ak_1} - 1 + \int_{-R+a}^{-R+a+\pi/2\varepsilon_1} (k_1 e^{ak_1})^{(p(-R+a)-1)/(p(r)-1)} \left[ \sin \left( \frac{\pi}{2} - \varepsilon_1(r - (-R+a)) \right) \right]^{1/(p(r)-1)} dr,
 \end{aligned} \tag{2.7}$$

$k_3$  and  $k_4$  satisfy

$$\begin{aligned}
 & e^{ak_4} - 1 + \int_{R-a-\pi/2\varepsilon_4}^{R-a} (k_4 e^{ak_4})^{q(R-a)-1/q(r)-1} \left[ \sin \left( \varepsilon_4(r - (R-a)) + \frac{\pi}{2} \right) \right]^{1/q(r)-1} dr \\
 & = e^{ak_3} - 1 + \int_{-R+a}^{-R+a+\pi/2\varepsilon_3} (k_3 e^{ak_3})^{q(-R+a)-1/q(r)-1} \left[ \sin \left( \frac{\pi}{2} - \varepsilon_3(r - (-R+a)) \right) \right]^{1/q(r)-1} dr,
 \end{aligned} \tag{2.8}$$

then  $\phi_1(x) \in C([-R, R])$ , and  $\phi_2(x) \in C([-R, R])$ . It is easy to see that  $\phi_i \geq 0$  and  $\phi_i \in C^1([-R, R])$ ,  $i = 1, 2$ . Obviously,  $\varepsilon_i = k_i^{-p^+} e^{-ak_i p^+}$  is continuous about  $k_i$ .

In the following, we will prove that  $(\phi_1, \phi_2)$  is a subsolution for  $(P_1)$ . By computation,

$$\begin{aligned}
 & -\Delta_{p(x)} \phi_1 = \\
 & \begin{cases} (k_2 e^{-k_2(x-R)})^{p(x)-1} [-k_2(p(x)-1) + p'(x) \ln k_2 - k_2 p'(x)(x-R)], & R-a < x \leq R, \\ \varepsilon_2 (k_2 e^{ak_2})^{(p(R-a)-1)} \cos \left( \varepsilon_2(x - (R-a)) + \frac{\pi}{2} \right), & R - \frac{\pi}{2\varepsilon_2} < x < R-a, \\ 0, & -R+a + \frac{\pi}{2\varepsilon_1} < x < R-a - \frac{\pi}{2\varepsilon_2}, \\ \varepsilon_1 (k_1 e^{ak_1})^{p(-R+a)-1} \left[ \cos \left( \frac{\pi}{2} - \varepsilon_1(x - (-R+a)) \right) \right], & -R+a < x < -R+a, \\ (k_1 e^{k_1(x+R)})^{p(x)-1} [-k_1(p(x)-1) - p'(x) \ln k_1 - k_1 p'(x)(x+R)], & -R \leq x < -R+a. \end{cases}
 \end{aligned} \tag{2.9}$$

If  $k_2$  is sufficiently large, we have

$$-\Delta_{p(x)} \phi_1 \leq -k_2 \left[ \inf p(x) - 1 - \sup |p'(x)| \left( \frac{\ln k_2}{k_2} + R - r \right) \right] \leq -k_2 a, \quad \forall x \in (R-a, R). \tag{2.10}$$

As  $a$  is a constant and only depends on  $p(x)$  and  $q(x)$ , when  $k_2$  is large enough, we have  $-k_2 a < 2b$ . Since  $\phi_1(x) \geq 0$  and  $f + h$  is monotone, we have

$$-\Delta_{p(x)} \phi_1 \leq 2b \leq f(0) + h(0) \leq f(\phi_2) + h(\phi_1), \quad R-a < x \leq R. \tag{2.11}$$

According to (H<sub>2</sub>), when  $k_i$  are large enough, we have

$$f(e^{ak_i} - 1) \geq 1, \quad g(e^{ak_i} - 1) \geq 1, \quad h(e^{ak_i} - 1) \geq 1, \quad \omega(e^{ak_i} - 1) \geq 1, \quad i = 1, 2, 3, 4, \quad (2.12)$$

where  $k_i$  are dependent on  $f, g, h, \omega, p$  and  $q$ , and they are independent on  $R$ . Since  $\varepsilon_2 = k_2^{-p^+} e^{-ak_2 p^+}$ , when  $x \in (R - a - \pi/2\varepsilon, R - a)$ , we have

$$-\Delta_{p(x)}\phi_1 = \varepsilon_2 (k_2 e^{ak_2})^{(p(R-1)-1)} \cos\left(\varepsilon(x - (R - a)) + \frac{\pi}{2}\right) \leq \varepsilon_2 k_2^{p^+} e^{ak_2 p^+} = 1. \quad (2.13)$$

Then we have

$$-\Delta_{p(x)}\phi_1 \leq 1 \leq f(\phi_2) + h(\phi_1), \quad R - a - \frac{\pi}{2\varepsilon_2} < x < R - a. \quad (2.14)$$

Obviously

$$-\Delta_{p(x)}\phi_1 = 0 \leq 1 \leq f(\phi_2) + h(\phi_1), \quad -R + a + \frac{\pi}{2\varepsilon_1} < x < R - a - \frac{\pi}{2\varepsilon_2}. \quad (2.15)$$

When  $k_2$  is large enough, from (2.7) we can see that  $k_1$  is large enough. Similar to the discussion of the above, we can conclude

$$-\Delta_{p(x)}\phi_1 \leq 1 \leq f(\phi_2) + h(\phi_1), \quad -R + a < x < -R + a + \frac{\pi}{2\varepsilon_1}, \quad (2.16)$$

$$-\Delta_{p(x)}\phi_1 \leq f(0) + h(0) \leq f(\phi_2) + h(\phi_1), \quad -R < x < -R + a. \quad (2.17)$$

Since  $\phi_i(x) \in C^1([-R, R])$ , combining (2.11), (2.14), (2.15), (2.16) and (2.17), we have

$$-\Delta_{p(x)}\phi_1 \leq f(\phi_2) + h(\phi_1), \quad \text{for a.e. } x \in (-R, R). \quad (2.18)$$

Similarly, when  $k_4$  is large enough, we have

$$-\Delta_{q(x)}\phi_2 \leq g(\phi_1) + \omega(\phi_2), \quad \text{for a.e. } x \in (-R, R). \quad (2.19)$$

Then  $(\phi_1, \phi_2)$  is a subsolution of  $(P_1)$ .

*Step 2.* We will construct a supersolution of  $(P_1)$ .

Let  $z_1$  be a solution of

$$-\left(|z_1'|^{p(x)-2} z_1'\right)' = 2\mu, \quad z_1(R) = 0 = z_1(-R), \quad (2.20)$$

where  $\mu$  is a positive constant and  $\mu > 1$ .

Obviously, there exists  $x_0 \in \Omega_R$  such that  $z_1(x) = \int_x^R |(r - x_0)2\mu|^{1/(p(r)-1)-1} 2\mu(r - x_0) dr$ . Note that  $x_0$  is dependent on  $\mu$ . Denote  $\beta = \beta(2\mu) = \max_{|x| \leq R} z_1(x)$ . It is easy to see that

$$\frac{1}{C} \mu^{1/p^+-1} \leq \beta(2\mu) \leq C \mu^{1/p^+-1}, \quad \text{where } C \geq 1 \text{ is a positive constant.} \quad (2.21)$$

Let us consider

$$\begin{aligned} -\Delta_{p(x)} z_1 &= 2\mu \quad \text{in } \Omega_R, \\ -\Delta_{q(x)} z_2 &= 2g(\beta(2\mu)) \quad \text{in } \Omega_R, \\ z_1 &= z_2 = 0 \quad \text{on } \partial\Omega_R. \end{aligned} \quad (2.22)$$

Similarly, we have

$$\frac{1}{C} [g(\beta(2\mu))]^{1/(q^+-1)} \leq \max_{|x| \leq R} z_2(x) \leq C [g(\beta(2\mu))]^{1/(q^+-1)}. \quad (2.23)$$

We will prove that  $(z_1, z_2)$  is a supersolution for  $(P_1)$ . From  $\lim_{t \rightarrow +\infty} \omega(t)/t^{q^- - 1} = 0$  and (2.23), when  $\mu$  is large enough, we can easily see that

$$-\Delta_{q(x)} z_2 = 2g(\beta(2\mu)) \geq g(z_1) + \omega(z_2). \quad (2.24)$$

Since  $\lim_{t \rightarrow +\infty} f[Cg(2t)^{1/(q^- - 1)}]/t^{p^- - 1} = 0$  and  $\lim_{t \rightarrow +\infty} h(t)/t^{p^- - 1} = 0$ , when  $\mu$  is large enough, according to (2.21) and (2.23), we have

$$2\mu \geq 2 \left( \frac{1}{C} \beta(2\mu) \right)^{p^- - 1} \geq f \left[ C(g(\beta(2\mu)))^{1/(q^- - 1)} \right] + h((\beta(2\mu))). \quad (2.25)$$

This means that

$$-\Delta_{p(x)} z_1 = 2\mu \geq f \left[ C(g(\beta(2\mu)))^{1/(q^- - 1)} \right] + h((\beta(2\mu))) \geq f(z_2) + h(z_1). \quad (2.26)$$

According to (2.24) and (2.26), we can conclude that  $(z_1, z_2)$  is a supersolution for  $(P_1)$ , when  $\mu$  is large enough.

*Step 3.* We will prove that  $\phi_1 \leq z_1$  and  $\phi_2 \leq z_2$ .

Obviously, when  $\mu$  is large enough, we can easily see that  $g(\beta(2\mu))$  is large enough, then

$$\begin{aligned} f(\phi_2) + h(\phi_1) &\leq \mu, \quad \forall x \in \Omega_R, \\ g(\phi_1) + \omega(\phi_2) &\leq g(\beta(2\mu)), \quad \forall x \in \Omega_R. \end{aligned} \quad (2.27)$$

Let us consider

$$-\Delta_{p(x)}\varpi = \mu \text{ in } \Omega_R. \quad (2.28)$$

It is easy to see that  $\phi_1$  is a subsolution of (2.28), when  $\mu$  is large enough. Obviously, we can see that  $z_1$  is a supersolution of (2.28), and

$$z_1(R) = \phi_1(R) = z_1(-R) = \phi_1(-R) = 0. \quad (2.29)$$

According to the comparison principle (see [12]), we can see that  $\phi_1 \leq z_1$ .  
Let us consider

$$-\Delta_{p(x)}\varpi = g(\beta(2\mu)) \text{ in } \Omega_R. \quad (2.30)$$

It is easy to see that  $\phi_2$  is a subsolution of (2.30), when  $\mu$  is large enough. Obviously, we can see that  $z_2$  is a supersolution of (2.30), and

$$z_2(R) = \phi_2(R) = z_2(-R) = \phi_2(-R) = 0. \quad (2.31)$$

According to the comparison principle (see [12]), we can see that  $\phi_2 \leq z_2$ .

Thus, we can conclude that  $\phi_1 \leq z_1$  and  $\phi_2 \leq z_2$ , when  $\mu$  is sufficiently large. This completes the proof. □

**Theorem 2.2.** *If  $(H_1)$ – $(H_5)$  hold, then  $(P)$  has infinitely many periodic solutions.*

*Proof.* Let  $R = nT$ . According to Theorem 2.1, we can conclude that there exists an integer  $n_0$  which is large enough such that  $(P_1)$  has a positive solution  $(u_n^\#(x), v_n^\#(x))$  for any integer  $n \geq n_0$ . Since  $p$  and  $q$  are even, and  $f, g, h$ , and  $\omega$  are odd, then  $(-u_n^\#(-x), -v_n^\#(-x))$  is a negative solution of  $(P_1)$ . We can define a  $C^1$  function  $(u_n(x), v_n(x))$  on  $[-nT, 3nT]$  as

$$\begin{aligned} u_n(x) &= \begin{cases} u_n^\#(x), & x \in [-nT, nT], \\ -u_n^\#(-(x - 2nT)), & x \in (nT, 3nT], \end{cases} \\ v_n(x) &= \begin{cases} v_n^\#(x), & x \in [-nT, nT], \\ -v_n^\#(-(x - 2nT)), & x \in (nT, 3nT]. \end{cases} \end{aligned} \quad (2.32)$$

We extend  $(u_n(x), v_n(x))$  as  $(u_n(x), v_n(x)) = (u_n(x + m4nT), v_n(x + m4nT))$ , where  $m$  is an integer such that  $x + m4nT \in [-nT, 3nT]$ . It is easy to see that  $u_n, v_n \in C^1(\mathbb{R})$ ,  $(u_n(x), v_n(x))$  is a solution of  $(P)$ , and the periodic of  $(u_n(x), v_n(x))$  is  $4nT$ . This completes the proof. □



## Acknowledgment

Partly supported by the National Science Foundation of China (10701066 & 10671084) and China Postdoctoral Science Foundation (20070421107) and the Natural Science Foundation of Henan Education Committee (2008-755-65 & 2009A120005) and the Natural Science Foundation of Jiangsu Education Committee (08KJD110007).

## References

- [1] E. Acerbi and G. Mingione, "Regularity results for stationary electro-rheological fluids," *Archive for Rational Mechanics and Analysis*, vol. 164, no. 3, pp. 213–259, 2002.
- [2] E. Acerbi and G. Mingione, "Gradient estimates for the  $p(x)$ -Laplacean system," *Journal für die Reine und Angewandte Mathematik*, vol. 584, pp. 117–148, 2005.
- [3] Y. Chen, S. Levine, and M. Rao, "Variable exponent, linear growth functionals in image restoration," *SIAM Journal on Applied Mathematics*, vol. 66, no. 4, pp. 1383–1406, 2006.
- [4] X. Fan, Y. Zhao, and D. Zhao, "Compact imbedding theorems with symmetry of Strauss-Lions type for the space  $W^{1,p(x)}(\Omega)$ ," *Journal of Mathematical Analysis and Applications*, vol. 255, no. 1, pp. 333–348, 2001.
- [5] X. Fan, "On the sub-supersolution method for  $p(x)$ -Laplacian equations," *Journal of Mathematical Analysis and Applications*, vol. 330, no. 1, pp. 665–682, 2007.
- [6] X. Fan, Q. Zhang, and D. Zhao, "Eigenvalues of  $p(x)$ -Laplacian Dirichlet problem," *Journal of Mathematical Analysis and Applications*, vol. 302, no. 2, pp. 306–317, 2005.
- [7] A. El Hamidi, "Existence results to elliptic systems with nonstandard growth conditions," *Journal of Mathematical Analysis and Applications*, vol. 300, no. 1, pp. 30–42, 2004.
- [8] P. Harjulehto, P. Hästö, and V. Latvala, "Harnack's inequality for  $p(\cdot)$ -harmonic functions with unbounded exponent  $p$ ," *Journal of Mathematical Analysis and Applications*, vol. 352, no. 1, pp. 345–359, 2009.
- [9] H. Hudzik, "On generalized Orlicz-Sobolev space," *Functiones et Approximatio Commentarii Mathematici*, vol. 4, pp. 37–51, 1976.
- [10] M. Mihăilescu and V. Rădulescu, "Continuous spectrum for a class of nonhomogeneous differential operators," *Manuscripta Mathematica*, vol. 125, no. 2, pp. 157–167, 2008.
- [11] M. Růžička, *Electrorheological Fluids: Modeling and Mathematical Theory*, vol. 1748 of *Lecture Notes in Mathematics*, Springer, Berlin, Germany, 2000.
- [12] Q. Zhang, "A strong maximum principle for differential equations with nonstandard  $p(x)$ -growth conditions," *Journal of Mathematical Analysis and Applications*, vol. 312, no. 1, pp. 24–32, 2005.
- [13] Q. Zhang, "Existence of solutions for  $p(x)$ -Laplacian equations with singular coefficients in  $\mathbb{R}^N$ ," *Journal of Mathematical Analysis and Applications*, vol. 348, no. 1, pp. 38–50, 2008.
- [14] Q. Zhang, "Existence of positive solutions for elliptic systems with nonstandard  $p(x)$ -growth conditions via sub-supersolution method," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 67, no. 4, pp. 1055–1067, 2007.
- [15] Q. Zhang, "Existence and asymptotic behavior of positive solutions to  $p(x)$ -Laplacian equations with singular nonlinearities," *Journal of Inequalities and Applications*, vol. 2007, Article ID 19349, 9 pages, 2007.
- [16] Q. Zhang, "Boundary blow-up solutions to  $p(x)$ -Laplacian equations with exponential nonlinearities," *Journal of Inequalities and Applications*, vol. 2008, Article ID 279306, 8 pages, 2008.
- [17] Q. Zhang, X. Liu, and Z. Qiu, "The method of subsuper solutions for weighted  $p(r)$ -Laplacian equation boundary value problems," *Journal of Inequalities and Applications*, vol. 2008, Article ID 621621, 19 pages, 2008.
- [18] V. V. Zhikov, "Averaging of functionals of the calculus of variations and elasticity theory," *Mathematics of the USSR-Izvestiya*, vol. 29, pp. 33–36, 1987.
- [19] D. D. Hai and R. Shivaji, "An existence result on positive solutions for a class of  $p$ -Laplacian systems," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 56, no. 7, pp. 1007–1010, 2004.
- [20] C. Chen, "On positive weak solutions for a class of quasilinear elliptic systems," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 62, no. 4, pp. 751–756, 2005.
- [21] J.-X. Feng and Z.-Q. Han, "Periodic solutions to differential systems with unbounded or periodic nonlinearities," *Journal of Mathematical Analysis and Applications*, vol. 323, no. 2, pp. 1264–1278, 2006.

- [22] R. Manásevich and J. Mawhin, "Periodic solutions for nonlinear systems with  $p$ -Laplacian-like operators," *Journal of Differential Equations*, vol. 145, no. 2, pp. 367–393, 1998.
- [23] J. Sun, Y. Ke, C. Jin, and J. Yin, "Existence of positive periodic solutions for the  $p$ -Laplacian system," *Applied Mathematics Letters*, vol. 20, no. 6, pp. 696–701, 2007.
- [24] Y. Tian and W. Ge, "Periodic solutions of non-autonomous second-order systems with a  $p$ -Laplacian," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 66, no. 1, pp. 192–203, 2007.