

Research Article

Rate of Convergence of a New Type Kantorovich Variant of Bleimann-Butzer-Hahn Operators

Lingju Chen¹ and Xiao-Ming Zeng²

¹ Department of Mathematics, Minjiang University, Fuzhou 350108, China

² Department of Mathematics, Xiamen University, Xiamen 361005, China

Correspondence should be addressed to Lingju Chen, lingjuchen@126.com
and Xiao-Ming Zeng, xmzeng@xmu.edu.cn

Received 28 September 2009; Accepted 16 November 2009

Recommended by Vijay Gupta

A new type Kantorovich variant of Bleimann-Butzer-Hahn operator J_n is introduced. Furthermore, the approximation properties of the operators J_n are studied. An estimate on the rate of convergence of the operators J_n for functions of bounded variation is obtained.

Copyright © 2009 L. Chen and X.-M. Zeng. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

In 1980, Bleimann et al. [1] introduced a sequence of positive linear Bernstein-type operators L_n (abbreviated in the following by BBH operators) defined on the infinite interval $I = [0, \infty)$ by

$$L_n(f, x) = \frac{1}{(1+x)^n} \sum_{k=0}^n \binom{n}{k} x^k f\left(\frac{k}{n+1-k}\right), \quad x \in I, \quad n \in \mathbb{N}, \quad (1.1)$$

where \mathbb{N} denotes the set of natural numbers.

Bleimann et al. [1] proved that $L_n(f, x) \rightarrow f(x)$ as $n \rightarrow \infty$ for $f \in C_b(I)$ (the space of all bounded continuous functions on I) and give an estimate on the rate of convergence of $L_n(f, x) \rightarrow f(x)$ measured with the second modulus of continuity of f .

In the present paper, we introduce a new type of Kantorovich variant of BBH operator J_n , also defined on I by

$$J_n(f, x) = \sum_{k=0}^n \binom{n}{k} p_x^k (1-p_x)^{n-k} \frac{\int_{I_k} f(t) dt}{\int_{I_k} dt}, \quad (1.2)$$

where $p_x = x/(1+x)$ ($x \geq 0$), $I_k = [k/(n+2-k), (k+1)/(n+1-k)]$, and dt is Lebesgue measure.

The operator (1.2) is different from another type of Kantorovich variant of BBH operator K_n :

$$K_n(f, x) = \frac{n+2}{(1+x)^n} \sum_{k=0}^n \binom{n}{k} x^k \int_{k/(n+2-k)}^{(k+1)/(n+1-k)} \frac{f(t)}{(1+t)^2} dt, \quad (1.3)$$

which was first considered by Abel and Ivan in [2]. The integrand function $f(t)/(1+t)^2$ in the operator (1.3) has been replaced with new integrand function $f(t)$ in the operator (1.2). In this paper we will study the approximation properties of J_n for the functions of bounded variation. The rate of convergence for functions of bounded variation was investigated by many authors such as Bojanić and Vuilleumier [3], Chêng [4], Guo and Khan [5], Zeng and Piriou [6], Gupta et al. [7], involving several different operators.

Throughout this paper the class of function Φ is defined as follows:

$$\Phi = \{f \mid f \text{ is of bounded variation on every finite subinterval of } I = [0, \infty)\}. \quad (1.4)$$

Our main result can be stated as follows.

Theorem 1.1. *Let $f \in \Phi$ and let $V_a^b(f)$ be the total variation of f on interval $[a, b]$. Then, for n sufficiently large, one has*

$$\left| J_n(f, x) - \frac{f(x+) - f(x-)}{2} \right| \leq \frac{5(1+x)}{2\sqrt{nx}} |f(x+) - f(x-)| + \frac{9(1+x)^2}{nx} \sum_{k=1}^n V_{x-x/\sqrt{k}}^{x+x/\sqrt{k}}(g_x) + O\left(\frac{1}{n}\right), \quad (1.5)$$

where

$$g_x(t) = \begin{cases} f(t) - f(x+), & x < t < \infty, \\ 0, & x = t, \\ f(t) - f(x-), & 0 \leq t < x. \end{cases} \quad (1.6)$$

2. Some Lemmas

In order to prove Theorem 1.1, we need the following lemmas for preparation. Lemma 2.1 is the well-known Berry-Esséen bound for the classical central limit theorem of probability theory. Its proof and further discussion can be founded in Feller [8, page 515].

Lemma 2.1. *Let $\{\xi\}_{i=1}^{\infty}$ be a sequence of independent and identically distributed random variables. And $0 < D\xi_1 < \infty$, $\beta_3 = E|\xi_1 - E\xi_1|^3 < +\infty$, then, there holds*

$$\max_{y \in \mathbb{R}} \left| P\left(\frac{1}{b_1\sqrt{n}} \sum_{k=1}^n (\xi_k - a_1) \leq y\right) - \frac{1}{\sqrt{2\pi}} \int_{-\infty}^y e^{-t^2/2} dt \right| < \frac{c}{\sqrt{n}} \frac{\beta_3}{b_1^3}, \quad (2.1)$$

where $a_1 = E\xi_1$, $b_1^2 = D\xi_1 = E(\xi_1 - E\xi_1)^2$, $1/\sqrt{2\pi} \leq c \leq 0.82$.

In addition, let $\{\xi\}_{i=1}^n$ be the random variables with two-point distribution

$$P_{\xi_i} = \begin{cases} x, & \xi_i = 1 \\ 1 - x, & \xi_i = 0, \end{cases} \quad (2.2)$$

where $i = 1, 2, \dots, n$. Then we can easily obtain that

$$a_1 = E\xi_1 = x, \quad b_1^2 = D\xi_1 = x(1-x), \quad \beta_3 = E|\xi_1 - E\xi_1|^3 \leq x(1-x)(2x^2 - 2x + 1). \quad (2.3)$$

Let $\eta_n = \sum_{i=1}^n \xi_i$, then we also have

$$P(\eta_n = k) = \binom{n}{k} x^k (1-x)^{n-k}, \quad k = 0, 1, \dots, n. \quad (2.4)$$

On the other hand, $J_n(f, x)$ can be written by following integral form:

$$J_n(f, x) = \sum_{k=0}^n p_{n,k} \left(\frac{x}{1+x} \right) \frac{\int_{I_k} f(t) dt}{\int_{I_k} dt} = \int_0^\infty f(t) H_n(x, t) dt, \quad (2.5)$$

where

$$H_n(x, t) = \sum_{k=0}^n p_{n,k} \left(\frac{x}{1+x} \right) \chi_k(t) \frac{1}{\int_{I_k} dt}, \quad \chi_k(t) = \begin{cases} 1, & t \in I_k, \\ 0, & t \notin I_k, \end{cases} \quad (2.6)$$

$I_k = [k/(n+2-k), (k+1)/(n+1-k)]$, $k = 0, 1, 2, \dots, n$. It is easy to verify that $\int_0^\infty H_n(x, u) du = 1$.

Lemma 2.2. *If $x \in (0, \infty)$ is fixed and n is sufficiently large, then*

(a) *for $0 \leq y < x$, there holds*

$$\int_0^y H_n(x, t) dt \leq \frac{1}{(x-y)^2} \frac{2x(1+x)^2}{n+1}, \quad (2.7)$$

(b) *for $x < z < \infty$, there holds*

$$\int_z^\infty H_n(x, t) dt \leq \frac{1}{(z-x)^2} \frac{2x(1+x)^2}{n+1}. \quad (2.8)$$

Proof. We first prove (a). Since $0 \leq y < x$, $t \in [0, y]$, then $(x-t)/(x-y) \geq 1$. Hence, we have

$$\int_0^y H_n(x, t) dt \leq \int_0^y \frac{(x-t)^2}{(x-y)^2} H_n(x, t) dt \leq \frac{1}{(x-y)^2} J_n((x-t)^2, x). \quad (2.9)$$

Direct calculation gives

$$J_n\left((x-t)^2, x\right) = \frac{x(1+x)^2}{n+1} + \frac{(1+x)^4}{3(n+1)(n+2)} + \frac{(1+x)^4(4x+1)}{3(n+1)(n+2)(n+3)} + o\left(n^{-4}\right). \quad (2.10)$$

Hence $\int_0^y H_n(x, t) dt \leq (1/(x-y)^2)(2x(1+x)^2/(n+1))$, for n sufficiently large.

The proof of (b) is similar. \square

Lemma 2.3 (see [9, Theorem 1] or, cf. [10]). *For every $x \in (0, 1)$, there holds*

$$p_{n,k}(x) = C_n^k x^k (1-x)^{n-k} \leq \frac{1}{\sqrt{2enx(1-x)}}. \quad (2.11)$$

3. Proof of Theorem 1.1

Let $f \in \Phi$, and $x \in I$, Bojanic-Cheng decomposition yields

$$f(t) = \frac{f(x+) - f(x-)}{2} + g_x(t) + \frac{f(x+) - f(x-)}{2} \operatorname{sgn}(t-x) + \delta_x(t) \left[f(x) - \frac{f(x+) - f(x-)}{2} \right], \quad (3.1)$$

where $g_x(t)$ is defined as in (1.6) and

$$\delta_x(t) = \begin{cases} 1, & t = x, \\ 0, & t \neq x. \end{cases} \quad (3.2)$$

Obviously, $J_n(\delta_x(t), x) = 0$. Thus it follows from (3.1) that

$$\left| J_n(f, x) - \frac{f(x+) - f(x-)}{2} \right| \leq |J_n(g_x, x)| + \left| J_n(\operatorname{sgn}(t-x), x) \frac{f(x+) - f(x-)}{2} \right|. \quad (3.3)$$

First of all, we estimate $|J_n(\operatorname{sgn}(t-x), x)|$

$$J_n(\operatorname{sgn}(t-x), x) = \sum_{k=0}^n \frac{(n+1-k)(n+2-k)}{n+2} p_{n,k} \left(\frac{x}{1+x} \right) \int_{I_k} \operatorname{sgn}(t-x) dt, \quad (3.4)$$

where $I_k = [k/(n+2-k), (k+1)/(n+1-k)]$.

Assuming that $x \in [k'/(n + 2 - k'), (k' + 1)/(n + 1 - k')]$, for some k' ($0 \leq k' \leq n$), then we have

$$\begin{aligned}
 J_n(\operatorname{sgn}(t - x), x) &= \sum_{k/(n+2-k) > x} p_{n,k} \left(\frac{x}{x+1} \right) - \sum_{(k+1)/(n+1-k) < x} p_{n,k} \left(\frac{x}{x+1} \right) \\
 &\quad + \frac{(n + 1 - k')(n + 2 - k')}{n + 2} p_{n,k'} \left(\frac{x}{1 + x} \right) \left(\int_x^{(k'+1)/(n+1-k)} dt - \int_{k'/(n+2-k)}^x dt \right) \\
 &= 1 - 2 \sum_{k/(n+2-k) \leq x} p_{n,k} \left(\frac{x}{x+1} \right) \\
 &\quad + 2 \frac{(n + 1 - k')(n + 2 - k')}{n + 2} p_{n,k'} \left(\frac{x}{1 + x} \right) \int_x^{(k'+1)/(n+1-k)} dt.
 \end{aligned} \tag{3.5}$$

Thus

$$|J_n(\operatorname{sgn}(t - x), x)| \leq \left| 1 - 2 \sum_{k/(n+2-k) \leq x} p_{n,k} \left(\frac{x}{x+1} \right) \right| + 2 p_{n,k'} \left(\frac{x}{1 + x} \right). \tag{3.6}$$

By Lemma 2.3 combining some direct computations, we can easily obtain

$$2 p_{n,k'} \left(\frac{x}{1 + x} \right) \leq \frac{2}{\sqrt{2en(x/(1 + x)) \cdot (1/(1 + x))}} \leq \frac{1 + x}{\sqrt{nx}}. \tag{3.7}$$

Set $y = x/(1 + x) < 1$, then by (2.4) and using Lemma 2.1, we have

$$\begin{aligned}
 &\left| 1 - 2 \sum_{k/(n+2-k) \leq x} p_{n,k} \left(\frac{x}{x+1} \right) \right| \\
 &= \left| 1 - 2 \sum_{k \leq (n+2)y} p_{n,k}(y) \right| = 2 \left| \frac{1}{2} - P(\eta_n \leq (n + 2)y) \right| \\
 &= 2 \left| \frac{1}{2} - P \left(\frac{\eta_n - ny}{\sqrt{ny(1 - y)}} \leq \frac{2y}{\sqrt{ny(1 - y)}} \right) \right| \\
 &= 2 \left| P \left(\frac{\eta_n - ny}{\sqrt{ny(1 - y)}} \leq \frac{2y}{\sqrt{ny(1 - y)}} \right) - \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{(2y-1)/\sqrt{ny(1-y)}} e^{-t^2/2} dt \right. \\
 &\quad \left. + \frac{1}{\sqrt{2\pi}} \int_0^{(2y-1)/\sqrt{ny(1-y)}} e^{-t^2/2} dt \right|
 \end{aligned}$$

$$\begin{aligned}
&\leq 2 \left| P \left(\frac{\eta_n - ny}{\sqrt{ny(1-y)}} \leq \frac{2y}{\sqrt{ny(1-y)}} \right) - \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{(2y-1)/\sqrt{ny(1-y)}} e^{-t^2/2} dt \right| \\
&\quad + \frac{2}{\sqrt{2\pi}} \int_0^{2y/\sqrt{ny(1-y)}} e^{-t^2/2} dt \\
&\leq \frac{2c\beta_3}{\sqrt{nb_1^3}} + \frac{2}{\sqrt{2\pi}} \int_0^{2y/\sqrt{ny(1-y)}} e^{-t^2/2} dt \\
&\leq \frac{2 \times 0.82 \times y(1-y)(2y^2 - 2y + 1)}{y(1-y)\sqrt{ny(1-y)}} + \frac{2}{\sqrt{2\pi}} \frac{2y}{\sqrt{ny(1-y)}} \\
&\leq \frac{4}{\sqrt{ny(1-y)}} = \frac{4(1+x)}{\sqrt{nx}}.
\end{aligned} \tag{3.8}$$

Thus, by (3.7), (3.8) we have

$$|J_n(\operatorname{sgn}(t-x), x)| \leq \frac{4(1+x)}{\sqrt{nx}} + \frac{1+x}{\sqrt{nx}} = \frac{5(1+x)}{\sqrt{nx}}. \tag{3.9}$$

Finally, we estimate $J_n(g_x, x)$.

First, interval $I = [0, \infty)$ can be decomposed into four parts as

$$D_1 = \left[0, x - \frac{x}{\sqrt{n}} \right], \quad D_2 = \left[x - \frac{x}{\sqrt{n}}, x + \frac{x}{\sqrt{n}} \right], \quad D_3 = \left[x + \frac{x}{\sqrt{n}}, 2x \right], \quad D_4 = [2x, +\infty]. \tag{3.10}$$

So $J_n(g_x, x)$ can be divided into four parts

$$J_n(g_x, x) = \int_0^{+\infty} g_x(t) H_n(x, t) dt = \Delta_{1,n}(g_x) + \Delta_{2,n}(g_x) + \Delta_{3,n}(g_x) + \Delta_{4,n}(g_x), \tag{3.11}$$

where $\Delta_{j,n}(g_x) = \int_{D_j} g_x(t) H_n(x, t) dt$.

Noticing $g_x(x) = 0$ and for $t \in D_2$, we have $g_x(t) = g_x(t) - g_x(x)$.

Thus

$$|\Delta_{2,n}(g_x)| \leq \int_{x-x/\sqrt{n}}^{x+x/\sqrt{n}} |g_x(t) - g_x(x)| H_n(x, t) dt \leq V_{x-x/\sqrt{n}}^{x+x/\sqrt{n}}(g_x) \leq \frac{1}{n} \sum_{k=1}^n V_{x-x/\sqrt{k}}^{x+x/\sqrt{k}}(g_x). \tag{3.12}$$

Next, let $y = x - x/\sqrt{n}$, $\lambda_n(x, t) = \int_0^t H_n(x, u) du$.

Now, we recall the Lebesgue-Stieltjes integral representation, and by using partial Lebesgue-Stieltjes integration, we get

$$\begin{aligned}
 |\Delta_{1,n}(g_x)| &= \left| \int_0^y g_x(t) d_t \lambda_n(x,t) \right| \\
 &= \left| g_x(y) \lambda_n(x,y) - \int_0^y \lambda_n(x,t) d_t g_x(t) \right| \\
 &= \left| (g_x(y) - g_x(x)) \lambda_n(x,y) - \int_0^y \lambda_n(x,t) d_t (g_x(t) - g_x(x)) \right| \\
 &\leq V_y^x(g_x) \lambda_n(x,y) + \int_0^y \lambda_n(x,t) d_t (-V_t^x(g_x)).
 \end{aligned} \tag{3.13}$$

An application of (a) in Lemma 2.2 yields

$$|\Delta_{1,n}(g_x)| \leq V_y^x(g_x) \frac{2x(1+x)^2}{(x-y)^2(n+1)} + \frac{2x(1+x)^2}{(n+1)} \int_0^y \frac{1}{(x-t)^2} d_t (-V_t^x(g_x)). \tag{3.14}$$

Furthermore, since

$$\int_0^y \frac{1}{(x-t)^2} d_t (-V_t^x(g_x)) = \frac{-V_y^x(g_x)}{(x-y)^2} + \frac{V_0^x(g_x)}{x^2} + 2 \int_0^y \frac{V_t^x(g_x)}{(x-t)^3} dt, \tag{3.15}$$

we have

$$|\Delta_{1,n}(g_x)| \leq \frac{2x(1+x)^2}{n+1} \left[\frac{V_0^x(g_x)}{x^2} + 2 \int_0^{x-x/\sqrt{n}} \frac{V_t^x(g_x)}{(x-t)^3} dt \right]. \tag{3.16}$$

Putting $t = x - x/\sqrt{u}$ in the last integral, we have

$$2 \int_0^{x-x/\sqrt{n}} \frac{V_t^x(g_x)}{(x-t)^3} dt = \frac{1}{x^2} \int_1^n V_{x-x/\sqrt{u}}^x(g_x) du \leq \frac{1}{x^2} \sum_{k=1}^n V_{x-x/\sqrt{k}}^x(g_x). \tag{3.17}$$

It follows from (3.16) and (3.17) that

$$|\Delta_{1,n}(g_x)| \leq \frac{2x(1+x)^2}{(n+1)x^2} \left(V_0^x(g_x) + \sum_{k=1}^n V_{x-x/\sqrt{k}}^x(g_x) \right) \leq \frac{4(1+x)^2}{(n+1)x} \sum_{k=1}^n V_{x-x/\sqrt{k}}^x(g_x). \tag{3.18}$$

By a similar method and using Lemma 2.2(b), we obtain

$$|\Delta_{3,n}(g_x)| \leq \frac{8(1+x)^2}{(n+1)x} \sum_{k=1}^n V_x^{x+x/\sqrt{k}}(g_x). \quad (3.19)$$

Now, the remainder of our work is to estimate $\Delta_{4,n}(g_x)$.

For $f(x)$ satisfying the growth condition $f(t) = O(t^r)$ for some positive integer r as $t \rightarrow +\infty$, we obviously have

$$|\Delta_{4,n}(g_x)| \leq \sum_{k/(n+2-k) > 2x} p_{n,k} \left(\frac{x}{1+x} \right) \frac{\int_{I_k} |g_x(t)| dt}{\int_{I_k} dt}. \quad (3.20)$$

Thus, for n sufficiently large, there exists a $M > 0$, such that the following inequalities hold:

$$\begin{aligned} |\Delta_{4,n}(g_x)| &\leq M \sum_{k/(n+2-k) > 2x} p_{n,k} \left(\frac{x}{1+x} \right) \frac{\int_{I_k} t^r dt}{\int_{I_k} dt} \\ &= M \sum_{k/(n+2-k) > 2(y/(1-y))} p_{n,k}(y) \frac{\int_{I_k} t^r dt}{\int_{I_k} dt} \\ &\leq M \sum_{k/(n+2-k) > 2(y/(1-y))} p_{n,k}(y) \left(\frac{k+1}{n+1-k} \right)^r, \end{aligned} \quad (3.21)$$

where $y = x/(1+x)$. By the definition of the Stirling numbers $S(r, s)$ of the second kind, we readily have

$$a^r = \sum_{s=1}^r S(r, s) a(a-1) \cdots (a-s+1), \quad r \in \mathbb{N}, \quad (3.22)$$

where the Stirling numbers $S(r, s)$ satisfy

$$S(n, 0) = \begin{cases} 1 & (n = 0), \\ 0 & (n \in \mathbb{N}). \end{cases} \quad (3.23)$$

Thus we can write

$$\sum_{k/(n+2-k) > 2(y/(1-y))} \left(\frac{k+1}{n+1-k} \right)^r p_{n,k}(y) = \sum_{s=1}^r S(r, s) A_s, \quad (3.24)$$

where

$$\begin{aligned}
 A_s &= \sum_{k/(n+2-k) > 2(y/(1-y))} \frac{(k+1)k \cdots (k-s+2)}{(n+1-k)^r} p_{n,k}(y) \\
 &= \sum_{k/(n+2-k) > 2(y/(1-y))} \frac{1}{(n+1-k)^r} \cdot \frac{n!(k+1)}{(k-s+1)!(n-k)!} y^k (1-y)^{n-k}.
 \end{aligned}
 \tag{3.25}$$

From $k/(n+2-k) > 2x, x/(1+x) = y$, we can easily find $k > (2n+4)y/(1+y)$. For a fixed $x > 0$, when $n > 2r + r/x$, we have $(k+1)/(k+1-s) < 2$. Thus there holds

$$A_s \leq 2 \sum_{k > (2n+4)y/(1+y)} \frac{1}{(n+1-k)^r} \cdot \frac{n!}{(k-s)!(n-k)!} y^k (1-y)^{n-k}.
 \tag{3.26}$$

Now using the similar method as that in the proof of Lemma 4 of [11], we deduce that

$$A_s \leq \frac{24r!n!y^{s-1}(1+y)^{r-s+2}}{(n+r-s)!(n+r-s+2)}, \quad \text{for } n > 2r + \frac{r}{x}.
 \tag{3.27}$$

From (3.21), (3.24), and (3.27), we obtain

$$\begin{aligned}
 |\Delta_{4,n}(g_x)| &\leq M \sum_{k/(n+2-k) > 2(y/(1-y))} p_{n,k}(y) \left(\frac{k+1}{n+1-k} \right)^r \\
 &= M \sum_{s=1}^r S(r,s) A_s = O\left(\frac{1}{n}\right).
 \end{aligned}
 \tag{3.28}$$

Finally, by combining (3.12), (3.18), (3.19), and (3.28), we deduce that

$$\begin{aligned}
 |J_n(g_x(t), x)| &\leq |\Delta_{1,n}(g_x)| + |\Delta_{2,n}(g_x)| + |\Delta_{3,n}(g_x)| + |\Delta_{4,n}(g_x)| \\
 &\leq \frac{4(1+x)^2}{(n+1)x} \sum_{k=1}^n V_{x-x/\sqrt{k}}^x(g_x) + \frac{1}{n} V_{x-x/\sqrt{k}}^{x+x/\sqrt{k}}(g_x) + \frac{8(1+x)^2}{(n+1)x} \sum_{k=1}^n V_x^{x+x/\sqrt{k}}(g_x) + O\left(\frac{1}{n}\right) \\
 &\leq \frac{1}{n} V_{x-x/\sqrt{k}}^{x+x/\sqrt{k}}(g_x) + \frac{8(1+x)^2}{(n+1)x} \sum_{k=1}^n V_{x-x/\sqrt{k}}^{x+x/\sqrt{k}}(g_x) + O\left(\frac{1}{n}\right) \\
 &\leq \frac{9(1+x)^2}{nx} \sum_{k=1}^n V_{x-x/\sqrt{k}}^{x+x/\sqrt{k}}(g_x) + O\left(\frac{1}{n}\right).
 \end{aligned}
 \tag{3.29}$$

Theorem 1.1 now follows from (3.3), (3.9), and (3.29).

Acknowledgments

This work is supported by the National Natural Science Foundation of China and the Fujian Provincial Science Foundation of China. The authors thank the associate editor and the referee(s) for their several important comments and suggestions which improve the quality of the paper.

References

- [1] G. Bleimann, P. L. Butzer, and L. Hahn, "A Bernstein-type operator approximating continuous functions on the semi-axis," *Indagationes Mathematicae*, vol. 83, no. 3, pp. 255–262, 1980.
- [2] U. Abel and M. Ivan, "A Kantorovich variant of the Bleimann, Butzer and Hahn operators," *Rendiconti del Circolo Matematico di Palermo. Serie II. Supplemento*, vol. 68, pp. 205–218, 2002.
- [3] R. Bojanić and M. Vuilleumier, "On the rate of convergence of Fourier-Legendre series of functions of bounded variation," *Journal of Approximation Theory*, vol. 31, no. 1, pp. 67–79, 1981.
- [4] F. Chêng, "On the rate of convergence of Bernstein polynomials of functions of bounded variation," *Journal of Approximation Theory*, vol. 39, no. 3, pp. 259–274, 1983.
- [5] S. S. Guo and M. K. Khan, "On the rate of convergence of some operators on functions of bounded variation," *Journal of Approximation Theory*, vol. 58, no. 1, pp. 90–101, 1989.
- [6] X.-M. Zeng and A. Piriou, "On the rate of convergence of two Bernstein-Bézier type operators for bounded variation functions," *Journal of Approximation Theory*, vol. 95, no. 3, pp. 369–387, 1998.
- [7] V. Gupta, U. Abel, and M. Ivan, "Rate of convergence of beta operators of second kind for functions with derivatives of bounded variation," *International Journal of Mathematics and Mathematical Sciences*, vol. 2005, no. 23, pp. 3827–3833, 2005.
- [8] W. Feller, *An Introduction to Probability Theory and Its Applications*, vol. 2, John Wiley & Sons, New York, NY, USA, 2nd edition, 1971.
- [9] X.-M. Zeng, "Bounds for Bernstein basis functions and Meyer-König and Zeller basis functions," *Journal of Mathematical Analysis and Applications*, vol. 219, no. 2, pp. 364–376, 1998.
- [10] X.-M. Zeng and J.-N. Zhao, "Exact bounds for some basis functions of approximation operators," *Journal of Inequalities and Applications*, vol. 6, no. 5, pp. 563–575, 2001.
- [11] H. M. Srivastava and V. Gupta, "Rate of convergence for the Bézier variant of the Bleimann-Butzer-Hahn operators," *Applied Mathematics Letters*, vol. 18, no. 8, pp. 849–857, 2005.