

Research Article

Convergence Theorems on Generalized Equilibrium Problems and Fixed Point Problems with Applications

Yan Hao,¹ Sun Young Cho,² and Xiaolong Qin³

¹ School of Mathematics, Physics and Information Science, Zhejiang Ocean University, Zhoushan 316004, China

² Department of Mathematics, Gyeongsang National University, Jinju 660-701, South Korea

³ Department of Mathematics, Hangzhou Normal University, Hangzhou 310036, China

Correspondence should be addressed to Yan Hao, zjhaoyan@yahoo.cn

Received 2 October 2009; Accepted 23 January 2010

Academic Editor: Jong Kim

Copyright © 2010 Yan Hao et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The purpose of this work is to introduce an iterative method for finding a common element of a solution set of a generalized equilibrium problem, of a solution set solutions of a variational inequality problem and of a fixed point set of a strict pseudocontraction. Strong convergence theorems are established in the framework of Hilbert spaces.

1. Introduction and Preliminaries

Let H be a real Hilbert space, C a nonempty closed and convex subset of H , and $B : C \rightarrow H$ a nonlinear mapping. Recall the following definitions.

(a) The mapping B is said to be *monotone* if

$$\langle Bx - By, x - y \rangle \geq 0, \quad \forall x, y \in C. \quad (1.1)$$

(b) B is said to be β -*strongly monotone* if there exists a constant $\beta > 0$ such that

$$\langle Bx - By, x - y \rangle \geq \beta \|x - y\|^2, \quad \forall x, y \in C. \quad (1.2)$$

(c) B is said to be β -inverse-strongly monotone if there exists a constant $\beta > 0$ such that

$$\langle Bx - By, x - y \rangle \geq \beta \|Bx - By\|^2, \quad \forall x, y \in C. \quad (1.3)$$

The classical variational inequality problem is to find $u \in C$ such that

$$\langle Bu, v - u \rangle \geq 0, \quad \forall v \in C. \quad (1.4)$$

In this paper, we use $VI(C, B)$ to denote the solution set of the problem (1.4). One can easily see that the variational inequality problem is equivalent to a fixed point problem. $u \in C$ is a solution to the problem (1.4) if and only if u is a fixed point of the mapping $P_C(I - \lambda B)$, where $\lambda > 0$ is a constant and I is the identity mapping.

Let $S : C \rightarrow C$ be a nonlinear mapping. In this paper, we use $F(S)$ to denote the fixed point set of S . Recall the following definitions.

(d) The mapping S is said to be *nonexpansive* if

$$\|Sx - Sy\| \leq \|x - y\|, \quad \forall x, y \in C. \quad (1.5)$$

(e) S is *strictly pseudocontractive* with a constant $k \in [0, 1)$ if

$$\|Sx - Sy\|^2 \leq \|x - y\|^2 + k\|(I - S)x - (I - S)y\|^2, \quad \forall x, y \in C. \quad (1.6)$$

For such a case, S is called a *k-strict pseudocontraction*.

(f) S is said to be *pseudocontractive* if

$$\|Sx - Sy\|^2 \leq \|x - y\|^2 + \|(I - S)x - (I - S)y\|^2, \quad \forall x, y \in C. \quad (1.7)$$

Clearly, the class of strict pseudocontractions falls into the one between a class of nonexpansive mappings and a class of pseudocontractions.

Recently, many authors considered the problem of finding a common element of the solution set of the variational inequality (1.4) and of fixed point set of a nonexpansive mapping in Hilbert spaces; see, for examples, [1–5] and the references therein.

In 2005, Iiduka and Takahashi [2] obtained the following theorem in a real Hilbert space.

Theorem IT. *Let C be a closed convex subset of a real Hilbert space H . Let B be an α -inverse-strongly monotone mapping of C into H and let S be a nonexpansive mapping of C into itself such that $F(S) \cap VI(C, B) \neq \emptyset$. Suppose $x_1 = x \in C$ and $\{x_n\}$ is given by*

$$x_{n+1} = \alpha_n x + (1 - \alpha_n) SP_C(x_n - \lambda_n Bx_n) \quad (1.8)$$

for every $n = 1, 2, \dots$, where $\{\alpha_n\}$ is a sequence in $[0, 1)$ and $\{\lambda_n\}$ is a sequence in $[a, b]$. If $\{\alpha_n\}$ and $\{\lambda_n\}$ are chosen so that $\{\lambda_n\} \in [a, b]$ for some a, b with $0 < a < b < 2\alpha$,

$$\lim_{n \rightarrow \infty} \alpha_n = 0, \quad \sum_{n=1}^{\infty} \alpha_n = \infty, \quad \sum_{n=1}^{\infty} |\alpha_{n+1} - \alpha_n| < \infty, \quad \sum_{n=1}^{\infty} |\lambda_{n+1} - \lambda_n| < \infty, \quad (1.9)$$

then $\{x_n\}$ converges strongly to $P_{F(S) \cap VI(C, B)} x$.

Let A be an inverse-strongly monotone mapping, and F a bifunction of $C \times C$ into \mathbb{R} , where \mathbb{R} is the set of real numbers. We consider the following equilibrium problem:

$$\text{find } z \in C \text{ such that } F(z, y) + \langle Az, y - z \rangle \geq 0, \quad \forall y \in C. \quad (1.10)$$

In this paper, the set of such $z \in C$ is denoted by $EP(F, A)$, that is,

$$EP(F, A) = \{z \in C : F(z, y) + \langle Az, y - z \rangle \geq 0, \forall y \in C\}. \quad (1.11)$$

In the case of $A \equiv 0$, the zero mapping, the problem (1.10) is reduced to

$$\text{Find } z \in C \text{ such that } F(z, y) \geq 0, \quad \forall y \in C. \quad (1.12)$$

In this paper, we use $EP(F)$ to denote the solution set of the problem (1.12), which was studied by many others; see, for examples, [1, 3, 6–23] and the reference therein. In the case of $F \equiv 0$, the problem (1.10) is reduced to the classical variational inequality (1.4). The problem (1.10) is very general in the sense that it includes, as special cases, optimization problems, variational inequalities, minimax problems, the Nash equilibrium problem in noncooperative games, and others; see, for instances, [15, 24].

To study the problems (1.10) and (1.12), we may assume that the bifunction $F : C \times C \rightarrow \mathbb{R}$ satisfies the following conditions:

- (A1) $F(x, x) = 0$ for all $x \in C$;
- (A2) F is monotone, that is, $F(x, y) + F(y, x) \leq 0$ for all $x, y \in C$;
- (A3) for each $x, y, z \in C$,

$$\limsup_{t \downarrow 0} F(tz + (1-t)x, y) \leq F(x, y); \quad (1.13)$$

- (A4) for each $x \in C$, $y \mapsto F(x, y)$ is convex and weakly lower semicontinuous.

Recently, S. Takahashi and W. Takahashi [21] considered the problem (1.12) by introducing an iterative method in a Hilbert space. To be more precise, they proved the following theorem.

Theorem TT 1. Let C be a nonempty closed convex subset of H . Let F be a bifunction from $C \times C$ to \mathbb{R} satisfying (A1)–(A4), and let S be a nonexpansive mapping of C into H such that $F(S) \cap \text{EP}(F) \neq \emptyset$. Let f be a contraction of H into itself, and let $\{x_n\}$ and $\{y_n\}$ be sequences generated by $x_1 \in H$ and

$$\begin{aligned} F(y_n, u) + \frac{1}{r_n} \langle u - y_n, y_n - x_n \rangle &\geq 0, \quad \forall u \in C, \\ x_{n+1} &= \alpha_n f(x_n) + (1 - \alpha_n) S y_n, \quad n \geq 1, \end{aligned} \quad (1.14)$$

where $\{\alpha_n\} \in [0, 1]$ and $\{r_n\} \subset (0, \infty)$ satisfy

$$\begin{aligned} \lim_{n \rightarrow \infty} \alpha_n &= 0, \quad \sum_{n=1}^{\infty} \alpha_n = \infty, \quad \sum_{n=1}^{\infty} |\alpha_{n+1} - \alpha_n| < \infty, \\ \liminf_{n \rightarrow \infty} r_n &> 0, \quad \sum_{n=1}^{\infty} |r_{n+1} - r_n| < \infty. \end{aligned} \quad (1.15)$$

Then $\{x_n\}$ and $\{y_n\}$ converge strongly to $z \in F(S) \cap \text{EP}(F)$, where $z = P_{F(S) \cap \text{EP}(F)} f(z)$.

Very recently, S. Takahashi and W. Takahashi [22] further considered the problem (1.10). Strong convergence theorems of common elements are established. More precisely, they obtained the following result.

Theorem TT 2. Let C be a closed convex subset of a real Hilbert space H , and let $F : C \times C \rightarrow \mathbb{R}$ be a bifunction satisfying (A1), (A2), (A3) and (A4). Let A be an α -inverse-strongly monotone mapping of C into H and let S be a nonexpansive mapping of C into itself such that $F(S) \cap \text{EP}(F, A) \neq \emptyset$. Let $u \in C$ and $x_1 \in C$ and let $\{z_n\} \subset C$ and $\{x_n\} \subset C$ be sequences generated by

$$\begin{aligned} F(z_n, y) + \langle Ax_n, y - z_n \rangle + \frac{1}{\lambda_n} \langle y - z_n, z_n - x_n \rangle &\geq 0, \quad \forall y \in C, \\ x_{n+1} &= \beta_n x_n + (1 - \beta_n) S[\alpha_n u + (1 - \alpha_n) z_n], \quad \forall n \geq 1, \end{aligned} \quad (1.16)$$

where $\{\alpha_n\} \subset [0, 1]$, $\{\beta_n\} \subset [0, 1]$, and $\{\lambda_n\} \subset [0, 2\alpha]$ satisfy

$$\begin{aligned} 0 < c \leq \beta_n \leq d < 1, \quad 0 < a \leq \lambda_n \leq b < 2\alpha, \\ \lim_{n \rightarrow \infty} (\lambda_n - \lambda_{n+1}) &= 0, \quad \lim_{n \rightarrow \infty} \alpha_n = 0, \quad \sum_{n=1}^{\infty} \alpha_n = \infty. \end{aligned} \quad (1.17)$$

Then, $\{x_n\}$ converges strongly to $z = P_{F(S) \cap \text{EP}(F, A)} u$.

In this paper, motivated by Theorem IT, Theorem TT1, and Theorem TT2, we introduce a general iterative method for the problem of finding a common element of a solution set of a generalized equilibrium problem (1.10), of a solution set of a variational inequality problem (1.4), and of a fixed point set of a strict pseudocontraction. Strong convergence theorems are established in the framework of Hilbert spaces. The results presented in this paper improve and extend the corresponding results announced by many others.

In order to prove our main results, we need the following lemmas.
The following lemmas can be found in [11, 24].

Lemma 1.1. *Let C be a nonempty closed convex subset of H and let $F : C \times C \rightarrow \mathbb{R}$ be a bifunction satisfying (A1)–(A4). Then, for any $r > 0$ and $x \in H$, there exists $z \in C$ such that*

$$F(z, y) + \frac{1}{r} \langle y - z, z - x \rangle \geq 0, \quad \forall y \in C. \quad (1.18)$$

Further, define a mapping T_r by

$$T_r x = \left\{ z \in C : F(z, y) + \frac{1}{r} \langle y - z, z - x \rangle \geq 0, \forall y \in C \right\} \quad (1.19)$$

for all $r > 0$ and $x \in H$. Then, the following hold.

- (1) T_r is single-valued;
- (2) T_r is firmly nonexpansive, that is, for any $x, y \in H$,

$$\|T_r x - T_r y\|^2 \leq \langle T_r x - T_r y, x - y \rangle; \quad (1.20)$$

- (3) $F(T_r) = \text{EP}(F)$;
- (4) $\text{EP}(F)$ is closed and convex.

Lemma 1.2 (see [25]). *Let C be a nonempty closed convex subset of a real Hilbert space H and $S : C \rightarrow C$ a k -strict pseudocontraction. Define $S_\alpha : C \rightarrow C$ by $S_\alpha x = \alpha x + (1 - \alpha)Sx$ for each $x \in C$. Then, as $\alpha \in [k, 1)$, S_α is nonexpansive and $F(S_\alpha) = F(S)$.*

Lemma 1.3 (see [26]). *Let $\{x_n\}$ and $\{y_n\}$ be bounded sequences in a Banach space X and let $\{\beta_n\}$ be a sequence in $[0, 1]$ with*

$$0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} \beta_n < 1. \quad (1.21)$$

Suppose that $x_{n+1} = (1 - \beta_n)y_n + \beta_n x_n$ for all integers $n \geq 0$ and

$$\limsup_{n \rightarrow \infty} (\|y_{n+1} - y_n\| - \|x_{n+1} - x_n\|) \leq 0. \quad (1.22)$$

Then $\lim_{n \rightarrow \infty} \|y_n - x_n\| = 0$.

The following lemma can be deduced from Bruck [8].

Lemma 1.4. *Let C be a closed convex subset of a strictly convex Banach space E . Let S_1, S_2 , and S_3 be three nonexpansive mappings on C . Suppose $\bigcap_{n=1}^3 F(S_n)$ is nonempty. Let a_1, a_2 , and a_3 be three constant in $(0, 1)$. Then the mapping S on C defined by*

$$Sx = a_1 S_1 x + a_2 S_2 x + a_3 S_3 x \quad (1.23)$$

for $x \in C$ is well defined, nonexpansive, and $F(S) = \bigcap_{n=1}^3 F(S_n)$ holds.

Lemma 1.5 (see [6]). *Let H be a real Hilbert space, C a nonempty closed and convex subset of E , and $S : C \rightarrow C$ a nonexpansive mapping. Then $I - S$ is demiclosed at zero.*

Lemma 1.6 (see [14]). *Let H be a real Hilbert space, C a nonempty closed and convex subset of E and $S : C \rightarrow C$ a k -strict pseudocontraction. Then $F(S)$ is closed and convex.*

Lemma 1.7 (see [27]). *Assume that $\{\alpha_n\}$ is a sequence of nonnegative real numbers such that*

$$\alpha_{n+1} \leq (1 - \gamma_n)\alpha_n + \delta_n, \quad (1.24)$$

where $\{\gamma_n\}$ is a sequence in $(0, 1)$ and $\{\delta_n\}$ is a sequence such that

- (a) $\sum_{n=1}^{\infty} \gamma_n = \infty$;
- (b) $\limsup_{n \rightarrow \infty} \delta_n / \gamma_n \leq 0$ or $\sum_{n=1}^{\infty} |\delta_n| < \infty$.

Then $\lim_{n \rightarrow \infty} \alpha_n = 0$.

2. Main Results

Theorem 2.1. *Let C be a nonempty closed and convex subset of a real Hilbert space H and F a bifunction from $C \times C$ to \mathbb{R} satisfying (A1)–(A4). Let A be an α -inverse-strongly monotone mapping of C into H and B a β -inverse-strongly monotone mapping of C into H . Let $S : C \rightarrow C$ be a k -strict pseudocontraction with a fixed point. Assume that $\Omega := \text{EP}(F, A) \cap F(S) \cap \text{VI}(C, B) \neq \emptyset$. Let $\{x_n\}$ be a sequence in C generated by*

$$\begin{aligned} x_1 &\in C, \quad \text{chosen arbitrarily,} \\ F(u_n, y) + \langle Ax_n, y - u_n \rangle + \frac{1}{r} \langle y - u_n, u_n - x_n \rangle &\geq 0, \quad \forall y \in C, \\ v_n &= P_C(x_n - \lambda Bx_n), \\ y_n &= \delta_n x_n + (1 - \delta_n) Sx_n, \\ x_{n+1} &= \alpha_n u + \beta_n x_n + \gamma_n (\mu_{(1,n)} y_n + \mu_{(2,n)} u_n + \mu_{(3,n)} v_n), \quad \forall n \geq 1, \end{aligned} \quad (2.1)$$

where u is a fixed element in C , $\{\alpha_n\}$, $\{\beta_n\}$, $\{\gamma_n\}$, $\{\mu_{(1,n)}\}$, $\{\mu_{(2,n)}\}$ and $\{\mu_{(3,n)}\}$ are sequences in $(0, 1)$, $\{\delta_n\}$ is sequence in $[k, 1)$, $r \in (0, 2\alpha]$ and $\lambda \in (0, 2\beta]$. Assume that the above control sequences satisfy the following restrictions

- (R1) $\alpha_n + \beta_n + \gamma_n = \mu_{(1,n)} + \mu_{(2,n)} + \mu_{(3,n)} = 1$, for all $n \geq 1$;
- (R2) $\lim_{n \rightarrow \infty} \alpha_n = 0$, $\sum_{n=1}^{\infty} \alpha_n = \infty$;
- (R3) $0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} \beta_n < 1$;
- (R4) $\lim_{n \rightarrow \infty} \delta_n = \delta \in [k, 1)$ and $\lim_{n \rightarrow \infty} \mu_{(i,n)} = \mu_i \in (0, 1)$ for all $1 \leq i \leq 3$.

Then the sequence $\{x_n\}$ defined by the iterative process (2.1) converges strongly to $\bar{x} = P_{\Omega} u$.

Proof. The proof is divided into six steps.

Step 1. Show that $P_{\Omega} u$ is well defined.

From Lemma 1.6, we see that $F(S)$ is closed and convex. On the other hand, we see that the mapping $I - rA$, where $r \in (0, 2\alpha]$, is nonexpansive. Indeed, for any $x, y \in C$, we have that

$$\begin{aligned} \|(I - rA)x - (I - rA)y\|^2 &= \|x - y - r(Ax - Ay)\|^2 \\ &= \|x - y\|^2 - 2r\langle x - y, Ax - Ay \rangle + r^2\|Ax - Ay\|^2 \\ &\leq \|x - y\|^2 - 2r\alpha\|Ax - Ay\|^2 + r^2\|Ax - Ay\|^2 \quad (2.2) \\ &= \|x - y\|^2 - r(2\alpha - r)\|Ax - Ay\|^2 \\ &\leq \|x - y\|^2. \end{aligned}$$

This shows that $I - rA$ is nonexpansive mapping. Similarly, we can prove that $I - \lambda B$, where $\lambda \in (0, 2\beta]$ is nonexpansive. It follows that $\text{VI}(C, B) = F(P_C(I - \lambda B))$, for all $\lambda > 0$ is closed and convex. From Lemma 1.1, we see that $\text{EP}(F, A) = F(T_r(I - rA))$. Since $T_r(I - rA)$ is nonexpansive, we obtain that $\text{EP}(F, A)$ is closed and convex. This shows that $P_\Omega u$ is well defined.

Step 2. Show that $\{x_n\}$ is bounded.

Put $T_n = \delta_n I + (1 - \delta_n)S$ for each $n \geq 1$. In view of Lemma 1.2 and (R4), we obtain that T_n is nonexpansive and $F(T_n) = F(S)$. Letting $x^* \in \Omega$, we obtain that

$$x^* = T_r(I - rA)x^* = T_n x^* = Sx^* = P_C(I - \lambda B)x^*, \quad \forall n \geq 1. \quad (2.3)$$

Note that $y_n = T_n x_n$ for each $n \geq 1$. It follows that $\|y_n - x^*\| \leq \|x_n - x^*\|$. Putting

$$e_n = \mu_{(1,n)}y_n + \mu_{(2,n)}u_n + \mu_{(3,n)}v_n, \quad \forall n \geq 1, \quad (2.4)$$

we have that

$$\begin{aligned} \|e_n - x^*\| &\leq \mu_{(1,n)}\|y_n - x^*\| + \mu_{(2,n)}\|u_n - x^*\| + \mu_{(3,n)}\|v_n - x^*\| \\ &= \mu_{(1,n)}\|y_n - x^*\| + \mu_{(2,n)}\|T_r(I - rA)x_n - T_r(I - rA)x^*\| \\ &\quad + \mu_{(3,n)}\|P_C(I - \lambda A)x_n - P_C(I - \lambda B)x^*\| \quad (2.5) \\ &\leq \mu_{(1,n)}\|x_n - x^*\| + \mu_{(2,n)}\|x_n - x^*\| + \mu_{(3,n)}\|x_n - x^*\| \\ &= \|x_n - x^*\|. \end{aligned}$$

It follows that

$$\begin{aligned}
 \|x_{n+1} - x^*\| &= \|\alpha_n u + \beta_n x_n + \gamma_n e_n - x^*\| \\
 &\leq \alpha_n \|u - x^*\| + \beta_n \|x_n - x^*\| + \gamma_n \|e_n - x^*\| \\
 &\leq \alpha_n \|u - x^*\| + \beta_n \|x_n - x^*\| + \gamma_n \|x_n - x^*\| \\
 &= \alpha_n \|u - x^*\| + (1 - \alpha_n) \|x_n - x^*\| \\
 &\leq \max\{\|u - x^*\|, \|x_1 - x^*\|\}.
 \end{aligned} \tag{2.6}$$

This shows that the sequence $\{x_n\}$ is bounded. Note that

$$\begin{aligned}
 \|u_n - x^*\| &= \|T_r(I - rA)x_n - x^*\| \leq \|x_n - x^*\|, \\
 \|v_n - x^*\| &= \|P_C(x_n - \lambda Bx_n) - x^*\| \leq \|x_n - x^*\|.
 \end{aligned} \tag{2.7}$$

This proves that the sequences $\{u_n\}$ and $\{v_n\}$ are bounded, too.

Step 3. Show that $x_{n+1} - x_n \rightarrow 0$ as $n \rightarrow \infty$.

Note that

$$\begin{aligned}
 \|y_{n+1} - y_n\| &= \|T_{n+1}x_{n+1} - T_nx_n\| \\
 &\leq \|T_{n+1}x_{n+1} - T_{n+1}x_n\| + \|T_{n+1}x_n - T_nx_n\| \\
 &\leq \|x_{n+1} - x_n\| + |\delta_{n+1} - \delta_n| M_1,
 \end{aligned} \tag{2.8}$$

where M_1 is an appropriate constant such that $M_1 = \sup_{n \geq 1} \{\|x_n - Sx_n\|\}$. On the other hand, we have

$$\begin{aligned}
 \|v_{n+1} - v_n\| &= \|P_C(x_{n+1} - \lambda Bx_{n+1}) - P_C(x_n - \lambda Bx_n)\| \leq \|x_{n+1} - x_n\|, \\
 \|u_{n+1} - u_n\| &= \|T_r(I - rA)x_{n+1} - T_r(I - rA)x_n\| \leq \|x_{n+1} - x_n\|.
 \end{aligned} \tag{2.9}$$

It follows from (2.1) and (2.9) that

$$\begin{aligned}
 \|e_{n+1} - e_n\| &= \|\mu_{(1,(n+1))}y_{n+1} + \mu_{(2,(n+1))}u_{n+1} + \mu_{(3,(n+1))}v_{n+1} \\
 &\quad - (\mu_{(1,n)}y_n + \mu_{(2,n)}u_n + \mu_{(3,n)}v_n)\| \\
 &\leq \mu_{(1,(n+1))} \|y_{n+1} - y_n\| + \|y_n\| |\mu_{(1,(n+1))} - \mu_{(1,n)}| \\
 &\quad + \mu_{(2,(n+1))} \|u_{n+1} - u_n\| + \|u_n\| |\mu_{(2,(n+1))} - \mu_{(2,n)}| \\
 &\quad + \mu_{(3,(n+1))} \|v_{n+1} - v_n\| + \|v_n\| |\mu_{(3,(n+1))} - \mu_{(3,n)}| \leq \|x_{n+1} - x_n\| \\
 &\quad + M_2 (|\mu_{(1,(n+1))} - \mu_{(1,n)}| + |\mu_{(2,(n+1))} - \mu_{(2,n)}| + |\mu_{(3,(n+1))} - \mu_{(3,n)}| + |\delta_{n+1} - \delta_n|),
 \end{aligned} \tag{2.10}$$

where M_2 is an appropriate constant such that

$$M_2 \geq \max \left\{ \sup_{n \geq 1} \|y_n\|, \sup_{n \geq 1} \|u_n\|, \sup_{n \geq 1} \|v_n\|, M_1 \right\}. \tag{2.11}$$

Put $l_n = (x_{n+1} - \beta_n x_n) / (1 - \beta_n)$, for each $n \geq 1$, that is,

$$x_{n+1} = (1 - \beta_n)l_n + \beta_n x_n, \quad \forall n \geq 1. \tag{2.12}$$

Now, we compute $\|l_{n+1} - l_n\|$. Notice that

$$\begin{aligned} l_{n+1} - l_n &= \frac{\alpha_{n+1}u + \gamma_{n+1}e_{n+1}}{1 - \beta_{n+1}} - \frac{\alpha_n u + \gamma_n e_n}{1 - \beta_n} \\ &= \frac{\alpha_{n+1}}{1 - \beta_{n+1}}u + \frac{1 - \beta_{n+1} - \alpha_{n+1}}{1 - \beta_{n+1}}e_{n+1} - \frac{\alpha_n}{1 - \beta_n}u - \frac{1 - \beta_n - \alpha_n}{1 - \beta_n}e_n \\ &= \frac{\alpha_{n+1}}{1 - \beta_{n+1}}(u - e_{n+1}) + \frac{\alpha_n}{1 - \beta_n}(e_n - u) + e_{n+1} - e_n. \end{aligned} \tag{2.13}$$

It follows that

$$\|l_{n+1} - l_n\| \leq \frac{\alpha_{n+1}}{1 - \beta_{n+1}} \|u - e_{n+1}\| + \frac{\alpha_n}{1 - \beta_n} \|e_n - u\| + \|e_{n+1} - e_n\|. \tag{2.14}$$

Substituting (2.10) into (2.14), we arrive at

$$\begin{aligned} &\|l_{n+1} - l_n\| - \|x_{n+1} - x_n\| \\ &\leq \frac{\alpha_{n+1}}{1 - \beta_{n+1}} \|u - e_{n+1}\| + \frac{\alpha_n}{1 - \beta_n} \|e_n - u\| \\ &\quad + M_2 (|\mu_{(1,(n+1))} - \mu_{(1,n)}| + |\mu_{(2,(n+1))} - \mu_{(2,n)}| + |\mu_{(3,(n+1))} - \mu_{(3,n)}| + |\delta_{n+1} - \delta_n|). \end{aligned} \tag{2.15}$$

It follows from the restrictions (R2)–(R4) that

$$\limsup_{n \rightarrow \infty} (\|l_{n+1} - l_n\| - \|x_{n+1} - x_n\|) < 0. \tag{2.16}$$

From Lemma 1.3, we obtain that

$$\lim_{n \rightarrow \infty} \|l_n - x_n\| = 0. \tag{2.17}$$

From (2.12), we see that

$$x_{n+1} - x_n = (1 - \beta_n)(l_n - x_n). \tag{2.18}$$

In view of (2.17), we get that

$$\lim_{n \rightarrow \infty} \|x_{n+1} - x_n\| = 0. \quad (2.19)$$

Step 4. Show that $x_n - e_n \rightarrow 0$ as $n \rightarrow \infty$.

From the iterative process (2.1), we have

$$\gamma_n(e_n - x_n) = x_{n+1} - x_n + \alpha_n(x_n - u). \quad (2.20)$$

This implies that

$$\gamma_n \|e_n - x_n\| \leq \|x_{n+1} - x_n\| + \alpha_n \|x_n - u\|. \quad (2.21)$$

It follows from the restrictions (R2) and (R3) that we arrive at

$$\lim_{n \rightarrow \infty} \|e_n - x_n\| = 0. \quad (2.22)$$

Step 5. Show that $\limsup_{n \rightarrow \infty} \langle u - \bar{x}, x_n - \bar{x} \rangle \leq 0$, where $\bar{x} = P_\Omega u$.

To show that, we can choose a sequence $\{x_{n_i}\}$ of $\{x_n\}$ such that

$$\limsup_{n \rightarrow \infty} \langle u - \bar{x}, x_n - \bar{x} \rangle = \lim_{i \rightarrow \infty} \langle u - \bar{x}, x_{n_i} - \bar{x} \rangle. \quad (2.23)$$

Since $\{x_{n_i}\}$ is bounded, we see that there exists a subsequence $\{x_{n_{i_j}}\}$ of $\{x_{n_i}\}$ which converges weakly to ξ . Without loss of generality, we may assume that $x_{n_i} \rightharpoonup \xi$.

Next, we show that $\xi \in \Omega = F(S) \cap VI(C, B) \cap EP(F, A)$. In fact, define a mapping $Q : C \rightarrow C$ by

$$Qx = \mu_1 Tx + \mu_2 T_r(I - rA)x + \mu_3 P_C(I - \lambda B), \quad \forall x \in C, \quad (2.24)$$

where $T = \delta I + (1 - \delta)S$. Note that T is nonexpansive and $F(T) = F(S)$. From Lemma 1.4, we see that Q is a nonexpansive mapping such that

$$F(Q) = F(T) \cap F(T_r(I - rA)) \cap F(P_C(I - \lambda B)) = F(S) \cap EP(F, A) \cap VI(C, B). \quad (2.25)$$

On the other hand, we have

$$\begin{aligned} \|e_n - Qx_n\| &= \|(\mu_{(1,n)}y_n + \mu_{(2,n)}T_r(I - rA)x_n + \mu_{(3,n)}P_C(I - \lambda B)x_n) \\ &\quad - (\mu_1 Tx_n + \mu_2 T_r(I - rA)x_n + \mu_3 P_C(I - \lambda B)x_n)\| \\ &\leq |\mu_{(1,n)}\delta_n - \mu_1\delta| \|x_n\| + |\mu_{(1,n)}(1 - \delta_n) - \mu_1(1 - \delta)| \|Sx_n\| \\ &\quad + \|T_r(I - rA)x_n\| |\mu_{(2,n)} - \mu_2| + |\mu_{(3,n)} - \mu_3| \|P_C(I - \lambda B)x_n\|. \end{aligned} \quad (2.26)$$

It follows from the condition (R4) that $e_n - Qx_n \rightarrow 0$. Note that

$$Qx_n - x_n = Qx_n - e_n + e_n - x_n. \quad (2.27)$$

From (2.22), we arrive at

$$\lim_{n \rightarrow \infty} \|Qx_n - x_n\| = 0. \quad (2.28)$$

It follows from Lemma 1.5 that

$$\xi \in F(Q) = \Omega. \quad (2.29)$$

Thanks to (2.23), we arrive at

$$\limsup_{n \rightarrow \infty} \langle u - \bar{x}, x_n - \bar{x} \rangle = \langle u - \bar{x}, \xi - \bar{x} \rangle \leq 0. \quad (2.30)$$

Step 6. Show that $x_n \rightarrow \bar{x}$ as $n \rightarrow \infty$.

Notice that

$$\begin{aligned} \|x_{n+1} - \bar{x}\|^2 &= \|\alpha_n u + \beta_n x_n + \gamma_n e_n - \bar{x}\|^2 \\ &= \langle \alpha_n(u - \bar{x}) + \beta_n(x_n - \bar{x}) + \gamma_n(e_n - \bar{x}), x_{n+1} - \bar{x} \rangle \\ &= \alpha_n \langle u - \bar{x}, x_{n+1} - \bar{x} \rangle + \beta_n \langle x_n - \bar{x}, x_{n+1} - \bar{x} \rangle + \gamma_n \langle e_n - \bar{x}, x_{n+1} - \bar{x} \rangle \\ &\leq \alpha_n \langle u - \bar{x}, x_{n+1} - \bar{x} \rangle + \beta_n \|x_n - \bar{x}\| \|x_{n+1} - \bar{x}\| + \gamma_n \|e_n - \bar{x}\| \|x_{n+1} - \bar{x}\| \\ &\leq \alpha_n \langle u - \bar{x}, x_{n+1} - \bar{x} \rangle + \beta_n \|x_n - \bar{x}\| \|x_{n+1} - \bar{x}\| + \gamma_n \|x_n - \bar{x}\| \|x_{n+1} - \bar{x}\| \\ &= (1 - \alpha_n) \|x_n - \bar{x}\| \|x_{n+1} - \bar{x}\| + \alpha_n \langle u - \bar{x}, x_{n+1} - \bar{x} \rangle \\ &\leq \frac{(1 - \alpha_n)}{2} (\|x_n - \bar{x}\|^2 + \|x_{n+1} - \bar{x}\|^2) + \alpha_n \langle u - \bar{x}, x_{n+1} - \bar{x} \rangle, \end{aligned} \quad (2.31)$$

which yields that

$$\|x_{n+1} - \bar{x}\|^2 \leq (1 - \alpha_n) \|x_n - \bar{x}\|^2 + 2\alpha_n \langle u - \bar{x}, x_{n+1} - \bar{x} \rangle. \quad (2.32)$$

In view of the restrictions (R2) and (2.30), we from Lemma 1.7 can conclude the desired conclusion easily. This completes the proof. \square

As corollaries of Theorem 2.1, we have the following results.

Corollary 2.2. Let C be a nonempty closed and convex subset of a real Hilbert space H and F a bifunction from $C \times C$ to \mathbb{R} satisfying (A1)–(A4). Let B be a β -inverse-strongly monotone mapping of C into H . Let $S : C \rightarrow C$ be a k -strict pseudocontraction with a fixed point. Assume that $\Omega := \text{EP}(F) \cap F(S) \cap \text{VI}(C, B) \neq \emptyset$. Let $\{x_n\}$ be a sequence in C generated by

$$\begin{aligned} x_1 &\in C, \quad \text{chosen arbitrarily,} \\ F(u_n, y) + \frac{1}{r} \langle y - u_n, u_n - x_n \rangle &\geq 0, \quad \forall y \in C, \\ v_n &= P_C(x_n - \lambda Bx_n), \\ y_n &= \delta_n x_n + (1 - \delta_n) Sx_n, \\ x_{n+1} &= \alpha_n u + \beta_n x_n + \gamma_n (\mu_{(1,n)} y_n + \mu_{(2,n)} u_n + \mu_{(3,n)} v_n), \quad \forall n \geq 1, \end{aligned} \tag{2.33}$$

where u is a fixed element in C , $\{\alpha_n\}$, $\{\beta_n\}$, $\{\gamma_n\}$, $\{\mu_{(1,n)}\}$, $\{\mu_{(2,n)}\}$, and $\{\mu_{(3,n)}\}$ are sequences in $(0, 1)$, $\{\delta_n\}$ is sequence in $[k, 1)$, $r \in (0, \infty)$, and $\lambda \in (0, 2\beta]$. Assume that the above control sequences satisfy the following restrictions:

- (R1) $\alpha_n + \beta_n + \gamma_n = \mu_{(1,n)} + \mu_{(2,n)} + \mu_{(3,n)} = 1$, for all $n \geq 1$;
- (R2) $\lim_{n \rightarrow \infty} \alpha_n = 0$, $\sum_{n=1}^{\infty} \alpha_n = \infty$;
- (R3) $0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} \beta_n < 1$;
- (R4) $\lim_{n \rightarrow \infty} \delta_n = \delta \in [k, 1)$ and $\lim_{n \rightarrow \infty} \mu_{(i,n)} = \mu_i \in (0, 1)$ for all $1 \leq i \leq 3$.

Then the sequence $\{x_n\}$ defined by the iterative process (2.1) converges strongly to $\bar{x} = P_{\Omega} u$.

Proof. In Theorem 2.1, put $A = 0$, the zero mapping. Then for any $\alpha > 0$, we see that the the following inequality holds.

$$\langle x - y, Ax - Ay \rangle \geq \alpha \|Ax - Ay\|^2. \tag{2.34}$$

Then, we can obtain the desired conclusion easily from Theorem 2.1. This completes the proof. \square

Corollary 2.3. Let C be a nonempty closed and convex subset of a real Hilbert space H and F a bifunction from $C \times C$ to \mathbb{R} satisfying (A1)–(A4). Let S_A be a k_α -strict pseudocontraction of C into H and K_B a k_β -strict pseudocontraction of C into H . Let $S : C \rightarrow C$ be a k -strict pseudocontraction with a fixed point. Assume that $\Omega := \text{EP}(F, I - S_A) \cap F(S) \cap F(S_B) \neq \emptyset$. Let $\{x_n\}$ be a sequence in C generated by

$$\begin{aligned} x_1 &\in C, \quad \text{chosen arbitrarily,} \\ F(u_n, y) + \langle x_n - S_A x_n, y - u_n \rangle + \frac{1}{r} \langle y - u_n, u_n - x_n \rangle &\geq 0, \quad \forall y \in C, \\ v_n &= (1 - \lambda)x_n + \lambda S_B x_n, \\ y_n &= \delta_n x_n + (1 - \delta_n) Sx_n, \\ x_{n+1} &= \alpha_n u + \beta_n x_n + \gamma_n (\mu_{(1,n)} y_n + \mu_{(2,n)} u_n + \mu_{(3,n)} v_n), \quad \forall n \geq 1, \end{aligned} \tag{2.35}$$

where u is a fixed element in C , $\{\alpha_n\}$, $\{\beta_n\}$, $\{\gamma_n\}$, $\{\mu_{(1,n)}\}$, $\{\mu_{(2,n)}\}$, and $\{\mu_{(3,n)}\}$ are sequences in $(0, 1)$, $\{\delta_n\}$ is sequence in $[k, 1)$, $r \in (0, 1 - k_\alpha]$, and $\lambda \in (0, 1 - k_\beta]$. Assume that the above control sequences satisfy the following restrictions:

- (R1) $\alpha_n + \beta_n + \gamma_n = \mu_{(1,n)} + \mu_{(2,n)} + \mu_{(3,n)} = 1$, for all $n \geq 1$;
- (R2) $\lim_{n \rightarrow \infty} \alpha_n = 0$, $\sum_{n=1}^{\infty} \alpha_n = \infty$;
- (R3) $0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} \beta_n < 1$;
- (R4) $\lim_{n \rightarrow \infty} \delta_n = \delta \in [k, 1)$ and $\lim_{n \rightarrow \infty} \mu_{(i,n)} = \mu_i \in (0, 1)$ for all $1 \leq i \leq 3$.

Then the sequence $\{x_n\}$ defined by the above iterative process converges strongly to $\bar{x} = P_\Omega u$.

Proof. Put $A = I - S_A$ and $B = I - S_B$. Then, we see that A is $((1 - k_\alpha)/2)$ -inverse-strongly monotone and B is $((1 - k_\beta)/2)$ -inverse-strongly monotone; see [7]. We have $F(S_B) = VI(C, B)$ and

$$P_C(x_n - \lambda_n Bx_n) = (1 - \lambda_n)x_n + \lambda_n S_B x_n. \quad (2.36)$$

It is easy to obtain the desired conclusion from Theorem 2.1. □

Remark 2.4. If $f : C \rightarrow C$ is a contractive mapping and we replace u by $f(x_n)$ in the recursion formula (2.1), we can obtain the so-called viscosity iteration method. We note that all theorems and corollaries of this paper carry over trivially to the so-called viscosity iteration method; see [28] for more details.

Acknowledgment

This project is supported by the National Natural Science Foundation of China (no. 10901140).

References

- [1] S. Chang, H. W. J. Lee, and C. K. Chan, "A new method for solving equilibrium problem fixed point problem and variational inequality problem with application to optimization," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 70, no. 9, pp. 3307–3319, 2009.
- [2] H. Iiduka and W. Takahashi, "Strong convergence theorems for nonexpansive mappings and inverse-strongly monotone mappings," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 61, no. 3, pp. 341–350, 2005.
- [3] X. Qin, Y. J. Cho, and S. M. Kang, "Viscosity approximation methods for generalized equilibrium problems and fixed point problems with applications," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 72, no. 1, pp. 99–112, 2010.
- [4] W. Takahashi and M. Toyoda, "Weak convergence theorems for nonexpansive mappings and monotone mappings," *Journal of Optimization Theory and Applications*, vol. 118, no. 2, pp. 417–428, 2003.
- [5] Y. Yao and J.-C. Yao, "On modified iterative method for nonexpansive mappings and monotone mappings," *Applied Mathematics and Computation*, vol. 186, no. 2, pp. 1551–1558, 2007.
- [6] F. E. Browder, "Convergence of approximants to fixed points of nonexpansive non-linear mappings in Banach spaces," *Archive for Rational Mechanics and Analysis*, vol. 24, pp. 82–90, 1967.
- [7] F. E. Browder and W. V. Petryshyn, "Construction of fixed points of nonlinear mappings in Hilbert space," *Journal of Mathematical Analysis and Applications*, vol. 20, pp. 197–228, 1967.
- [8] R. E. Bruck, Jr., "Properties of fixed-point sets of nonexpansive mappings in Banach spaces," *Transactions of the American Mathematical Society*, vol. 179, pp. 251–262, 1973.

- [9] L.-C. Ceng, S. Al-Homidan, Q. H. Ansari, and J.-C. Yao, "An iterative scheme for equilibrium problems and fixed point problems of strict pseudo-contraction mappings," *Journal of Computational and Applied Mathematics*, vol. 223, no. 2, pp. 967–974, 2009.
- [10] L.-C. Ceng and J.-C. Yao, "Hybrid viscosity approximation schemes for equilibrium problems and fixed point problems of infinitely many nonexpansive mappings," *Applied Mathematics and Computation*, vol. 198, no. 2, pp. 729–741, 2008.
- [11] P. L. Combettes and S. A. Hirstoaga, "Equilibrium programming in Hilbert spaces," *Journal of Nonlinear and Convex Analysis*, vol. 6, no. 1, pp. 117–136, 2005.
- [12] V. Colao, G. Marino, and H.-K. Xu, "An iterative method for finding common solutions of equilibrium and fixed point problems," *Journal of Mathematical Analysis and Applications*, vol. 344, no. 1, pp. 340–352, 2008.
- [13] C. Jaiboon, P. Kumam, and U. W. Humphries, "Weak convergence theorem by an extragradient method for variational inequality, equilibrium and fixed point problems," *Bulletin of the Malaysian Mathematical Sciences Society*, vol. 32, no. 2, pp. 173–185, 2009.
- [14] G. Marino and H.-K. Xu, "Weak and strong convergence theorems for strict pseudo-contractions in Hilbert spaces," *Journal of Mathematical Analysis and Applications*, vol. 329, no. 1, pp. 336–346, 2007.
- [15] A. Moudafi and M. Théra, "Proximal and dynamical approaches to equilibrium problems," in *Ill-Posed Variational Problems and Regularization Techniques (Trier, 1998)*, vol. 477 of *Lecture Notes in Economics and Mathematical Systems*, pp. 187–201, Springer, Berlin, Germany, 1999.
- [16] S. Plubtieng and R. Punpaeng, "A new iterative method for equilibrium problems and fixed point problems of nonexpansive mappings and monotone mappings," *Applied Mathematics and Computation*, vol. 197, no. 2, pp. 548–558, 2008.
- [17] X. Qin, Y. J. Cho, and S. M. Kang, "Convergence theorems of common elements for equilibrium problems and fixed point problems in Banach spaces," *Journal of Computational and Applied Mathematics*, vol. 225, no. 1, pp. 20–30, 2009.
- [18] X. Qin, M. Shang, and Y. Su, "Strong convergence of a general iterative algorithm for equilibrium problems and variational inequality problems," *Mathematical and Computer Modelling*, vol. 48, no. 7-8, pp. 1033–1046, 2008.
- [19] X. Qin and Y. Su, "Strong convergence theorems for relatively nonexpansive mappings in a Banach space," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 67, no. 6, pp. 1958–1965, 2007.
- [20] A. Tada and W. Takahashi, "Weak and strong convergence theorems for a nonexpansive mapping and an equilibrium problem," *Journal of Optimization Theory and Applications*, vol. 133, no. 3, pp. 359–370, 2007.
- [21] S. Takahashi and W. Takahashi, "Viscosity approximation methods for equilibrium problems and fixed point problems in Hilbert spaces," *Journal of Mathematical Analysis and Applications*, vol. 331, no. 1, pp. 506–515, 2007.
- [22] S. Takahashi and W. Takahashi, "Strong convergence theorem for a generalized equilibrium problem and a nonexpansive mapping in a Hilbert space," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 69, no. 3, pp. 1025–1033, 2008.
- [23] W. Takahashi and K. Zembayashi, "Strong and weak convergence theorems for equilibrium problems and relatively nonexpansive mappings in Banach spaces," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 70, no. 1, pp. 45–57, 2009.
- [24] E. Blum and W. Oettli, "From optimization and variational inequalities to equilibrium problems," *The Mathematics Student*, vol. 63, no. 1–4, pp. 123–145, 1994.
- [25] H. Zhou, "Convergence theorems of fixed points for κ -strict pseudo-contractions in Hilbert spaces," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 69, no. 2, pp. 456–462, 2008.
- [26] T. Suzuki, "Strong convergence of Krasnoselskii and Mann's type sequences for one-parameter non-expansive semigroups without Bochner integrals," *Journal of Mathematical Analysis and Applications*, vol. 305, no. 1, pp. 227–239, 2005.
- [27] H.-K. Xu, "Iterative algorithms for nonlinear operators," *Journal of the London Mathematical Society*, vol. 66, no. 1, pp. 240–256, 2002.
- [28] T. Suzuki, "Moudafi's viscosity approximations with Meir-Keeler contractions," *Journal of Mathematical Analysis and Applications*, vol. 325, no. 1, pp. 342–352, 2007.