

Research Article

A Study on the p -Adic q -Integral Representation on \mathbb{Z}_p Associated with the Weighted q -Bernstein and q -Bernoulli Polynomials

T. Kim,¹ A. Bayad,² and Y.-H. Kim¹

¹ Division of General Education-Mathematics, Kwangwoon University, Seoul 139-701, Republic of Korea

² Département de Mathématiques, Université d'Evry Val d'Essonne, Boulevard François Mitterrand, 91025 Evry Cedex, France

Correspondence should be addressed to A. Bayad, abayad@maths.univ-evry.fr

Received 6 December 2010; Accepted 15 January 2011

Academic Editor: Vijay Gupta

Copyright © 2011 T. Kim et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We investigate some interesting properties of the weighted q -Bernstein polynomials related to the weighted q -Bernoulli numbers and polynomials by using p -adic q -integral on \mathbb{Z}_p .

1. Introduction and Preliminaries

Let p be a fixed prime number. Throughout this paper, \mathbb{Z}_p , \mathbb{Q}_p , and \mathbb{C}_p will denote the ring of p -adic integers, the field of p -adic rational numbers, and the completion of the algebraic closure of \mathbb{Q}_p , respectively. Let \mathbb{N} be the set of natural numbers, and let $\mathbb{Z}_+ = \mathbb{N} \cup \{0\}$. Let ν_p be the normalized exponential valuation of \mathbb{C}_p with $|p|_p = p^{-\nu_p(p)} = 1/p$. Let q be regarded as either a complex number $q \in \mathbb{C}$ or a p -adic number $q \in \mathbb{C}_p$. If $q \in \mathbb{C}$, then we always assume $|q| < 1$. If $q \in \mathbb{C}_p$, we assume that $|1 - q|_p < 1$. In this paper, we define the q -number as $[x]_q = (1 - q^x)/(1 - q)$ (see [1–13]).

Let $C[0, 1]$ be the set of continuous functions on $[0, 1]$. For $\alpha \in \mathbb{N}$ and $n, k \in \mathbb{Z}_+$, the weighted q -Bernstein operator of order n for $f \in C[0, 1]$ is defined by

$$\mathbb{B}_{n,q}^{(\alpha)}(f | x) = \sum_{k=0}^n f\left(\frac{k}{n}\right) \binom{n}{k} [x]_{q^\alpha}^k [1-x]_{q^\alpha}^{n-k} = \sum_{k=0}^n f\left(\frac{k}{n}\right) B_{k,n}^{(\alpha)}(x, q). \quad (1.1)$$

Here $B_{k,n}^{(\alpha)}(x, q)$ is called the weighted q -Bernstein polynomials of degree n (see [2, 5, 6]).

Let $UD(\mathbb{Z}_p)$ be the space of uniformly differentiable functions on \mathbb{Z}_p . For $f \in UD(\mathbb{Z}_p)$, the p -adic q -integral on \mathbb{Z}_p , which is called the bosonic q -integral on \mathbb{Z}_p , is defined by

$$I_q(f) = \int_{\mathbb{Z}_p} f(x) d\mu_q(x) = \lim_{N \rightarrow \infty} \frac{1}{[p^N]_q} \sum_{x=0}^{p^N-1} f(x) q^x, \quad (1.2)$$

(see [10]).

The Carlitz's q -Bernoulli numbers are defined by

$$\beta_{0,q} = 1, \quad q(q\beta + 1)^k - \beta_{k,q} = \begin{cases} 1, & \text{if } k = 1, \\ 0, & \text{if } k > 1, \end{cases} \quad (1.3)$$

with the usual convention about replacing β^k by $\beta_{k,q}$ (see [3, 9, 10]). In [3], Carlitz also defined the expansion of Carlitz's q -Bernoulli numbers as follows:

$$\beta_{0,q}^h = \frac{h}{[h]_q}, \quad q^h(q\beta^h + 1)^n - \beta_{n,q}^h = \begin{cases} 1, & \text{if } n = 1, \\ 0, & \text{if } n > 1, \end{cases} \quad (1.4)$$

with the usual convention about replacing $(\beta^h)^n$ by $\beta_{n,q}^h$.

The weighted q -Bernoulli numbers are constructed in previous paper [6] as follows: for $\alpha \in \mathbb{N}$,

$$\tilde{\beta}_{0,q}^{(\alpha)} = 1, \quad q(q^\alpha \tilde{\beta}^{(\alpha)} + 1)^n - \tilde{\beta}_{n,q}^{(\alpha)} = \begin{cases} \frac{\alpha}{[\alpha]_q}, & \text{if } n = 1, \\ 0, & \text{if } n > 1, \end{cases} \quad (1.5)$$

with the usual convention about replacing $(\tilde{\beta}^{(\alpha)})^n$ by $\tilde{\beta}_{n,q}^{(\alpha)}$. Let $f_n(x) = f(x+n)$. By the definition (1.2) of p -adic q -integral on \mathbb{Z}_p , we easily get

$$\begin{aligned} qI_q(f_1) &= q \lim_{N \rightarrow \infty} \frac{1}{[p^N]_q} \sum_{x=0}^{p^N-1} f(x+1) q^x, \\ &= \lim_{N \rightarrow \infty} \frac{1}{[p^N]_q} \sum_{x=0}^{p^N-1} f(x) q^x + \lim_{N \rightarrow \infty} \frac{f(p^N) q^{p^N} - f(0)}{[p^N]_q} \\ &= \int_{\mathbb{Z}_p} f(x) d\mu_q(x) + (q-1)f(0) + \frac{q-1}{\log q} f'(0), \end{aligned} \quad (1.6)$$

Continuing this process, we obtain easily the relation

$$q^n \int_{\mathbb{Z}_p} f_n(x) d\mu_q(x) - \int_{\mathbb{Z}_p} f(x) d\mu_q(x) = (q-1) \sum_{l=0}^{n-1} q^l f(l) + \frac{q-1}{\log q} \sum_{l=0}^{n-1} q^l f'(l), \quad (1.7)$$

where $n \in \mathbb{N}$ and $f'(l) = df(l)/dx$ (see [6]).

Then by (1.2), applying to the function $x \rightarrow [x]_{q^\alpha}^n$, we can see that

$$\tilde{\beta}_{n,q}^{(\alpha)} = \int_{\mathbb{Z}_p} [x]_{q^\alpha}^n d\mu_q(x) = -\frac{n\alpha}{[\alpha]_q} \sum_{m=0}^\infty q^{m\alpha+m} [m]_{q^\alpha}^{n-1} + (1-q) \sum_{m=0}^\infty q^m [m]_{q^\alpha}^n. \tag{1.8}$$

The weighted q -Bernoulli polynomials are also defined by the generating function as follows:

$$F_q^{(\alpha)}(t, x) = -t \frac{\alpha}{[\alpha]_q} \sum_{m=0}^\infty q^{m\alpha+m} e^{[m+x]_{q^\alpha} t} + (1-q) \sum_{m=0}^\infty q^m e^{[m+x]_{q^\alpha} t} = \sum_{n=0}^\infty \tilde{\beta}_{n,q}^{(\alpha)}(x) \frac{t^n}{n!}, \tag{1.9}$$

(see[6]). Thus, we note that

$$\begin{aligned} \tilde{\beta}_{n,q}^{(\alpha)}(x) &= \sum_{l=0}^n \binom{n}{l} [x]_{q^\alpha}^{n-l} q^{\alpha l x} \tilde{\beta}_{l,q}^{(\alpha)} \\ &= -\frac{n\alpha}{[\alpha]_q} \sum_{m=0}^\infty q^{m\alpha+m} [m+x]_{q^\alpha}^{n-1} + (1-q) \sum_{m=0}^\infty q^m [m+x]_{q^\alpha}^n. \end{aligned} \tag{1.10}$$

From (1.2) and the previous equalities, we obtain the Witt’s formula for the weighted q -Bernoulli polynomials as follows:

$$\tilde{\beta}_{n,q}^{(\alpha)}(x) = \int_{\mathbb{Z}_p} [x+y]_{q^\alpha}^n d\mu_q(y) = \sum_{l=0}^n \binom{n}{l} q^{\alpha l x} [x]_{q^\alpha}^{n-l} \int_{\mathbb{Z}_p} [y]_{q^\alpha}^l d\mu_q(y). \tag{1.11}$$

By using (1.2) and the weighted q -Bernoulli polynomials, we easily get

$$q^n \tilde{\beta}_{m,q}^{(\alpha)}(n) - \tilde{\beta}_{m,q}^{(\alpha)} = (q-1) \sum_{l=0}^{n-1} q^l [l]_{q^\alpha}^m + \frac{m\alpha}{[\alpha]_q} \sum_{l=0}^{n-1} q^{\alpha l + l} [l]_{q^\alpha}^{m-1}, \tag{1.12}$$

where $n, \alpha \in \mathbb{N}$ and $m \in \mathbb{Z}_+$ (see [6]).

In this paper, we consider the weighted q -Bernstein polynomials to express the bosonic q -integral on \mathbb{Z}_p and investigate some properties of the weighted q -Bernstein polynomials associated with the weighted q -Bernoulli polynomials by using the expression of p -adic q -integral on \mathbb{Z}_p of those polynomials.

2. Weighted q -Bernstein Polynomials and q -Bernoulli Polynomials

In this section, we assume that $\alpha \in \mathbb{N}$ and $q \in \mathbb{C}_p$ with $|1-q|_p < 1$.

Now we consider the p -adic weighted q -Bernstein operator as follows:

$$\mathbb{B}_{n,q}^{(\alpha)}(f \mid x)(fx) = \sum_{k=0}^n f\left(\frac{k}{n}\right) \binom{n}{k} [x]_{q^\alpha}^k [1-x]_{q^\alpha}^{n-k} = \sum_{k=0}^n f\left(\frac{k}{n}\right) B_{k,n}^{(\alpha)}(x, q). \tag{2.1}$$

The p -adic q -Bernstein polynomials with weight α of degree n are given by

$$B_{k,n}^{(\alpha)}(x, q) = \binom{n}{k} [x]_{q^\alpha}^k [1-x]_{q^\alpha}^{n-k}, \quad (2.2)$$

where $x \in \mathbb{Z}_p$, $\alpha \in \mathbb{N}$, and $n, k \in \mathbb{Z}_+$ (see [6, 7]). Note that $B_{k,n}^{(\alpha)}(x, q) = B_{n-k,n}^{(\alpha)}(1-x, 1/q)$. That is, the weighted q -Bernstein polynomials are symmetric.

From the definition of the weighted q -Bernoulli polynomials, we have

$$\tilde{\beta}_{n,q^{-1}}^{(\alpha)}(1-x) = (-1)^n q^{\alpha n} \tilde{\beta}_{n,q}^{(\alpha)}(x). \quad (2.3)$$

By the definition of p -adic q -integral on \mathbb{Z}_p , we get

$$\begin{aligned} \int_{\mathbb{Z}_p} [1-x]_{q^\alpha}^n d\mu_q(x) &= q^{\alpha n} (-1)^n \int_{\mathbb{Z}_p} [-1+x]_{q^\alpha}^n d\mu_q(x) \\ &= \int_{\mathbb{Z}_p} (1-[x]_{q^\alpha})^n d\mu_q(x). \end{aligned} \quad (2.4)$$

From (2.3) and (2.4), we have

$$\int_{\mathbb{Z}_p} [1-x]_{q^\alpha}^n d\mu_q(x) = \sum_{l=0}^n \binom{n}{l} (-1)^l \tilde{\beta}_{l,q}^{(\alpha)} = q^{\alpha n} (-1)^n \tilde{\beta}_{n,q}^{(\alpha)} (-1) = \tilde{\beta}_{n,q}^{(\alpha)}(2). \quad (2.5)$$

Therefore, we obtain the following lemma.

Lemma 2.1. *For $n \in \mathbb{Z}_+$, one has*

$$\begin{aligned} \int_{\mathbb{Z}_p} [1-x]_{q^\alpha}^n d\mu_q(x) &= \sum_{l=0}^n \binom{n}{l} (-1)^l \tilde{\beta}_{l,q}^{(\alpha)} = q^{\alpha n} (-1)^n \tilde{\beta}_{n,q}^{(\alpha)} (-1) = \tilde{\beta}_{n,q}^{(\alpha)}(2), \\ \tilde{\beta}_{n,q^{-1}}^{(\alpha)}(1-x) &= (-1)^n q^{\alpha n} \tilde{\beta}_{n,q}^{(\alpha)}(x). \end{aligned} \quad (2.6)$$

By (2.2), (2.3), and (2.4), we get

$$q^2 \tilde{\beta}_{n,q}^{(\alpha)}(2) = n \frac{\alpha}{[\alpha]_q} q^{1+\alpha} + q^2 - q + \tilde{\beta}_{n,q}^{(\alpha)}, \quad \text{if } n > 1. \quad (2.7)$$

Thus, we have

$$\tilde{\beta}_{n,q}^{(\alpha)}(2) = \frac{1}{q^2} \tilde{\beta}_{n,q}^{(\alpha)} + \frac{n\alpha}{[\alpha]_q} q^{\alpha-1} + 1 - \frac{1}{q}, \quad \text{if } n > 1. \quad (2.8)$$

Therefore, by (2.8), we obtain the following proposition.

Proposition 2.2. For $n \in \mathbb{N}$ with $n > 1$, one has

$$\tilde{\beta}_{n,q}^{(\alpha)}(2) = \frac{1}{q^2} \tilde{\beta}_{n,q}^{(\alpha)} + \frac{n\alpha}{[\alpha]_q} q^{\alpha-1} + 1 - \frac{1}{q}. \tag{2.9}$$

By using Proposition 2.2 and Lemma 2.1, we obtain the following corollary.

Corollary 2.3. For $n \in \mathbb{N}$ with $n > 1$, one has

$$\int_{\mathbb{Z}_p} [1-x]_{q^{-\alpha}}^n d\mu_q(x) = q^2 \tilde{\beta}_{n,q^{-1}}^{(\alpha)} + \frac{n\alpha}{[\alpha]_q} + 1 - q, \tag{2.10}$$

$$\int_{\mathbb{Z}_p} [1-x]_{q^{-\alpha}}^n d\mu_q(x) = \frac{n\alpha}{[\alpha]_q} + 1 - q + q^2 \int_{\mathbb{Z}_p} [x]_{q^{-\alpha}}^n d\mu_{q^{-1}}(x) = \int_{\mathbb{Z}_p} (1 - [x]_{q^\alpha})^n d\mu_q(x). \tag{2.11}$$

Taking the bosonic q -integral on \mathbb{Z}_p for one weighted q -Bernstein polynomials in (2.1), we have

$$\begin{aligned} \int_{\mathbb{Z}_p} B_{k,n}^{(\alpha)}(x, q) d\mu_q(x) &= \binom{n}{k} \int_{\mathbb{Z}_p} [x]_{q^\alpha}^k [1-x]_{q^{-\alpha}}^{n-k} d\mu_q(x) \\ &= \binom{n}{k} \sum_{l=0}^{n-k} \binom{n-k}{l} (-1)^l \int_{\mathbb{Z}_p} [x]_{q^\alpha}^{k+l} d\mu_q(x) \\ &= \binom{n}{k} \sum_{l=0}^{n-k} \binom{n-k}{l} (-1)^l \tilde{\beta}_{k+l,q}^{(\alpha)}. \end{aligned} \tag{2.12}$$

By the symmetry of q -Bernstein polynomials, we get

$$\begin{aligned} \int_{\mathbb{Z}_p} B_{k,n}^{(\alpha)}(x, q) d\mu_q(x) &= \int_{\mathbb{Z}_p} B_{n-k,n}^{(\alpha)}\left(1-x, \frac{1}{q}\right) d\mu_q(x) \\ &= \binom{n}{k} \sum_{l=0}^k \binom{k}{l} (-1)^{k+l} \int_{\mathbb{Z}_p} [1-x]_{q^{-\alpha}}^{n-l} d\mu_q(x). \end{aligned} \tag{2.13}$$

For $n > k + 1$, by (2.11) and (2.13), we have

$$\begin{aligned} \int_{\mathbb{Z}_p} B_{k,n}^{(\alpha)}(x, q) d\mu_q(x) &= \binom{n}{k} \sum_{l=0}^k \binom{k}{l} (-1)^{k+l} \left(\frac{n\alpha}{[\alpha]_q} + 1 - q + q^2 \int_{\mathbb{Z}_p} [x]_{q^{-\alpha}}^{n-l} d\mu_{q^{-1}}(x) \right) \\ &= \begin{cases} \frac{n\alpha}{[\alpha]_q} + 1 - q + q^2 \tilde{\beta}_{n,q^{-1}}^{(\alpha)}, & \text{if } k = 0, \\ \binom{n}{k} q^2 \sum_{l=0}^k \binom{k}{l} (-1)^{k+l} \tilde{\beta}_{n-l,q^{-1}}^{(\alpha)}, & \text{if } k > 0. \end{cases} \end{aligned} \quad (2.14)$$

By comparing the coefficients on the both sides of (2.12) and (2.14), we obtain the following theorem.

Theorem 2.4. For $n, k \in \mathbb{Z}_+$ with $n > k + 1$, one has

$$\sum_{l=0}^{n-k} \binom{n-k}{l} (-1)^l \tilde{\beta}_{k+l,q}^{(\alpha)} = q^2 \sum_{l=0}^k \binom{k}{l} (-1)^{k+l} \tilde{\beta}_{n-l,q^{-1}}^{(\alpha)}, \quad \text{if } k \neq 0. \quad (2.15)$$

In particular, when $k = 0$, one has

$$\frac{n\alpha}{[\alpha]_q} + 1 - q + q^2 \tilde{\beta}_{n,q^{-1}}^{(\alpha)} = \sum_{l=0}^n \binom{n}{l} (-1)^l \tilde{\beta}_{l,q}^{(\alpha)}. \quad (2.16)$$

Let $m, n, k \in \mathbb{Z}_+$ with $m + n > 2k + 1$. Then we see that

$$\begin{aligned} &\int_{\mathbb{Z}_p} B_{k,n}^{(\alpha)}(x, q) B_{k,m}^{(\alpha)}(x, q) d\mu_q(x) \\ &= \binom{n}{k} \binom{m}{k} \int_{\mathbb{Z}_p} [x]_{q^{-\alpha}}^{2k} [1-x]_{q^{-\alpha}}^{n+m-2k} d\mu_q(x) \\ &= \binom{n}{k} \binom{m}{k} \sum_{l=0}^{2k} \binom{2k}{l} (-1)^{l+2k} \int_{\mathbb{Z}_p} [1-x]_{q^{-\alpha}}^{n+m-l} d\mu_q(x). \quad (2.17) \\ &= \binom{n}{k} \binom{m}{k} \sum_{l=0}^{2k} \binom{2k}{l} (-1)^{l+2k} \left(\frac{n\alpha}{[\alpha]_q} + 1 - q + q^2 \int_{\mathbb{Z}_p} [x]_{q^{-\alpha}}^{n+m-l} d\mu_{q^{-1}}(x) \right) \\ &= \binom{n}{k} \binom{m}{k} \sum_{l=0}^{2k} \binom{2k}{l} (-1)^{l+2k} \left(\frac{n\alpha}{[\alpha]_q} + 1 - q + q^2 \tilde{\beta}_{n+m-l,q^{-1}}^{(\alpha)} \right). \end{aligned}$$

Therefore, by (2.17), we obtain the following theorem.

Theorem 2.5. For $m, n, k \in \mathbb{Z}_+$ with $m + n > 2k + 1$, one has

$$\int_{\mathbb{Z}_p} B_{k,n}^{(\alpha)}(x, q) B_{k,m}^{(\alpha)}(x, q) d\mu_q(x) = \begin{cases} \frac{n\alpha}{[\alpha]_q} + 1 - q + q^2 \tilde{\beta}_{n+m, q^{-1}}^{(\alpha)}, & \text{if } k = 0, \\ \binom{n}{k} \binom{m}{k} q^2 \sum_{l=0}^{2k} \binom{2k}{l} (-1)^{l+2k} \tilde{\beta}_{n+m-l, q^{-1}}^{(\alpha)}, & \text{if } k \neq 0. \end{cases} \tag{2.18}$$

For $m, n, k \in \mathbb{Z}_+$, we have

$$\begin{aligned} & \int_{\mathbb{Z}_p} B_{k,n}^{(\alpha)}(x, q) B_{k,m}^{(\alpha)}(x, q) d\mu_q(x) \\ &= \binom{n}{k} \binom{m}{k} \int_{\mathbb{Z}_p} [x]_{q^\alpha}^{2k} [1-x]_{q^{-\alpha}}^{n+m-2k} d\mu_q(x) \\ &= \binom{n}{k} \binom{m}{k} \sum_{l=0}^{n+m-2k} \binom{n+m-2k}{l} (-1)^l \int_{\mathbb{Z}_p} [x]_{q^\alpha}^{2k+l} d\mu_q(x) \\ &= \binom{n}{k} \binom{m}{k} \sum_{l=0}^{n+m-2k} \binom{n+m-2k}{l} (-1)^l \tilde{\beta}_{l+2k, q}^{(\alpha)}. \end{aligned} \tag{2.19}$$

Therefore, by (2.18) and (2.19), we obtain the following theorem.

Theorem 2.6. For $m, n, k \in \mathbb{Z}_+$ with $m + n > 2k + 1$, one has

$$\frac{n\alpha}{[\alpha]_q} + 1 - q + q^2 \tilde{\beta}_{n+m-l, q^{-1}}^{(\alpha)} = \sum_{l=0}^{n+m} \binom{n+m}{l} (-1)^l \tilde{\beta}_{l, q}^{(\alpha)}. \tag{2.20}$$

Furthermore, for $k \neq 0$, one has

$$\sum_{l=0}^{n+m-2k} \binom{n+m-2k}{l} (-1)^l \tilde{\beta}_{l+2k, q}^{(\alpha)} = q^2 \sum_{l=0}^{2k} \binom{2k}{l} (-1)^{l+2k} \tilde{\beta}_{n+m-l, q^{-1}}^{(\alpha)}. \tag{2.21}$$

By the induction hypothesis, we obtain the following theorem.

Theorem 2.7. For $s \in \mathbb{N}$ and $k, n_1, \dots, n_s \in \mathbb{Z}_+$ with $n_1 + n_2 + \dots + n_s > sk + 1$, one has

$$\int_{\mathbb{Z}_p} \left(\prod_{i=1}^s B_{k, n_i}^{(\alpha)}(x, q) \right) d\mu_q(x) = \begin{cases} \frac{n\alpha}{[\alpha]_q} + 1 - q + q^2 \tilde{\beta}_{n_1+\dots+n_s, q^{-1}}^{(\alpha)}, & \text{if } k = 0, \\ \left(\prod_{i=1}^s \binom{n_i}{k} \right) \sum_{l=0}^{sk} \binom{sk}{l} (-1)^{l+sk} \tilde{\beta}_{n_1+\dots+n_s-l, q^{-1}}^{(\alpha)}, & \text{if } k \neq 0. \end{cases} \tag{2.22}$$

For $s \in \mathbb{N}$, let $k, n_1, \dots, n_s \in \mathbb{Z}_+$ with $n_1 + n_2 + \dots + n_s > sk + 1$. Then we show that

$$\int_{\mathbb{Z}_p} \left(\prod_{i=1}^s B_{k, n_i}^{(\alpha)}(x, q) \right) d\mu_q(x) = \left(\prod_{i=1}^s \binom{n_i}{k} \right) \sum_{l=0}^{n_1 + \dots + n_s - sk} \binom{n_1 + \dots + n_s - sk}{l} (-1)^l \tilde{\beta}_{l+sk, q}^{(\alpha)}. \quad (2.23)$$

Therefore, by Theorem 2.7 and (2.23), we obtain the following theorem.

Theorem 2.8. For $s \in \mathbb{N}$, let $k, n_1, \dots, n_s \in \mathbb{Z}_+$ with $n_1 + n_2 + \dots + n_s > sk + 1$. Then one sees that for $k = 0$

$$\sum_{l=0}^{n_1 + \dots + n_s} \binom{n_1 + \dots + n_s}{l} (-1)^l \tilde{\beta}_{l, q}^{(\alpha)} = \frac{n\alpha}{[\alpha]_q} + 1 - q + q^2 \tilde{\beta}_{n_1 + \dots + n_s, q^{-1}}^{(\alpha)}. \quad (2.24)$$

For $k \neq 0$, one has

$$\sum_{l=0}^{sk} \binom{sk}{l} (-1)^{l+sk} \tilde{\beta}_{n_1 + \dots + n_s - l, q^{-1}}^{(\alpha)} = \sum_{l=0}^{n_1 + \dots + n_s - sk} \binom{n_1 + \dots + n_s - sk}{l} (-1)^l \tilde{\beta}_{l+sk, q}^{(\alpha)}. \quad (2.25)$$

References

- [1] M. Acikgoz and Y. Simsek, "On multiple interpolation functions of the Nörlund-type q -Euler polynomials," *Abstract and Applied Analysis*, vol. 2009, Article ID 382574, 14 pages, 2009.
- [2] A. Bayad, J. Choi, T. Kim, Y.-H. Kim, and L. C. Jang, " q -extension of Bernstein polynomials with weighted $(\alpha; \beta)$," *Journal of Computational and Applied Mathematics*. In press.
- [3] L. Carlitz, "Expansions of q -Bernoulli numbers," *Duke Mathematical Journal*, vol. 25, pp. 355–364, 1958.
- [4] A. S. Hegazi and M. Mansour, "A note on q -Bernoulli numbers and polynomials," *Journal of Nonlinear Mathematical Physics*, vol. 13, no. 1, pp. 9–18, 2006.
- [5] L.-C. Jang, W.-J. Kim, and Y. Simsek, "A study on the p -adic integral representation on \mathbb{Z}_p associated with Bernstein and Bernoulli polynomials," *Advances in Difference Equations*, vol. 2010, Article ID 163217, 6 pages, 2010.
- [6] T. Kim, "On the weighted q -Bernoulli numbers and polynomials," <http://arxiv.org/abs/1011.5305>.
- [7] T. Kim, "A note on q -Bernstein polynomials," *Russian Journal of Mathematical Physics*, vol. 18, no. 1, 2011.
- [8] T. Kim, "Barnes-type multiple q -zeta functions and q -Euler polynomials," *Journal of Physics A*, vol. 43, no. 25, Article ID 255201, 11 pages, 2010.
- [9] T. Kim, " q -Bernoulli numbers and polynomials associated with Gaussian binomial coefficients," *Russian Journal of Mathematical Physics*, vol. 15, no. 1, pp. 51–57, 2008.
- [10] T. Kim, "On a q -analogue of the p -adic log gamma functions and related integrals," *Journal of Number Theory*, vol. 76, no. 2, pp. 320–329, 1999.
- [11] B. A. Kupershmidt, "Reflection symmetries of q -Bernoulli polynomials," *Journal of Nonlinear Mathematical Physics*, vol. 12, supplement 1, pp. 412–422, 2005.
- [12] H. Ozden, I. N. Cangul, and Y. Simsek, "Remarks on q -Bernoulli numbers associated with Daehee numbers," *Advanced Studies in Contemporary Mathematics*, vol. 18, no. 1, pp. 41–48, 2009.
- [13] S.-H. Rim, J.-H. Jin, E.-J. Moon, and S.-J. Lee, "On multiple interpolation functions of the q -Genocchi polynomials," *Journal of Inequalities and Applications*, vol. 2010, Article ID 351419, 13 pages, 2010.