

Research Article

Limit Distribution of Inventory Level of Perishable Inventory Model

Hailing Dong¹ and Guochao Jiang²

¹ *School of Mathematics and Computational Science, Shenzhen University, Nanhai Avenue 3688, Guangdong, Shenzhen 518060, China*

² *Department of Urban Planning and Economic Management, Harbin Institute of Technology Shenzhen Graduate School, Shenzhen 518055, China*

Correspondence should be addressed to Hailing Dong, hailingdong@gmail.com

Received 27 January 2011; Accepted 14 June 2011

Academic Editor: Paulo Batista Gonçalves

Copyright © 2011 H. Dong and G. Jiang. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

This paper studies a perishable inventory model, which assumes that each perishable item has finite lifetime, and only one item is consumed each time. The lifetimes of perishable items are independent random variables with the general distribution and so are the consumption internal. Under this assumption, by using backward equations and limit distribution of Markov skeleton processes, this paper obtains the existence conditions and the explicit expression of the limit distribution of the inventory level of perishable inventory model.

1. Introduction

Perishable goods are common in our daily life. In this paper, perishable goods refer to the items that have finite lifetime, like putrescible foods, easily-expired medicines, volatile liquids, and so on. Perishable inventory model can be widely used in blood banks, chemical and food industry.

In the past few decades, researchers have paid much attention to perishable inventory model. The inventory problem of perishable items was first studied by Whittin [1] who considered fashion goods perishing at the end of a prescribed storage period. Ghare and Schrader [2] proposed an inventory model, in which the rate of perishable is a constant, and the consumption internals have exponential distribution. Based on the inventory model proposed by Ghare and Schrader [2], series of studies are carried out (see Raafat [3], Goyal and Giri [4], and their references). Recently, Li et al. [5] considered some factors, like demand,

deteriorating rate, price discount, allow shortage or not, inflation, time value of money, and so on, as important factors in the perishable inventory study, then they divided current perishable items inventory study literatures into two categories from the perspective of study scope and reviewed the literature for each category. Karmakar and Choudhury [6] focused on the modeling of perishable items with shortages and reviewed the corresponding inventory models. Other representative works can be seen in [7–15].

An interesting and important study of perishable inventory model is about the inventory level process. Ravichandran [16] obtained the explicit expression of the stationary distribution of the inventory level in operating the (S, s) policy, with positive lead time and poisson demand. Chiu [17] developed the expected inventory level to determine a best (Q, r) ordering policy under a positive order lead time and fixed life perishability. Liu and Yang [18] analyzed an (s, S) continuous review model and obtains the matrix-geometric solutions for the steady-state probability distribution of the inventory level, with finite lifetimes and positive lead times. Sivakumar [19] obtained the joint probability distribution of the inventory level and the number of demands in the orbit, where the life time of each items is assumed to be exponential. Other related papers can be seen in [20–22] and so on.

In order to facilitate the mathematical treatment, most of these papers assume that the lifetime of item or consumption interval equals to constant or has exponential distribution, so that the inventory level of perishable inventory model can be reduced to a Markov process. However, in practice, the lifetime of item or consumption interval is not necessarily exponential, but a wide range of distribution. In this case, the inventory level of perishable inventory model is not and hardly been converted into a Markov process, which leads to a bottleneck on the mathematical treatment. To the best of our knowledge, no previous studies obtained the existence conditions of the limit distribution of the inventory level. Thus, we intend to work at it.

Markov skeleton process provides an effective solution to the problem. Markov skeleton processes which are proposed by Hou et al. [23] in 1997 are more extensive than Markov processes. Markov skeleton process has been in-depth studied (representative works, see [24–27]). This paper proves that the inventory level of perishable inventory model is a positive recurrent Doob skeleton process which is a special case of Markov skeleton processes. Hence, by applying backward equations and limit distribution of Markov skeleton processes, this paper obtains the existence conditions and the explicit expression of the limit distribution of the inventory level. Moreover, this paper obtains the probability of the inventory level greater than 0 and the probability of the inventory level less than or equal to 0, which can then be used for the evaluation of inventory system performance.

This paper is organized as follows. Section 2 introduces Markov skeleton processes and presents its backward equations and limit distribution. Section 3 introduces a perishable inventory model and applies Markov skeleton processes approach to study the limit distribution of inventory level process.

2. Markov Skeleton Processes

In this section, we introduce Markov skeleton processes and present its backward equations and limit distribution.

2.1. Definition of Markov Skeleton Processes

Definition 2.1 (see [26]). A stochastic process $X = \{X(t, \omega), 0 \leq t < \infty\}$ which takes values on a polish space (E, \mathcal{E}) is called a Markov skeleton process if there exists a sequence of optional stopping times $\{\tau_n\}_{n \geq 0}$, satisfying

- (i) $\tau_n \uparrow +\infty$ with $\tau_0 = 0$, and for each $n \geq 0$, $\tau_n < \infty$;
- (ii) for all $n = 0, 1, \dots$, $\tau_{n+1} = \tau_n + \theta_{\tau_n} \cdot \tau_1$;
- (iii) for every τ_n and any bounded $\mathcal{E}^{[0, \infty)}$ -measurable function f defined on $E^{[0, \infty)}$

$$E[f(X(\tau_n + \cdot)) | \mathcal{F}_{\tau_n}^X] = E[f(X(\tau_n + \cdot)) | X(\tau_n)] \quad \text{P-a.s.}, \quad (2.1)$$

where $\Omega_{\tau_n} = (\omega : \tau_n(\omega) < \infty)$, and $\mathcal{F}_{\tau_n}^X = \{A : \forall t \geq 0, A \cap (\omega : \tau_n \leq t) \in \mathcal{F}_t^X\}$ is the σ -algebra on Ω_{τ_n} . $\{\tau_n\}_{n=0}^\infty$ is called skeleton time sequence of the Markov skeleton process X . Furthermore, if on Ω_{τ_n}

$$E[f(X(\tau_n + \cdot)) | \mathcal{F}_{\tau_n}^X] = E[(f(X(\tau_n + \cdot)) | X(\tau_n))] = E_{X(\tau_n)}[f(X(\cdot))] \quad (2.2)$$

P-a.s. holds, where $E_x(\cdot)$ denotes the expectation corresponding to $P(\cdot | X(0) = x)$, then X is called a time homogeneous Markov skeleton process.

Definition 2.2 (see [26]). A time homogeneous Markov skeleton process $X = \{X(t, \omega), \leq t < \infty\} \rightarrow \{X(t, \omega), 0 \leq t < \infty\}$ is called normal, if there exists a function $h(x, t, A)$ on $E \times \mathbf{R}^+ \times \varepsilon$, such that

- (i) for fixed x and t , $h(x, t, \cdot)$ is a finite measure on ε ,
- (ii) for fixed $A \in \varepsilon$, $h(\cdot, \cdot, A)$ is $\varepsilon \times \mathcal{B}(\mathbf{R}^+)$ measurable function on $E \times \mathbf{R}^+$,
- (iii) for any $t \geq 0$, $A \in \varepsilon$,

$$h(X(\tau_n), t, A) = P\{X(\tau_n + t) \in A, \tau_{n+1} - \tau_n > t | X(\tau_n)\} \quad \text{P-a.s.} \quad (2.3)$$

2.2. Backward Equations of Markov Skeleton Processes

Theorem 2.3 (see [26]). Suppose that $X = \{X(t); t \geq 0\}$ is a normal Markov skeleton process with $\{\tau_n\}_{n=0}^\infty$ as its skeleton time sequence, then for any $x \in E$, $t \geq 0$, $A \in \varepsilon$,

$$p(x, t, A) = h(x, t, A) + \int_E \int_0^t \sum_{n=1}^\infty q^{(n)}(x, ds, dy) h(y, t - s, A). \quad (2.4)$$

Thus, $p(x, t, A)$ is a minimal nonnegative solution to the following nonnegative equation system: $\forall x \in E$, $t \geq 0$, $A \in \varepsilon$,

$$p(x, t, A) = h(x, t, A) + \int_E \int_0^t q(x, ds, dy) p(y, t - s, A). \quad (2.5)$$

Formula (2.5) is called the backward equations of Markov skeleton processes.

2.3. Limit Distribution of Markov Skeleton Processes

Definition 2.4 (see [27]). Suppose that $X(t)$ is a normal Markov skeleton process with $\{\tau_n\}_{n=0}^{\infty}$ as its skeleton time sequence. If there exists probability measure $\pi(\cdot)$ on (E, \mathcal{E}) , such that for any $A \in \mathcal{E}$,

$$P(X(\tau_1) \in A \mid X(0) = x, \tau_1 = s) = P(X(\tau_1) \in A) = \pi(A), \quad (2.6)$$

then $X(t)$ is called a Doob skeleton process, $\pi(\cdot)$ is called the characteristic measure of $X(t)$, and $\{\tau_n, n = 1, 2, \dots\}$ is the Doob skeleton time sequence of $X(t)$.

For any $n \in \mathbb{N}$, $t \geq 0$, $A \in \mathcal{E}$,

$$q^{(n)}(X(\tau_m), t, A) = P\{X(\tau_{m+n}) \in A, \tau_{m+n} \leq t \mid X(\tau_m)\} \quad (2.7)$$

and $q^{(1)}(x, t, A)$ is abbreviated to $q(x, t, A)$,

$$\begin{aligned} p(x, t, A) &= P\{X(t) \in A \mid X(0) = x\}, \\ F(x, t) &= P(\tau_1 \leq t \mid X(0) = x), \quad \forall x \in E, t \geq 0, \\ F(t) &= \int_E F(x, t) \pi(dx), \quad t \geq 0, \\ h(t, A) &= \int_E h(x, t, A) \pi(dx), \\ m &= \int_0^{\infty} t dF(t). \end{aligned} \quad (2.8)$$

Definition 2.5 (see [27]). Suppose that $X(t)$ is a Doob skeleton processes. If $m < \infty$ and for any $x \in E$, $F(x, 0) \equiv 0$, $F(x, \infty) \equiv 1$, then $X(t)$ is called a positive recurrent Doob skeleton process.

Theorem 2.6 (see [27]). Suppose that $X(t)$ is a positive recurrent Doob skeleton process. If $F(t)$ is not lattice distribution, then for $\forall A \in \mathcal{E}$, the limit distribution $p(\cdot)$ of $X(t)$ exists,

$$p(A) = \lim_{t \rightarrow \infty} p(x, t, A) = \frac{\int_0^{\infty} h(t, A) dt}{m}, \quad (2.9)$$

and $p(\cdot)$ is a probability distribution in (E, \mathcal{E}) .

3. Limit Distribution of Inventory Level of Perishable Inventory Model

The perishable inventory model studied in this paper has been proposed and investigated in [26], which obtained the backward equations of the inventory level of this model. Different from [26], this paper study the limit distribution of the inventory level.

3.1. Perishable Inventory Model

First, we present the details of the perishable inventory model as follows (see [26]).

- (1) Assume that lifetimes of inventory commodities are i.i.d random variables, with a common distribution function $F(t)$, where $F(t)$ is continuous and satisfies $\int_0^\infty t dF(t) = \mu_1$.
- (2) Sell one item each time, and the sale times of each item are i.i.d random variables, with a common distribution function $G(t)$, where $G(t)$ is continuous and satisfies $\int_0^\infty t dG(t) = \mu_2$. Assume that the sale times are also independent of the commodities' lifetimes.
- (3) The maximum capacity of the warehouse is a fixed value S_{\max} . When the inventory level becomes S_{\min} ($S_{\min} < 0$) (i.e., the quantity of out of stock arrives S_{\min}), new commodities are replenished to increase the inventory level until it reaches S_{\max} .

Let $S(t)$ denote inventory level at time t . When $F(t)$ and $G(t)$ are not exponential distributions, $\{S(t), t \geq 0\}$ is not a Markov process. In this case, we introduce supplementary variables as follows: $\theta(t)$ denotes the lifetime of the item in stock at time t , and $\hat{\theta}(t)$ denotes the time interval between the last sale before t and time t .

As one item is consumed and the other item perishes at the same time is a rare event, so we don't consider this case and suppose $F(t)$ and $G(t)$ are continuous distribution. Let τ_n denote the n th discontinuous point of $(S(t), \theta(t), \hat{\theta}(t))$, that is, one item is consumed or perishes at τ_n . At τ_n , $(S(t), \theta(t), \hat{\theta}(t))$ has Markov property, so by Definition 2.1, $(S(t), \theta(t), \hat{\theta}(t))$ is a Markov skeleton process with τ_n as its Markov skeleton time sequence.

3.2. Limit Distribution of Inventory Level

In this subsection, we obtain the limit distribution of inventory level.

Suppose that $T_0 = 0$, and T_n denotes the n th times when the process $(S(t), \theta(t), \hat{\theta}(t))$ returns to state $(S_{\max}, 0, 0)$. Let

$$\begin{aligned} T_{n+1} &= T_n + \theta_{T_n} \cdot T_1, \quad n = 1, 2, \dots, \\ Y_0 &= T_1, \quad Y_i = T_{i+1} - T_i, \quad i \geq 1, \end{aligned} \tag{3.1}$$

then Y_i is the replenishment interval. By Definition 2.2,

$$h_{S_{\max}, j}(t) = P\{S(t) = j, t \leq T_1 \mid S(0) = S_{\max}, \theta(0) = \hat{\theta}(0) = 0\}. \tag{3.2}$$

Let $M(t)$ denote the distribution function of Y_i , and M denote the expectation of Y_i , then,

$$\begin{aligned} M &= E\left\{T_1 \mid S(0) = S_{\max}, \theta(0) = \hat{\theta}(0) = 0\right\} \\ &= \int_0^{\infty} P\left\{T_1 \geq t \mid S(0) = S_{\max}, \theta(0) = \hat{\theta}(0) = 0\right\} dt \\ &= \int_0^{\infty} \sum_{j=S_{\min}+1}^{S_{\max}} h_j(t) dt. \end{aligned} \quad (3.3)$$

Theorem 3.1. *If $\mu_1 < +\infty$, $\mu_2 < +\infty$, $S(t)$ is a positive recurrent Doob skeleton process with $\{T_n\}$ as its Doob skeleton time sequence.*

Proof. As T_n denotes the beginning of the n th replenishment and T_n^- denotes the moment before the n th replenishment, we have $S(T_n^-) = S_{\min}$, $S(T_n) = S_{\max}$, then $S(t)$ satisfies Markov property at T_n , which assures that $S(t)$ is a Markov skeleton process. At the beginning of every replenishment, $S(T_n) \equiv S_{\max}$, so we have

$$P(S(T_1) = j \mid S(0) = i, T_1 = s) = \delta_0(j - S_{\max}). \quad (3.4)$$

Thus, $S(t)$ is Doob skeleton process by Definition 2.4. If $\mu_1 < +\infty$, $\mu_2 < +\infty$, we obtain $M = E[Y_i] < +\infty$. Therefore, $S(t)$ is a positive recurrent Doob skeleton process with $\{T_n\}$ as its Doob skeleton time sequence. \square

Theorem 3.2. *If $\mu_1 < +\infty$, $\mu_2 < +\infty$, and $F(x)$, $G(x)$ are not lattice distribution, then, for $j \in \{S_{\min} + 1, S_{\min} + 2, \dots, S_{\max}\}$,*

$$p_j(t) = \lim_{t \rightarrow \infty} p(S(t) = j \mid S(0) = i) = \frac{\int_0^{\infty} h_j(t) dt}{M} = \frac{\int_0^{\infty} h_j(t) dt}{\int_0^{\infty} \left[\sum_{j=S_{\min}+1}^{S_{\max}} h_j(t) \right] dt}; \quad (3.5)$$

for $A = \{1, 2, \dots, S_{\max}\}$,

$$p(A) = \lim_{t \rightarrow \infty} p(S(t) \in A \mid S(0) = i) = \frac{\int_0^{\infty} h(t, A) dt}{M} = \frac{\int_0^{\infty} \left[\sum_{j=1}^{S_{\max}} h_j(t) \right] dt}{\int_0^{\infty} \left[\sum_{j=S_{\min}+1}^{S_{\max}} h_j(t) \right] dt}; \quad (3.6)$$

for $B = \{S_{\min} + 1, S_{\min} + 2, \dots, 0\}$,

$$p(B) = \lim_{t \rightarrow \infty} p(S(t) \in B \mid S(0) = i) = \frac{\int_0^{\infty} h(t, B) dt}{M} = \frac{\int_0^{\infty} \left[\sum_{j=S_{\min}+1}^0 h_j(t) \right] dt}{\int_0^{\infty} \left[\sum_{j=S_{\min}+1}^{S_{\max}} h_j(t) \right] dt}, \quad (3.7)$$

and $p(\cdot)$ is a probability distribution in (E, \mathcal{E}) .

Proof. If $F(x)$, $G(x)$ are not lattice distribution, then $M(t)$ is not lattice distribution. According to Theorems 2.6, 3.1, and formula (3.3), we get formulas (3.5)–(3.7). Thus, the proof of the theorem is completed. \square

3.3. The Explicit Expression of $h_j(t)$

Next, we intend to give the explicit expression of $h_j(t)$ by applying backward equations of Markov skeleton processes.

By formula (3.4), we have $\pi(dx) = \delta_0(x - S_{\max})$. Then,

$$h_j(t) = \sum_{i=0}^{\infty} h_{ij}(t) \delta_0(i - S_{\max}) = h_{S_{\max},j}(t), \quad (3.8)$$

where $h_{S_{\max},j}(t)$ is defined in (3.2).

Let $\hat{\tau}_0 = 0$, $\hat{\tau}_n = \tau_n \wedge T_1$, $n = 1, 2, \dots$, where τ_n denotes the n th discontinuous point of $(S(t), \theta_1(t), \theta_2(t))$, then

$$\hat{\tau}_n \uparrow T_1, \quad n \uparrow +\infty. \quad (3.9)$$

According to Theorem 3.1, $(S(t), \theta_1(t), \theta_2(t))$, $t < T_1$ is a Markov skeleton process with $\hat{\tau}_n$ as its skeleton time sequence.

For $i > 0$, let

$$\begin{aligned} \hat{h}(i, \theta, \hat{\theta}, j, A, \hat{A}, t) &= P\{S(t) = j, \theta(t) \in A, \hat{\theta}(t) \in \hat{A}, t < \hat{\tau}_1 \mid S(0) = i, \theta(0) = \theta, \hat{\theta}(0) = \hat{\theta}\}; \\ \hat{q}(i, \theta, \hat{\theta}, ds, j, A, \hat{A}) &= P\{S(ds) = j, \theta(ds) \in A, \hat{\theta}(ds) \in \hat{A} \mid S(0) = i, \theta(0) = \theta, \hat{\theta}(0) = \hat{\theta}\}; \\ \hat{p}(i, \theta, \hat{\theta}, j, A, \hat{A}, t) &= P\{S(t) = j, \theta(t) \in A, \hat{\theta}(t) \in \hat{A}, t < T_1 \mid S(0) = i, \theta(0) = \theta, \hat{\theta}(0) = \hat{\theta}\}. \end{aligned} \quad (3.10)$$

Thus, $h_{S_{\max},j}(t)$ can be expressed as follows:

$$h_{S_{\max},j}(t) = \hat{p}(S_{\max}, 0, 0, j, t). \quad (3.11)$$

Lemma 3.3. When $F(t)$ and $G(t)$ are continuous, we have

$$\hat{h}(S_{\max}, 0, 0, j, t) = \begin{cases} 0, & j \neq S_{\max}; \\ (1 - F(t))^{S_{\max}} (1 - G(t)), & j = S_{\max}, \end{cases}$$

$$\hat{q}(S_{\max}, 0, 0, ds, j) = \begin{cases} C_{S_{\max}}^1 (1 - F(s))^{S_{\max}-1} (1 - G(s)) F(ds), & j = S_{\max} - 1, \\ \text{one item perishes, and no item is consumed;} & \\ C_{S_{\max}}^1 (1 - F(s))^{S_{\max}} G(ds), & j = S_{\max} - 1, \\ \text{one item is consumed, and no item perishes;} & \\ 0, & j \neq S_{\max} - 1. \end{cases} \quad (3.12)$$

Proof. By the definition of $\hat{h}(S_{\max}, 0, 0, j, t)$, there is no state transition of $S(t)$ up to t . If $j \neq S_{\max}$, $\hat{h}(S_{\max}, 0, 0, j, t) = 0$. If $j = S_{\max}$, which means that no item is consumed or perishes up to t , then $\hat{h}(S_{\max}, 0, 0, j, t) = (1 - F(t))^{S_{\max}}(1 - G(t))$.

By the definition of $\hat{q}(S_{\max}, 0, 0, ds, j)$, $S(t)$ will transfer from state S_{\max} to state j at ds . As $F(t), G(t)$ are continuous, when $j \neq S_{\max} - 1$, $\hat{q}(S_{\max}, 0, 0, ds, j) = 0$.

If one item perishes at time ds , and no item is consumed up to s , then $j = S_{\max} - 1$, $\hat{q}(S_{\max}, 0, 0, ds, j) = C_{S_{\max}}^1 (1 - F(s))^{S_{\max}-1} (1 - G(s)) F(ds)$.

If one item is consumed at time ds , and no item perishes up to s , then $j = S_{\max} - 1$, $\hat{q}(S_{\max}, 0, 0, ds, j) = C_{S_{\max}}^1 (1 - F(s))^{S_{\max}} G(ds)$. \square

According to Theorem 2.3 and Lemma 3.3, we have the following.

Theorem 3.4. $\hat{p}(S_{\max}, 0, 0, j, t)$ is the minimal nonnegative solution to the following nonnegative linear equation,

$$\begin{aligned} \hat{p}(S_{\max}, 0, 0, j, t) &= \delta_{S_{\max}, j} (1 - F(t))^{S_{\max}} (1 - G(t)) \\ &+ \int_0^t C_{S_{\max}}^1 (1 - F(s))^{S_{\max}-1} (1 - G(s)) F(ds) \hat{p}(S_{\max} - 1, s, s, j, t - s) \\ &+ \int_0^t C_{S_{\max}}^1 (1 - F(s))^{S_{\max}} G(ds) \hat{p}(S_{\max} - 1, s, 0, j, t - s). \end{aligned} \quad (3.13)$$

Thus, combining formulas (3.8), (3.11), and Theorem 3.4, the explicit expression of $h_j(t)$ is obtained; $h_j(t) = \hat{p}(S_{\max}, 0, 0, j, t)$ is the minimal nonnegative solution to formula (3.13).

Acknowledgments

This work was supported by the National Natural Science Foundation for Young Scholars of China (no. 11001179). The authors thank the editor and the anonymous referees for their constructive comments and suggestions for improving the quality of the paper.

References

- [1] T. M. Whitin, *Theory of Inventory Management*, Princeton University Press, Princeton, NJ, USA, 1957.
- [2] P. M. Ghare and G. P. Schrader, "A model for an exponentially decaying inventory," *Journal of Industrial Engineering*, vol. 14, no. 5, pp. 238–243, 1963.
- [3] F. Raafat, "Survey of literature on continuously deteriorating inventory models," *Journal of the Operational Research Society*, vol. 42, no. 1, pp. 27–37, 1991.
- [4] S. K. Goyal and B. C. Giri, "Recent trends in modeling of deteriorating inventory," *European Journal of Operational Research*, vol. 134, no. 1, pp. 1–16, 2001.
- [5] R. X. Li, H. J. Lan, and J. R. Mawhinney, "A review on deteriorating inventory study," *Journal Service Science & Management*, vol. 3, pp. 117–129, 2010.
- [6] B. Karmakar and K. D. Choudhury, "A review on inventory models for deteriorating items with shortages," *Assam University Journal of Science & Technology : Physical Sciences and Technology*, vol. 6, pp. 51–59, 2010.
- [7] C. L. Williams and B. E. Patuwo, "A perishable inventory model with positive order lead times," *European Journal of Operational Research*, vol. 116, no. 2, pp. 352–373, 1999.

- [8] Y. Adachi, T. Nose, and S. Kuriyama, "Optimal inventory control policy subject to different selling prices of perishable commodities," *International Journal of Production Economics*, vol. 60, pp. 389–394, 1999.
- [9] P. S. Deng, R. H.-J. Lin, and P. Chu, "A note on the inventory models for deteriorating items with ramp type demand rate," *European Journal of Operational Research*, vol. 178, no. 1, pp. 112–120, 2007.
- [10] G. P. Samanta and A. Roy, "A production inventory model with deteriorating items and shortages," *Yugoslav Journal of Operations Research*, vol. 14, no. 2, pp. 219–230, 2004.
- [11] S. R. Singh, N. Kumar, and R. Kumari, "Two warehouse inventory model for deteriorating items with shortages under inflation and time value of money," *International Journal of Computational and Applied Mathematics*, vol. 4, no. 1, pp. 83–94, 2009.
- [12] N. H. Shah and K. T. Shukla, "Deteriorating inventory model for waiting time partial backlogging," *Applied Mathematical Sciences*, vol. 3, no. 9–12, pp. 421–428, 2009.
- [13] N. H. Shah and Y. K. Shah, "Literature survey on inventory model for deteriorating items," *Economic Annals*, vol. 44, pp. 221–237, 2000.
- [14] Y. K. Shah and M. C. Jaiswal, "An order-level inventory model for a system with constant rate of deterioration," *Opsearch*, vol. 14, no. 3, pp. 174–184, 1977.
- [15] S.-P. Wang, "An inventory replenishment policy for deteriorating items with shortages and partial backlogging," *Computers & Operations Research*, vol. 29, no. 14, pp. 2043–2051, 2002.
- [16] N. Ravichandran, "Stochastic analysis of a continuous review perishable inventory system with positive lead time and Poisson demand," *European Journal of Operational Research*, vol. 84, no. 2, pp. 444–457, 1995.
- [17] H. N. Chiu, "An approximation to the continuous review inventory model with perishable items and lead times," *European Journal of Operational Research*, vol. 87, no. 1, pp. 93–108, 1995.
- [18] L. Liu and T. Yang, "An (s, S) random lifetime inventory model with a positive lead time," *European Journal of Operational Research*, vol. 113, no. 1, pp. 52–63, 1999.
- [19] B. Sivakumar, "A perishable inventory system with retrial demands and a finite population," *Journal of Computational and Applied Mathematics*, vol. 224, no. 1, pp. 29–38, 2009.
- [20] H. K. Alfares, "Inventory model with stock-level dependent demand rate and variable holding cost," *International Journal of Production Economics*, vol. 108, no. 1-2, pp. 259–265, 2007.
- [21] T. Roy and K. S. Chaudhuri, "A production-inventory model under stock-dependent demand, Weibull distribution deterioration and shortage," *International Transactions in Operational Research*, vol. 16, no. 3, pp. 325–346, 2009.
- [22] K. Kanchanasuntorn and A. Techanitisawad, "An approximate periodic model for fixed-life perishable products in a two-echelon inventory-distribution system," *International Journal of Production Economics*, vol. 100, no. 1, pp. 101–115, 2006.
- [23] Z. Hou, Z. Liu, and J. Zou, "QNQL processes: (H, Q) -processes and their applications," *Chinese Science Bulletin*, vol. 42, no. 11, pp. 881–886, 1997.
- [24] Z.-T. Hou, "Markov skeleton processes and applications to queueing systems," *Acta Mathematicae Applicatae Sinica. English Series*, vol. 18, no. 4, pp. 537–552, 2002.
- [25] Z. Hou, C. Yuan, J. Zou et al., "Transient distribution of the length of $GI/G/N$ queueing systems," *Stochastic Analysis and Applications*, vol. 21, no. 3, pp. 567–592, 2003.
- [26] Z. T. Hou and G. X. Liu, *Markov Skeleton Processes and Their Applications*, Science Press and International Press, 2005.
- [27] H. L. Dong, Z. T. Hou, and G. C. Jiang, "Limit distribution of Markov skeleton processes," *Acta Mathematicae Applicatae Sinica*, vol. 33, no. 2, pp. 290–296, 2010 (Chinese).



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

