

Research Article

A Defect-Correction Mixed Finite Element Method for Stationary Conduction-Convection Problems

Zhiyong Si and Yinnian He

Faculty of Science, Xi'an Jiaotong University, Xi'an 710049, China

Correspondence should be addressed to Yinnian He, heyn@mail.xjtu.edu.cn

Received 29 July 2010; Revised 15 November 2010; Accepted 5 January 2011

Academic Editor: Katica R. (Stevanovic) Hedrih

Copyright © 2011 Z. Si and Y. He. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

A defect-correction mixed finite element method (MFEM) for solving the stationary conduction-convection problems in two-dimension is given. In this method, we solve the nonlinear equations with an added artificial viscosity term on a grid and correct this solution on the same grid using a linearized defect-correction technique. The stability is given and the error analysis in L^2 and H^1 -norm of u , T and the L^2 -norm of p are derived. The theory analysis shows that our method is stable and has a good precision. Some numerical results are also given, which show that the defect-correction MFEM is highly efficient for the stationary conduction-convection problems.

1. Introduction

In this paper, we consider the stationary conduction-convection problems in two dimension whose coupled equations governing viscous incompressible flow and heat transfer for the incompressible fluid are Boussinesq approximations to the stationary Navier-Stokes equations.

(\mathcal{P}) Find $(u, p, T) \in X \times M \times W$ such that

$$\begin{aligned} -\nu \Delta u + (u \cdot \nabla)u + \nabla p &= \lambda j T, & x \in \Omega, \\ \operatorname{div} u &= 0, & x \in \Omega, \\ -\Delta T + \lambda u \cdot \nabla T &= 0, & x \in \Omega, \\ u &= 0, \quad T = T_0, & x \in \partial\Omega, \end{aligned} \tag{1.1}$$

where Ω is a bounded domain in \mathbb{R}^2 assumed to have a Lipschitz continuous boundary $\partial\Omega$. $u = (u_1(x), u_2(x))^T$ represents the velocity vector, $p(x)$ the pressure, $T(x)$ the temperature, $\lambda > 0$ the Grashoff number, $j = (0, 1)^T$ the two-dimensional vector, and $\nu > 0$ the viscosity.

As we know the conduction-convection problem contains the velocity vector field, the pressure field and the temperature field, so finding the numerical solution of conduction-convection problems is very difficult. The conduction-convection problems is an important system of equations in atmospheric dynamics and dissipative nonlinear system of equations, so lots of works are devoted to this problem [1–6]. There are also some works devoted to the nonstationary conduction-convection problems [7–10]. In [8], Luo et al. gave an optimizing reduced PLSMFE for the nonstationary conduction-convection problems. They combined PLSMEF method with POD to deal with the problems. In [11], an analysis of conduction natural convection conjugate heat transfer in the gap between concentric cylinders under solar irradiation was studied. In [12], a Newton iterative mixed finite element method for the stationary conduction-convection problems was shown by Si et al. In [13], Si and He gave a coupled Newton iterative mixed finite element method for the stationary conduction-convection problems.

The defect-correction method is an iterative improvement technique for increasing the accuracy of a numerical solution without applying a grid refinement. Due to its good efficiency, there are many works devoted to this method, for example, [14–28]. In [18], a method making it possible to apply the idea of iterated defect correction to finite element methods was given. A method for solving the time-dependent Navier-Stokes equations, aiming at higher Reynolds' number, was presented in [23]. In [27], an accurate approximations for self-adjoint elliptic eigenvalues was presented. In [28], Stetter exposed the common structural principle of all these techniques and exhibit the principal modes of its implementation in a discretization context.

In this paper we present a defect-correction MFEM for the stationary conduction convection problems. In this method, we solve the nonlinear equations with an added artificial viscosity term on a finite element grid and correct this solution on the same grid using a linearized defect-correction technique. Actually, the defect-correction MFEM incorporates the artificial viscosity term as a stabilizing factor, making both the nonlinear system easier to resolve and the linearized system easier to precondition. The stability and error analysis of the coupled the defect-correction MFEM show that this method is stable and has a good precision. Some numerical experiments show that our analysis is proper and our method is effective. And it can be used for solving the convection-conduction problems with much small viscosity.

This paper is organized as follows. In Section 2, the functional settings and some assumptions are given. Section 3 is devoted to the defect-correction MFEM. Section 4 gives the stability analysis. Section 5 presents the error analysis. In Section 6, some numerical results and the numerical analysis to validate the effectiveness of the method are laid out.

2. Functional Setting for the Conduction Convection Problems

In this section, we aim to describe some of the notations and results which will be frequently used in this paper. The Sobolev spaces used in this context are standard [29]. For the mathematical setting of the conduction-convection problems and MFEM of conduction-convection problems (1.1), we introduce the Hilbert spaces

$$\begin{aligned} X &= H_0^1(\Omega)^2, & W &= H^1(\Omega), \\ M &= L_0^2(\Omega) \doteq \left\{ \varphi \in L^2(\Omega); \int_{\Omega} \varphi \, dx = 0 \right\}. \end{aligned} \tag{2.1}$$

\mathfrak{T}_h is the uniformly regular family of triangulation of $\overline{\Omega}$, indexed by a parameter $h = \max_{K \in \mathfrak{T}_h} \{h_K; h_K = \text{diam}(K)\}$. We introduce the finite element subspace $X_h \subset X$, $M_h \subset M$, $W_h \subset W$ as follows

$$\begin{aligned} X_h &= \left\{ v_h \in X \cap C^0(\overline{\Omega})^2; v_h|_K \in P_\ell(K)^2, \forall K \in \mathfrak{T}_h \right\}, \\ M_h &= \left\{ q_h \in M \cap C^0(\overline{\Omega}); q_h|_K \in P_k(K), \forall K \in \mathfrak{T}_h \right\}, \\ W_h &= \left\{ \phi_h \in W \cap C^0(\overline{\Omega}); \phi_h|_K \in P_l(K), \forall K \in \mathfrak{T}_h \right\}, \end{aligned} \quad (2.2)$$

where $P_\ell(K)$ is the space of piecewise polynomials of degree ℓ on K , and $\ell \geq 1, k \geq 1, l \geq 1$ are three integers. $W_{0h} = W_h \cap H_0^1(\Omega)$, and (X_h, M_h) satisfies the discrete LBB condition

$$\sup_{v_h \in X_h} \frac{d(\varphi_h, v_h)}{\|\nabla v_h\|_0} \geq \beta \|\varphi_h\|_0, \quad \forall \varphi_h \in M_h, \quad (2.3)$$

where $d(\varphi, v) = (\varphi, \text{div } v)$.

With the above notations, the Galerkin mixed variation and the mixed FEM problem for the conduction-convection problems (\mathcal{P}) are defined, respectively, as follows.

(\mathcal{P}') Find $(u, p, T) \in X \times M \times W$ such that

$$\begin{aligned} va(u, v) - d(p, v) + d(\varphi, u) + b(u, u, v) &= \lambda(jT, v), \quad \forall v \in X, \varphi \in M, \\ \bar{a}(T, \varphi) + \lambda \bar{b}(u, T, \varphi) &= 0, \quad \forall \varphi \in W_0. \end{aligned} \quad (2.4)$$

(\mathcal{P}'') Find $(u_h, p_h, T_h) \in X_h \times M_h \times W_h$ such that

$$\begin{aligned} va(u_h, v_h) - d(p_h, v_h) + d(\varphi_h, u_h) + b(u_h, u_h, v_h) &= \lambda(jT_h, v_h), \quad \forall v_h \in X_h, \varphi_h \in M_h, \\ \bar{a}(T_h, \varphi_h) + \lambda \bar{b}(u_h, T_h, \varphi_h) &= 0, \quad \forall \varphi_h \in W_{0h}, \end{aligned} \quad (2.5)$$

where $a(u, v) = (\nabla u, \nabla v)$, $d(\varphi, v) = (\varphi, \text{div } v)$, $\bar{a}(T, \varphi) = (\nabla T, \nabla \varphi)$, and

$$\begin{aligned} b(u, v, w) &= \frac{1}{2} \left[\int_{\Omega} \sum_{i,k=1}^2 u_i \frac{\partial v_k}{\partial x_i} w_k dx - \sum_{i,k=1}^2 u_i \frac{\partial w_k}{\partial x_i} v_k dx \right], \quad \forall u, v, w \in X, \\ \bar{b}(u, T, \varphi) &= \frac{1}{2} \left[\int_{\Omega} \sum_{i=1}^2 u_i \frac{\partial T}{\partial x_i} \varphi dx - \sum_i u_i \frac{\partial \varphi}{\partial x_i} T dx \right], \quad \forall u \in X, T, \varphi \in W. \end{aligned} \quad (2.6)$$

The following assumptions and results are recalled (see [7, 29–31]).

(A₁) There exists a constant C_0 which only depends on Ω , such that

- (i) $\|u\|_0 \leq C_0 \|\nabla u\|_0$, $\|u\|_{0,4} \leq C_0 \|\nabla u\|_0$, for all $u \in H_0^1(\Omega)^2$ (or $H_0^1(\Omega)$),
- (ii) $\|u\|_{0,4} \leq C_0 \|u\|_1$, for all $u \in H^1(\Omega)^2$,
- (iii) $\|u\|_{0,4} \leq \sqrt{2} \|\nabla u\|_0^{1/2} \|u\|_0^{1/2}$, for all $u \in H_0^1(\Omega)^2$ (or $H_0^1(\Omega)$).

(A₂) Assuming $\partial\Omega \in C^{k,\alpha}$ ($k \geq 0$, $\alpha > 0$), then, for $T_0 \in C^{k,\alpha}(\partial\Omega)$, there exists an extension in $C_0^{k,\alpha}(\mathbb{R}^2)$ (denote T_0 also), such that

$$\|T_0\|_{k,q} \leq \varepsilon, \quad k \geq 0, \quad 1 \leq q \leq \infty, \quad (2.7)$$

where ε is an arbitrary positive constant.

(A₃) $b(\cdot, \cdot, \cdot)$ and $\bar{b}(\cdot, \cdot, \cdot)$ have the following properties.

(i) For all $u \in X$, $v, w \in X$ (or $T, \varphi \in H_0^1(\Omega)$), there holds that

$$b(u, v, v) = 0, \quad b(u, v, w) = -b(u, w, v), \quad (2.8)$$

$$\bar{b}(u, T, T) = 0, \quad \bar{b}(u, T, \varphi) = -\bar{b}(u, \varphi, T). \quad (2.9)$$

(ii) For all $u \in X$, $v \in H^1(\Omega)^2$ (or $T \in H^1(\Omega)$), for all $w \in X$ (or $\varphi \in H_0^1(\Omega)$), there holds that

$$|b(u, v, w)| \leq N \|\nabla u\|_0 \|\nabla v\|_0 \|\nabla w\|_0, \quad (2.10)$$

$$|\bar{b}(u, T, \varphi)| \leq \bar{N} \|\nabla u\|_0 \|\nabla T\|_0 \|\nabla \varphi\|_0, \quad (2.11)$$

where

$$N = \frac{\sup_{u,v,w} |b(u, v, w)|}{(\|\nabla u\|_0 \|\nabla v\|_0 \|\nabla w\|_0)}, \quad (2.12)$$

$$\bar{N} = \frac{\sup_{u,T,\varphi} |\bar{b}(u, T, \varphi)|}{(\|\nabla u\|_0 \|\nabla T\|_0 \|\nabla \varphi\|_0)}.$$

We recall the following existence, uniqueness and regularity result of (\mathcal{P}') (see [7, Chapter 4]).

Theorem 2.1 (see [7]). *Under the assumption of (A₁)~(A₃), letting $A \equiv 2\nu^{-1}\lambda(3C_0 + 1)\|T_0\|_1$, $B \equiv 2\|\nabla T_0\|_0 + 2(C_0^2\lambda)^{-1}A$, there exist $0 < \delta_1, \delta_2 \leq 1$ such that*

$$\nu^{-1}NA \leq 1 - \delta_1, \quad \delta_1^{-1}\nu^{-1}C_0^2\lambda^2B\bar{N} \leq 1 - \delta_2. \quad (2.13)$$

Then, there exists a unique solution $(u, p, T) \in X \times M \times W$ for (\mathcal{P}') , and

$$\|\nabla u\|_0 \leq A, \quad \|\nabla T\|_0 \leq B. \quad (2.14)$$

Some estimates of the trilinear form b are given in the following lemma and the proof can be found in [30, 32–34].

Lemma 2.2. *The trilinear form b satisfies the following estimate:*

$$|b(u_h, v_h, w)| + |b(v_h, u_h, w)| + |b(w, u_h, v_h)| \leq C_0 |\log h|^{1/2} \|\nabla v_h\|_0 \|\nabla u_h\|_0 \|w\|_0, \quad (2.15)$$

for all $u_h, v_h \in V_h, w \in L^2(\Omega)^2$.

Lemma 2.3. *Suppose that (A_1) – (A_3) are valid and ε is a positive constant, such that*

$$\frac{32C_0^2 \lambda^2 \bar{N} \varepsilon}{3\nu} < 1, \quad \|\nabla T_0\|_0 \leq \frac{\varepsilon}{4}, \quad \|T_0\|_0 \leq \frac{C_0 \varepsilon}{4}, \quad (2.16)$$

then (\mathcal{P}'') has a unique solution $(u_h, p_h, T_h) \in X_h \times M_h \times W_h$, such that $T|_{\partial\Omega} = T_0$ and

$$\|\nabla u_h\|_0 \leq \frac{5C_0^2 \lambda \varepsilon}{3\nu}, \quad \|\nabla T_h\|_0 \leq \varepsilon. \quad (2.17)$$

Proof. The proof of the existence and the uniqueness of the solution has been given by Luo [7]. Let $T_h = \omega_h + T_0, \varphi_h = \omega_h$ in (2.5), we can get

$$\bar{a}(\omega_h, \omega_h) = -\lambda \bar{b}(u_h, T_0, \omega_h) - \bar{a}(T_0, \omega_h). \quad (2.18)$$

Using (2.11) and (2.16), we deduce

$$\|\nabla \omega_h\|_0 \leq \|\nabla T_0\|_0 + \lambda \bar{N} \varepsilon \|\nabla u_h\|_0. \quad (2.19)$$

Letting $v_h = u_h, \varphi_h = p_h$ in the first equation of (2.5), we get

$$\nu \|\nabla u_h\|_0^2 = |\lambda(jT_h, u_h)| \leq \lambda C_0 \|T_h\|_0 \|\nabla u_h\|_0. \quad (2.20)$$

By (2.16), we can obtain

$$\begin{aligned} \|\nabla u_h\|_0 &\leq \nu^{-1} \lambda C_0 \|T_h\|_0 \\ &\leq \nu^{-1} \lambda C_0 (\|\omega_h\|_0 + \|T_0\|_0) \\ &\leq \nu^{-1} \lambda C_0^2 \|\nabla \omega_h\|_0 + \nu^{-1} \lambda C_0 \|T_0\|_0 \\ &\leq \nu^{-1} \lambda C_0 \|T_0\|_0 + \nu^{-1} \lambda C_0^2 \|\nabla T_0\|_0 + \nu^{-1} \lambda^2 C_0^2 \bar{N} \varepsilon \|\nabla u_h\|_0. \end{aligned} \quad (2.21)$$

Using (2.16) again, we get

$$\|\nabla u_h\|_0 \leq \frac{5C_0^2 \lambda \varepsilon}{3\nu}. \quad (2.22)$$

By (2.19), we deduce

$$\begin{aligned}
\|\nabla T_h\|_0 &\leq \|\nabla \omega_h\|_0 + \|\nabla T_0\|_0 \\
&\leq 2\|\nabla T_0\|_0 + \lambda \overline{N} \varepsilon \|\nabla u_h\|_0 \\
&\leq 2\|\nabla T_0\|_0 + \frac{5C_0^2 \lambda^2 \overline{N} \varepsilon^2}{3\nu} \\
&\leq \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon.
\end{aligned} \tag{2.23}$$

□

We introduce the Laplace operator

$$\mathcal{A}u = -\Delta u, \quad \forall u \in D(\mathcal{A}) = H^2(\Omega)^2 \cap X. \tag{2.24}$$

Lemma 2.4 (see [35, 36]). *For all $u, w \in X, v \in D(A)$ there holds that*

$$|b(u, v, w)| + |b(v, u, w)| + |b(w, u, v)| \leq C \|\mathcal{A}v\|_0 \|w\|_0 \|\nabla u\|_0. \tag{2.25}$$

3. The Defect-Correction Method

The aim of this section is to give a method for solving the nonlinear system (2.5) on a coarser mesh than one uses when employing the standard FEM; the coarse-mesh solution is corrected using the same grid in our method. The defect-correction method in which we consider incorporates an artificial viscosity parameter σh as a stabilizing factor in the solution algorithm. For a fixed grid parameter h the method requires the solution of one nonlinear system and a few linear correction steps. It is described in the following paragraphs. We consider the following problems which is identical to (2.5) except for an artificial viscosity term.

(\mathcal{P}^*) Find $(u_h^0, p_h^0, T_h^0) \in X_h \times M_h \times W_h$ such that

$$\begin{aligned}
(\nu + \sigma h)a(u_h^0, v_h) - d(p_h^0, v_h) + d(\varphi_h, u_h^0) + b(u_h^0, u_h^0, v_h) &= \lambda(jT_h^0, v_h), \\
\forall v_h \in X_h, \varphi_h \in M_h,
\end{aligned} \tag{3.1}$$

$$(1 + \sigma h)\bar{a}(T_h^0, \varphi_h) + \lambda\bar{b}(u_h^0, T_h^0, \varphi_h) = 0, \quad \forall \varphi_h \in W_{0h}.$$

We define the residual or named defect $R(u_h^0, p_h^0, T_h^0), Q(u_h^0, p_h^0, T_h^0)$ for the momentum systems as follows:

$$\begin{aligned}
(R(u_h^0, p_h^0, T_h^0), v_h) &= \lambda(jT_h^0, v_h) - \nu a(u_h^0, v_h) + d(p_h^0, v_h) \\
&\quad - d(\varphi_h, u_h^0) - b(u_h^0, u_h^0, v_h), \\
(Q(u_h^0, p_h^0, T_h^0), \varphi_h) &= -\bar{a}(T_h^0, \varphi_h) - \lambda\bar{b}(u_h^0, T_h^0, \varphi_h).
\end{aligned} \tag{3.2}$$

Define the correction $(\varepsilon_h^0, \varrho_h^0, \tau_h^0)$ satisfying the following linear problem:

$$\begin{aligned}
 & (\nu + \sigma h)a(\varepsilon_h^0, v_h) - d(\varrho_h^0, v_h) + d(\varphi_h, \varepsilon_h^0) + b(\varepsilon_h^0, u_h^0, v_h) + b(u_h^0, \varepsilon_h^0, v_h) \\
 & \quad = \left(R(u_h^0, p_h^0, T_h^0), v_h \right), \quad \forall v_h \in X_h, \varphi_h \in M_h, \\
 & (1 + \sigma h)\bar{a}(\tau_h^0, \varphi_h) + \lambda\bar{b}(u_h^0, \tau_h^0, \varphi_h) + \lambda\bar{b}(\varepsilon_h^0, T_h^0, \varphi_h) \\
 & \quad = \left(Q(u_h^0, p_h^0, T_h^0), \varphi_h \right), \quad \forall \varphi_h \in W_{0h}.
 \end{aligned} \tag{3.3}$$

Define $u_h^1 = u_h^0 + \varepsilon^0$, $p_h^1 = p_h^0 + \varrho_h^0$, $T_h^1 = T_h^0 + \tau_h^0$, which are hoped to be better solutions of the problems. In order to obtain the equations for (u_h^1, p_h^1, T_h^1) , we use the residual equation (3.2) to rewrite the linear problems (3.3); we obtain

$$(\rho^\dagger) \begin{cases} (\nu + \sigma h)a(u_h^1, v_h) - d(p_h^1, v_h) + d(\varphi_h, u_h^1) + b(u_h^0, u_h^1, v_h) + b(u_h^1, u_h^0, v_h) \\ \quad = \lambda(jT_h^1, v_h) + \sigma ha(u_h^0, v_h) + b(u_h^0, u_h^0, v_h), \quad \forall v_h \in X_h, \varphi_h \in M_h, \\ (1 + \sigma h)\bar{a}(T_h^1, \varphi_h) + \lambda\bar{b}(u_h^1, T_h^0, \varphi_h) + \lambda\bar{b}(u_h^0, T_h^1, \varphi_h) \\ \quad = \sigma h\bar{a}(T_h^0, \varphi_h) + \lambda\bar{b}(u_h^0, T_h^0, \varphi_h), \quad \forall \varphi_h \in W_{0h}. \end{cases} \tag{3.4}$$

In general, this method can be described as follows.

Step 1. Solve the nonlinear systems (3.1) for (u_h^0, p_h^0, T_h^0) .

Step 2. For $i = 1, 2, \dots, m$, solve the linear equations

$$(\rho^\dagger) \begin{cases} (\nu + \sigma h)a(u_h^i, v_h) - d(p_h^i, v_h) + d(\varphi_h, u_h^i) + b(u_h^{i-1}, u_h^i, v_h) + b(u_h^i, u_h^{i-1}, v_h) \\ \quad = (T_h^i, v_h) + \sigma ha(u_h^{i-1}, v_h) + b(u_h^{i-1}, u_h^{i-1}, v_h), \quad \forall v_h \in X_h, \varphi_h \in M_h, \\ (1 + \sigma h)\bar{a}(T_h^i, \varphi_h) + \lambda\bar{b}(u_h^i, T_h^{i-1}, \varphi_h) + \lambda\bar{b}(u_h^{i-1}, T_h^i, \varphi_h) \\ \quad = \sigma h\bar{a}(T_h^{i-1}, \varphi_h) + \lambda\bar{b}(u_h^{i-1}, T_h^{i-1}, \varphi_h), \quad \forall \varphi_h \in W_{0h}. \end{cases} \tag{3.5}$$

For each i the residual is given by

$$\begin{aligned}
 & \left(R(u_h^i, p_h^i, T_h^i), v_h \right) = \lambda(jT_h^i, v_h) - \nu a(u_h^i, v_h) + d(p_h^i, v_h) \\
 & \quad - d(\varphi_h, u_h^i) - b(u_h^i, u_h^i, v_h), \\
 & \left(Q(u_h^i, p_h^i, T_h^i), \varphi_h \right) = -\bar{a}(T_h^i, \varphi_h) - \lambda\bar{b}(u_h^i, T_h^i, \varphi_h).
 \end{aligned} \tag{3.6}$$

The correction $(\varepsilon_h^i, \varrho_h^i, \tau_h^i)$ is given by

$$\begin{aligned}
& (\nu + \sigma h)a(\varepsilon_h^i, v_h) - d(\varrho_h^i, v_h) + d(\varphi_h, \varepsilon_h^i) + b(\varepsilon_h^i, u_h^i, v_h) + b(u_h^i, \varepsilon_h^i, v_h) \\
& = \left(R(u_h^i, p_h^i, T_h^i), v_h \right), \quad \forall v_h \in X_h, \varphi_h \in M_h, \\
& (1 + \sigma h)\bar{a}(\tau_h^i, \varphi_h) + \lambda \bar{b}(u_h^i, \tau_h^i, \varphi_h) + \lambda \bar{b}(\varepsilon_h^i, T_h^i, \varphi_h) \\
& = \left(Q(u_h^i, p_h^i, T_h^i), \varphi_h \right), \quad \forall \varphi_h \in W_{0h}.
\end{aligned} \tag{3.7}$$

Remark 3.1. From the numerical experiments, we see that one or two correction steps is adequate. And this is as same as [24].

4. Stability Analysis

In this section, we give the stability analysis. It is given by the following theorems.

Theorem 4.1. *Under the assumptions of Lemma 2.3, then (u_h^0, T_h^0) defined by (\mathcal{D}^*) satisfies*

$$\left\| \nabla u_h^0 \right\|_0 \leq \frac{5C_0^2 \lambda \varepsilon}{3(\nu + \sigma h)}, \quad \left\| \nabla T_h^0 \right\|_0 \leq \varepsilon. \tag{4.1}$$

Moreover, if

$$\frac{25C_0^2 N \lambda \varepsilon}{3(\nu + \sigma h)^2} < 1, \tag{4.2}$$

(\mathcal{D}^*) admits a unique solution.

Proof. We define the set

$$\mathcal{B}_M = \left\{ \tilde{v}_h \in X_h; \left\| \nabla \tilde{v}_h \right\|_0 \leq \frac{5C_0^2 \lambda \varepsilon}{3(\nu + \sigma h)} \right\}. \tag{4.3}$$

Let \tilde{u}_h be in \mathcal{B}_M . Then

$$(1 + \sigma h)\bar{a}(T_h^0, \varphi_h) + \lambda \bar{b}(\tilde{u}_h, T_h^0, \varphi_h) = 0, \quad \forall \varphi_h \in W_{0h} \tag{4.4}$$

has a unique solution $T_h^0 \in W_h$ such that $T_h|_{\partial\Omega} = T_0$. For a given T_h^0 , we consider the following problem:

$$\begin{aligned}
& (\nu + \sigma h)a(u_h^{0*}, v_h) - d(p_h^{0*}, v_h) + d(\varphi_h, u_h^{0*}) + b(u_h^{0*}, u_h^{0*}, v_h) = \lambda(jT_h^0, v_h), \\
& \forall v_h \in X_h, \varphi_h \in M_h.
\end{aligned} \tag{4.5}$$

By the theory of the Navier-Stokes equations, we get (4.5) has a unique solution $(u_h^{0*}, p_h^{0*}) \in X_h \times M_h$ (see [31]). It means that (4.4) and (4.5) give a unique $u_h^{0*} \in X_h$ for a given $\tilde{u}_h \in X_h$, we denote

$$u_h^{0*} = \ell_h \tilde{u}_h. \quad (4.6)$$

Setting $T_h^0 = \omega_h^0 + T_0$, $\varphi_h = \omega_h^0$ in (4.4) and using (2.9), we can obtain

$$(1 + \sigma h) \bar{a}(\omega_h^0, \omega_h^0) = -\lambda \bar{b}(\tilde{u}_h, T_0, \omega_h^0) - (1 + \sigma h) \bar{a}(T_0, \omega_h^0). \quad (4.7)$$

Using (2.7), (2.11), and (2.16), we can get

$$\begin{aligned} (1 + \sigma h) \|\nabla \omega_h^0\|_0 &\leq \lambda \bar{N} \|\nabla \tilde{u}_h\|_0 \|\nabla T_0\|_0 + (1 + \sigma h) \|\nabla T_0\|_0, \\ \|\nabla \omega_h^0\|_0 &\leq \frac{\lambda \bar{N} \varepsilon}{4} \|\nabla \tilde{u}_h\|_0 + \|\nabla T_0\|_0. \end{aligned} \quad (4.8)$$

Using the triangle inequality, we have

$$\begin{aligned} \|\nabla T_h^0\|_0 &\leq \|\nabla T_0\|_0 + \|\nabla \omega_h^0\|_0 \\ &\leq \frac{\lambda \bar{N} \varepsilon}{4} \|\nabla \tilde{u}_h\|_0 + 2\|\nabla T_0\|_0 \\ &\leq \frac{5C_0^2 \bar{N} \lambda \varepsilon^2}{12(\nu + \sigma h)} + \frac{\varepsilon}{2} \leq \varepsilon. \end{aligned} \quad (4.9)$$

Letting $v_h = u_h^{0*}$, $\varphi_h = p_h^0$ in (4.5) and using (2.8), we get

$$(\nu + \sigma h) a(u_h^{0*}, u_h^{0*}) = \lambda (j_{T_h^0}^0, u_h^{0*}). \quad (4.10)$$

Letting $T_h^0 = \omega_h^0 + T_0$ and using (2.9), we have

$$\begin{aligned} (\nu + \sigma h) \|\nabla u_h^{0*}\|_0 &\leq C_0^2 \lambda \|\nabla \omega_h^0\|_0 + C_0 \lambda \|T_0\|_0 \\ &\leq C_0^2 \lambda^2 \bar{N} \|\nabla \tilde{u}_h\|_0 \|\nabla T_0\|_0 + C_0 \lambda (1 + C_0) \|\nabla T_0\|_0 \\ &\leq C_0^2 \lambda \varepsilon. \end{aligned} \quad (4.11)$$

Namely,

$$\|\nabla u_h^{0*}\|_0 \leq \frac{5C_0^2 \lambda \varepsilon}{3(\nu + \sigma h)}. \quad (4.12)$$

Hence, we proved that ℓ_h maps \mathcal{B}_M to \mathcal{B}_M . It follows from Brouwer's fixed-point theorem that there exists a solution to system (\mathcal{P}^*) .

To prove the uniqueness, assume that $(u_h^{01}, p_h^{01}, T_h^{01}), (u_h^{02}, p_h^{02}, T_h^{02}) \in X_h \times M_h \times W_h$, and $T_h^{01}|_{\partial\Omega} = T_h^{02}|_{\partial\Omega} = T_0$ are two solutions of (\mathcal{P}^*) . Then, we obtain that

$$\begin{aligned} & (\nu + \sigma h)a(u_h^{01} - u_h^{02}, v_h) - d(p_h^{01} - p_h^{02}, v_h) + d(\varphi_h, u_h^{01} - u_h^{02}) + b(u_h^{01} - u_h^{02}, u_h^{01}, v_h) \\ & \quad + b(u_h^{02}, u_h^{01} - u_h^{02}, v_h) = \lambda(j(T_h^{01} - T_h^{02}), v_h), \quad \forall v_h \in X_h, \varphi_h \in M_h, \\ & (1 + \sigma h)\bar{a}(T_h^{01} - T_h^{02}, \varphi_h) + \lambda\bar{b}(u_h^{02}, T_h^{01} - T_h^{02}, \varphi_h) + \lambda\bar{b}(u_h^{01} - u_h^{02}, T_h^{01}, \varphi_h) = 0, \quad \forall \varphi_h \in W_{0h}. \end{aligned} \quad (4.13)$$

Let $v_h = u_h^{01} - u_h^{02}, \varphi_h = p_h^{01} - p_h^{02}$ in the first equation of (4.13), we can get

$$(\nu + \sigma h)\|\nabla(u_h^{01} - u_h^{02})\|_0 \leq N\|\nabla(u_h^{01} - u_h^{02})\|_0\|\nabla u_h^{01}\|_0 + C_0^2\lambda\|\nabla(T_h^{01} - T_h^{02})\|_0. \quad (4.14)$$

Setting $\varphi_h = T_h^{01} - T_h^{02}$ in the second equation of (4.13), we obtain

$$(1 + \sigma h)\|\nabla(T_h^{01} - T_h^{02})\|_0 \leq \lambda\bar{N}\|\nabla(u_h^{01} - u_h^{02})\|_0\|\nabla T_h^{01}\|_0. \quad (4.15)$$

By (4.14) and (4.15), we deduce

$$\begin{aligned} & (\nu + \sigma h)\|\nabla(u_h^{01} - u_h^{02})\|_0 \\ & \quad \leq N\|\nabla u_h^{01}\|_0\|\nabla(u_h^{01} - u_h^{02})\|_0 + C_0^2\lambda^2\bar{N}\|\nabla T_h^{01}\|_0\|\nabla(u_h^{01} - u_h^{02})\|_0 \\ & \quad \leq \frac{5C_0^2N\lambda\varepsilon}{3(\nu + \sigma h)}\|\nabla(u_h^{01} - u_h^{02})\|_0 + \frac{C_0^2\lambda\bar{N}\varepsilon}{4\nu}\|\nabla(u_h^{01} - u_h^{02})\|_0. \end{aligned} \quad (4.16)$$

Using (4.2), we obtain

$$\|\nabla(u_h^{01} - u_h^{02})\|_0 \leq \frac{2}{5}\|\nabla(u_h^{01} - u_h^{02})\|_0. \quad (4.17)$$

Namely,

$$\|\nabla(u_h^{01} - u_h^{02})\|_0 = 0. \quad (4.18)$$

By (4.15), we see that $\|\nabla(T_h^{01} - T_h^{02})\|_0 = 0$. Therefore, it follows that (\mathcal{P}^*) has a unique solution.

Then, we give the prove of (4.1) without using (4.2). Letting $v_h = u_h^0, \varphi_h = p_h^0$ in the first equation of (3.1) and using (2.8), we get

$$(\nu + \sigma h)a(u_h^0, u_h^0) = \lambda(j(T_h^0, u_h^0)). \quad (4.19)$$

Letting $T_h^0 = \omega_h^0 + T_0$, we have

$$(\nu + \sigma h) \left\| \nabla u_h^0 \right\|_0 \leq C_0^2 \lambda \left\| \nabla \omega_h^0 \right\|_0 + C_0 \lambda \|T_0\|_0. \quad (4.20)$$

Letting $T_h^0 = \omega_h^0 + T_0$, $\psi_h = \omega_h^0$ in the second equation of (3.1) and using (2.9), we can obtain

$$(1 + \sigma h) \bar{a}(\omega_h^0, \omega_h^0) = -\lambda \bar{b}(u_h^0, T_0, \omega_h^0) - (1 + \sigma h) \bar{a}(T_0, \omega_h^0). \quad (4.21)$$

Using (2.7), (2.11), and (2.16), we can get

$$\begin{aligned} (1 + \sigma h) \left\| \nabla \omega_h^0 \right\|_0 &\leq \lambda \bar{N} \left\| \nabla u_h^0 \right\|_0 \| \nabla T_0 \|_0 + (1 + \sigma h) \| \nabla T_0 \|_0, \\ \left\| \nabla \omega_h^0 \right\|_0 &\leq \frac{\lambda \bar{N} \varepsilon}{4} \left\| \nabla u_h^0 \right\|_0 + \| \nabla T_0 \|_0. \end{aligned} \quad (4.22)$$

By (4.20) and (4.22), we can deduce

$$\begin{aligned} \left\| \nabla u_h^0 \right\|_0 &\leq (\nu + \sigma h)^{-1} \lambda C_0^2 \left\| \nabla \omega_h^0 \right\|_0 + (\nu + \sigma h)^{-1} C_0 \lambda \|T_0\|_0 \\ &\leq (\nu + \sigma h)^{-1} \left(\lambda C_0 \|T_0\|_0 + C_0^2 \lambda \| \nabla T_0 \|_0 + \frac{C_0^2 \lambda^2 \bar{N} \varepsilon}{4} \left\| \nabla u_h^0 \right\|_0 \right). \end{aligned} \quad (4.23)$$

Using (2.16), we get

$$\left\| \nabla u_h^0 \right\|_0 \leq \frac{5C_0^2 \lambda \varepsilon}{3(\nu + \sigma h)}. \quad (4.24)$$

Using (2.7), (2.11), (2.16), and (4.20), we can get

$$\begin{aligned} \left\| \nabla \omega_h^0 \right\|_0 &\leq \lambda \bar{N} \left\| \nabla u_h^0 \right\|_0 \| \nabla T_0 \|_0 + \| \nabla T_0 \|_0 \\ &\leq \frac{\lambda \bar{N} \varepsilon}{4} \left\| \nabla u_h^0 \right\|_0 + \| \nabla T_0 \|_0 \\ &\leq \frac{3\varepsilon}{4}, \\ \left\| \nabla T_h^0 \right\|_0 &\leq \left\| \nabla \omega_h^0 \right\|_0 + \| \nabla T_0 \|_0 \\ &\leq \varepsilon. \end{aligned} \quad (4.25)$$

Therefore, we finish the proof. \square

Theorem 4.2. *Under the assumptions of Lemma 2.3, and*

$$\frac{25C_0^2 N \lambda \varepsilon}{3(\nu + \sigma h)^2} < 1, \quad (4.26)$$

(u_h^1, T_h^1) defined by (3.4) satisfies

$$\|\nabla u_h^1\|_0 \leq \delta, \quad \|\nabla T_h^1\|_0 \leq \frac{5\varepsilon}{6} + \lambda \bar{N} \delta \varepsilon + \sigma h \varepsilon, \quad (4.27)$$

where $\delta \doteq (103C_0^2 \lambda \varepsilon / 48 + \sigma h (5C_0^2 \lambda \varepsilon / 3\nu)) / (7/10)(\nu + \sigma h)$.

Proof. Letting $v_h = u_h^1$, $\varphi_h = p_h^1$ in the first equation of (3.4) and using (2.8), we get

$$(\nu + \sigma h)a(u_h^1, u_h^1) + b(u_h^1, u_h^0, u_h^1) = b(u_h^0, u_h^0, u_h^1) + \sigma h a(u_h^0, u_h^1) + \lambda(jT_h^0, u_h^1). \quad (4.28)$$

Letting $T_h^0 = \omega_h^0 + T_0$ and using (2.10), we have

$$\begin{aligned} (\nu + \sigma h)\|\nabla u_h^1\|_0 &\leq N\|\nabla u_h^1\|_0\|\nabla u_h^0\|_0 + \sigma h\|\nabla u_h^0\|_0 + N\|\nabla u_h^0\|_0^2 \\ &\quad + C_0^2 \lambda \|\nabla \omega_h^1\|_0 + C_0 \lambda \|T_0\|_0. \end{aligned} \quad (4.29)$$

Let $T_h^1 = \omega_h^1 + T_0$, $\varphi_h = \omega_h^1$ in the second equation of (3.4), we can obtain

$$\begin{aligned} (1 + \sigma h)\bar{a}(\omega_h^1, \omega_h^1) &= -\lambda \bar{b}(u_h^0, T_0, \omega_h^1) - \lambda \bar{b}(u_h^1, T_h^0, \omega_h^1) + \lambda \bar{b}(u_h^0, T_h^0, \omega_h^1) \\ &\quad + \sigma h \bar{a}(T_h^0, \omega_h^1) - (1 + \sigma h)\bar{a}(T_0, \omega_h^1). \end{aligned} \quad (4.30)$$

Using (2.11) and (2.16), we can get

$$\begin{aligned} (1 + \sigma h)\|\nabla \omega_h^1\|_0 &\leq \lambda \bar{N}\|\nabla u_h^0\|_0\|\nabla T_0\|_0 + \lambda \bar{N}\|\nabla u_h^1\|_0\|\nabla T_h^0\|_0 \\ &\quad + \lambda \bar{N}\|\nabla u_h^0\|_0\|\nabla T_h^0\|_0 + \sigma h\|\nabla T_h^0\|_0 + (1 + \sigma h)\|\nabla T_0\|_0, \end{aligned} \quad (4.31)$$

$$\begin{aligned} \|\nabla \omega_h^1\|_0 &\leq \lambda \bar{N}\|\nabla u_h^0\|_0\|\nabla T_0\|_0 + \lambda \bar{N}\|\nabla u_h^1\|_0\|\nabla T_h^0\|_0 \\ &\quad + \lambda \bar{N}\|\nabla u_h^0\|_0\|\nabla T_h^0\|_0 + \sigma h\|\nabla T_h^0\|_0 + \|\nabla T_0\|_0. \end{aligned} \quad (4.32)$$

Using (4.29), we get

$$\begin{aligned}
& (\nu + \sigma h) \left\| \nabla u_h^1 \right\|_0 \\
& \leq N \left\| \nabla u_h^1 \right\|_0 \left\| \nabla u_h^0 \right\|_0 + \sigma h \left\| \nabla u_h^0 \right\|_0 + N \left\| \nabla u_h^0 \right\|_0^2 \\
& \quad + C_0^2 \lambda \left(\lambda \bar{N} \left\| \nabla u_h^0 \right\|_0 \left\| \nabla T_0 \right\|_0 + \lambda \bar{N} \left\| \nabla u_h^1 \right\|_0 \left\| \nabla T_h^0 \right\|_0 + \lambda \bar{N} \left\| \nabla u_h^0 \right\|_0 \left\| \nabla T_h^0 \right\|_0 \right. \\
& \quad \left. + \left\| \nabla T_h^0 \right\|_0 + \left\| \nabla T_0 \right\|_0 \right) + C_0 \lambda \left\| T_0 \right\|_0. \tag{4.33} \\
& (\nu + \sigma h - N \left\| \nabla u_h^0 \right\|_0 - C_0^2 \lambda^2 \bar{N} \left\| \nabla T_h^0 \right\|_0) \left\| \nabla u_h^1 \right\|_0 \\
& \leq N \left\| \nabla u_h^0 \right\|_0^2 + \sigma h \left\| \nabla u_h^0 \right\|_0 + C_0^2 \lambda^2 \bar{N} \left\| \nabla u_h^0 \right\|_0 \left\| \nabla T_0 \right\|_0 + C_0^2 \lambda^2 \bar{N} \left\| \nabla u_h^0 \right\|_0 \left\| \nabla T_h^0 \right\|_0 \\
& \quad + C_0^2 \lambda \left\| \nabla T_h^0 \right\|_0 + C_0^2 \lambda \left\| \nabla T_0 \right\|_0 + C_0 \lambda \left\| T_0 \right\|_0.
\end{aligned}$$

Using (2.16), (4.26), and Theorem 4.2, we can obtain

$$\begin{aligned}
\frac{7}{10} (\nu + \sigma h) \left\| \nabla u_h^1 \right\|_0 & \leq \frac{25 N C_0^4 \lambda^2 \varepsilon^2}{9 (\nu + \sigma h)^2} + \sigma h \frac{5 C_0^2 \lambda \varepsilon}{3 (\nu + \sigma h)} + \frac{10 \bar{N} C_0^4 \lambda^3 \varepsilon^2}{3 (\nu + \sigma h)} + \frac{3 C_0^2 \lambda^2 \varepsilon^2}{2} \\
& \leq \frac{103 C_0^2 \lambda \varepsilon}{48} + \sigma h \frac{5 C_0^2 \lambda \varepsilon}{3 (\nu + \sigma h)}. \tag{4.34}
\end{aligned}$$

Namely,

$$\left\| \nabla u_h^1 \right\|_0 \leq \frac{103 C_0^2 \lambda \varepsilon / 48 + \sigma h (5 C_0^2 \lambda \varepsilon / 3 \nu)}{(7/10) (\nu + \sigma h)} \doteq \delta. \tag{4.35}$$

Using (2.16), (4.31), and (4.35), we can get

$$\begin{aligned}
\left\| \nabla \omega_h^1 \right\|_0 & \leq \frac{10 \bar{N} C_0^2 \lambda^2 \varepsilon^2}{3 (\nu + \sigma h)} + \lambda \bar{N} \delta \varepsilon + \sigma h \varepsilon + \frac{\varepsilon}{4} \\
& \leq \frac{\varepsilon}{3} + \frac{\varepsilon}{4} + \lambda \bar{N} \delta \varepsilon + \sigma h \varepsilon \\
& = \frac{7 \varepsilon}{12} + \lambda \bar{N} \delta \varepsilon + \sigma h \varepsilon. \tag{4.36}
\end{aligned}$$

Using the triangle inequality, we can get

$$\left\| \nabla T_h^1 \right\|_0 \leq \left\| \nabla \omega_h^1 \right\|_0 + \left\| \nabla T_0 \right\|_0 \leq \frac{5 \varepsilon}{6} + \lambda \bar{N} \delta \varepsilon + \sigma h \varepsilon. \tag{4.37}$$

Therefore, we finish the proof. \square

5. Error Analysis

In this section, we establish the H^1 and L^2 -bounds of the error $u - u_h^i$, $T - T_h^i$, $i = 0, 1$ and L^2 -bound of the error $p - p_h^i$, $i = 0, 1$. In order to obtain the error estimates, we define the Galerkin projection $(R_h, Q_h) = (R_h(u, p), Q_h(u, p)) : (X, M) \rightarrow (X_h, M_h)$, such that

$$a(R_h - u, v_h) - d(Q_h - p, v_h) + d(q_h, R_h - u) = 0, \quad \forall (u, p) \in (X, M), (v_h, q_h) \in (X_h, M_h). \quad (5.1)$$

Lemma 5.1 (see [37, 38]). *The Galerkin projection (R_h, Q_h) satisfies*

$$\|R_h - u\|_0 + h(\|\nabla(R_h - u)\|_0 + \|Q_h - p\|_0) \leq Ch^{r+1}(\nu\|u\|_{r+1} + \|p\|_r), \quad r = 1, 2. \quad (5.2)$$

Lemma 5.2 (see [7]). *There exists $\tilde{r}_h : W \rightarrow W_h$ for all $\psi \in W$ holds that*

$$(\nabla(\psi - \tilde{r}_h\psi), \nabla\varphi_h) = 0, \quad \forall \varphi_h \in W_h, \quad (5.3)$$

$$\int_{\Omega} (\psi - \tilde{r}_h\psi) dx = 0, \quad \|\nabla\tilde{r}_h\psi\|_0 \leq \|\nabla\psi\|_0. \quad (5.4)$$

When $\psi \in W^{k,q}(\Omega)$ ($1 \leq q \leq \infty$), there holds

$$\|\psi - \tilde{r}_h\psi\|_{-s,q} \leq Ch^{k+s} |\psi|_{k,q}, \quad -1 \leq s \leq m, \quad 0 \leq k \leq r+1. \quad (5.5)$$

There exists $\bar{r}_h : W_0 \rightarrow W_{0h}$ for all $\psi \in W_0$ holds that

$$(\nabla(\psi - \bar{r}_h\psi), \nabla\varphi_h) = 0, \quad \forall \varphi_h \in W_{0h}, \quad \|\nabla\bar{r}_h\psi\|_0 \leq \|\nabla\psi\|_0. \quad (5.6)$$

When $\psi \in W^{r,q}(\Omega)$ ($1 \leq q \leq \infty$), there holds

$$\|\psi - \bar{r}_h\psi\|_{-s,q} \leq Ch^{k+s} |\psi|_{r,q}, \quad -1 \leq s \leq r, \quad 0 \leq k \leq r+1. \quad (5.7)$$

Lemma 5.3 (see [7]). *If $(A_1) \sim (A_3)$ hold and $(u, p, T) \in H^{r+1}(\Omega) \times H^r(\Omega) \times H^{r+1}(\Omega)$ and (u_h, P_h, T_h) are the solution of problem (\mathcal{P}') and (\mathcal{P}'') , respectively, then there holds that*

$$\|\nabla(u - u_h)\|_0 + \|p - p_h\|_0 + \|\nabla(T - T_h)\|_0 \leq Ch^r (\|u\|_{r+1} + \|p\|_r + \|T\|_{r+1}). \quad (5.8)$$

Lemma 5.4. Under the assumptions of Lemma 2.3, (u_h, p_h, T_h) is the solution of (3.1), (u_h^0, p_h^0, T_h^0) defined by (3.4), then there hold

$$\begin{aligned} \left\| \nabla(u_h - u_h^0) \right\|_0 &\leq \frac{50\sigma C_0^2 \lambda h}{21\nu} + \frac{10C_0^2 \lambda \sigma \varepsilon h}{7\nu}, \\ \left\| \nabla(T_h - T_h^0) \right\|_0 &\leq 2\sigma h \varepsilon + \frac{\sigma h \varepsilon}{14\nu}, \\ \beta \left\| p_h - p_h^0 \right\|_0 &\leq \frac{95\sigma h C_0^2 \lambda \varepsilon}{21\nu} + \frac{19\sigma h C_0^2 \lambda \varepsilon}{7} + \sigma h C_0^2 \lambda + \frac{\sigma h C_0^2 \lambda}{14\nu}. \end{aligned} \quad (5.9)$$

Proof. Subtracting (3.1) from (2.5) we get the error equations, namely $(u_h - u_h^0, p_h - p_h^0, T_h - T_h^0)$ satisfy

$$\begin{aligned} \nu a(u_h - u_h^0, v_h) - \sigma h a(u_h^0, v_h) + d(\varphi_h, u_h - u_h^0) - d(p_h - p_h^0, v_h) + b(u_h^0, u_h - u_h^0, v_h) \\ + b(u_h - u_h^0, u_h, v_h) = \lambda(j(T_h - T_h^0), v_h), \quad \forall v_h \in X_h, \varphi_h \in M_h, \\ \bar{a}(T_h - T_h^0, \varphi_h) - \sigma h \bar{a}(T_h^0, \varphi_h) + \lambda \bar{b}(u_h - u_h^0, T_h, \varphi_h) + \bar{b}(u_h^0, T_h - T_h^0, \varphi_h) = 0, \quad \forall \varphi_h \in W_{0h}. \end{aligned} \quad (5.10)$$

Letting $v_h = u_h - u_h^0, \varphi_h = p_h - p_h^0$ in the first equation of (5.10) and using (2.11), (2.8), and (A₁), we can get

$$\nu \left\| \nabla(u_h - u_h^0) \right\|_0 \leq \sigma h \left\| \nabla u_h^0 \right\|_0 + N \left\| \nabla(u_h - u_h^0) \right\|_0 \|\nabla u_h\|_0 + C_0^2 \lambda \left\| \nabla(T_h - T_h^0) \right\|_0. \quad (5.11)$$

Hence, we deduce

$$(\nu - N \|\nabla u_h\|_0) \left\| \nabla(u_h - u_h^0) \right\|_0 \leq \sigma h \left\| \nabla u_h^0 \right\|_0 + C_0^2 \lambda \left\| \nabla(T_h - T_h^0) \right\|_0. \quad (5.12)$$

Letting $\varphi_h = T_h - T_h^0$ in the second equation of (5.10) and using (2.9), we obtain

$$\bar{a}(T_h - T_h^0, T_h - T_h^0) + \sigma h \bar{a}(T_h^0, T_h - T_h^0) + \lambda \bar{b}(u_h - u_h^0, T_h, T_h - T_h^0) = 0. \quad (5.13)$$

Using (2.11), we can get

$$\left\| \nabla(T_h - T_h^0) \right\|_0 \leq \sigma h \left\| \nabla T_h^0 \right\|_0 + \lambda \bar{N} \left\| \nabla(u_h - u_h^0) \right\|_0 \|\nabla T_h\|_0. \quad (5.14)$$

By (5.12), we deduce

$$(\nu - N \|\nabla u_h\|_0) \left\| \nabla(u_h - u_h^0) \right\|_0 \leq \sigma h \left\| \nabla u_h^0 \right\|_0 + C_0^2 \lambda \sigma h \left\| \nabla T_h^0 \right\|_0 + C_0^2 \lambda^2 \bar{N} \left\| \nabla(u_h - u_h^0) \right\|_0 \|\nabla T_h\|_0. \quad (5.15)$$

Using (4.1), we can obtain

$$\begin{aligned} \left(\nu - N\|\nabla u_h\|_0 - \frac{3\nu}{32} \right) \|\nabla(u_h - u_h^0)\|_0 &\leq \sigma h \|\nabla u_h^0\|_0 + C_0^2 \lambda \sigma h \|\nabla T_h^0\|_0 \\ &\leq \frac{5\sigma h C_0^2 \lambda \varepsilon}{3(\nu + \sigma h)} + C_0^2 \lambda \sigma h \varepsilon. \end{aligned} \quad (5.16)$$

By using (2.16) and (2.17), there holds

$$\nu - N\|\nabla u_h\|_0 - \frac{3\nu}{32} \geq \frac{7\nu}{10}. \quad (5.17)$$

Therefore, we can deduce

$$\|\nabla(u_h - u_h^0)\|_0 \leq \frac{50\sigma h C_0^2 \lambda \varepsilon}{21\nu(\nu + \sigma h)} + \frac{10C_0^2 \lambda \sigma h \varepsilon}{7\nu}. \quad (5.18)$$

By (5.14) and (5.18), we can have

$$\begin{aligned} \|\nabla(T_h - T_h^0)\|_0 &\leq \sigma h \varepsilon + \lambda \bar{N} \varepsilon \left(\frac{50\sigma h C_0^2 \lambda \varepsilon}{21\nu(\nu + \sigma h)} + \frac{10C_0^2 \lambda \sigma h \varepsilon}{7\nu} \right) \\ &\leq 2\sigma h \varepsilon + \frac{\sigma h \varepsilon}{14\nu}. \end{aligned} \quad (5.19)$$

Letting $\varphi_h = 0$, $v_h = u_h - u_h^0$ in the first equation of (5.10) and using (2.3), we have

$$\begin{aligned} \beta \|p_h - p_h^0\|_0 &\leq \nu \|\nabla(u_h - u_h^0)\|_0 + \sigma h \|\nabla u_h^0\|_0 + N \|\nabla(u_h - u_h^0)\|_0 \|\nabla u_h\|_0 + C_0^2 \lambda \|\nabla(T_h - T_h^0)\|_0 \\ &\leq \frac{50\sigma h C_0^2 \lambda \varepsilon}{21(\nu + \sigma h)} + \frac{10\sigma h C_0^2 \lambda \varepsilon}{7} + \frac{5\sigma h C_0^2 \lambda \varepsilon}{3\nu} + \frac{10\sigma h C_0^2 \lambda \varepsilon}{21\nu} + \frac{2\sigma h C_0^2 \lambda \varepsilon}{7} \\ &\quad + \sigma h C_0^2 \lambda \varepsilon + \sigma h C_0^2 \lambda + \frac{\sigma h C_0^2 \lambda}{14\nu} \\ &\leq \frac{95\sigma h C_0^2 \lambda \varepsilon}{21(\nu + \sigma h)} + \frac{19\sigma h C_0^2 \lambda \varepsilon}{7} + \sigma h C_0^2 \lambda + \frac{\sigma h C_0^2 \lambda}{14\nu}. \end{aligned} \quad (5.20)$$

Hence, we finish the proof. \square

Theorem 5.5. *Under the assumptions of Lemmas 2.3 and 5.3, the following inequality*

$$\left\| \nabla(u - u_h^0) \right\|_0 + \left\| p - p_h^0 \right\|_0 + \left\| \nabla(T - T_h^0) \right\|_0 \leq Ch^r (\|u\|_{r+1} + \|p\|_r + \|T\|_{r+1}) + Ch, \quad (5.21)$$

holds, where C is a positive constant numbers.

Proof. By Lemmas 5.3, 5.4, and the triangle inequality this theorem is obviously true. \square

Lemma 5.6. *For all $u \in H^2(\Omega) \cap X$, $\omega \in W_0$, $\varphi \in H^2(\Omega) \cap W_0$, there hold that*

$$\left| \bar{b}(u - R_h, \omega, \varphi) \right| \leq C \|u - R_h\|_0 \|\mathcal{A}\omega\|_0 \|\nabla\varphi\|_0, \quad (5.22)$$

$$\left| \bar{b}(u, T - \tilde{r}_h T, \varphi) \right| \leq C \|\mathcal{A}u\|_0 \|T - \tilde{r}_h T\|_0 \|\nabla\varphi\|_0. \quad (5.23)$$

Proof. Letting $\bar{\omega} = (\omega, 0)^T$, we have

$$\bar{b}(u - R_h, \omega, \varphi) = b(u - R_h, \bar{\omega}, \varphi). \quad (5.24)$$

Using (2.25), we can deduce (5.22). Because $T - \tilde{r}_h T \in W_0$, (5.23) holds. \square

Theorem 5.7. *Under the assumptions of Lemmas 2.3 and 5.3, the following inequality:*

$$\left\| u - u_h^0 \right\|_0 + \left\| T - T_h^0 \right\|_0 \leq Ch^{r+1} (\|u\|_{r+1} + \|p\|_r + \|T\|_{r+1}) + Ch, \quad (5.25)$$

holds, where C is a positive constant.

Proof. Subtracting (3.1) from (2.4) we get the error equations, namely,

$$\begin{aligned} & \nu a(u - u_h^0, v_h) - \sigma h a(u_h^0, v_h) + b(u - u_h^0, u_h^0, v_h) + b(u, u - u_h^0, v_h) - d(p - p_h^0, v_h) \\ & \quad + d(\varphi_h, u - u_h^0) = \lambda(j(T - T_h^0), v_h), \quad \forall v_h \in X_h, \varphi_h \in M_h, \\ & \bar{a}(T - T_h^0, \varphi_h) - \sigma h \bar{a}(T_h^0, \varphi_h) + \lambda \bar{b}(u - u_h^0, T, \varphi_h) + \lambda \bar{b}(u_h^0, T - T_h^0, \varphi_h) = 0, \quad \forall \varphi_h \in W_{0h}. \end{aligned} \quad (5.26)$$

Letting $e_h^0 = R_h - u_h^0$, $\eta_h^0 = Q_h - p_h^0$, $\xi_h^0 = \tilde{r}_h T - T_h^0$ and using (5.1) and (5.3), we can get

$$\begin{aligned} & \nu a(e_h^0, v_h) - \sigma h a(u_h^0, v_h) + b(u - u_h^0, u_h^0, v_h) + b(u, u - u_h^0, v_h) - d(\eta_h^0, v_h) + d(\varphi_h, e_h^0) \\ & \quad = \lambda(j(T - T_h^0), v_h), \quad \forall v_h \in X_h, \varphi_h \in M_h, \\ & \bar{a}(\xi_h, \varphi_h) - \sigma h \bar{a}(T_h^0, \varphi_h) + \lambda \bar{b}(u - u_h^0, T, \varphi_h) + \lambda \bar{b}(u_h^0, T - T_h^0, \varphi_h) = 0, \quad \forall \varphi_h \in W_{0h}. \end{aligned} \quad (5.27)$$

Taking $v_h = e_h^0$, $\varphi_h = \eta_h^0$ in the first equation of (5.27), we obtain

$$\begin{aligned} & \nu a(e_h^0, e_h^0) - \sigma h a(u_h^0, e_h^0) + b(e_h^0, u_h^0, e_h^0) + b(u - R_h, u_h^0, e_h^0) + b(u, u - R_h, e_h^0) \\ & = \lambda(j(T - T_h^0), e_h^0), \quad \forall v_h \in X_h, \varphi_h \in M_h. \end{aligned} \quad (5.28)$$

Using (2.10) and (A₁), we deduce

$$\begin{aligned} (\nu - N \|\nabla u_h^0\|_0) \|\nabla e_h^0\|_0^2 & \leq |b(u - R_h, u, e_h^0)| + |b(u_h^0, u - R_h, e_h^0)| \\ & \quad + |\lambda(j(T - T_h^0), e_h^0)| + |\sigma h a(u_h^0, e_h^0)| \\ & \leq N(\|\nabla u\|_0 + \|\nabla u_h^0\|_0) \|\nabla(u - R_h)\|_0 \|\nabla e_h^0\|_0 \\ & \quad + C_0^2 \lambda \|\nabla(T - T_h^0)\|_0 \|\nabla e_h^0\|_0 + \sigma h \|\nabla u_h^0\|_0 \|\nabla e_h^0\|_0. \end{aligned} \quad (5.29)$$

Using Theorem 2.1, (2.16), (4.1), and (5.2), we can obtain

$$\|\nabla e_h^0\|_0 \leq Ch^{r+1}(\|u\|_{r+1} + \|p\|_r + \|T\|_{r+1}) + Ch. \quad (5.30)$$

Taking $\psi_h = \xi_h^0$ in the second equation of (5.27) and using (2.9) we have

$$\bar{a}(\xi_h^0, \xi_h^0) - \sigma h \bar{a}(T_h^0, \xi_h^0) + \lambda \bar{b}(u_h^0, T - \tilde{r}_h T, \xi_h^0) + \lambda \bar{b}(u - R_h, T, \xi_h^0) + \lambda \bar{b}(e_h^0, T, \xi_h^0) = 0. \quad (5.31)$$

By (2.9), we have

$$\begin{aligned} & \bar{b}(u_h^0, T - \tilde{r}_h T, \xi_h^0) + \bar{b}(u - R_h, T, \xi_h^0) + \bar{b}(e_h^0, T, \xi_h^0) \\ & = -\bar{b}(e_h^0, T - \tilde{r}_h T, \xi_h^0) - \bar{b}(u - R_h, T - \tilde{r}_h T, \xi_h^0) + \bar{b}(u, T - \tilde{r}_h T, \xi_h^0) \\ & \quad + \bar{b}(u - R_h, T, \xi_h^0) + \bar{b}(e_h^0, T - \tilde{r}_h T, \xi_h^0) + \bar{b}(e_h^0, \xi_h^0, \xi_h^0) + \bar{b}(e_h^0, T_h^0, \xi_h^0). \end{aligned} \quad (5.32)$$

Letting $T = \omega + T_0$, $\omega \in W_0$ and using Lemma 5.6, we can get

$$\begin{aligned}
& \left| \bar{b}(u_h^0, T - \tilde{r}_h T, \xi_h^0) + \bar{b}(u - R_h, T, \xi_h^0) + \bar{b}(e_h, T, \xi_h^0) \right| \\
& \leq \bar{N} \left\| \nabla e_h^0 \right\|_0 \left\| \nabla(T - \tilde{r}_h T) \right\|_0 \left\| \nabla \xi_h^0 \right\|_0 + \bar{N} \left\| \nabla(u - R_h) \right\|_0 \left\| \nabla(T - \tilde{r}_h T) \right\|_0 \left\| \nabla \xi_h^0 \right\|_0 \\
& \quad + \bar{N} \left\| \nabla e_h^0 \right\|_0 \left\| \nabla(T - \tilde{r}_h T) \right\|_0 \left\| \nabla \xi_h^0 \right\|_0 + \bar{N} \left\| \nabla e_h^0 \right\|_0 \left\| \nabla T_h^0 \right\|_0 \left\| \nabla \xi_h^0 \right\|_0 \\
& \quad + C \|\mathcal{A}u\|_0 \|T - \tilde{r}_h T\|_0 \left\| \nabla \xi_h^0 \right\|_0 + C \|u - R_h\|_0 \|\mathcal{A}\omega\|_0 \left\| \nabla \xi_h^0 \right\|_0 \\
& \quad + C \left\| \nabla(u - R_h) \right\|_0 \left\| \nabla T_0 \right\|_0 \left\| \nabla \xi_h^0 \right\|_0.
\end{aligned} \tag{5.33}$$

By assumption (A₂), letting $\varepsilon < h$ and using Lemma 5.1 and (2.16), (4.26), and (5.33), we can deduce

$$\left\| \nabla \xi_h^0 \right\|_0 \leq Ch^{r+1} (\|u\|_{r+1} + \|p\|_r + \|T\|_{r+1}) + Ch. \tag{5.34}$$

Hence, we have

$$\begin{aligned}
\left\| T - T_h^0 \right\|_0 & \leq \|T - \tilde{r}_h T\|_0 + \left\| \xi_h^0 \right\|_0 \\
& \leq \|T - \tilde{r}_h T\|_0 + C_0 \left\| \nabla \xi_h^0 \right\|_0 \\
& \leq Ch^{r+1} (\|u\|_{r+1} + \|p\|_r + \|T\|_{r+1}) + Ch.
\end{aligned} \tag{5.35}$$

By (2.10) and (2.25), we can deduce

$$\begin{aligned}
& \left| b(u - R_h, u, e_h^0) \right| + \left| b(u_h^0, u - R_h, e_h^0) \right| \\
& \leq \left| b(u - R_h, u, e_h^0) \right| + \left| b(u, u - R_h, e_h^0) \right| \\
& \quad + \left| b(u - R_h, u - R_h, e_h^0) \right| + \left| b(e_h, u - R_h, e_h^0) \right| \\
& \leq C \|\mathcal{A}u\|_0 \|u - R_h\|_0 \left\| \nabla e_h^0 \right\|_0 \\
& \quad + N \left(\left\| \nabla(u - R_h) \right\|_0 + \left\| \nabla e_h^0 \right\|_0 \right) \left\| \nabla(u - R_h) \right\|_0 \left\| \nabla e_h^0 \right\|_0 \\
& \leq Ch^2 \left\| \nabla e_h^0 \right\|_0.
\end{aligned} \tag{5.36}$$

Using (5.29), we can obtain

$$\begin{aligned} \left(\nu - N \left\| \nabla u_h^0 \right\|_0 \right) \left\| \nabla e_h^0 \right\|_0^2 &\leq Ch^{r+1} (\|u\|_{r+1} + \|p\|_r + \|T\|_{r+1}) \left\| \nabla e_h^0 \right\|_0 \\ &+ C_0 \lambda \left\| T - T_h^0 \right\|_0 \left\| \nabla e_h^0 \right\|_0 + \sigma h \left\| \nabla u_h^0 \right\|_0 \left\| \nabla e_h^0 \right\|_0. \end{aligned} \quad (5.37)$$

By using (2.16) and (4.1), there holds

$$\nu - N \left\| \nabla u_h^0 \right\|_0 \geq \frac{4\nu}{5}. \quad (5.38)$$

Hence, we can deduce from (5.37)

$$\left\| \nabla e_h^0 \right\|_0 \leq Ch^{r+1} (\|u\|_{r+1} + \|p\|_r + \|T\|_{r+1}) + Ch. \quad (5.39)$$

Therefore, we can deduce

$$\begin{aligned} \|u - u_h^0\|_0 &\leq \|u - R_h\|_0 + \left\| e_h^0 \right\|_0 \\ &\leq \|u - R_h\|_0 + C_0 \left\| \nabla e_h^0 \right\|_0 \\ &\leq Ch^{r+1} (\|u\|_{r+1} + \|p\|_r + \|T\|_{r+1}) + Ch. \end{aligned} \quad (5.40)$$

□

Theorem 5.8. *Under the assumptions of Lemmas 2.3 and 5.3, then there holds*

$$\begin{aligned} \left\| \nabla(u - u_h^1) \right\|_0 + \left\| \nabla(T - T_h^1) \right\|_0 &\leq Ch^r (\|u\|_{r+1} + \|p\|_r + \|T\|_{r+1}) + Ch^2, \\ \left\| u - u_h^1 \right\|_0 + \left\| T - T_h^1 \right\|_0 + h \left\| p - p_h^1 \right\|_0 &\leq Ch^{r+1} (\|u\|_{r+1} + \|p\|_r + \|T\|_{r+1}) + Ch^2, \end{aligned} \quad (5.41)$$

where C is a positive constant.

Proof. Subtracting (3.4) from (2.4) we get the error equations, namely,

$$\begin{aligned} \nu a(u - u_h^1, v_h) - \sigma h a(u_h^1, v_h) + b(u, u, v_h) - b(u_h^1, u_h^0, v_h) \\ - b(u_h^0, u_h^1, v_h) - d(p - p_h^1, v_h) + d(\varphi_h, u - u_h^1) \\ = \lambda \left(j(T - T_h^1), v_h \right) - b(u_h^0, u_h^0, v_h) - \sigma h a(u_h^0, v_h), \quad \forall v_h \in X_h, \varphi_h \in M_h, \\ \bar{a}(T - T_h^1, \varphi_h) - \sigma h \bar{a}(T_h^1, \varphi_h) + \lambda \bar{b}(u, T, \varphi_h) - \lambda \bar{b}(u_h^1, T_h^0, \varphi_h) - \lambda \bar{b}(u_h^0, T_h^1, \varphi_h) \\ = -\sigma h \bar{a}(T_h^0, \varphi_h) - \lambda \bar{b}(u_h^0, T_h^0, \varphi_h), \quad \forall \varphi_h \in W_{0h}. \end{aligned} \quad (5.42)$$

Letting $e_h^1 = R_h - u_h^1$, $\eta_h^1 = Q_h - p_h^1$, $\xi_h^1 = \tilde{r}_h T - T_h^1$, using (5.1) and (5.3) and adding and subtracting appropriate terms in the above expression yields

$$\begin{aligned}
& (\nu + \sigma h)a(e_h^1, v_h) + b(u_h^0, u - u_h^1, v_h) + b(u - u_h^1, u_h^0, v_h) - d(\eta_h^1, v_h) + d(\varphi_h, e_h^1) \\
& = \lambda(j(T - T_h^0), v_h) + \sigma ha(R_h - u_h^0, v_h) - b(u - u_h^0, u - u_h^0, v_h), \quad \forall v_h \in X_h, \varphi_h \in M_h, \\
& (1 + \sigma h)\bar{a}(\xi_h^1, \varphi_h) + \lambda\bar{b}(u_h^0, T - T_h^1, \varphi_h) - \lambda\bar{b}(u - u_h^1, T_h^0, \varphi_h) \\
& = \sigma h\bar{a}(\tilde{r}_h T - T_h^0, \varphi_h) - \lambda\bar{b}(u - u_h^0, T - T_h^0, \varphi_h), \quad \forall \varphi_h \in W_{0h}.
\end{aligned} \tag{5.43}$$

Letting $v_h = e_h^1$, $\varphi_h = \eta_h^1$ in the first equation of (5.43), we can deduce

$$\begin{aligned}
& (\nu + \sigma h)a(e_h^1, e_h^1) + b(u_h^0, u - R_h, e_h^1) + b(u - R_h, u_h^0, e_h^1) + b(e_h^1, u_h^0, e_h^1) \\
& = \lambda(j(T - T_h^0), e_h^1) + \sigma ha(R_h - u_h^0, e_h^1) - b(u - u_h^0, u - u_h^0, e_h^1)
\end{aligned} \tag{5.44}$$

By (2.10) and (2.25), we can deduce

$$\begin{aligned}
& (\nu + \sigma h - N\|\nabla u_h^0\|_0)\|\nabla e_h^1\|_0 \leq C\|\mathcal{A}u_h^0\|_0\|u - R_h\|_0 + \sigma h\|\nabla(R_h - u_h^0)\|_0 \\
& \quad + N\|\nabla(u - u_h^0)\|_0^2 + \lambda C_0\|T - T_h^0\|_0.
\end{aligned} \tag{5.45}$$

Using (2.16), (4.1), (4.26), (5.21), and (5.2), we can obtain

$$\|\nabla e_h^1\|_0 \leq Ch^{r+1}(\|u\|_{r+1} + \|p\|_r + \|T\|_{r+1}) + Ch^2. \tag{5.46}$$

Using (5.2) and triangle inequality, we can have

$$\begin{aligned}
\|\nabla(u - u_h^1)\|_0 & \leq \|\nabla(u - R_h)\|_0 + \|\nabla e_1\|_0 \\
& \leq Ch^r(\|u\|_{r+1} + \|p\|_r + \|T\|_{r+1}) + Ch^2, \\
\|u - u_h^1\|_0 & \leq \|u - R_h\|_0 + \|e_1\|_0 \\
& \leq \|u - R_h\|_0 + C_0\|\nabla e_1\|_0 \\
& \leq Ch^{r+1}(\|u\|_{r+1} + \|p\|_r + \|T\|_{r+1}) + Ch^2.
\end{aligned} \tag{5.47}$$

Letting $\psi_h = \xi_h^1$ in the second equation of (5.43) and using (2.8), we can deduce

$$\begin{aligned} & (1 + \sigma h)\bar{a}(\xi_h^1, \xi_h^1) + \lambda\bar{b}(u_h^0, T - \tilde{r}_h T, \xi_h^1) - \lambda\bar{b}(u - u_h^1, T_h^0, \xi_h^1) \\ & = \sigma h\bar{a}(T - T_h^0, \xi_h^1) - \lambda\bar{b}(u - u_h^0, T - T_h^0, \xi_h^1). \end{aligned} \quad (5.48)$$

Letting $T_h^0 = \omega_h^0 + T_0$ and using (2.11), (5.22), and (5.23), we have

$$\begin{aligned} (1 + \sigma h)\|\nabla \xi_h^1\|_0 & \leq C\lambda\|\mathcal{A}u_h^0\|_0\|T - \tilde{r}_h T\|_0 + C\lambda\|u - u_h^1\|_0\|\mathcal{A}\omega_h^0\|_0 \\ & \quad + \bar{N}\lambda\|\nabla(u - u_h^1)\|_0\|\nabla T_0\|_0 + \sigma h\|\nabla(T - T_h^0)\|_0 \\ & \quad + \lambda\bar{N}\|\nabla(u - u_h^0)\|_0\|\nabla(T - T_h^0)\|_0. \end{aligned} \quad (5.49)$$

Using (5.5), (5.21), (5.47), we can obtain

$$\|\nabla \xi_h^1\|_0 \leq Ch^{r+1}(\|u\|_{r+1} + \|p\|_r + \|T\|_{r+1}) + Ch^2. \quad (5.50)$$

Using (5.2) and triangle inequality, we can have

$$\begin{aligned} \|\nabla(T - T_h^1)\|_0 & \leq \|\nabla(T - \tilde{r}_h T)\|_0 + \|\nabla \xi_h^1\|_0 \\ & \leq Ch^r(\|u\|_{r+1} + \|p\|_r + \|T\|_{r+1}) + Ch^2, \\ \|T - T_h^1\|_0 & \leq \|T - \tilde{r}_h T\|_0 + \|\xi_h^1\|_0 \leq \|T - \tilde{r}_h T\|_0 + C_0\|\nabla \xi_h^1\|_0 \\ & \leq Ch^{r+1}(\|u\|_{r+1} + \|p\|_r + \|T\|_{r+1}) + Ch^2. \end{aligned} \quad (5.51)$$

Taking $\varphi_h = 0$, $v_h = R_h - u_h^1$ in the first equation of (5.43) and using (2.3), we have

$$\begin{aligned} \beta\|\eta_h^1\|_0 & \leq (v + \sigma h)\|\nabla e_h^1\|_0 + \sigma h\|\nabla(u - u_h^0)\|_0 + 2N\|\nabla u_h^0\|_0\|\nabla(u - u_h^1)\|_0 \\ & \quad + C_0\lambda\|T - T_h^0\|_0 + N\|\nabla(u - u_h^0)\|_0^2. \end{aligned} \quad (5.52)$$

By (4.1), (5.21), and (5.47), we can deduce

$$\|\eta_h^1\|_0 \leq Ch^r(\|u\|_{r+1} + \|p\|_r + \|T\|_{r+1}) + Ch^2. \quad (5.53)$$




Figure 1: Physics model of the cavity flows.




Figure 2: The numerical Isotherms (a) and the numerical Isobar (b) for $\nu = 1/2000$ by the defect-correction MFEM with $h = \sqrt{2}/40$, $\sigma = 0.4$.

Using (5.2) and triangle inequality, we can have

$$\|p - p_h^1\|_0 \leq \|p - Q_h\|_0 + \|\eta_h^1\|_0 \leq Ch^r (\|u\|_{r+1} + \|p\|_r + \|T\|_{r+1}) + Ch^2. \quad (5.54)$$

□

6. Numerical Experiments

In this section, we present some numerical examples with a physical model of square cavity stationary flow. We choose different ν for comparison. The side length of the square cavity and the boundary conditions are given by Figure 1. From Figure 1, we can see that the $T = 0$




Figure 3: The numerical streamline for $\nu = 1/2000$ by the defect-correction MFEM with $h = \sqrt{2}/40$, $\sigma = 0.4$.




Figure 4: The numerical Isotherms (a) and the numerical Isobar (b) for $\nu = 1/5000$ by the defect-correction MFEM with $h = \sqrt{2}/100$, $\sigma = 0.4$.

on left and lower boundaries, $\partial T/\partial n = 0$ on upper boundary, and $T = 4y(1 - y)$ on right boundary of the cavity. We use $P_2 - P_1 - P_2$ finite element here.

Firstly, we choose $\nu = 1/2000$, $\sigma = 0.4$ and divide the cavity into $M \times N = 40 \times 40$, that is, $h = \sqrt{2}/40$. Figure 2 gives the numerical isotherms (a) and the numerical isobar (b). Figure 3 gives the numerical streamline. From the numerical results, we can see that our method is stable and has a good precision.

Secondly, we choose $\nu = 1/5000$, $\sigma = 0.4$ to show our our method suiting for solving the conduction convection problems with small viscosity. It is well known that it is more and more difficult to solve the problem by numerical method as ν changing smaller and smaller.




Figure 5: The numerical streamline for $\nu = 1/5000$ by the defect-correction MFEM with $h = \sqrt{2}/100$, $\sigma = 0.4$.




Figure 6: The numerical Isotherms (a) and the numerical Isobar (b) for $\nu = 1/6000$ by the defect-correction MFEM with $h = \sqrt{2}/100$, $\sigma = 0.4$.

Hence, we divide the cavity into $M \times N = 100 \times 100$, namely $h = \sqrt{2}/100$. Figure 4 gives the numerical isotherms (a) and the numerical isobar (b), and Figure 5 shows the numerical streamline. At last, we choose $\nu = 1/6000$, $\sigma = 0.4$. Figure 6 gives the numerical isotherms (a) and the numerical isobar (b), and Figure 7 shows the numerical streamline.

Just as Remark 3.1, we only use one correction step in our numerical experiments. From the numerical, we can see that when $\nu = 0.5 \times 10^{-3}$ the numerical streamline is very regular. The pressure is small near the wall. But the numerical streamline changes more and more immethodical with ν changing smaller and smaller. And the pressure changes bigger




Figure 7: The numerical streamline for $\nu = 1/6000$ by the defect-correction MFEM with $h = \sqrt{2}/100$, $\sigma = 0.4$.

near the wall. In conclusion, the defect-correction MFEM is highly efficient for the stationary conduction-convection problems and it can be used for solving the convection-conduction problems with much small viscosity.

Acknowledgments





The authors would like to thank the editor and the referees for their criticism, valuable comments, which led to the improvement of this paper. This work is supported by the NSF of China (no. 10971166) and the National High Technology Research and Development program of China (863 program, no. 2009AA01A135).

References

- [1] J. A. M. García, J. M. G. Cabeza, and A. C. Rodríguez, "Two-dimensional non-linear inverse heat conduction problem based on the singular value decomposition," *International Journal of Thermal Sciences*, vol. 48, no. 6, pp. 1081–1093, 2009.
- [2] Z. D. Luo and X. M. Lu, "A least-squares Galerkin/Petrov mixed finite element method for stationary conduction-convection problems," *Mathematica Numerica Sinica*, vol. 25, no. 2, pp. 231–244, 2003.
- [3] M. S. Mesquita and M. J. S. de Lemos, "Optimal multigrid solutions of two-dimensional convection-conduction problems," *Applied Mathematics and Computation*, vol. 152, no. 3, pp. 725–742, 2004.
- [4] C. P. Naveira, M. Lachi, R. M. Cotta, and J. Padet, "Hybrid formulation and solution for transient conjugated conduction-external convection," *International Journal of Heat and Mass Transfer*, vol. 52, no. 1-2, pp. 112–123, 2009.
- [5] Q. W. Wang, M. Yang, and W. Q. Tao, "Natural convection in a square enclosure with an internal isolated vertical plate," *Wärme- und Stoffübertragung*, vol. 29, no. 3, pp. 161–169, 1994.
- [6] M. Yang, W. Q. Tao, Q. W. Wang, and S. S. Lue, "On identical problems of natural convection in enclosures and applications of the identity character," *Journal of Thermal Science*, vol. 2, no. 2, pp. 116–125, 1993.
- [7] Z. D. Luo, *The Bases and Applications of Mixed Finite Element Methods*, Science Press, Beijing, China, 2006.

- [8] Z. Luo, J. Chen, I. M. Navon, and J. Zhu, "An optimizing reduced PLSMFE formulation for non-stationary conduction-convection problems," *International Journal for Numerical Methods in Fluids*, vol. 60, no. 4, pp. 409–436, 2009.
- [9] J. N. Reddy and D. K. Gartling, *The Finite Element Method Transfer and Fluid Dynamics*, CRC Press, Washington, DC, USA, 2nd edition, 2000.
- [10] L. Q. Tang and T. T. H. Tsang, "A least-squares finite element method for time-dependent incompressible flows with thermal convection," *International Journal for Numerical Methods in Fluids*, vol. 17, no. 4, pp. 271–289, 1993.
- [11] D. C. Kim and Y. D. Choi, "Analysis of conduction-natural convection conjugate heat transfer in the gap between concentric cylinders under solar irradiation," *International Journal of Thermal Sciences*, vol. 48, no. 6, pp. 1247–1258, 2009.
- [12] Z. Si, T. Zhang, and K. Wang, "A Newton iterative mixed finite element method for stationary conduction-convection problems," *International Journal of Computational Fluid Dynamics*, vol. 24, no. 3-4, pp. 135–141, 2010.
- [13] Z. Si and Y. He, "A coupled Newton iterative mixed fixed element method for stationary conduction-convection problems," *Computing*, vol. 89, no. 1-2, pp. 1–25, 2010.
- [14] O. Axelsson and M. Nikolova, "Adaptive refinement for convection-diffusion problems based on a defect-correction technique and finite difference method," *Computing*, vol. 58, no. 1, pp. 1–30, 1997.
- [15] V. Ervin and W. Layton, "An analysis of a defect-correction method for a model convection-diffusion equation," *SIAM Journal on Numerical Analysis*, vol. 26, no. 1, pp. 169–179, 1989.
- [16] V. J. Ervin, W. J. Layton, and J. M. Maubach, "Adaptive defect-correction methods for viscous incompressible flow problems," *SIAM Journal on Numerical Analysis*, vol. 37, no. 4, pp. 1165–1185, 2000.
- [17] V. J. Ervin, J. S. Howell, and H. Lee, "A two-parameter defect-correction method for computation of steady-state viscoelastic fluid flow," *Applied Mathematics and Computation*, vol. 196, no. 2, pp. 818–834, 2008.
- [18] R. Frank, J. Hertling, and J. P. Monnet, "The application of iterated defect correction to variational methods for elliptic boundary value problems," *Computing*, vol. 30, no. 2, pp. 121–135, 1983.
- [19] J. L. Gracia and E. O'Riordan, "A defect correction parameter-uniform numerical method for a singularly perturbed convection diffusion problem in one dimension," *Numerical Algorithms*, vol. 41, no. 4, pp. 359–385, 2006.
- [20] W. Heinrichs, "Defect correction for convection-dominated flow," *SIAM Journal on Scientific Computing*, vol. 17, no. 5, pp. 1082–1091, 1996.
- [21] W. Kramer, R. Minero, H. J. H. Clercx, and R. M. M. Mattheij, "A finite volume local defect correction method for solving the transport equation," *Computers and Fluids*, vol. 38, no. 3, pp. 533–543, 2009.
- [22] O. Koch and E. B. Weinmüller, "Iterated defect correction for the solution of singular initial value problems," *SIAM Journal on Numerical Analysis*, vol. 38, no. 6, pp. 1784–1799, 2001.
- [23] A. Labovschii, "A defect correction method for the time-dependent Navier-Stokes equations," *Numerical Methods for Partial Differential Equations*, vol. 25, no. 1, pp. 1–25, 2009.
- [24] W. Layton, H. K. Lee, and J. Peterson, "A defect-correction method for the incompressible Navier-Stokes equations," *Applied Mathematics and Computation*, vol. 129, no. 1, pp. 1–19, 2002.
- [25] R. Minero, M. J. H. Anthonissen, and R. M. M. Mattheij, "A local defect correction technique for time-dependent problems," *Numerical Methods for Partial Differential Equations*, vol. 22, no. 1, pp. 128–144, 2006.
- [26] R. Martin and H. Guillard, "A second order defect correction scheme for unsteady problems," *Computers and Fluids*, vol. 25, no. 1, pp. 9–27, 1996.
- [27] L. Shen and A. Zhou, "A defect correction scheme for finite element eigenvalues with applications to quantum chemistry," *SIAM Journal on Scientific Computing*, vol. 28, no. 1, pp. 321–338, 2006.
- [28] H. J. Stetter, "The defect correction principle and discretization methods," *Numerische Mathematik*, vol. 29, no. 4, pp. 425–443, 1978.
- [29] R. A. Adams, *Sobolev Spaces*, Pure and Applied Mathematics, vol. 6, Academic Press, New York, NY, USA, 1975.
- [30] Y. He and J. Li, "Convergence of three iterative methods based on the finite element discretization for the stationary Navier-Stokes equations," *Computer Methods in Applied Mechanics and Engineering*, vol. 198, no. 15-16, pp. 1351–1359, 2009.
- [31] R. Temam, *Navier-Stokes Equations: Theory and Numerical Analysis*, vol. 2 of *Studies in Mathematics and Its Applications*, North-Holland, Amsterdam, The Netherlands, 3rd edition, 1984.

- [32] Y. He, "Two-level method based on finite element and Crank-Nicolson extrapolation for the time-dependent Navier-Stokes equations," *SIAM Journal on Numerical Analysis*, vol. 41, no. 4, pp. 1263–1285, 2003.
- [33] Y. He and W. Sun, "Stability and convergence of the Crank-Nicolson/Adams-Bashforth scheme for the time-dependent Navier-Stokes equations," *SIAM Journal on Numerical Analysis*, vol. 45, no. 2, pp. 837–869, 2007.
- [34] A. T. Hill and E. Süli, "Approximation of the global attractor for the incompressible Navier-Stokes equations," *IMA Journal of Numerical Analysis*, vol. 20, no. 4, pp. 633–667, 2000.
- [35] V. Girault and P.-A. Raviart, *Finite Element Methods for Navier-Stokes Equations: Theory and Algorithms*, vol. 5 of *Springer Series in Computational Mathematics*, Springer, Berlin, Germany, 1986.
- [36] Y. He and K. Li, "Two-level stabilized finite element methods for the steady Navier-Stokes problem," *Computing*, vol. 74, no. 4, pp. 337–351, 2005.
- [37] Y. He and A. Wang, "A simplified two-level method for the steady Navier-Stokes equations," *Computer Methods in Applied Mechanics and Engineering*, vol. 197, no. 17-18, pp. 1568–1576, 2008.
- [38] J. Li and Z. Chen, "A new local stabilized nonconforming finite element method for the Stokes equations," *Computing*, vol. 82, no. 2-3, pp. 157–170, 2008.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

