

Journal of Inequalities in Pure and Applied Mathematics

NOTE ON SOME HADAMARD-TYPE INEQUALITIES

M. KLARIČIĆ BAKULA AND J. PEČARIĆ

Department of Mathematics
Faculty of Natural Sciences, Mathematics and Education
University of Split , Teslina 12
21 000 Split, Croatia.


EMail: milica@pmfst.hr

Faculty of Textile Technology
University of Zagreb
Pierottijeva 6, 10000 Zagreb
Croatia.

EMail: pecaric@hazu.hr

URL: <http://mahazu.hazu.hr/DepMPCS/indexJP.html>

©2000 Victoria University
ISSN (electronic): 1443-5756
018-04



volume 5, issue 3, article 74,
2004.

*Received 20 January, 2004;
accepted 05 April, 2004.*

Communicated by: [C. Giordano](#)

Abstract

Contents



Home Page

Go Back

Close

Quit

Abstract


Some Hadamard-type inequalities involving the product of two convex functions are obtained. Our results generalize the corresponding results of B.G.Pachpatte.

2000 Mathematics Subject Classification: 26D15, 26D20.

Key words: Integral inequalities, Hadamard's inequality

Contents

1	Introduction	3
2	Results	6
	References	



Note on Some Hadamard-Type Inequalities

M. Klaričić Bakula and J. Pečarić

Title Page

Contents



Go Back

Close

Quit

Page 2 of 20

1. Introduction

Let f be a convex function on $[a, b] \subset \mathbb{R}$. The following double inequality:

$$(1.1) \quad f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a) + f(b)}{2}$$

is known in the literature as Hadamard's inequality [1, p. 137], [2, p. 10] for convex functions.

Recently B.G.Pachpatte [3] considered some new integral inequalities, analogous to that of Hadamard, involving the product of two convex functions. In [3] the following theorem has been proved:

Theorem 1.1. *Let f and g be nonnegative, convex functions on $[a, b] \subset \mathbb{R}$. Then*


(i)

$$(1.2) \quad \frac{1}{b-a} \int_a^b f(x) g(x) dx \leq \frac{1}{3} M(a, b) + \frac{1}{6} N(a, b),$$

(ii)

$$(1.3) \quad 2f\left(\frac{a+b}{2}\right) g\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) g(x) dx + \frac{1}{6} M(a, b) + \frac{1}{3} N(a, b),$$

where $M(a, b) = f(a)g(a) + f(b)g(b)$ and $N(a, b) = f(a)g(b) + f(b)g(a)$. Inequalities (1.2) and (1.3) are sharp in the sense that equalities hold for some $f(x)$ and $g(x)$ on $[a, b]$.



Note on Some Hadamard-Type Inequalities

M. Klaričić Bakula and J. Pečarić

Title Page

Contents



Go Back

Close

Quit

Page 3 of 20

In the following Theorem 1.2 we give a variant of the corresponding Theorem 2 in [3].

Theorem 1.2. *Let f and g be nonnegative, convex functions on $[a, b] \subset \mathbb{R}$. Then*

(i)

$$(1.4) \quad \frac{3}{2(b-a)^2} \int_a^b \int_a^b \int_0^1 f(tx + (1-t)y) g(tx + (1-t)y) dt dx dy \\ \leq \frac{1}{b-a} \int_a^b f(x) g(x) dx + \frac{1}{8} [M(a, b) + N(a, b)];$$


(ii)

$$(1.5) \quad \frac{3}{b-a} \int_a^b \int_0^1 f\left(tx + (1-t)\left(\frac{a+b}{2}\right)\right) \\ \times g\left(tx + (1-t)\left(\frac{a+b}{2}\right)\right) dt dx \\ \leq \frac{1}{b-a} \int_a^b f(x) g(x) dx + \frac{1}{2} [M(a, b) + N(a, b)],$$

where $M(a, b)$ and $N(a, b)$ are as in Theorem 1.1.

It should be noted that in [3, Theorem 2] inequalities (3) and (4) are established. Inequality (3) from [3, Theorem 2] is a variant of our inequality (1.4) in which

$$\frac{1}{8} \left[\frac{M(a, b) + N(a, b)}{(b-a)^2} \right]$$



Note on Some Hadamard-Type Inequalities

M. Klaričić Bakula and J. Pečarić

Title Page

Contents



Go Back

Close

Quit

Page 4 of 20

stands in place of the term $\frac{1}{8} [M(a, b) + N(a, b)]$. Analogously, inequality (4) from [3, Theorem 2] is a variant of our inequality (1.5) in which


$$\frac{1}{4} \left(\frac{1 + b - a}{b - a} \right) [M(a, b) + N(a, b)]$$

stands in place of the term $\frac{1}{2} [M(a, b) + N(a, b)]$.

However, one can compare inequalities (3) and (4) with (1.4) and (1.5), respectively, to find out that estimates given by (1.4) and (1.5) are better (worse) than those given by (3) and (4) in [3, Theorem 2] in case of $b - a < 1$ ($b - a > 1$).

But on careful inspection of the proof in [3, Theorem 2], the reader can notice some errors in Pachpatte's calculation, so inequalities (3) and (4) in [3, Theorem 2] are in fact incorrect.

The aim of this paper is to prove some simple generalizations of Theorem 1.1 and Theorem 1.2, which additionally involve weight functions and also non-linear transformations of the base interval $[a, b]$. Those generalizations are established in Theorem 2.1 and Theorem 2.2. The above cited Theorem 1.1 is a special case of Theorem 2.1, while the above Theorem 1.2 is a special case of our Theorem 2.2 (see Remark 2.2).



Note on Some Hadamard-Type Inequalities

M. Klaričić Bakula and J. Pečarić

Title Page

Contents



Go Back

Close

Quit

Page 5 of 20

2. Results

Throughout the rest of the paper we shall use the following notation

$$[h; x, y] = \frac{h(y) - h(x)}{y - x}, \quad x \neq y$$

$$\tilde{h}(t) = th(\alpha + \beta - t),$$

$$\hat{h}(t) = th(t)$$

where $h : [\alpha, \beta] \rightarrow \mathbb{R}$ is a function, $[\alpha, \beta] \subset \mathbb{R}$, $x, y, t \in [\alpha, \beta]$. Note that from the above equalities we get

$$[\tilde{h}; \alpha, \beta] = \frac{\beta h(\alpha) - \alpha h(\beta)}{\beta - \alpha},$$


$$[\hat{h}; \alpha, \beta] = \frac{\beta h(\beta) - \alpha h(\alpha)}{\beta - \alpha},$$

and, by simple calculation,

$$(2.1) \quad [\hat{h}; \alpha, \beta] - [\tilde{h}; \alpha, \beta] = (\alpha + \beta) [h; \alpha, \beta].$$

The following results are valid:

Theorem 2.1. *Let f be a nonnegative convex function on $[m_1, M_1]$, g a nonnegative convex function on $[m_2, M_2]$, $u : [a, b] \rightarrow [m_1, M_1]$ and $v : [a, b] \rightarrow [m_2, M_2]$ continuous functions, and $p : [a, b] \rightarrow \mathbb{R}$ a positive integrable function. Then*



Note on Some Hadamard-Type Inequalities

M. Klaričić Bakula and J. Pečarić

Title Page

Contents



Go Back

Close

Quit


Page 6 of 20

(i)

$$(2.2) \quad \frac{1}{P} \int_a^b p(x) f(u(x)) g(v(x)) dx$$
$$\leq [f; m_1, M_1] [g; m_2, M_2] \frac{1}{P} \int_a^b p(x) u(x) v(x) dx$$
$$+ [f; m_1, M_1] [\tilde{g}; m_2, M_2] \frac{1}{P} \int_a^b p(x) u(x) dx$$
$$+ [\tilde{f}; m_1, M_1] [g; m_2, M_2] \frac{1}{P} \int_a^b p(x) v(x) dx$$
$$+ [\tilde{f}; m_1, M_1] [\tilde{g}; m_2, M_2].$$

(ii)

$$(2.3) \quad f\left(\frac{m_1 + M_1}{2}\right) g\left(\frac{m_2 + M_2}{2}\right)$$
$$\leq \frac{1}{4P} \left[\int_a^b p(x) f(u(x)) g(v(x)) dx \right.$$
$$+ \left. \int_a^b p(x) f(M_1 + m_1 - u(x)) g(M_2 + m_2 - v(x)) dx \right]$$
$$+ \frac{1}{4P} \left[-2 [f; m_1, M_1] [g; m_2, M_2] \int_a^b p(x) u(x) v(x) dx \right.$$
$$+ \left. ([\hat{g}; m_2, M_2] - [\tilde{g}; m_2, M_2]) [f; m_1, M_1] \int_a^b p(x) u(x) dx \right]$$



**Note on Some Hadamard-Type
Inequalities**

M. Klaričić Bakula and J. Pečarić

Title Page

Contents



Go Back

Close

Quit

Page 7 of 20

$$\begin{aligned}
 & + \left([\widehat{f}; m_1, M_1] - [\widetilde{f}; m_1, M_1] \right) [g; m_2, M_2] \int_a^b p(x) v(x) dx \\
 & + \frac{1}{4} \left([\widetilde{f}; m_1, M_1] [\widehat{g}; m_2, M_2] + [\widehat{f}; m_1, M_1] [\widetilde{g}; m_2, M_2] \right),
 \end{aligned}$$

where $P = \int_a^b p(x) dx$.

Proof. For any $x \in [a, b]$ we can write


$$(2.4) \quad u(x) = \frac{M_1 - u(x)}{M_1 - m_1} m_1 + \frac{u(x) - m_1}{M_1 - m_1} M_1$$

and

$$(2.5) \quad v(x) = \frac{M_2 - v(x)}{M_2 - m_2} m_2 + \frac{v(x) - m_2}{M_2 - m_2} M_2.$$

Since f and g are convex functions we have

$$\begin{aligned}
 f(u(x)) & \leq \frac{M_1 - u(x)}{M_1 - m_1} f(m_1) + \frac{u(x) - m_1}{M_1 - m_1} f(M_1) \\
 & = \frac{u(x)}{M_1 - m_1} (f(M_1) - f(m_1)) + \frac{M_1 f(m_1) - m_1 f(M_1)}{M_1 - m_1} \\
 (2.6) \quad & = [f; m_1, M_1] u(x) + [\widetilde{f}; m_1, M_1]
 \end{aligned}$$



Note on Some Hadamard-Type Inequalities

M. Klaričić Bakula and J. Pečarić

Title Page

Contents



Go Back

Close

Quit

Page 8 of 20

and

$$\begin{aligned}
 g(v(x)) &\leq \frac{M_2 - v(x)}{M_2 - m_2} g(m_2) + \frac{v(x) - m_2}{M_2 - m_2} g(M_2) \\
 &= \frac{v(x)}{M_2 - m_2} (g(M_2) - g(m_2)) + \frac{M_2 g(m_2) - m_2 g(M_2)}{M_2 - m_2} \\
 (2.7) \quad &= [g; m_2, M_2] v(x) + [\tilde{g}; m_2, M_2].
 \end{aligned}$$


Functions f and g are nonnegative by assumption, so after multiplying (2.6) and (2.7) we obtain

$$\begin{aligned}
 (2.8) \quad f(u(x)) g(v(x)) &\leq [f; m_1, M_1] [g; m_2, M_2] u(x) v(x) + [f; m_1, M_1] [\tilde{g}; m_2, M_2] u(x) \\
 &\quad + [g; m_2, M_2] [\tilde{f}; m_1, M_1] v(x) + [\tilde{f}; m_1, M_1] [\tilde{g}; m_2, M_2].
 \end{aligned}$$

Now, multiplying (2.8) by weight $p(x)$, integrating over $[a, b]$ and dividing by $P > 0$ we get (i).

To obtain (ii) we can write

$$\begin{aligned}
 \frac{m_1 + M_1}{2} &= \frac{1}{2} \left(\frac{M_1 - u(x)}{M_1 - m_1} m_1 + \frac{u(x) - m_1}{M_1 - m_1} M_1 \right. \\
 &\quad \left. + \frac{u(x) - m_1}{M_1 - m_1} m_1 + \frac{M_1 - u(x)}{M_1 - m_1} M_1 \right), \\
 \frac{m_2 + M_2}{2} &= \frac{1}{2} \left(\frac{M_2 - v(x)}{M_2 - m_2} m_2 + \frac{v(x) - m_2}{M_2 - m_2} M_2 \right. \\
 &\quad \left. + \frac{v(x) - m_2}{M_2 - m_2} m_2 + \frac{M_2 - v(x)}{M_2 - m_2} M_2 \right).
 \end{aligned}$$



Note on Some Hadamard-Type Inequalities

M. Klaričić Bakula and J. Pečarić

Title Page

Contents



Go Back

Close

Quit


Page 9 of 20

Using the Hadamard inequality (1.1) and the convexity of functions f and g , we get

$$\begin{aligned} & f\left(\frac{m_1 + M_1}{2}\right) g\left(\frac{m_2 + M_2}{2}\right) \\ & \leq \frac{1}{4} \left[f\left(\frac{M_1 - u(x)}{M_1 - m_1} m_1 + \frac{u(x) - m_1}{M_1 - m_1} M_1\right) \right. \\ & \quad \left. + f\left(\frac{u(x) - m_1}{M_1 - m_1} m_1 + \frac{M_1 - u(x)}{M_1 - m_1} M_1\right) \right] \\ & \quad \times \left[g\left(\frac{M_2 - v(x)}{M_2 - m_2} m_2 + \frac{v(x) - m_2}{M_2 - m_2} M_2\right) \right. \\ & \quad \left. + g\left(\frac{v(x) - m_2}{M_2 - m_2} m_2 + \frac{M_2 - v(x)}{M_2 - m_2} M_2\right) \right]. \end{aligned}$$

According to (2.4) and (2.5), after some simple calculus we obtain

$$\begin{aligned} (2.9) \quad & f\left(\frac{m_1 + M_1}{2}\right) g\left(\frac{m_2 + M_2}{2}\right) \\ & \leq \frac{1}{4} [f(u(x))g(v(x)) + f(M_1 + m_1 - u(x))g(M_2 + m_2 - v(x))] \\ & \quad + \frac{1}{4} \left[f\left(\frac{M_1 - u(x)}{M_1 - m_1} m_1 + \frac{u(x) - m_1}{M_1 - m_1} M_1\right) \right. \\ & \quad \times g\left(\frac{v(x) - m_2}{M_2 - m_2} m_2 + \frac{M_2 - v(x)}{M_2 - m_2} M_2\right) \end{aligned}$$



Note on Some Hadamard-Type Inequalities

M. Klaričić Bakula and J. Pečarić

Title Page

Contents



Go Back

Close

Quit

Page 10 of 20


$$+ f \left(\frac{u(x) - m_1}{M_1 - m_1} m_1 + \frac{M_1 - u(x)}{M_1 - m_1} M_1 \right) \\ \times g \left(\frac{M_2 - v(x)}{M_2 - m_2} m_2 + \frac{v(x) - m_2}{M_2 - m_2} M_2 \right) \Bigg].$$

Using the convexity of functions f and g , from inequality (2.9) we get

$$(2.10) \quad f \left(\frac{m_1 + M_1}{2} \right) g \left(\frac{m_2 + M_2}{2} \right) \\ \leq \frac{1}{4} [f(u(x))g(v(x)) + f(M_1 + m_1 - u(x))g(M_2 + m_2 - v(x))] \\ + \frac{1}{4} \left[\left(\frac{M_1 - u(x)}{M_1 - m_1} f(m_1) + \frac{u(x) - m_1}{M_1 - m_1} f(M_1) \right) \right. \\ \times \left(\frac{v(x) - m_2}{M_2 - m_2} g(m_2) + \frac{M_2 - v(x)}{M_2 - m_2} g(M_2) \right) \\ \left. + \left(\frac{u(x) - m_1}{M_1 - m_1} f(m_1) + \frac{M_1 - u(x)}{M_1 - m_1} f(M_1) \right) \right. \\ \left. \times \left(\frac{M_2 - v(x)}{M_2 - m_2} g(m_2) + \frac{v(x) - m_2}{M_2 - m_2} g(M_2) \right) \right].$$

With respect to the notation introduced at the beginning of this section, inequality (2.10) becomes

$$f \left(\frac{m_1 + M_1}{2} \right) g \left(\frac{m_2 + M_2}{2} \right)$$



Note on Some Hadamard-Type Inequalities

M. Klaričić Bakula and J. Pečarić

Title Page

Contents




Go Back

Close

Quit

Page 11 of 20



Note on Some Hadamard-Type Inequalities

M. Klaričić Bakula and J. Pečarić

Title Page

Contents



Go Back

Close

Quit

Page 12 of 20

$$\begin{aligned}
 &\leq \frac{1}{4} \left\{ f(u(x)) g(v(x)) \right. \\
 &\quad \left. + f(M_1 + m_1 - u(x)) g(M_2 + m_2 - v(x)) \right\} \\
 &+ \frac{1}{4} \left\{ \left([f; m_1, M_1] u(x) + [\tilde{f}; m_1, M_1] \right) \right. \\
 &\quad \times \left([\hat{g}; m_2, M_2] - [g; m_2, M_2] v(x) \right) \\
 &\quad \left. + \left([\hat{f}; m_1, M_1] - [f; m_1, M_1] u(x) \right) \right. \\
 &\quad \left. \times \left([g; m_2, M_2] v(x) + [\tilde{g}; m_2, M_2] \right) \right\} \\
 &= \frac{1}{4} \left\{ f(u(x)) g(v(x)) \right. \\
 &\quad \left. + f(M_1 + m_1 - u(x)) g(M_2 + m_2 - v(x)) \right\} \\
 &+ \frac{1}{4} \left\{ -2 [f; m_1, M_1] [g; m_2, M_2] u(x) v(x) \right. \\
 &\quad \left. + \left([\hat{g}; m_2, M_2] - [\tilde{g}; m_2, M_2] \right) [f; m_1, M_1] u(x) \right. \\
 &\quad \left. + \left([\hat{f}; m_1, M_1] - [\tilde{f}; m_1, M_1] \right) [g; m_2, M_2] v(x) \right. \\
 &\quad \left. + [\tilde{f}; m_1, M_1] [\hat{g}; m_2, M_2] + [\hat{f}; m_1, M_1] [\tilde{g}; m_2, M_2] \right\}
 \end{aligned}
 \tag{2.11}$$


Now we multiply both sides of (2.11) by $p(x)$, integrate over $[a, b]$ and divide by P . We thus obtain (ii) and the proof is completed. \square

Remark 2.1. Pachpatte's results (1.2) and (1.3) can be obtained from (2.2) and (2.3) respectively if we put $p(x) = 1, u(x) = v(x) = x$ for all $x \in [a, b]$ (then we have $m_1 = m_2 = a$ and $M_1 = M_2 = b$). In the case of $g(x) \equiv 1$ inequality (i) becomes the right side of Hadamard's inequality (1.1).

Theorem 2.2. Let f be a nonnegative convex function on $[m_1, M_1]$, g a nonnegative convex function on $[m_2, M_2]$, $u : [a, b] \rightarrow [m_1, M_1]$ and $v : [a, b] \rightarrow [m_2, M_2]$ continuous functions, and $p, q : [a, b] \rightarrow \mathbb{R}$ positive integrable functions. Then

(i)

$$\begin{aligned} & \frac{1}{PQ} \int_a^b \int_a^b \int_0^1 p(x) q(y) f(tu(x) + (1-t)u(y)) \\ & \quad \times g(tv(x) + (1-t)v(y)) dt dx dy \\ & \leq \frac{1}{3PQ} \left[Q \int_a^b f(u(x)) g(v(x)) p(x) dx \right. \\ & \quad \left. + P \int_a^b f(u(y)) g(v(y)) q(y) dy \right] \\ & \quad + \frac{1}{3PQ} \int_a^b p(x) f(u(x)) dx \int_a^b q(y) g(v(y)) dy; \end{aligned}$$



Note on Some Hadamard-Type Inequalities

M. Klaričić Bakula and J. Pečarić

Title Page

Contents



Go Back

Close

Quit

Page 13 of 20

(ii)

$$\begin{aligned} & \frac{1}{P} \int_a^b \int_0^1 p(x) f(tu(x) + (1-t)\bar{u}) g(tv(x) + (1-t)\bar{v}) dt dx \\ & \leq \frac{1}{3P} \int_a^b p(x) f(u(x)) g(v(x)) dx + \frac{1}{3} f(\bar{u}) g(\bar{v}) \\ & + \frac{1}{6P} \left[g(\bar{v}) \int_a^b p(x) f(u(x)) dx + f(\bar{u}) \int_a^b p(x) g(v(x)) dx \right], \end{aligned}$$

where $\bar{u} = \frac{1}{P} \int_a^b p(x) u(x) dx$, $\bar{v} = \frac{1}{Q} \int_a^b q(x) v(x) dx$.

Proof. Since f and g are convex functions, for $t \in [0, 1]$ we have

$$(2.12) \quad f(tu(x) + (1-t)u(y)) \leq tf(u(x)) + (1-t)f(u(y))$$


$$(2.13) \quad g(tv(x) + (1-t)v(y)) \leq tg(v(x)) + (1-t)g(v(y)).$$

Functions f and g are nonnegative, so multiplying (2.12) and (2.13) we get

$$\begin{aligned} (2.14) \quad & f(tu(x) + (1-t)u(y)) g(tv(x) + (1-t)v(y)) \\ & \leq t^2 f(u(x)) g(v(x)) + (1-t)^2 f(u(y)) g(v(y)) \\ & + t(1-t) [f(u(x)) g(v(y)) + f(u(y)) g(v(x))]. \end{aligned}$$

Integrating (2.14) over $[0, 1]$ we obtain

$$\begin{aligned} (2.15) \quad & \int_0^1 f(tu(x) + (1-t)u(y)) g(tv(x) + (1-t)v(y)) dt \\ & \leq \frac{1}{3} [f(u(x)) g(v(x)) + f(u(y)) g(v(y))] \end{aligned}$$



Note on Some Hadamard-Type Inequalities

M. Klaričić Bakula and J. Pečarić

Title Page

Contents



Go Back

Close

Quit

Page 14 of 20


$$+ \frac{1}{6} [f(u(x))g(v(y)) + f(u(y))g(v(x))].$$

Now we multiply (2.15) by $p(x)q(y)$, integrate over $[a, b] \times [a, b]$ and divide by PQ , where

$$P = \int_a^b p(x) dx, \quad Q = \int_a^b q(x) dx,$$

so we get

$$\begin{aligned} & \frac{1}{PQ} \int_a^b \int_a^b \int_0^1 p(x) q(y) f(tu(x) + (1-t)u(y)) \\ & \quad \times g(tv(x) + (1-t)v(y)) dt dx dy \\ & \leq \frac{1}{3PQ} \left[\int_a^b p(x) f(u(x)) g(v(x)) dx \int_a^b q(y) dy \right. \\ & \quad \left. + \int_a^b q(y) f(u(y)) g(v(y)) dy \int_a^b p(x) dx \right] \\ & \quad + \frac{1}{6PQ} \left[\int_a^b p(x) f(u(x)) dx \int_a^b q(y) g(v(y)) dy \right. \\ & \quad \left. + \int_a^b p(y) f(u(y)) dy \int_a^b q(x) g(v(x)) dx \right] \\ & = \frac{1}{3PQ} \left[\int_a^b p(x) f(u(x)) g(v(x)) dx \int_a^b q(y) dy \right. \end{aligned}$$



Note on Some Hadamard-Type Inequalities

M. Klaričić Bakula and J. Pečarić

Title Page

Contents



Go Back

Close

Quit

Page 15 of 20

$$(2.16) \quad \left[\int_a^b q(y) f(u(y)) g(v(y)) dy \int_a^b p(x) dx \right] + \frac{1}{3PQ} \int_a^b p(x) f(u(x)) dx \int_a^b q(y) g(v(y)) dy.$$

This is the desired inequality (i).


To prove inequality (ii), in (2.12) and (2.13) we substitute $u(y)$ and $v(y)$ with \bar{u} and \bar{v} respectively.

Then we obtain

$$(2.17) \quad f(tu(x) + (1-t)\bar{u}) g(tv(x) + (1-t)\bar{v}) \leq t^2 f(u(x)) g(v(x)) + (1-t)^2 f(\bar{u}) g(\bar{v}) + t(1-t) [f(u(x)) g(\bar{v}) + f(\bar{u}) g(v(x))].$$

Integrating (2.17) in respect to t over $[0, 1]$ we obtain

$$(2.18) \quad \int_0^1 f(tu(x) + (1-t)\bar{u}) g(tv(x) + (1-t)\bar{v}) dt \leq \frac{1}{3} [f(u(x)) g(v(x)) + f(\bar{u}) g(\bar{v})] + \frac{1}{6} [f(u(x)) g(\bar{v}) + f(\bar{u}) g(v(x))].$$



Note on Some Hadamard-Type Inequalities

M. Klaričić Bakula and J. Pečarić

Title Page

Contents



Go Back

Close

Quit

Page 16 of 20

Similarly as before, from (2.18) we get

$$\begin{aligned} & \frac{1}{P} \int_a^b \int_0^1 p(x) f(tu(x) + (1-t)\bar{u}) g(tv(x) + (1-t)\bar{v}) dt dx \\ & \leq \frac{1}{3P} \int_a^b p(x) f(u(x)) g(v(x)) dx + \frac{1}{3} f(\bar{u}) g(\bar{v}) \\ & \quad + \frac{1}{6P} \left[g(\bar{v}) \int_a^b p(x) f(u(x)) dx + f(\bar{u}) \int_a^b p(x) g(v(x)) dx \right]. \end{aligned}$$


This completes the proof. \square

Remark 2.2. If in (i) we put $u(x) = v(x) = x$ for all $x \in [a, b]$, it becomes

$$\begin{aligned} (2.19) \quad & \frac{1}{PQ} \int_a^b \int_a^b \int_0^1 p(x) q(y) f(tx + (1-t)y) g(tx + (1-t)y) dt dx dy \\ & \leq \frac{1}{3PQ} \left[Q \int_a^b p(x) f(x) g(x) dx + P \int_a^b q(y) f(y) g(y) dy \right] \\ & \quad + \frac{1}{3PQ} \int_a^b p(x) f(x) dx \int_a^b q(y) g(y) dy, \end{aligned}$$

so by using a generalization of Hadamard's inequality [1, p.138]

$$(2.20) \quad f\left(\frac{a+b}{2}\right) \leq \frac{1}{P} \int_a^b p(x) f(x) dx \leq \frac{f(a) + f(b)}{2}$$



Note on Some Hadamard-Type Inequalities

M. Klaričić Bakula and J. Pečarić

Title Page

Contents



Go Back

Close

Quit

Page 17 of 20


which holds for $p(a+t) = p(b-t)$, $0 \leq t \leq \frac{1}{2}(a+b)$, we obtain from (2.19) the following inequality

$$\begin{aligned}
 & \frac{1}{PQ} \int_a^b \int_a^b \int_0^1 p(x) q(y) f(tx + (1-t)y) g(tx + (1-t)y) dt dx dy \\
 & \leq \frac{1}{3PQ} \left[Q \int_a^b p(x) f(x) g(x) dx + P \int_a^b q(y) f(y) g(y) dy \right] \\
 & \quad + \frac{1}{3} \frac{f(a) + f(b)}{2} \frac{g(a) + g(b)}{2} \\
 & = \frac{1}{3PQ} \left[Q \int_a^b p(x) f(x) g(x) dx + P \int_a^b q(y) f(y) g(y) dy \right] \\
 (2.21) \quad & + \frac{1}{12} [M(a, b) + N(a, b)].
 \end{aligned}$$

Now it is easy to observe that if $p(x) = q(x) = 1$ for all $x \in [a, b]$ inequality (2.21) becomes the corrected Pachpatte's result (1.4).

If we do the same in (ii) we get

$$\begin{aligned}
 & \frac{1}{P} \int_a^b \int_0^1 p(x) f\left(tx + (1-t)\frac{a+b}{2}\right) g\left(tx + (1-t)\frac{a+b}{2}\right) dt dx \\
 & \leq \frac{1}{3P} \int_a^b p(x) f(x) g(x) dx + \frac{1}{3} f\left(\frac{a+b}{2}\right) g\left(\frac{a+b}{2}\right) \\
 & + \frac{1}{6P} \left[g\left(\frac{a+b}{2}\right) \int_a^b p(x) f(x) dx + f\left(\frac{a+b}{2}\right) \int_a^b p(x) g(x) dx \right].
 \end{aligned}$$



Note on Some Hadamard-Type Inequalities

M. Klaričić Bakula and J. Pečarić

Title Page

Contents



Go Back

Close

Quit

Page 18 of 20


Using again (2.20) we obtain

$$\begin{aligned} & \frac{1}{P} \int_a^b \int_0^1 p(x) f\left(tx + (1-t)\frac{a+b}{2}\right) g\left(tx + (1-t)\frac{a+b}{2}\right) dt dx \\ & \leq \frac{1}{3P} \int_a^b p(x) f(x) g(x) dx + \frac{1}{3} \frac{f(a) + f(b)}{2} \frac{g(a) + g(b)}{2} \\ & \quad + \frac{1}{6} \left[\frac{g(a) + g(b)}{2} \frac{f(a) + f(b)}{2} + \frac{f(a) + f(b)}{2} \frac{g(a) + g(b)}{2} \right] \\ & = \frac{1}{3P} \int_a^b p(x) f(x) g(x) dx + \frac{1}{6} (M(a, b) + N(a, b)) \end{aligned}$$

Furthermore, in the case $p(x) = 1$ for all $x \in [a, b]$ we get

$$\begin{aligned} & \frac{1}{b-a} \int_a^b \int_0^1 f\left(tx + (1-t)\frac{a+b}{2}\right) g\left(tx + (1-t)\frac{a+b}{2}\right) dt dx \\ & \leq \frac{1}{3(b-a)} \int_a^b f(x) g(x) dx + \frac{1}{6} (M(a, b) + N(a, b)), \end{aligned}$$

which is the corrected Pachpatte's result (1.5).



Note on Some Hadamard-Type Inequalities

M. Klaričić Bakula and J. Pečarić

Title Page

Contents



Go Back


Close

Quit

Page 19 of 20

References

- [1] J.E. PEČARIĆ, F. PROSCHAN AND Y.L. TONG, *Convex Functions, Partial Orderings, and Statistical Applications*, Academic Press, Inc. (1992).
- [2] D.S. MITRINOVIĆ, J.E. PEČARIĆ AND A.M. FINK, *Classical and New Inequalities in Analysis*, Kluwer Academic Publishers, 1993.
- [3] B.G. PACHPATTE, On some inequalities for convex functions, *RGMIA Res. Rep. Coll.*, **6**(E) (2003). [ONLINE [http://rgmia.vu.edu.au/v6\(E\).html](http://rgmia.vu.edu.au/v6(E).html)].



Note on Some Hadamard-Type Inequalities

M. Klaričić Bakula and J. Pečarić

Title Page

Contents



Go Back

Close

Quit

Page 20 of 20