


## A CLASS OF INEQUALITIES RELATED TO THE ANGLE BISECTORS AND THE SIDES OF A TRIANGLE

SHAN-HE WU AND ZHI-HUA ZHANG

Department of Mathematics  
Longyan College  
Longyan Fujian 364012  
People's Republic of China.  
*EMail:* [wushanhe@yahoo.com.cn](mailto:wushanhe@yahoo.com.cn)

Zixing Educational Research Section  
Chenzhou, Hunan 423400  
People's Republic of China.  
*EMail:* [zxzh1234@163.com](mailto:zxzh1234@163.com)



---

volume 7, issue 3, article 108,  
2006.

*Received 18 February, 2006;  
accepted 23 March, 2006.*

*Communicated by: A. Lupaş*

---

Abstract

Contents



Home Page

Go Back

Close

Quit

## Abstract

In the present paper a class of geometric inequalities concerning the angle bisectors and the sides of a triangle are established. Moreover an interesting open problem is proposed.

*2000 Mathematics Subject Classification:* 26D15.

*Key words:* Inequality; Triangle; Angle bisector; Cyclic sum; Best coefficient.

The research was supported by the Natural Science Foundation of Fujian Province Education Department of China under grant No. JA05324.

## Contents

1	Introduction and Main Results .....	3
2	Proof of Theorem 1.1 .....	5
3	An Application .....	12
	References	



---

### A Class of Inequalities Related to the Angle Bisectors and the Sides of a Triangle

Shan-He Wu and Zhi-Hua Zhang

---

Title Page

Contents



Go Back

Close

Quit

Page 2 of 16

# 1. Introduction and Main Results


For a given triangle  $ABC$  we assume that  $A, B, C$  denote its angles,  $a, b, c$  denote the lengths of its corresponding sides,  $w_a, w_b, w_c$  denote respectively the bisectors of angles  $A, B, C$ . Let  $R, r$  and  $s$  be the circumradius, the inradius and the semi-perimeter of a triangle respectively. In addition we will customarily use the symbols  $\sum$  (cyclic sum) and  $\prod$  (cyclic product), such as

$$\sum f(a) = f(a) + f(b) + f(c), \quad \prod f(a) = f(a)f(b)f(c).$$

The angle bisectors of triangles have many interesting properties. In particular, inequalities for angle bisectors is a very attractive subject and plays an important role in the study of geometry. A large number of related results can be found in the well-known monographs [1] – [3]. In recent years, we have given considerable attention to these inequalities (see [4] – [8]). Recently, the following interesting double inequality concerning the angle bisectors and the sides, which was presented by X.-Zh. Yang, T.-Y. Ma and W.-L. Dong in [9, 10], has come to our attention:

$$(1.1) \quad \frac{3\sqrt{3}}{2} + \left(\frac{8}{3} - \frac{3\sqrt{3}}{2}\right) \left(1 - \frac{2r}{R}\right) \leq \sum \frac{w_a}{a} \leq \frac{3\sqrt{3}}{2} + 2\sqrt{3} \left(\frac{R}{2r} - 1\right).$$

The above result also motivates us to investigate some similar inequalities. We give here sharp lower and upper bounds for the sum  $\sum \frac{a}{w_a}$ . Moreover, in Section 3 the obtained result will be used for establishing an analogue of inequality (1.1).



---

**A Class of Inequalities Related to the Angle Bisectors and the Sides of a Triangle**

Shan-He Wu and Zhi-Hua Zhang

---

Title Page

Contents



Go Back

Close


Quit

Page 3 of 16

**Theorem 1.1.** *In any triangle  $ABC$  the following double inequalities hold*

$$(1.2) \quad \frac{1}{2} \left( \frac{s}{r} + \sqrt{3} \right) \leq \sum \frac{a}{w_a} \leq \frac{\sqrt{2}}{2} \left( \frac{s}{r} + 2\sqrt{6} - 3\sqrt{3} \right),$$

*with equality if and only if the triangle is an equilateral. Furthermore,  $\frac{1}{2}$  and  $\frac{\sqrt{2}}{2}$  are the best coefficients in (1.2).*



---

**A Class of Inequalities Related  
to the Angle Bisectors and the  
Sides of a Triangle**

Shan-He Wu and Zhi-Hua Zhang

---

Title Page

Contents



Go Back

Close

Quit

Page 4 of 16

## 2. Proof of Theorem 1.1

To prove Theorem 1.1, we shall use the following known results [2, p. 3, p. 241] (see also [1])

**Lemma 2.1.** *In any triangle  $ABC$  we have the following inequalities*

$$(2.1) \quad s^4 \leq s^2(4R^2 + 20Rr - 2r^2) - r(4R + r)^3,$$

$$(2.2) \quad 2R^2 + 10Rr - r^2 - 2(R - 2r)\sqrt{R^2 - 2Rr} \\ \leq s^2 \leq 2R^2 + 10Rr - r^2 + 2(R - 2r)\sqrt{R^2 - 2Rr},$$

*with equality if and only if the triangle is isosceles.*

$$(2.3) \quad s \leq \frac{1}{\sqrt{3}}(4R + r),$$

*with equality if and only if the triangle is equilateral.*


*In any acute triangle  $ABC$  we have*

$$(2.4) \quad s^2 \geq 4R^2 + 4Rr + r^2,$$

*with equality if and only if the triangle is equilateral.*

**Proof of Theorem 1.1.** By the formula for angle bisector of triangle  $ABC$   $w_a = \frac{2bc}{b+c} \cos \frac{A}{2}$ , we have

$$\sum \frac{a}{w_a} = \sum (\csc B + \csc C) \sin \frac{A}{2} \\ = \left( \sum \csc A \right) \left( \sum \sin \frac{A}{2} \right) - \frac{1}{2} \sum \sec \frac{A}{2}.$$



**A Class of Inequalities Related to the Angle Bisectors and the Sides of a Triangle**

Shan-He Wu and Zhi-Hua Zhang

Title Page

Contents



Go Back

Close

Quit

Page 5 of 16

Based on the above result, it follows from the identity  $\prod \cot \frac{A}{2} = \frac{s}{r}$  that the inequality (1.2) is equivalent to the following inequality

$$(2.5) \quad \frac{1}{2} \prod \cot \frac{A}{2} + \frac{\sqrt{3}}{2} \leq \left( \sum \csc A \right) \left( \sum \sin \frac{A}{2} \right) - \frac{1}{2} \sum \sec \frac{A}{2} \\ \leq \frac{\sqrt{2}}{2} \prod \cot \frac{A}{2} + \frac{4\sqrt{3} - 3\sqrt{6}}{2}.$$

Using a substitution  $A \rightarrow \pi - 2A$ ,  $B \rightarrow \pi - 2B$ ,  $C \rightarrow \pi - 2C$  in (2.5), then the inequality (2.5) can be translated to

$$(2.6) \quad \frac{1}{2} \prod \tan A + \frac{\sqrt{3}}{2} \leq \left( \sum \csc 2A \right) \left( \sum \cos A \right) - \frac{1}{2} \sum \csc A \\ \leq \frac{\sqrt{2}}{2} \prod \tan A + \frac{4\sqrt{3} - 3\sqrt{6}}{2}.$$


Now, in order to prove the inequality (2.5), it is enough to prove that the inequality (2.6) to be valid for any acute triangle.

Note that the following known identities for a triangle [2, p. 55-60]:

$$\sum \csc 2A = \frac{s^4 + s^2(2r^2 - 8Rr - 4R^2) + 16R^3r + 20R^2r^2 + 8Rr^3 + r^4}{4rs(s^2 - 4R^2 - 4Rr - r^2)},$$

$$\prod \tan A = \frac{2rs}{s^2 - 4R^2 - 4Rr - r^2},$$

$$\sum \csc A = \frac{s^2 + 4Rr + r^2}{2rs},$$



**A Class of Inequalities Related to the Angle Bisectors and the Sides of a Triangle**

Shan-He Wu and Zhi-Hua Zhang

Title Page

Contents



Go Back

Close

Quit

Page 6 of 16

$$\sum \cos A = \frac{R+r}{R},$$

taking these identities into (2.6), we find that the inequality (2.6) is equivalent to

$$(2.7) \quad 4Rr^2s^2 + 2\sqrt{3}(s^2 - 4R^2 - 4Rr - r^2)srR \\ \leq H \leq 4\sqrt{2}Rr^2s^2 + (8\sqrt{3} - 6\sqrt{6})(s^2 - 4R^2 - 4Rr - r^2)srR,$$


where

$$H = (R+r)[s^4 + s^2(2r^2 - 8Rr - 4R^2) \\ + 16R^3r + 20R^2r^2 + 8Rr^3 + r^4] \\ - R(s^2 + 4Rr + r^2)(s^2 - 4R^2 - 4Rr - r^2).$$

Let us now prove the inequality (2.7) to be valid for any acute triangle.

Using the inequalities (2.1), (2.3) and (2.4), we have

$$(2.8) \quad H - 4\sqrt{2}Rr^2s^2 - (8\sqrt{3} - 6\sqrt{6})(s^2 - 4R^2 - 4Rr - r^2)srR \\ \leq (R+r)[s^2(4R^2 + 20Rr - 2r^2) - r(4R+r)^3 \\ + s^2(2r^2 - 8Rr - 4R^2) + 16R^3r + 20R^2r^2 + 8Rr^3 + r^4] \\ - R(s^2 + 4Rr + r^2)(s^2 - 4R^2 - 4Rr - r^2) - 4\sqrt{2}Rr^2s^2 \\ - (8 - 6\sqrt{2})(s^2 - 4R^2 - 4Rr - r^2)(4R+r)rR \\ = s^2 \left[ (-40 + 24\sqrt{2})R + (6 + 2\sqrt{2})r \right] Rr \\ + (160 - 96\sqrt{2})R^4r + (152 - 120\sqrt{2})R^3r^2$$



**A Class of Inequalities Related to the Angle Bisectors and the Sides of a Triangle**

Shan-He Wu and Zhi-Hua Zhang

Title Page

Contents



Go Back

Close

Quit

Page 7 of 16

$$\begin{aligned}
& + \left(52 - 48\sqrt{2}\right) R^2 r^3 + \left(6 - 6\sqrt{2}\right) R r^4 \\
& = F(s, R, r).
\end{aligned}$$


From Euler's inequality  $R \geq 2r$ , we observe that  $(-40 + 24\sqrt{2})R + (6 + 2\sqrt{2})r < 0$ .

*Case 1.* When  $R > (\sqrt{2} + 1)r$ , by inequality (2.4) we have

$$\begin{aligned}
F(s, R, r) & \leq (4R^2 + 4Rr + r^2) \left[ (-40 + 24\sqrt{2})R + (6 + 2\sqrt{2})r \right] Rr \\
& \quad + \left(160 - 96\sqrt{2}\right) R^4 r + \left(152 - 120\sqrt{2}\right) R^3 r^2 \\
& \quad + \left(52 - 48\sqrt{2}\right) R^2 r^3 + \left(6 - 6\sqrt{2}\right) R r^4 \\
& = \left(16 - 16\sqrt{2}\right) R^3 r^2 + \left(36 - 16\sqrt{2}\right) R^2 r^3 + \left(12 - 4\sqrt{2}\right) R r^4 \\
& = -4Rr^2 \left[ R - (\sqrt{2} + 1)r \right] \left[ (4\sqrt{2} - 4)R + (4\sqrt{2} - 5)r \right] \\
& < 0.
\end{aligned}$$

*Case 2.* When  $2r \leq R \leq (\sqrt{2} + 1)r$ , by inequality (2.2) we get

$$\begin{aligned}
F(s, R, r) & \leq \left[ 2R^2 + 10Rr - r^2 - 2(R - 2r)\sqrt{R^2 - 2Rr} \right] \\
& \quad \times \left[ (-40 + 24\sqrt{2})R + (6 + 2\sqrt{2})r \right] Rr + \left(160 - 96\sqrt{2}\right) R^4 r \\
& \quad + \left(152 - 120\sqrt{2}\right) R^3 r^2 + \left(52 - 48\sqrt{2}\right) R^2 r^3 + \left(6 - 6\sqrt{2}\right) R r^4 \\
& = 4Rr^3(R - 2r)[F_1(R, r) + F_2(R, r)],
\end{aligned}$$



**A Class of Inequalities Related to the Angle Bisectors and the Sides of a Triangle**

Shan-He Wu and Zhi-Hua Zhang

Title Page

Contents



Go Back

Close

Quit

Page 8 of 16



where

$$F_1(R, r) = \left[ \left( 20 - 12\sqrt{2} \right) \left( \frac{R}{r} \right) - 3 - \sqrt{2} \right] \sqrt{\left( \frac{R}{r} \right) \left( \frac{R}{r} - 2 \right)},$$

$$F_2(R, r) = \left( 20 - 12\sqrt{2} \right) \left( \frac{R}{r} \right)^2 + \left( -19 + 7\sqrt{2} \right) \left( \frac{R}{r} \right) + \sqrt{2}.$$

We deduce from  $2 \leq R/r \leq \sqrt{2} + 1$  that


$$\begin{aligned} F_1(R, r) &\leq \left[ \left( 20 - 12\sqrt{2} \right) \left( \sqrt{2} + 1 \right) - 3 - \sqrt{2} \right] \sqrt{\left( \sqrt{2} + 1 \right) \left( \sqrt{2} + 1 - 2 \right)} \\ &= 7\sqrt{2} - 7, \end{aligned}$$

$$\begin{aligned} F_2(R, r) &\leq \left( 20 - 12\sqrt{2} \right) \left( \sqrt{2} + 1 \right)^2 + \left( -19 + 7\sqrt{2} \right) \left( \sqrt{2} + 1 \right) + \sqrt{2} \\ &= 7 - 7\sqrt{2}, \end{aligned}$$

which leads to  $F(s, R, r) \leq 0$ .

Consequently

$$(2.9) \quad H - 4\sqrt{2}Rr^2s^2 - \left( 8\sqrt{3} - 6\sqrt{6} \right) \left( s^2 - 4R^2 - 4Rr - r^2 \right) srR \leq 0.$$



**A Class of Inequalities Related to the Angle Bisectors and the Sides of a Triangle**

Shan-He Wu and Zhi-Hua Zhang

Title Page

Contents



Go Back

Close

Quit

Page 9 of 16

On the other hand, utilizing the inequalities (2.3) and (2.4), we have

$$\begin{aligned}
 & H - 4Rr^2s^2 - 2\sqrt{3}(s^2 - 4R^2 - 4Rr - r^2)srR \\
 & \geq (R + r) [s^4 + s^2(2r^2 - 8Rr - 4R^2) + 16R^3r + 20R^2r^2 + 8Rr^3 + r^4] \\
 & \quad - R(s^2 + 4Rr + r^2)(s^2 - 4R^2 - 4Rr - r^2) \\
 & \quad - 4\sqrt{2}Rr^2s^2 - 2(s^2 - 4R^2 - 4Rr - r^2)(4R + r)R \\
 & = r(s^2 - 4R^2 - 4Rr - 3r^2)^2 + 4r(R + r)F(s, R, r),
 \end{aligned}$$

where


$$F(s, R, r) = -s^2(3R - 2r) + 12R^3 + 4R^2r - Rr^2 - 2r^3.$$

By Euler's inequality  $R \geq 2r$ , we conclude that  $3R - 2r > 0$ . Using the inequality (2.2) yields

$$\begin{aligned}
 F(s, R, r) & \geq -(3R - 2r) \left[ 2R^2 + 10Rr - r^2 + 2(R - 2r)\sqrt{R^2 - 2Rr} \right] \\
 & \quad + 12R^3 + 4R^2r - Rr^2 - 2r^3 \\
 & = 2(R - 2r) \left[ R(3R - 5r) + r^2 - (3R - 2r)\sqrt{R^2 - 2Rr} \right] \\
 & = 2(R - 2r) \frac{[Rr^2(3R - 2r) + r^4]}{R(3R - 5r) + r^2 + (3R - 2r)\sqrt{R^2 - 2Rr}} \\
 & \geq 0.
 \end{aligned}$$

Consequently

$$(2.10) \quad H - 4Rr^2s^2 - 2\sqrt{3}(s^2 - 4R^2 - 4Rr - r^2)srR \geq 0.$$



**A Class of Inequalities Related to the Angle Bisectors and the Sides of a Triangle**

Shan-He Wu and Zhi-Hua Zhang

Title Page

Contents



Go Back

Close

Quit

Page 10 of 16

Combining (2.9) and (2.10) yields the inequality (2.7), then from (2.7), the inequality (1.2) follows immediately. Moreover, from the process of proving inequality (1.2), it is easy to observe that the equalities hold in (1.2) if and only if the triangle is equilateral.

Next, we need to show that the coefficients  $\frac{1}{2}$  and  $\frac{\sqrt{2}}{2}$  in (1.2) are best possible in the strong sense.


Consider the inequality (1.2) in a general form as

$$(2.11) \quad \lambda \left( \frac{s}{r} + \frac{2\sqrt{3}}{\lambda} - 3\sqrt{3} \right) \leq \sum \frac{a}{w_a} \leq k \left( \frac{s}{r} + \frac{2\sqrt{3}}{k} - 3\sqrt{3} \right).$$

Putting  $a = 1, b = 1, c = 2t$  in (2.11) yields that

$$(2.12) \quad \begin{aligned} \lambda(1+t) + (2\sqrt{3} - 3\sqrt{3}\lambda) t \sqrt{\frac{1-t}{1+t}} \\ \leq \frac{4t^2 + (1+2t)\sqrt{2-2t}}{2+2t} \\ \leq k(1+t) + (2\sqrt{3} - 3\sqrt{3}k) t \sqrt{\frac{1-t}{1+t}}. \end{aligned}$$

In (2.12), passing the limit as  $t \rightarrow 0$  and  $t \rightarrow 1$  respectively, we find that  $\lambda \leq \frac{1}{2}$  and  $k \geq \frac{\sqrt{2}}{2}$ . Thus the best possible values for  $\lambda$  and  $k$  in (2.11) is that  $\lambda_{\max} = \frac{1}{2}$ ,  $k_{\min} = \frac{\sqrt{2}}{2}$ . This completes the proof of Theorem 1.1.  $\square$



A Class of Inequalities Related to the Angle Bisectors and the Sides of a Triangle

Shan-He Wu and Zhi-Hua Zhang

Title Page

Contents



Go Back

Close

Quit

Page 11 of 16

### 3. An Application

As an application of Theorem 1.1, we establish an analogue of the inequality (1.1), as follows.

**Theorem 3.1.** *In any triangle ABC the following double inequalities hold*

$$(3.1) \quad 2\sqrt{3} + \frac{3}{2} \left(1 - \frac{2r}{R}\right) \leq \sum \frac{a}{w_a} \leq 2\sqrt{3} + 2\sqrt{2} \left(\frac{R}{2r} - 1\right),$$


with equality if and only if the triangle is equilateral. Furthermore,  $2\sqrt{2}$  is the best coefficient in the right-hand side of inequality (3.1).

*Proof.* Applying Theorem 1.1 and Blundon's inequality [11]  $s \leq 2R + (3\sqrt{3} - 4)r$ , it follows that

$$\sum \frac{a}{w_a} \leq \frac{\sqrt{2}}{2} \left(\frac{s}{r} + 2\sqrt{6} - 3\sqrt{3}\right) \leq 2\sqrt{3} + 2\sqrt{2} \left(\frac{R}{2r} - 1\right).$$

On the other hand, by using Theorem 1.1 and Gerretsen's inequality [12]  $s^2 \geq 16Rr - 5r^2$ , we have

$$\begin{aligned} \sum \frac{a}{w_a} - 2\sqrt{3} - \frac{3}{2} \left(1 - \frac{2r}{R}\right) &\geq \frac{1}{2} \left(\frac{s}{r} + \sqrt{3}\right) - 2\sqrt{3} - \frac{3}{2} \left(1 - \frac{2r}{R}\right) \\ &= \frac{1}{2Rr} [sR - 3(R - 2r)r - 6\sqrt{3}Rr] \end{aligned}$$



A Class of Inequalities Related to the Angle Bisectors and the Sides of a Triangle

Shan-He Wu and Zhi-Hua Zhang

Title Page

Contents



Go Back

Close

Quit

Page 12 of 16

$$\begin{aligned}
&\geq \frac{1}{2Rr} \left[ R\sqrt{16Rr - 5r^2} - 3(R - 2r)r - 3\sqrt{3}Rr \right] \\
&= \frac{R^2(16Rr - 5r^2) - [3(R - 2r)r + 3\sqrt{3}Rr]^2}{2Rr \left[ R\sqrt{16Rr - 5r^2} + 3(R - 2r)r + 3\sqrt{3}Rr \right]} \\
&= \frac{16(R - 2r)^3 + (55 - 18\sqrt{3})(R - 2r)^2 + (64 - 36\sqrt{3})(R - 2r)}{2R \left[ R\sqrt{16Rr - 5r^2} + 3(R - 2r)r + 3\sqrt{3}Rr \right]} \\
&\geq 0,
\end{aligned}$$

so that

$$\sum \frac{a}{w_a} \geq 2\sqrt{3} + \frac{3}{2} \left( 1 - \frac{2r}{R} \right).$$

The inequality (3.1) is proved. It follows directly from Theorem 1.1 that the equalities hold in (3.1) if and only if the triangle is equilateral.

Let us now show that the coefficient  $2\sqrt{2}$  in the right-hand side of inequality (3.1) is best possible.


Consider the inequality (3.1) in a general form as

$$(3.2) \quad \sum \frac{a}{w_a} \leq 2\sqrt{3} + \mu \left( \frac{R}{2r} - 1 \right).$$

Putting  $a = 1$ ,  $b = 1$ ,  $c = 2t$  in (3.2) yields that

$$(3.3) \quad \frac{4t^2 + (1 + 2t)\sqrt{2 - 2t}}{2 + 2t} \leq \frac{\mu}{4\sqrt{1 - t^2}} + (2\sqrt{3} - \mu) t \sqrt{\frac{1 - t}{1 + t}}.$$

Passing the limit as  $t \rightarrow 0$  in (3.3), we get  $\mu \geq 2\sqrt{2}$ . Thus the best possible value for  $\mu$  in (3.2) should be  $\mu_{\min} = 2\sqrt{2}$ . The proof of Theorem 3.1 is complete.  $\square$



**A Class of Inequalities Related to the Angle Bisectors and the Sides of a Triangle**

Shan-He Wu and Zhi-Hua Zhang

Title Page

Contents



Go Back

Close

Quit


Page 13 of 16

It is worth noticing that the coefficient  $\frac{3}{2}$  is not best possible for the left-hand side of inequality (3.1), this may lead us to further discussion of the following significant problem.

**Open Problem.** *Determine the best coefficient  $\mu$  for which the inequality below holds*

$$(3.4) \quad \sum \frac{a}{w_a} \geq 2\sqrt{3} + \mu \left(1 - \frac{2r}{R}\right).$$

It seems that the problem is complicated and difficult. Indeed, it is unable to be solved in a same way as the foregoing technique.




---

**A Class of Inequalities Related to the Angle Bisectors and the Sides of a Triangle**

Shan-He Wu and Zhi-Hua Zhang

---

Title Page

Contents



Go Back


Close

Quit

Page 14 of 16

## References

- [1] O. BOTTEMA, R.Z. DJORDJEVIĆ, R.R. JANIĆ, D.S. MITRINOVIĆ AND P.M. VASIĆ, *Geometric Inequalities.*, Groningen, Wolters-Noordhoff, 1969.
- [2] D.S. MITRINOVIĆ, J.E. PEČARIĆ AND V. VOLENEC, *Recent Advances in Geometric Inequalities*, Dordrecht, Netherlands, Kluwer Academic Publishers, 1989.
- [3] D.S. MITRINOVIĆ, J.E. PEČARIĆ, V. VOLENEC AND J. CHEN, Addenda to the monograph: Recent Advances in Geometric Inequalities (I), *Journal of Ningbo University*, **4**(2) (1991), 1–145.
- [4] SH.-H. WU, An inverse inequality for the angle bisectors and its analogue. *Journal of Suzhou Railway Teachers College*, **17**(8) (2000), 42–44.(in Chinese)
- [5] SH.-H. WU, A strengthened inequality for the angle bisectors and its analogue. *Journal of Longyan Teachers College*, **20**(10) (2002), 98–99.(in Chinese)
- [6] SH.-H. WU, Some inequalities related to the angle bisectors and the radius of escribed circle, *Journal of Guizhou Educational College*, **13**(8) (2002), 16–18. (in Chinese)
- [7] SH.-H. WU, An inverse inequality for the angle bisectors, *Maths Teaching in Middle School*, **6** (2001), 62–63. (in Chinese)



---

A Class of Inequalities Related  
to the Angle Bisectors and the  
Sides of a Triangle

Shan-He Wu and Zhi-Hua Zhang

---

Title Page

Contents




Go Back

Close

Quit

Page 15 of 16

- [8] SH.-H. WU, A simple proof of an inequality concerning the angle bisectors, *Fujian High-School Mathematics*, **3** (1997), 19–20. (in Chinese)
- [9] X.-ZH. YANG, Sharpness of an inequality involving the angle bisectors, *Middle School Mathematics*, **12** (1995), 20–22. (in Chinese)
- [10] T.-Y. MA AND W.-L. DONG, The inverse version of an inequality concerning the angle bisectors, *Bulletin of Maths (Wuhan)*, **11** (1995), 19–20. (in Chinese)
- [11] W.J. BLUNDON, Inequalities associated with the triangle, *Canad. Math. Bull.*, **8** (1965), 615–626.
- [12] J.C. GERRETSEN, *Ongelijkheden in de Driehoek*, *Nieuw Tijdschr. Wisk.*, **41** (1953), 1–7.




---

**A Class of Inequalities Related to the Angle Bisectors and the Sides of a Triangle**

Shan-He Wu and Zhi-Hua Zhang

---

Title Page

Contents



Go Back

Close

Quit

Page 16 of 16