


POWER-MONOTONE SEQUENCES AND INTEGRABILITY OF TRIGONOMETRIC SERIES

J. NÉMETH

Department of Mathematics
Bolyai Institute, University of Szeged,
Aradi Vértanúk Tere 1.
6720 Szeged, Hungary
EMail: nemethj@math.u-szeged.hu



volume 4, issue 1, article 3,
2003.

*Received 13 June, 2002;
accepted 20 September, 2002.*

Communicated by: A. Babenko

[Abstract](#)

[Contents](#)



[Home Page](#)

[Go Back](#)

[Close](#)

[Quit](#)

Abstract

The theorem proved in this paper is a generalization of some results, concerning integrability of trigonometric series, due to R.P. Boas, L. Leindler, etc. This result can be considered as an example showing the utility of the notion of power-monotone sequences.

2000 Mathematics Subject Classification: 42A32.

Key words: Trigonometrical series, Integrability, Quasi power-monotone sequence.

The author was partially supported by the Hungarian National Foundation for Scientific Research under Grant # T 029094.

Contents

1	Introduction	3
2	Result	6
3	Lemmas	7
4	Proofs	9
	References	



Power-Monotone Sequences and Integrability of Trigonometric Series

J. Németh

Title Page

Contents



Go Back

Close

Quit

Page 2 of 14

1. Introduction

Several authors have studied the integrability of the series

$$(1.1) \quad g(x) = \sum_{n=1}^{\infty} b_n \sin nx$$

requiring certain conditions on the sequence $\{b_n\}$ (see [1] – [6] and [9] – [14]).

For example R.P. Boas in [2] proved the following result for (1.1):

Theorem 1.1. *If $b_n \downarrow 0$ then for $0 \leq \gamma \leq 1$, $x^{-\gamma}g(x) \in L[0, \pi]$ if and only if $\sum_{n=1}^{\infty} n^{\gamma-1}b_n$ converges.*

This theorem had previously been proved for $\gamma = 0$ by W.H. Young [14] and was later extended by P. Heywood [6] for $1 < \gamma < 2$.

Further generalization was given by Aljančić, R. Bojanić and M. Tomić in [1], by using the so called slowly varying functions.


A positive, continuous function defined on $[0, \infty)$ is called slowly varying if $\frac{L(tx)}{L(x)} \rightarrow 1$, if $x \rightarrow \infty$ for all $t > 0$.

They proved among others the following results:

Theorem 1.2. *Let $L(x)$ be a convex, non-decreasing slowly varying function. If $b_n \downarrow 0$, then $L(1/x)g(x) \in L(0, \pi]$ if and only if $\sum_{n=1}^{\infty} n^{-1}L(n)b_n$ converges.*

Theorem 1.3. *Let $L(x)$ be a slowly varying function. If $b_n \downarrow 0$ and $0 < \gamma < 2$, then $x^{-\gamma}L\left(\frac{1}{x}\right)g(x) \in L[0, \pi]$ if and only if $\sum_{n=1}^{\infty} n^{\gamma-1}L(n)b_n$ converges.*

Later the monotonicity condition on $\{b_n\}$ was changed to more general ones by S.M. Shah [11] and L. Leindler [9]. Before formulating a result of this type we need a definition due to L. Leindler.



Title Page

Contents



Go Back

Close

Quit

Page 3 of 14

A sequence $\mathbf{c} := \{c_n\}$ of positive numbers tending to zero is of rest bounded variation, or briefly $R_0^+ BVS$, if it has the property

$$(1.2) \quad \sum_{n=m}^{\infty} |c_n - c_{n+1}| \leq K(\mathbf{c})c_m$$

for all natural number m , where $K(\mathbf{c})$ is a constant depending only on \mathbf{c} .

Using this notion L. Leindler ([9]) proved

Theorem 1.4. *Let $\{b_n\} \in R_0^+ BVS$. If $0 \leq \gamma \leq 1$ and*

$$(1.3) \quad \sum_{n=1}^{\infty} n^{\gamma-1} b_n < \infty,$$


then $x^{-\gamma}g(x) \in L(0; \pi)$.

The aim of the present note is to give further generalization of above mentioned theorems by using the concept of the so called quasi β -power-monotone sequence changing the function x^γ to more general one. We deal only with the sufficiency of the conditions because only this point of the proofs has interest in showing up the utility of the quasi β -power-monotone sequences, the proof of the necessity, in general, goes on the same way as in the earlier cited papers.

First we need some definitions before formulating our result and the lemmas used in the proof.

Following L. Leindler we shall say that a sequence $\gamma := \{\gamma_n\}$ of positive terms is quasi β -power-monotone increasing (decreasing) if there exists a natural number $N := N(\beta, \gamma)$ and constant $K := K(\beta, \gamma) > 1$ such that

$$(1.4) \quad Kn^\beta \gamma_n \geq m^\beta \gamma_m \quad (n^\beta \gamma_n \leq Km^\beta \gamma_m)$$



Title Page

Contents



Go Back

Close

Quit

Page 4 of 14

holds for any $n \geq m \geq N$.

Here and in the sequel, K and K_i denote positive constants that are not necessarily the same of each occurrence.

If (1.4) holds with $\beta = 0$ then we omit the attribute “ β -power”.

Furthermore, we shall say that a sequence $\gamma := \{\gamma_n\}$ of positive terms is quasi geometrically increasing (decreasing) if there exist natural numbers $\mu := \mu(\gamma)$, $N := N(\gamma)$ and a constant $K := K(\gamma) \geq 1$ such that

$$(1.5) \quad \gamma_{n+\mu} \geq 2\gamma_n \quad \text{and} \quad \gamma_n \leq K\gamma_{n+1} \quad (\gamma_{n+\mu} \leq \frac{1}{2}\gamma_n \quad \text{and} \quad \gamma_{n+1} \leq K\gamma_n)$$

hold for all $n \geq N$.

A sequence $\{\gamma_n\}$ is said to be bounded by blocks if the inequalities


$$(1.6) \quad \alpha_1 \Gamma_m^{(k)} \leq \gamma_n \leq \alpha_2 \Gamma_M^{(k)}, \quad 0 < \alpha_1 \leq \alpha_2 < \infty$$

hold for any $2^k \leq n \leq 2^{k+1}$, $k = 1, 2, \dots$, where

$$\Gamma_m^{(k)} := \min(\gamma_{2^k}, \gamma_{2^{k+1}}) \quad \text{and} \quad \Gamma_M^{(k)} := \max(\gamma_{2^k}, \gamma_{2^{k+1}}).$$

Finally, for a given sequence $\{\gamma_n\}$, $\gamma(x)$ will denote the following function:

$$\gamma(x) = \gamma_n \quad \text{if} \quad x = \frac{1}{n}, n \geq 1; \quad \text{linear on the interval} \quad \left[\frac{1}{n+1}, \frac{1}{n} \right].$$



Title Page

Contents



Go Back

Close

Quit

Page 5 of 14


2. Result

Theorem 2.1. Let $\{b_n\} \in R_0^+ BVS$ and $\{\gamma_n\}$ such that $\gamma_n n^{-2+\varepsilon}$ is quasi-monotone decreasing for some $\varepsilon > 0$. If

$$(2.1) \quad \sum_{n=1}^{\infty} \frac{\gamma_n}{n} b_n < \infty$$

then $\gamma(x)g(x) \in L(0; \pi]$.

Remark 2.1. This result is a generalization of Theorem 1.4 since in (1.3) n^γ is replaced by γ_n and the case $0 \leq \gamma \leq 1$ is extending to $0 \leq \gamma < 2$. Furthermore the sufficiency parts of Theorem 1.2 and 1.3 are also special cases of our Theorem in a few respects: namely the monotonicity of $\{b_n\}$ is changed to the property of $R_0^+ BVS$ and as we will prove later for any slowly varying function $L(x)$ and for $0 \leq \gamma < 2$ the sequence $\{n^\gamma L(n)n^{-2+\varepsilon}\}$ is quasi-monotone decreasing for some $\varepsilon(> 0)$ therefore $n^\gamma L(n)$ can be replaced by $\{\gamma_n\}$ ($0 \leq \gamma < 2$). Moreover using our result it turns out that the convexity and monotonicity conditions on $L(x)$ can be dropped in the case of Theorem 1.2. For example our result contains statement of the type $\sum_{n=2}^{\infty} \frac{1}{n \log n} b_n < \infty \Rightarrow \frac{1}{\log \frac{1}{x}} g(x) \in L[0; \pi]$, too.



Power-Monotone Sequences
and Integrability of
Trigonometric Series

J. Németh

Title Page

Contents



Go Back

Close

Quit

Page 6 of 14

3. Lemmas

We need the following lemmas.

Lemma 3.1. ([8]) *A positive sequence $\{\gamma_n\}$ bounded by blocks is quasi ε -power-monotone decreasing with a certain positive exponent ε if and only if the sequence $\{\gamma_{2^n}\}$ is quasi geometrically decreasing.*

Lemma 3.2. ([7]) *For any positive sequence $\gamma := \{\gamma_n\}$ the inequalities*

$$(3.1) \quad \sum_{n=m}^{\infty} \gamma_n \leq K \gamma_m \quad (m = 1, 2, \dots; K \geq 1),$$

or

$$(3.2) \quad \sum_{n=1}^m \gamma_n \leq K \gamma_m \quad (m = 1, 2, \dots; K \geq 1),$$

hold if and only if γ is quasi geometrically decreasing or increasing, respectively.


Lemma 3.3. *If $\{\gamma_n\}$ has the same property as in Theorem 2.1, then*

$$(3.3) \quad \gamma_n \leq K \cdot \sum_{k=1}^n \frac{\gamma_k}{k} \quad \text{for all } n,$$

and

$$(3.4) \quad \gamma_n \leq K \cdot n^2 \quad \text{for all } n.$$

These statements immediately follow from the definition of $\{\gamma_n\}$.



Title Page

Contents



Go Back

Close

Quit

Page 7 of 14


Lemma 3.4. If $\{b_n\} \in R_0^+ BVS$ and (2.1) is satisfied then

$$(3.5) \quad \sum_{n=1}^{\infty} |b_n - b_{n+1}| \sum_{k=1}^n \frac{\gamma_k}{k} < \infty.$$

Proof. Using the definition of $R_0^+ BVS$ (see (1.2)) we have from (2.1) that

$$\sum_{n=1}^{\infty} \frac{\gamma_n}{n} \sum_{k=n}^{\infty} |b_k - b_{k+1}| < \infty.$$

Now changing the order of summation we get (3.5). □



Power-Monotone Sequences
and Integrability of
Trigonometric Series

J. Németh

Title Page

Contents



Go Back

Close

Quit

Page 8 of 14

4. Proofs

Proof of Theorem 2.1. Since $\{b_n\}$ is of bounded variation, the function $g(x)$ is continuous except perhaps at 0 ([15, p. 4]), so we are concerned only with a neighbourhood of 0.

We shall write $c(x) := 1 - \cos x$. Then by Abel transformation ([3, p. 5]) we have

$$\begin{aligned}\frac{1}{2}g(x) &= \frac{1}{\sin x} \sum_{n=0}^{\infty} b_{n+1} [c\{(n+2)x\} - c(nx)] \\ &= \frac{1}{\sin x} \left[-b_2c(x) + b_3c(2x) + \sum_{n=3}^{\infty} (b_{n-1} - b_{n+1})c(nx) \right].\end{aligned}$$

Since $\sin x \sim x$ as $x \rightarrow 0$, so it is enough to prove the existence of

$$(4.1) \quad \int_0^1 \gamma(x) \cdot \frac{1}{x} \sum_{n=3}^{\infty} |b_{n-1} - b_{n+1}| c(nx) dx.$$


and the integrability of $\gamma(x)c(x)/x$ and $\gamma(x)c(2x)/x$.

Applying Levi's theorem, the existence of (4.1) will follow from

$$(4.2) \quad \sum_{n=3}^{\infty} |b_{n-1} - b_{n+1}| \int_0^1 \gamma(x) \frac{1}{x} c(nx) dx < \infty.$$

Divide the integral $\int_0^1 \gamma(x) \frac{1}{x} c(nx) dx$ into two parts for a fixed n :

$$(4.3) \quad \int_0^1 \gamma(x) \frac{1}{x} c(nx) dx = \int_0^{1/n} \gamma(x) \frac{1}{x} c(nx) dx + \int_{1/n}^1 \gamma(x) \frac{1}{x} c(nx) dx$$



Title Page

Contents



Go Back

Close

Quit

Page 9 of 14


$$= I_1 + I_2.$$

In the estimate of I_1 we use that from the property of $\{\gamma_n\}$ assumed in Theorem 2.1 it follows that $\{\frac{\gamma_{2^n}}{4^n}\}$ is geometrically decreasing and so by Lemma 3.1 (3.1) can be applied for this sequence. So, in the last step using (3.3) also, we get that

$$\begin{aligned}
 (4.4) \quad I_1 &= \int_0^{1/n} \gamma(x) \frac{1}{x} c(nx) dx \\
 &= \int_0^{1/n} \gamma(x) \frac{1}{x} (1 - \cos nx) dx \\
 &= \int_0^{1/n} \gamma(x) x n^2 \frac{1 - \cos nx}{n^2 x^2} dx \\
 &\leq K_1 n^2 \int_0^{1/n} \gamma(x) x dx \\
 &\leq K_2 n^2 \sum_{k=n}^{\infty} \gamma\left(\frac{1}{k}\right) \frac{1}{k^3} \leq K_3 n^2 \sum_{l=\lfloor \log n \rfloor}^{\infty} \gamma\left(\frac{1}{2^l}\right) \frac{1}{4^l} \\
 &\leq K_4 \gamma\left(\frac{1}{n}\right) = K_4 \gamma_n \leq K_5 \sum_{k=1}^n \frac{\gamma_k}{k}.
 \end{aligned}$$

Now we estimate I_2 :

$$\begin{aligned}
 (4.5) \quad I_2 &= \int_{1/n}^1 \gamma(x) \frac{1}{x} c(nx) dx \\
 &\leq 2 \int_{1/n}^1 \gamma(x) \frac{1}{x} dx \leq K \sum_{k=1}^n \gamma\left(\frac{1}{k}\right) \frac{1}{k} = K \sum_{k=1}^n \frac{\gamma_k}{k}.
 \end{aligned}$$



Title Page

Contents



Go Back

Close

Quit

Page 10 of 14

Since (4.3), (4.4), (4.5) with (3.5) give (4.2), so the integral (4.1) exists. Finally the integrability of $\gamma(x)c(x)/x$ and $\gamma(x)c(2x)/x$ can be proved by the same way that was used in the estimate of I_1 in (4.4), applying still (3.4).

Thus the proof of Theorem 2.1 is complete. □

Proof of Remark 2.1. The only fact we need to show that the sequence $\{n^\gamma L(n)n^{-2+\varepsilon}\}$ is quasi monotone decreasing for some $\varepsilon(> 0)$, where $0 \leq \gamma < 2$ and $L(x)$ is an arbitrary slowly varying function. According to Lemma 3.1 it is enough to prove that the sequence $\{n^{\gamma-2}L(n)\}$ is bounded by blocks and that $\{2^{n(\gamma-2)}L(2^n)\}$ is a quasi geometrically decreasing sequence. It is obvious that for the sequence $\{n^{\gamma-2}L(n)\}$ (1.6) is equivalent to the existence of positive constants K_1, K_2 such that

$$(4.6) \quad K_1 \leq \frac{L(2^k + \ell)}{L(2^k)} \leq K_2$$

hold for arbitrary k and $1 \leq \ell \leq 2^k$. But since from the definition of $L(x)$

$$(4.7) \quad \lim_{x \rightarrow \infty} \frac{L(tx)}{L(x)} = 1$$

is uniformly satisfied in the ratio t on the interval $[1, 2]$ (see [1, p. 69]) therefore (4.6) holds.

In order to prove that for the sequence $2^{n(\gamma-2)}L(2^n)$ the properties (1.5) hold it is enough to show that there exist natural numbers μ and N and a constant $K \geq 1$ such that

$$(4.8) \quad (2^{\gamma-2})^{n+\mu}L(2^{n+\mu}) \leq \frac{1}{2}(2^{\gamma-2})^nL(2^n)$$



Title Page

Contents



Go Back

Close

Quit

Page 11 of 14

and


$$(4.9) \quad (2^{\gamma-2})^{n+1} L(2^{n+1}) \leq K(2^{\gamma-2})^n L(2^n)$$

hold if $n > N$.

However, (4.8) is equivalent to

$$(4.10) \quad \frac{L(2^{n+\mu})}{L(2^n)} \leq \frac{1}{2} (2^{2-\gamma})^\mu$$

and if $(2^{2-\gamma})^\mu > 2$ then by using (4.7), (4.10) holds if n is large enough, which gives (4.8). Finally since (4.9) can be obtained by using a similar argument as before, Remark 2.1 is proved. \square



Power-Monotone Sequences
and Integrability of
Trigonometric Series

J. Németh

Title Page

Contents



Go Back


Close

Quit

Page 12 of 14

References

- [1] S. ALJANČIĆ, R. BOJANIĆ AND M. TOMIĆ, Sur, l'intégrabilité de certaines séries trigono- métriques, *Acad. Serbe Sci. Publ. Math.*, **8** (1955), 67–84.
- [2] R.P. BOAS, Jr., Integrability of trigonometrical series III, *Quart. J. Math. (Oxford)*, **3**(2) (1952), 217–221.
- [3] R.P. BOAS, Jr., *Integrability theorems for trigonometric transforms*, Springer-Verlag, Berlin-Heidelberg, 1967.
- [4] YUNG-MING CHEN, On the integrability of functions defined by trigonometrical series, *Math. Z.*, **66** (1956), 9–12.
- [5] YUNG-MING CHEN, Some asymptotic properties of Fourier constants and integrability theorems, *Math. Z.*, **68** (1957), 227–244.
- [6] P. HEYWOOD, On the integrability of functions defined by trigonometric series, *Quart. J. Math. (Oxford)*, **5**(2) (1954), 71–76.
- [7] L. LEINDLER, On the converses of inequality of Hardy and Littlewood, *Acta Sci. Math. (Szeged)*, **58** (1993), 191–196.
- [8] L. LEINDLER AND J. NÉMETH, On the connection between quasi power-monotone and quasi geometrical sequences with application to integrability theorems for power series, *Acta Math. Hungar.*, **68** (1-2) (1995), 7–19.
- [9] L. LEINDLER, A new class of numerical sequences and its applications to sine and cosine series, *Analysis Math.*, **28** (2002), 279–286.



Title Page

Contents




Go Back

Close

Quit

Page 13 of 14

- [10] G.G. LORENTZ, Fourier Koeffizienten und Funktionenklassen, *Math. Z.*, **51** (1948), 135–149.
- [11] S.M. SHAH, Trigonometric series with quasi-monotone coefficients, *Proc. Amer. Math. Soc.*, **13** (1962), 266–273.
- [12] S. O'SHEA, Note on an integrability theorem for sine series, *Quart. J. Math. (Oxford)*, **8**(2) (1957), 279–281.
- [13] G. SUNOUCHI, Integrability of trigonometric series, *J. Math. Tokyo*, **1** (1953), 99–103.
- [14] W.H. YOUNG, On the Fourier series of bounded variation, *Proc. London Math. Soc.*, **12** (1913) 41–70.
- [15] A. ZYGMUND, *Trigonometric series*, Vol. 1, Cambridge Univ. Press, New York, 1959.



Power-Monotone Sequences
and Integrability of
Trigonometric Series

J. Németh

Title Page

Contents



Go Back

Close

Quit

Page 14 of 14