

Journal of Inequalities in Pure and Applied Mathematics

π AND SOME OTHER CONSTANTS

ANTHONY SOFO

School of Computer Science and Mathematics
Victoria University
PO Box 14428, MCMC 8001
VIC, Australia.

*E*Mail: anthony.sofa@vu.edu.au
*U*RL: <http://rgmia.vu.edu.au/sofo/>

©2000 Victoria University
ISSN (electronic): 1443-5756
123-05



volume 6, issue 5, article 138,
2005.

*Received 18 March, 2005;
accepted 26 October, 2005.*

Communicated by: J. Sándor

Abstract

Contents




Home Page

Go Back

Close

Quit



Abstract

We consider a particular definite integral and reduce it to hypergeometric form. Then we develop identities for some numerical constants and the number π .

2000 Mathematics Subject Classification: Primary 33B15; Secondary 33C05.

Key words: Hypergeometric summation, Definite integration, Binomial type series.

This paper is based on the talk given by the author within the “International Conference of Mathematical Inequalities and their Applications, I”, December 06-08, 2004, Victoria University, Melbourne, Australia [<http://rgmia.vu.edu.au/conference>]

Contents

1	Introduction	3
2	The Integral	8
3	Illustrative Examples	19
4	Some Estimates	24
	References	

π and Some Other Constants

Anthony Sofo

Title Page

Contents



Go Back

Close

Quit

Page 2 of 31

1. Introduction

π , is a real number and defined as the ratio of the circumference of a circle to its diameter. π 's digits have many interesting properties and π has a rich history dating back to the time of the Babylonians and Egyptians, circa 2000 B.C. The Bible has two references, I Kings 7:23 and Chronicles 4:2, to Pi and gives it an estimate of about 3. The Babylonians gave an estimate of π as $3\frac{1}{8}$ and the Egyptians also obtained $3\frac{13}{81}$ as an estimate. We know that $3\frac{1}{8} < \pi < 3\frac{13}{81}$.

Many researchers have increasingly calculated the number of decimal places for the value of π . Apparently in September 2002, Dr. Kanada and his team, from the University of Tokyo, calculated π to 1.2411 trillion digits, indeed a world record. Many, many formulae also exist for the representation of π , and a collection of these formulae is listed below.

Vieta (~1593), see [7], gave an infinite product of nested radicals for the reciprocal of π , namely


$$\frac{2}{\pi} = \sqrt{\frac{1}{2}} \sqrt{\frac{1}{2} + \frac{1}{2} \sqrt{\frac{1}{2}}} \sqrt{\frac{1}{2} + \frac{1}{2} \sqrt{\frac{1}{2} + \frac{1}{2} \sqrt{\frac{1}{2}}}} \cdots$$

Wallis (~1650), see [7], gave

$$\frac{\pi}{4} = \prod_{r=1}^{\infty} \left(1 - \frac{1}{(2r+1)^2} \right).$$

Leibnitz (~1670), see [7], gave the very slow converging series

$$\frac{\pi}{4} = \sum_{r=0}^{\infty} \frac{(-1)^r}{2r+1}.$$



π and Some Other Constants

Anthony Sofo

Title Page

Contents



Go Back

Close

Quit

Page 3 of 31

Newton (~1666) admitted being ashamed at having computed π to fifteen decimal places, by the formula

$$\begin{aligned}\pi &= \frac{3\sqrt{3}}{4} + 24 \int_0^{\frac{1}{4}} \sqrt{x - x^2} dx \\ &= \frac{3\sqrt{3}}{4} + 2 - \frac{3}{4} \left\{ \frac{1}{5} + \frac{1}{7 \cdot 2^4} + \frac{1}{9 \cdot 2^7} + \frac{5}{11 \cdot 2^{12}} + \dots \right\} \\ &= \frac{3\sqrt{3}}{4} + 2 - \frac{3}{4} \sum_{k=0}^{\infty} \binom{2k}{k} \frac{1}{16^k (k+1)(2k+5)}\end{aligned}$$

Euler (~1750) gave many representations of π including:

$$\frac{\pi}{2} = \lim_{n \rightarrow \infty} \left[\frac{1}{n} + \frac{1}{6n^2} + 4n \sum_{j=1}^n \frac{1}{n^2 + j^2} \right].$$


Ramanujan (~1914) has also given many representations of π and its reciprocal, including:

$$\frac{1}{\pi} = \frac{2\sqrt{2}}{3^4 \cdot 11^2} \sum_{r=0}^{\infty} \frac{(4r)! (1103 + (2 \cdot 5 \cdot 7 \cdot 13 \cdot 29) r)}{(r!)^4 (2^2 \cdot 3^2 \cdot 11)^{4r}}.$$

For a fuller account of Ramanujan's work the interested reader is referred to the books of Berndt [4].

Comtet (1974) gave

$$\pi^4 = \frac{2^3 \cdot 3^4 \cdot 5}{17} \sum_{r=1}^{\infty} \frac{1}{r^4 \binom{2r}{r}}.$$



π and Some Other Constants

Anthony Sofo

Title Page

Contents



Go Back

Close

Quit

Page 4 of 31

D. and G. Chudnovsky (1989) gave

$$\frac{1}{\pi} = 12 \sum_{r=0}^{\infty} \frac{(-1)^r (6r)!}{(r!)^3 (3r)!} \cdot \frac{13 \cdot 1045493 + 2 \cdot 3^2 \cdot 7 \cdot 11 \cdot 19 \cdot 127 \cdot 163r}{(2^{18} \cdot 3^3 \cdot 5^3 \cdot 23^3 \cdot 29^3)^{r+\frac{1}{2}}}.$$

Bailey, Borwein and Plouffe (1996) gave

$$(1.1) \quad \pi = \sum_{r=0}^{\infty} \frac{1}{16^r} \left[\frac{4}{8r+1} - \frac{2}{8r+4} - \frac{1}{8r+5} - \frac{1}{8r+6} \right].$$

Bellard (1997) gave

$$\pi = \frac{1}{3^2 \cdot 5^2 \cdot 11 \cdot 13 \cdot 23} \left[\sum_{r=1}^{\infty} \frac{3P(r)}{\binom{7r}{2r} 2^{r-1}} - 2^4 \cdot 5 \cdot 254741 \right],$$

where

$$P(r) = -13 \cdot 29 \cdot 2351653r^5 + 193 \cdot 16193509r^4 - 5^2 \cdot 7 \cdot 79 \cdot 212873r^3 \\ + 5 \cdot 206392559r^2 - 2 \cdot 98441137r + 2^3 \cdot 13 \cdot 43 \cdot 2459.$$

Lupas [10], gave

$$\pi = 4 + \sum_{k=1}^{\infty} (-16)^k \frac{\binom{2k}{k} (40k^2 + 16k + 1)}{\binom{4k}{2k}^2 2k (4k + 1)^2}.$$

The original Lupas formula contained a minor misprint which has been corrected here.



π and Some Other Constants

Anthony Sofo

Title Page

Contents



Go Back

Close

Quit

Page 5 of 31

Borwein and Girgensohn (2003), wrote

$$\pi = \ln 4 + 10 \sum_{r=1}^{\infty} \frac{1}{2^r r \binom{3r}{r}}.$$

Sofa [16] has given

$$\frac{\pi}{2} = \sqrt{2} + \ln(\sqrt{2} - 1) + \sum_{r=0}^{\infty} \binom{4r}{2r} \frac{1}{16^r (2r+1)(4r+1)}$$

and

$$\pi^2 = \frac{1308}{135} + \frac{12}{5} \sum_{r=1}^{\infty} \frac{4^r}{r^2 \binom{2r}{r} (r+1)(2r+1)(2r+3)}.$$

Many other representations of π exist, including the famous Machin-type formulae such as


$$\frac{\pi}{4} = 4 \arctan\left(\frac{1}{5}\right) - \arctan\left(\frac{1}{239}\right).$$

There are also many other connections of π and other mathematical constants, including:

$$e^{i\pi} + 1 = 0,$$

$$\pi^3 + 8\pi = 56 - 8 \sum_{r=1}^{\infty} \frac{(-1)^r}{r(r+1)(2r+1)^3},$$

$$\frac{\pi^2}{6} = 3 \ln^2 \phi + \sum_{r=0}^{\infty} \frac{(-1)^r (r!)^2}{(2r)! (2r+1)^2}, \text{ where } \phi \text{ is the Golden ratio}$$



π and Some Other Constants

Anthony Sofa

Title Page

Contents



Go Back

Close

Quit

Page 6 of 31


and

$$\pi^2 = -12e^3 \sum_{r=1}^{\infty} \frac{1}{r^2} \cos \left(\frac{9}{r\pi + \sqrt{(r\pi)^2 - 3^2}} \right).$$

A selection of some series expansion representations of π including some of the above is given by Sebah and Gourdon [15].

There are other nice articles and books relating to π including [2, 3, 6, 7, 8, 11, 12].

The aim of this paper is to derive representations of π , as well as some other constants, by the consideration of a particular definite integral. The following integral will now be investigated.



π and Some Other Constants

Anthony Sofo

Title Page

Contents



Go Back

Close

Quit

Page 7 of 31

2. The Integral

Theorem 2.1. For k, m and α real positive numbers and $a \geq 1$, then

$$(2.1) \quad I(a, k, m, \alpha) = \int_0^{\frac{1}{a}} \frac{x^m}{(1-x^k)^\alpha} dx$$

$$(2.2) \quad = \sum_{r=0}^{\infty} \frac{(\alpha)_r}{r! (rk + m + 1) a^{rk+m+1}}$$

$$(2.3) \quad = T_0 {}_2F_1 \left[\begin{matrix} \frac{m+1}{k}, \alpha \\ \frac{m+1+k}{k} \end{matrix} \middle| \frac{1}{a^k} \right]$$

$$(2.4) \quad = \frac{1}{k} \left[B\left(1-\alpha, \frac{m+1}{k}\right) - B\left(1-a^{-k}; 1-\alpha, \frac{m+1}{k}\right) \right],$$

where


$$(2.5) \quad T_0 = \frac{1}{(m+1) a^{m+1}},$$

$(b)_s$ is Pochhammer's symbol defined by

$$(2.6) \quad \begin{cases} (b)_0 = 1 \\ (b)_s = b(b+1) \cdots (b+s-1) = \frac{\Gamma(b+s)}{\Gamma(b)}, \end{cases}$$

$\Gamma(b)$ is the classical Gamma function, ${}_2F_1[\cdot, \cdot]$ is the Gauss Hypergeometric function, $B(s, t)$ is the classical Beta function and

$$B(z; s, t) = \int_0^z u^{s-1} (1-u)^{t-1} du$$



π and Some Other Constants

Anthony Sofo

Title Page

Contents



Go Back

Close

Quit

Page 8 of 31

is the incomplete Beta function.

Proof.

$$\begin{aligned} I(a, k, m, \alpha) &= \int_0^{\frac{1}{a}} \frac{x^m}{(1-x^k)^\alpha} dx \\ &= \int_0^{\frac{1}{a}} \sum_{r=0}^{\infty} (-1)^r \binom{-\alpha}{r} x^{kr+m}. \end{aligned}$$

where we have utilised

$$\frac{1}{(1+z)^\beta} = \sum_{r=0}^{\infty} \binom{-\beta}{r} z^r$$

and from

$$\binom{-\beta}{r} = (-1)^r \binom{\beta+r-1}{r} = \frac{(-1)^r (\beta)_r}{r!}$$


we have

$$I(a, k, m, \alpha) = \int_0^{\frac{1}{a}} \sum_{r=0}^{\infty} \frac{(\alpha)_r}{r!} x^{kr+m}.$$

Reversing the order of integration and summation and substituting the integration limits we obtain the result (2.2). The result (2.4) is obtained by the use of the substitution $u = 1 - x^k$.

Binomial sums are intrinsically associated with generalised hypergeometric functions and if from (2.2) we let

$$(2.7) \quad T_r = \frac{(\alpha)_r}{r! (rk + m + 1) a^{rk+m+1}},$$



π and Some Other Constants

Anthony Sofo

Title Page

Contents



Go Back

Close

Quit

Page 9 of 31

then we get the ratio

$$(2.8) \quad \frac{T_{r+1}}{T_r} = \frac{(\alpha + r) \left(r + \frac{m+1}{k}\right)}{a^k (r+1) \left(r + \frac{m+1+k}{k}\right)}$$

where T_0 is given by (2.5). From (2.5) and (2.2) we can write


$$\begin{aligned} I(a, k, m, \alpha) &= T_0 {}_2F_1 \left[\begin{matrix} \frac{m+1}{k}, \alpha \\ \frac{m+1+k}{k} \end{matrix} \middle| \frac{1}{a^k} \right] \\ &= \frac{1}{k} \left[B \left(1 - \alpha, \frac{m+1}{k} \right) - B \left(1 - a^{-k}; 1 - \alpha, \frac{m+1}{k} \right) \right], \end{aligned}$$

which is the result (2.3). We can now match (2.2) and (2.3) so that

$$\begin{aligned} \sum_{r=0}^{\infty} \frac{(\alpha)_r}{r! (rk + m + 1) a^{rk+m+1}} \\ &= T_0 {}_2F_1 \left[\begin{matrix} \frac{m+1}{k}, \alpha \\ \frac{m+1+k}{k} \end{matrix} \middle| \frac{1}{a^k} \right] \\ &= \frac{1}{k} \left[B \left(1 - \alpha, \frac{m+1}{k} \right) - B \left(1 - a^{-k}; 1 - \alpha, \frac{m+1}{k} \right) \right], \end{aligned}$$

and the infinite series converges for $|a^{-k}| < 1$. □

In a previous paper, Sofo [17] has utilised (2.1) for the case $a = 1$ and



π and Some Other Constants

Anthony Sofo

Title Page

Contents



Go Back

Close

Quit

Page 10 of 31

developed identities for π and other constants, such as

$$\pi = \frac{15p! \left(\sqrt{30 + 6\sqrt{5}} + 1 - \sqrt{5} \right)}{4 \binom{7}{30}_p} \sum_{r=0}^{\infty} \frac{\binom{7}{30}_r}{r! (30r + 30p + 7)},$$

for $p = 0, 1, 2, 3, \dots$

Remark 1. Bailey, Borwein, Borwein and Plouffe see [7] utilised (2.1) for $a = \sqrt{2}$, $\alpha = 1$, $k = 8$ and $m = \beta - 1$, $\beta < 8$ to prove the new formula (1.1). Subsequently Hirschhorn [9] has shown that (1.1) can be obtained from standard integration procedures.


The following lemma will be useful in the consideration of the integral (2.1).

Lemma 2.2. For $p = 0, 1, 2, \dots$ the following well-known identities are given by Beyer [5]

$$(2.9) \quad \sin^{2p+1} x = \frac{(-1)^p}{2^{2p}} \sum_{j=0}^p (-1)^j \binom{2p+1}{p} \sin((2p+1-2j)x)$$

and

$$(2.10) \quad \sin^{2p+2} x = \frac{1}{2^{2p+1}} \left[\binom{2p+1}{p} - (-1)^p \sum_{j=0}^p (-1)^j \binom{2p+2}{p} \cos((2p+2-2j)x) \right].$$



π and Some Other Constants

Anthony Sofo

Title Page

Contents



Go Back

Close

Quit

Page 11 of 31

The integral (2.1) can be simplified as follows. Consider the case $k = 2$; then from (2.1)

$$(2.11) \quad I(a, m, 2, \alpha) = \int_0^{\frac{1}{a}} \frac{x^m}{(1-x^2)^\alpha} dx.$$

The following lemma concerns the integral (2.11).

Lemma 2.3.


(i) For $m = 2p + 1$, $2\alpha = 2q + 1$; $p = 0, 1, 2, \dots$, and $q = 0, 1, 2, \dots$

$$(2.12) \quad I(a, m, 2, \alpha)$$

$$\begin{aligned} &= \int_0^{\frac{1}{a}} \frac{x^{2p+1}}{(1-x^2)^{\frac{2q+1}{2}}} dx = \int_0^{\theta^*} \frac{\sin^{2p+1} \theta}{\cos^{2q} \theta} d\theta \\ &= \frac{1}{2} \left[B\left(\frac{1-2q}{2}, p+1\right) - B\left(1-a^{-2}; \frac{1-2q}{2}, p+1\right) \right] \\ &= \frac{1}{2q-1} \cdot \frac{\left(\frac{1}{a}\right)^{2p+2}}{\cos^{2q-1} \theta^*} + \sum_{j=1}^{q-1} \frac{(-1)^j \left(\frac{1}{a}\right)^{2p+2}}{\cos^{2(q-j)-1} \theta^*} \prod_{i=1}^j \frac{2(p-q+i)-1}{2(q-i)+1} \\ &\quad + (-1)^q \prod_{i=1}^q \frac{2(p-q+i)-1}{2(q-i)+1} \left[\frac{(-1)^p}{2^{2p}} \right. \\ &\quad \left. \times \sum_{s=0}^p \frac{(-1)^s}{2p-2s+1} \binom{2p+1}{s} \{1 - \cos((2p-2s+1)\theta^*)\} \right]. \end{aligned}$$

where $x = \sin \theta$ and $\theta^* = \arcsin\left(\frac{1}{a}\right)$.

Note, the middle term in the right hand side of (2.12) is identically zero for $q = 1$ and only the last sum applies for $q = 0$.



Title Page

Contents



Go Back

Close

Quit

(ii) For $m = 2p + 2$, $2\alpha = 2q + 1$; $p = 0, 1, 2, \dots$, and $q = 0, 1, 2, \dots$

(2.13) $I(a, m, 2, \alpha)$


$$\begin{aligned}
 &= \int_0^{\frac{1}{a}} \frac{x^{2p+2}}{(1-x^2)^{\frac{2q+1}{2}}} dx = \int_0^{\theta^*} \frac{\sin^{2p+2} \theta}{\cos^{2q} \theta} d\theta \\
 &= \frac{1}{2} \left[B\left(\frac{1-2q}{2}, p + \frac{3}{2}\right) - B\left(1-a^{-2}; \frac{1-2q}{2}, p + \frac{3}{2}\right) \right] \\
 &= \frac{1}{2q-1} \cdot \frac{\left(\frac{1}{a}\right)^{2p+3}}{\cos^{2q-1} \theta^*} + \sum_{j=1}^{q-1} \frac{(-1)^j \left(\frac{1}{a}\right)^{2p+3}}{\cos^{2(q-j)-1} \theta^*} \prod_{i=1}^j \frac{2(p-q+i+1)}{2(q-i)+1} \\
 &\quad + (-1)^q \prod_{i=1}^q \frac{2(p-q+i+1)}{2(q-i)+1} \left\{ \frac{1}{2^{2p+1}} \left[\binom{2p+1}{p} \theta^* \right. \right. \\
 &\quad \left. \left. - (-1)^p \sum_{s=0}^p \frac{(-1)^s}{2p-2s+2} \binom{2p+2}{s} \sin((2p-2s+2)\theta^*) \right] \right\}.
 \end{aligned}$$

where $x = \sin \theta$ and $\theta^* = \arcsin\left(\frac{1}{a}\right)$.

Note, the middle term in the right hand side of (2.13) is identically zero for $q = 1$ and only the last two terms apply for $q = 0$.

Proof. (i) From

$$I(a, m, 2, \alpha) = \int_0^{\theta^*} \frac{\sin^{2p+1} \theta}{\cos^{2q} \theta} d\theta,$$



π and Some Other Constants

Anthony Sofo

Title Page

Contents




Go Back

Close

Quit

Page 13 of 31



Title Page

Contents



Go Back

Close

Quit

Page 14 of 31

integrating by parts once leads to

$$\begin{aligned}
 I(a, m, 2, \alpha) &= \frac{1}{2q-1} \left[\frac{\sin^{2p+2} \theta}{\cos^{2q-1} \theta} \right]_0^{\theta^*} - (2p+3-2q) \int_0^{\theta^*} \frac{\sin^{2p+1} \theta}{\cos^{2q-2} \theta} d\theta \\
 &= \frac{\left(\frac{1}{a}\right)^{2p+2}}{(2q-1) \cos^{2q-1} \theta^*} - \frac{2p-2q+3}{2q-1} \int_0^{\theta^*} \frac{\sin^{2p+1} \theta}{\cos^{2q-2} \theta} d\theta,
 \end{aligned}$$

and repeated integration by parts gives us

$$\begin{aligned}
 (2.14) \quad I(a, m, 2, \alpha) &= \frac{\left(\frac{1}{a}\right)^{2p+2}}{(2q-1) \cos^{2q-1} \theta^*} \\
 &+ \sum_{j=1}^{q-1} \frac{(-1)^j \left(\frac{1}{a}\right)^{2p+2}}{\cos^{2(q-j)-1} \theta^*} \prod_{i=1}^j \frac{2(p-q+i)+1}{2(q-i)+1} \\
 &\quad + (-1)^q \prod_{i=1}^q \frac{2(p-q+i)+1}{2(q-i)+1} \int_0^{\theta^*} \sin^{2p+1} \theta d\theta.
 \end{aligned}$$

Substituting (2.9), from Lemma 2.2, into (2.14) and integrating, results in (2.12) hence part (i) of the lemma is proved.

(ii) The proof of part (ii) of the lemma follows the same footsteps as part (i).

□

The following lemma is given and will be useful in the simplification of the left hand side of (2.17) and (2.18).

Lemma 2.4. For $r = 0, 1, 2, \dots$ and $q = 1, 2, 3, \dots$ then

$$(2.15) \quad \frac{(q + \frac{1}{2})_r}{r!} = \frac{1}{4^r} \binom{2r}{r} \prod_{\rho=0}^{q-1} \frac{2r + 2\rho + 1}{2\rho + 1},$$

and for $q = 0$,

$$(2.16) \quad \frac{(\frac{1}{2})_r}{r!} = \frac{1}{4^r} \binom{2r}{r}.$$

Proof. For $q = 0$, then

$$\frac{(\frac{1}{2})_r}{r!} = \frac{\Gamma(r + \frac{1}{2})}{r! \Gamma(\frac{1}{2})} = \frac{1}{4^r} \binom{2r}{r}.$$

This result is well known and is also given by Wilf [18].

For $q \geq 1$, let


$$P(q) := \frac{(q + \frac{1}{2})_r}{r!} = \frac{1}{4^r} \binom{2r}{r} \prod_{\rho=0}^{q-1} \frac{2r + 2\rho + 1}{2\rho + 1}$$

then

$$P(1) = \frac{(\frac{3}{2})_r}{r!} = \frac{(2r + 1) \Gamma(\frac{1}{2}) (\frac{1}{2})_r}{2 \Gamma(\frac{3}{2}) r!},$$

and from (2.16)

$$P(1) = \frac{2r + 1}{4^r} \binom{2r}{r},$$



π and Some Other Constants

Anthony Sofo

Title Page

Contents



Go Back

Close

Quit

Page 15 of 31

which satisfies the right hand of (2.15) for $q = 1$.

Consider

$$\begin{aligned} P(q+1) &= \frac{(q + \frac{3}{2})_r}{r!} = \frac{\Gamma(q + r + \frac{3}{2})}{r! \Gamma(q + \frac{3}{2})} \\ &= \left(\frac{2q + 2r + 1}{2q + 1} \right) \frac{\Gamma(q + r + \frac{1}{2})}{r! \Gamma(q + \frac{1}{2})} \\ &= \left(\frac{2q + 2r + 1}{2q + 1} \right) \frac{(q + \frac{1}{2})_r}{r!} \end{aligned}$$

and from (2.15)

$$\begin{aligned} P(q+1) &= \left(\frac{2q + 2r + 1}{2q + 1} \right) \frac{1}{4^r} \binom{2r}{r} \prod_{\rho=0}^{q-1} \frac{2r + 2\rho + 1}{2\rho + 1} \\ &= \frac{1}{4^r} \binom{2r}{r} \prod_{\rho=0}^q \frac{2r + 2\rho + 1}{2\rho + 1} \end{aligned}$$


hence the lemma is proved. □

For $k = 2$ the following theorem now applies.


Theorem 2.5.

(i) For $m = 2p + 1$, $2\alpha = 2q + 1$; $p = 0, 1, 2, \dots$, $q = 0, 1, 2, \dots$,

$$(2.17) \quad \sum_{r=0}^{\infty} \frac{(q + \frac{1}{2})_r}{r! (2r + 2p + 2) a^{2r+2p+2}}$$



Title Page	
Contents	
◀◀	▶▶
◀	▶
Go Back	
Close	
Quit	
Page 16 of 31	



Title Page

Contents



Go Back

Close

Quit

Page 17 of 31

$$\begin{aligned}
 &= \sum_{r=0}^{\infty} \binom{2r}{r} \frac{\prod_{\rho=0}^{q-1} \frac{2r+2\rho+1}{2\rho+1}}{4^r (2r+2p+2) a^{2r+2p+2}} \\
 &= T_0^* {}_2F_1 \left[\begin{matrix} p+1, \frac{1}{2}(2q+1) \\ p+2 \end{matrix} \middle| \frac{1}{a^2} \right] \\
 &= \frac{1}{2} \left[B \left(\frac{1-2q}{2}, p+1 \right) - B \left(1-a^{-2}; \frac{1-2q}{2}, p+1 \right) \right] \\
 &= \frac{\left(\frac{1}{a}\right)^{2p+2}}{(2q-1) \cos^{2q-1} \theta^*} + \sum_{j=1}^{q-1} \frac{(-1)^j \left(\frac{1}{a}\right)^{2p+2}}{\cos^{2(q-j)-1} \theta^*} \prod_{i=1}^j \frac{2(p-q+i)+1}{2(q-i)+1} \\
 &\quad + (-1)^q \prod_{i=1}^q \frac{2(p-q+i)+1}{2(q-i)+1} \\
 &\quad \times \left[\frac{(-1)^p}{2^{2p}} \sum_{s=0}^p \frac{(-1)^s}{2p-2s+1} \binom{2p+1}{s} \{1 - \cos((2p-2s+1)\theta^*)\} \right]
 \end{aligned}$$


where

$$T_0^* = \frac{1}{(2p+2) a^{2p+2}}$$

and $\theta^* = \arcsin\left(\frac{1}{a}\right)$.

(ii) For $m = 2p + 2$, $2\alpha = 2q + 1$; $p = 0, 1, 2, \dots$, $q = 0, 1, 2, \dots$,

$$(2.18) \quad \sum_{r=0}^{\infty} \frac{\left(q + \frac{1}{2}\right)_r}{r! (2r+2p+3) a^{2r+2p+3}}$$



Title Page

Contents



Go Back

Close

Quit

Page 18 of 31

$$\begin{aligned}
 &= \sum_{r=0}^{\infty} \binom{2r}{r} \frac{\prod_{\rho=0}^{q-1} \frac{2r+2\rho+1}{2\rho+1}}{4^r (2r+2p+3) a^{2r+2p+3}} \\
 &= T_0^{\nabla} {}_2F_1 \left[\begin{matrix} \frac{1}{2}(2p+3), \frac{1}{2}(2q+1) \\ \frac{1}{2}(2p+5) \end{matrix} \middle| \frac{1}{a^2} \right] \\
 &= \frac{1}{2} \left[B \left(\frac{1-2q}{2}, p + \frac{3}{2} \right) - B \left(1-a^{-2}; \frac{1-2q}{2}, p + \frac{3}{2} \right) \right] \\
 &= \frac{\left(\frac{1}{a}\right)^{2p+3}}{(2q-1) \cos^{2q-1} \theta^*} + \sum_{j=1}^{q-1} \frac{(-1)^j \left(\frac{1}{a}\right)^{2p+3}}{\cos^{2(q-j)-1} \theta^*} \prod_{i=1}^j \frac{2(p-q+i+1)}{2(q-i)+1} \\
 &\quad + (-1)^q \prod_{i=1}^q \frac{2(p-q+i+1)}{2(q-i)+1} \left\{ \frac{1}{2^{2p+1}} \left[\binom{2p+1}{p} \theta^* \right. \right. \\
 &\quad \left. \left. - (-1)^p \sum_{s=0}^p \frac{(-1)^s}{2p-2s+2} \binom{2p+2}{s} \sin((2p-2s+2)\theta^*) \right] \right\},
 \end{aligned}$$

where

$$T_0^{\nabla} = \frac{1}{(2p+3) a^{2p+3}}$$

and $\theta^* = \arcsin\left(\frac{1}{a}\right)$.

The proof of Theorem 2.5 follows directly from Lemma 2.3 , (2.14) and Lemma 2.4.

Some examples will now be given expressing π and other constants in terms of an infinite series.

3. Illustrative Examples

Example 3.1. From (2.17) with $q = 2$, $a = 2$ and $\theta^* = \frac{\pi}{6}$, we have


$$\begin{aligned} & \frac{2}{\sqrt{3}} \left(\frac{8}{3} - 2(2p-1) \right) + 8(-1)^p (2p-1)(2p+1) \\ & \times \sum_{s=0}^p \frac{(-1)^s}{2p-2s+1} \binom{2p+1}{s} \left\{ 1 - \cos(2p-2s+1) \frac{\pi}{6} \right\} \\ & = \sum_{r=0}^{\infty} \binom{2r}{r} \frac{(2r+1)(2r+3)}{16^r (r+p+1)}. \end{aligned}$$

Hence, for $p = 7$,

$$\sqrt{3} = \frac{3 \cdot 2^{22}}{7 \cdot 163 \cdot 6367} - \frac{3^2 \cdot 11}{2^6 \cdot 7 \cdot 163 \cdot 6376} \sum_{r=0}^{\infty} \binom{2r}{r} \frac{(2r+1)(2r+3)}{16^r (r+8)}.$$

Example 3.2. From (2.18) with $q = 2$, $a = \frac{2}{\sqrt{3}}$ and $\theta^* = \frac{\pi}{3}$, we have

$$\begin{aligned} & 8 - 4p + \frac{16p(p+1)}{(\sqrt{3})^{2p+3}} \left[\binom{2p+1}{p} \frac{\pi}{3} \right. \\ & \left. - (-1)^p \sum_{s=0}^p \frac{(-1)^s}{2p-2s+2} \binom{2p+2}{s} \sin(2p-2s+2) \frac{\pi}{3} \right] \\ & = \sum_{r=0}^{\infty} \binom{2r}{r} \frac{(2r+1)(2r+3)}{(2r+2p+3)} \left(\frac{3}{16} \right)^r \end{aligned}$$



π and Some Other Constants

Anthony Sofo

Title Page

Contents



Go Back

Close

Quit

Page 19 of 31

or rearranging,

$$\frac{\pi}{3} = \frac{3^{p+\frac{3}{2}}}{16p(p+1) \binom{2p+1}{p}} \left\{ 4p - 8 + \sum_{r=0}^{\infty} \binom{2r}{r} \frac{(2r+1)(2r+3)}{(2r+2p+3)} \left(\frac{3}{16}\right)^r \right\} + \frac{(-1)^p}{\binom{2p+1}{p}} \sum_{s=0}^p \frac{(-1)^s}{2p-2s+2} \binom{2p+2}{s} \sin(2p-2s+2) \frac{\pi}{3},$$

and for $p = 2$

$$\frac{20\sqrt{3}\pi}{81} = 1 + \frac{1}{16} \sum_{r=0}^{\infty} \binom{2r}{r} \frac{(2r+1)(2r+3)}{2r+7} \left(\frac{3}{16}\right)^r.$$

For $p = 2, a = \sqrt{2}$

$$\pi = \frac{8}{3} + \frac{1}{3\sqrt{2}} \sum_{r=0}^{\infty} \binom{2r}{r} \frac{(2r+1)(2r+3)}{2r+5} \left(\frac{1}{8}\right)^r.$$

Example 3.3. For $q = 3, p = 2$ and $a = \sqrt{2}$,

$$\pi = \frac{52}{15} - \frac{1}{30\sqrt{2}} \sum_{r=0}^{\infty} \binom{2r}{r} \frac{(2r+1)(2r+3)(2r+5)}{2r+7} \left(\frac{1}{8}\right)^r.$$

Example 3.4. For $q = 4, p = 3$ and $a = 2$,

$$\pi = \frac{1712\sqrt{3}}{945} + \frac{1}{8960} \sum_{r=0}^{\infty} \binom{2r}{r} \frac{(2r+1)(2r+3)(2r+5)(2r+7)}{(2r+9)16^r}.$$



π and Some Other Constants

Anthony Sofo

Title Page

Contents




Go Back

Close

Quit

Page 20 of 31



Title Page

Contents



Go Back

Close

Quit

Page 21 of 31

Example 3.5. For $q = 5$, $p = 0$ and $a = \sqrt{5}$,

$$\sqrt{5} = \frac{2^{11}}{3 \cdot 5 \cdot 193 \cdot 2731} \sum_{r=0}^{\infty} \binom{2r}{r} \frac{(2r+1)(2r+5)(2r+7)(2r+9)(2r+11)}{20^r}.$$

Example 3.6. For $q = 6$, $p = 5$ and $a = \sqrt{2}$,

$$\pi = \frac{2^3 \cdot 1289}{5 \cdot 7 \cdot 9 \cdot 11} + \frac{1}{5 \cdot 16 \cdot 829 \cdot \sqrt{2}} \sum_{r=0}^{\infty} \left[\binom{2r}{r} \times \frac{(2r+1)(2r+3)(2r+5)(2r+7)(2r+9)(2r+11)}{(2r+13)8^r} \right].$$

Example 3.7. For $q = 4$ ($\alpha = \frac{9}{2}$), $p = 59$ ($m = 120$) and $a = 2$,

$$\pi = \frac{\Omega_1}{\Omega_2 \sqrt{3}} + \frac{1}{\Omega_3} \sum_{r=0}^{\infty} \binom{2r}{r} \frac{(2r+1)(2r+3)(2r+5)(2r+7)}{(2r+121)16^r},$$

where

$$\begin{aligned} \Omega_1 &= 15604102274295581508678435968572864501995513795052733 \\ &= (46042305118509401202197)(338907929004245243145594887689), \end{aligned}$$

$$\begin{aligned} \Omega_2 &= 2^6 \cdot 3^5 \cdot 5^2 \cdot 7^2 \cdot 11 \cdot 13 \cdot 17 \cdot 19 \cdot 23 \cdot 29 \cdot 31 \cdot 37 \cdot 41 \cdot 43 \cdot 47 \cdot 53 \cdot 59 \\ &\quad \cdot 61 \cdot 67 \cdot 71 \cdot 73 \cdot 79 \cdot 83 \cdot 89 \cdot 97 \cdot 101 \cdot 103 \cdot 107 \cdot 109 \cdot 113 \end{aligned}$$

and

$$\Omega_3 = 2^{11} \cdot 3^3 \cdot 5 \cdot 7 \cdot 13 \cdot 17 \cdot 19 \cdot 23 \cdot 29 \cdot 31 \cdot 37 \cdot 59 \\ \cdot 61 \cdot 67 \cdot 71 \cdot 73 \cdot 79 \cdot 83 \cdot 89 \cdot 97 \cdot 101 \cdot 103 \cdot 107 \cdot 109 \cdot 113.$$

In this case the first term of the sum gives π accurate to over forty decimal places.

Other particular values of constants may be obtained from (2.13).

Example 3.8. For $a = 2$, $m = 10$, $k = 1$ and $\alpha = 7$

$$\ln 2 = \frac{27947}{2^7 \cdot 3^2 \cdot 5 \cdot 7} + \frac{1}{2^{12} \cdot 3 \cdot 5 \cdot 7} \sum_{r=0}^{\infty} \binom{r+6}{r} \frac{1}{(r+11) 2^r}.$$

It is of some interest to note that from (2.18) for $m = 0$ and $\alpha = w > 1$, integer, we may eventually write, after integration by parts, and using (2.2)

$$\ln \left(\frac{a+1}{a-1} \right) = \frac{2^w (w-1)!}{(2w-3)!!} \left[\sum_{r=0}^{\infty} \binom{r+w-1}{r} \frac{1}{(2r+1) a^{2r+1}} \right. \\ \left. - \frac{1}{a(2w-1)} \sum_{j=1}^{w-1} \frac{1}{2^j} \left(\frac{a^2}{a^2-1} \right)^{w-j} \prod_{\nu=1}^j \frac{2w-2\nu+1}{w-\nu} \right],$$

where $(2w-3)!! = (2w-3)(2w-5)\cdots 5 \cdot 3 \cdot 1$, and $a > 1$.

For $a = 11$ and $w = 7$, we have

$$\ln \left(\frac{6}{5} \right) = \frac{2^{11}}{3 \cdot 7 \cdot 11^2} \sum_{r=0}^{\infty} \binom{r+6}{r} \frac{1}{(2r+1) 121^r} - \frac{11 \cdot 179 \cdot 17047711}{2^9 \cdot 3^8 \cdot 5^6}.$$



π and Some Other Constants

Anthony Sofo

Title Page

Contents



Go Back


Close

Quit

Page 22 of 31

Remark 2. In the degenerative case of $w = 1$ then we obtain the well-known formula as listed in Abramowitz and Stegun [1], namely

$$\ln \left(\frac{a+1}{a-1} \right) = 2 \sum_{r=0}^{\infty} \frac{1}{(2r+1) a^{2r+1}}.$$



π and Some Other Constants

Anthony Sofo

Title Page

Contents



Go Back

Close

Quit

Page 23 of 31

4. Some Estimates

It is useful to be able to obtain some estimates of the representation of the series (2.2). This is done in the following theorems.

Theorem 4.1. *Given that*


$$(4.1) \quad S(a, k, \alpha, m) = \sum_{r=0}^{\infty} \frac{(\alpha)_r}{r! (rk + m + 1) a^{rk+m+1}},$$

then

$$(4.2) \quad \frac{1}{(m+1)a^{m+1}} < S(a, k, \alpha, m) \leq \begin{cases} \frac{1}{((mq+1)a^{mq+1})^{\frac{1}{q}} k^{\frac{1}{p}}} \left[B\left(1 - \alpha p, \frac{1}{k}\right) - B\left(1 - a^{-k}; 1 - \alpha p, \frac{1}{k}\right) \right]^{\frac{1}{p}}, \\ \left(\frac{a^k}{a^k - 1}\right)^{\alpha} \frac{1}{(m+1)a^{m+1}}, \quad a > 1, \end{cases}$$

for real numbers p and q where $p > 1$, $\frac{1}{p} + \frac{1}{q} = 1$, $B(s, t)$ is the classical Beta function and $B(z; s, t)$ is the incomplete Beta function as described in Theorem 2.1.

Proof. Let $f(x) = \frac{1}{(1-x^k)^{\alpha}}$ and $g(x) = x^m$. Since $|f(x)|^p$ and $|g(x)|^q$ are integrable functions defined on $x \in [0, \frac{1}{a}]$, then by Hölder's integral inequality



π and Some Other Constants

Anthony Sofo

Title Page

Contents



Go Back

Close

Quit

Page 24 of 31

[14]

$$S(a, k, \alpha, m) \leq \left(\int_0^{\frac{1}{a}} x^{mq} dx \right)^{\frac{1}{q}} \left(\int_0^{\frac{1}{a}} \frac{dx}{(1-x^k)^{\alpha p}} \right)^{\frac{1}{p}}.$$

Now

$$\int_0^{\frac{1}{a}} \frac{dx}{(1-x^k)^{\alpha p}} = \frac{1}{k} \left[B\left(1 - \alpha p, \frac{1}{k}\right) - B\left(1 - a^{-k}; 1 - \alpha p, \frac{1}{k}\right) \right]$$

by the substitution $u = 1 - x^k$, and hence

$$S(a, k, \alpha, m) \leq \frac{1}{((mq+1)a^{mq+1})^{\frac{1}{q}} k^{\frac{1}{p}}} \times \left[B\left(1 - \alpha p, \frac{1}{k}\right) - B\left(1 - a^{-k}; 1 - \alpha p, \frac{1}{k}\right) \right]^{\frac{1}{p}}.$$


The lower bound on $S(a, k, \alpha, m)$ is $\frac{1}{(m+1)a^{m+1}}$ since the sum (2.2) is one of positive terms.

The second part of the inequality (4.2) is obtained from

$$\int_{x_0}^{x_1} |f(x)g(x)| dx \leq \text{ess sup}_{x \in [x_0, x_1]} |f(x)| \int_{x_0}^{x_1} |g(x)| dx.$$

Since $f(x)$ is monotonic on $x \in [0, \frac{1}{a}]$,

$$\text{ess sup}_{x \in [0, \frac{1}{a}]} \left\{ \frac{1}{(1-x^k)^\alpha} \right\} = \left(\frac{a^k}{a^k - 1} \right)^\alpha$$



π and Some Other Constants

Anthony Sofo

Title Page

Contents



Go Back

Close

Quit

Page 25 of 31

and

$$\int_0^{\frac{1}{a}} x^m dx = \frac{1}{(m+1)a^{m+1}},$$

hence

$$\frac{1}{(m+1)a^{m+1}} < S(a, k, \alpha, m) \leq \left(\frac{a^k}{a^k - 1}\right)^\alpha \frac{1}{(m+1)a^{m+1}}.$$

The result (4.2) follows and the theorem is proved. □


The next theorem develops an inequality of (4.1) based on the pre-Grüss result.

Theorem 4.2. For $a > 1$,

$$(4.3) \quad \left| S(a, k, \alpha, m) - \frac{1}{k(m+1)a^m} \right. \\ \left. \times \left\{ B\left(1 - \alpha, \frac{1}{k}\right) - B\left(1 - a^{-k}; 1 - \alpha, \frac{1}{k}\right) \right\} \right| \\ \leq \frac{m}{2(m+1)a^{m+1}\sqrt{2m+1}} \left[\left(\frac{a^k}{a^k - 1}\right)^\alpha - 1 \right].$$

Proof. Define

$$T(g, f) := \frac{1}{x_1 - x_0} \int_{x_0}^{x_1} f(x) g(x) dx \\ - \frac{1}{x_1 - x_0} \int_{x_0}^{x_1} f(x) dx \cdot \frac{1}{x_1 - x_0} \int_{x_0}^{x_1} g(x) dx$$



π and Some Other Constants

Anthony Sofo

Title Page

Contents



Go Back

Close

Quit

Page 26 of 31

for $f(x)$ and $g(x)$ integrable functions, as given in Theorem 4.1, and defined on $x \in [0, \frac{1}{a}]$, then the pre-Grüss inequality [13] states that

$$|T(g, f)| \leq \frac{\Gamma - \gamma}{2} [T(g, g)]^{\frac{1}{2}}$$

for $\gamma \leq f(x) \leq \Gamma$.

Now, for $x \in [0, \frac{1}{a}]$


$$\gamma = 1 \leq f(x) = \frac{1}{(1-x^k)^\alpha} \leq \left(\frac{a^k}{a^k-1}\right)^\alpha = \Gamma,$$

$$\begin{aligned} T(g, f) &= a \int_0^{\frac{1}{a}} \frac{x^m}{(1-x^k)^\alpha} dx - a^2 \int_0^{\frac{1}{a}} \frac{dx}{(1-x^k)^\alpha} \int_0^{\frac{1}{a}} x^m dx \\ &= aS(a, k, \alpha, m) - \frac{a^2}{(m+1)a^{m+1}k} \\ &\quad \times \left[B\left(1-\alpha, \frac{1}{k}\right) - B\left(1-a^{-k}; 1-\alpha, \frac{1}{k}\right) \right]. \end{aligned}$$

In a similar fashion

$$[T(g, g)]^{\frac{1}{2}} = \frac{m}{(m+1)a^m\sqrt{2m+1}}.$$

Combining $T(g, f)$ and $[T(g, g)]^{\frac{1}{2}}$ gives us the result (4.3) after a little algebraic simplification. \square



π and Some Other Constants

Anthony Sofo

Title Page

Contents



Go Back

Close

Quit

Page 27 of 31

Open Problem 1. From Example 3.2, we have that

$$U_{\infty} = \frac{3^{p+\frac{3}{2}}}{16p(p+1) \binom{2p+1}{p}} \left\{ 4p - 8 + \sum_{r=0}^{\infty} \binom{2r}{r} \frac{(2r+1)(2r+3)}{(2r+2p+3)} \left(\frac{3}{16}\right)^r \right\} \\ + \frac{(-1)^p}{\binom{2p+1}{p}} \sum_{s=0}^p \frac{(-1)^s}{2p-2s+2} \binom{2p+2}{s} \sin(2p-2s+2) \frac{\pi}{3} = \frac{\pi}{3}.$$

Now let us consider the following. For a finite positive integer R let


$$U_R = \frac{3^{p+\frac{3}{2}}}{16p(p+1) \binom{2p+1}{p}} \left\{ 4p - 8 + \sum_{r=0}^R \binom{2r}{r} \frac{(2r+1)(2r+3)}{(2r+2p+3)} \left(\frac{3}{16}\right)^r \right\} \\ + \frac{(-1)^p}{\binom{2p+1}{p}} \sum_{s=0}^p \frac{(-1)^s}{2p-2s+2} \binom{2p+2}{s} \sin(2p-2s+2) \frac{\pi}{3},$$

$$U_R = V + W,$$

in fact

$$U_R < \frac{\pi}{3}.$$

For a fixed positive integer R , it can be shown, by standard analysis methods,



π and Some Other Constants

Anthony Sofo

Title Page

Contents



Go Back

Close

Quit

Page 28 of 31

that

$$\lim_{p \rightarrow \infty} V = \lim_{p \rightarrow \infty} \frac{3^{p+\frac{3}{2}}}{16p(p+1) \binom{2p+1}{p}} \times \left\{ 4p - 8 + \sum_{r=0}^R \binom{2r}{r} \frac{(2r+1)(2r+3)}{(2r+2p+3)} \left(\frac{3}{16}\right)^r \right\} = 0$$

and as $p \rightarrow \infty$ uniformly


$$W = \frac{(-1)^p}{\binom{2p+1}{p}} \sum_{s=0}^p \frac{(-1)^s}{2p-2s+2} \binom{2p+2}{s} \sin(2p-2s+2) \frac{\pi}{3} \approx \frac{\pi}{3}.$$

This implies that for $p \rightarrow \infty$, $W \approx U_\infty$ and

$$W = \frac{(-1)^p}{\binom{2p+1}{p}} \sum_{s=0}^p \frac{(-1)^s}{2p-2s+2} \binom{2p+2}{s} \sin(2p-2s+2) \frac{\pi}{3} \approx \frac{\pi}{3}.$$

An open problem is to prove, or provide a contradiction to,

$$\lim_{p \rightarrow \infty} W = \lim_{p \rightarrow \infty} \left\{ \frac{(-1)^p}{\binom{2p+1}{p}} \sum_{s=0}^p \frac{(-1)^s}{2p-2s+2} \binom{2p+2}{s} \sin(2p-2s+2) \frac{\pi}{3} \right\} = \frac{\pi}{3}.$$



π and Some Other Constants

Anthony Sofo

Title Page

Contents



Go Back


Close

Quit

Page 29 of 31

References

- [1] M. ABRAMOWITZ AND I. STEGUN, *Handbook of Mathematical functions*, Dover Publications Inc. New York, 1970.
- [2] J. ARNDT AND C. HAENEL, *Pi-Unleashed*, Springer-Verlag, Germany, 2001.
- [3] P. BECKMANN, *A History of Pi*, 3rd Edition, New York, 1989.
- [4] B. BERNDT, *Ramanujans Notebooks*, Springer Verlag, New York. Part I, 1985. Part II, 1989. Part III, 1991. Part IV, 1994. Part V, 1998.
- [5] W.H. BEYER, *CRC Standard Mathematical Tables*, 28th Ed., Boca Raton FL., CRC Press, 1987.
- [6] D. BLATNER, *The Joy of Pi*. Walker, New York, 1997.
- [7] J. BORWEIN AND P. BORWEIN, *Pi and the AGM: A Study in Analytic Number Theory and Computational Complexity*, Wiley, New York, 1987, reprinted 1998.
- [8] S.R. FINCH, *Mathematical Constants. Encyclopedia of Mathematics and its Applications*, 94. Cambridge University Press, USA, 2003.
- [9] M. HIRSCHHORN, A new formula for π . *Australian Math. Soc. Gazette*, **25** (1999), 82–83.
- [10] A. LUPAŞ, *Formulae for some classical constants*, Schriftenreihe des fachbereichs Mathematik, Gerhard Mercator Universitat Duisburg, 2000, 70–76.



π and Some Other Constants

Anthony Sofo

Title Page

Contents




Go Back

Close

Quit

Page 30 of 31

- [11] *Pi*, Wolfram Research, 2003. <http://mathworld.wolfram.com/Pi>
- [12] *Pi Formulas*, Wolfram Research, 2003. <http://mathworld.wolfram.com/PiFormulas>
- [13] M. MATIĆ, J.E. PEČARIĆ AND N. UJEVIĆ, On new estimation of the remainder in generalised Taylor's formula, *Math. Ineq. and App.*, **2** (1999), 343–361.
- [14] D.S. MITRINOVIĆ, J.E. PEČARIĆ AND A.M. FINK, *Classical and New Inequalities in Analysis*, Kluwer Academic Publishers, Boston, 1993.
- [15] P. SEBAH AND X. GOURDON, *Collection of Series for π* , 2002. <http://numbers.computation.free.fr/constants/Pi/pi.html>
- [16] A. SOFO, *Computational Techniques for the Summation of Series*, Kluwer Publishing Co., New York, 2003.
- [17] A. SOFO, Some Representations of π , *Australian Math. Soc. Gazette*, **31** (2004), 184–189.
- [18] H.S. WILF, *Generatingfunctionology*, Academic Press Inc., New York, 1994.



π and Some Other Constants

Anthony Sofo

Title Page

Contents



Go Back

Close

Quit

Page 31 of 31