


# Journal of Inequalities in Pure and Applied Mathematics

## NOTES ON QI TYPE INTEGRAL INEQUALITIES

LAZHAR BOUGOFFA

King Khalid University  
Faculty of Science  
Department of Mathematics  
P. O. Box 9004  
Abha Saudi Arabia.  
EMail: [abogafah@kku.edu.sa](mailto:abogafah@kku.edu.sa)

©2000 Victoria University  
ISSN (electronic): 1443-5756  
091-03



---

volume 4, issue 4, article 77,  
2003.

*Received 01 July, 2003;*  
*accepted 08 October, 2003.*

*Communicated by: F. Qi*

---

[Abstract](#)

[Contents](#)



[Home Page](#)

[Go Back](#)

[Close](#)

[Quit](#)

## Abstract

By utilizing a reversed Hölder inequality and a reversed convolution inequality, some new integral inequalities which originate from an open problem posed in [F. Qi, *Several integral inequalities*, J. Inequal. Pure Appl. Math. 1(2) (2000), Art. 19. Available online at [http://jipam.vu.edu.au/v1n2.html/001\\_00.html](http://jipam.vu.edu.au/v1n2.html/001_00.html). RGMIA Res. Rep. Coll. 2 (1999), no. 7, Art. 9, 1039–1042. Available online at <http://rgmia.vu.edu.au/v2n7.html>] are established.


**2000 Mathematics Subject Classification:** Primary 26D15; Secondary 39B62, 60E15.

**Key words:** Qi type integral inequality, Reversed Hölder inequality, Reversed convolution inequality.

The author would like to thank the anonymous referees for their valuable help to renew the first section, add and correct several references, and improve the typesetting of this paper.

## Contents

1	Introduction .....	3
2	Two Lemmas .....	5
3	Main Results and Proofs .....	6
	References	



### Notes on Qi Type Integral Inequalities

Lazhar Bougoffa

Title Page

Contents



Go Back

Close

Quit

Page 2 of 10

# 1. Introduction

In [6] F. Qi posed the following open problem:

*Under what conditions does the inequality*

$$(1.1) \quad \int_a^b [f(x)]^t dx \geq \left( \int_a^b f(x) dx \right)^{t-1}$$

*hold for  $t > 1$ ?*


In response, affirmative answers, generalizations, reversed forms, and interpretations of inequality (1.1) have been obtained by several mathematicians.

The paper [9] first gave an affirmative answer to this open problem using the integral version of Jensen's inequality and a lemma of convexity. The second affirmative answer to (1.1) was given by Towghi in [8]. Motivated by (1.1), Pogány [5] proved the following

$$(1.2) \quad \int_a^b [f(x)]^\alpha dx \geq \left( \int_a^b f(x) dx \right)^\beta$$

and its reversed form under assumptions of the bounds, depending on  $b - a$ ,  $\alpha$  and  $\beta$ , and convexity of  $f$ , which contains an answer to the above open problem and some reversed forms of (1.1).

In [3, 4], by employing a functional inequality which is an abstract generalization of the classical Jessen's inequality, the following functional inequality (1.4) was established, from which inequality (1.1), some integral inequalities, and an interesting discrete inequality involving sums can be deduced:



Title Page

Contents



Go Back

Close

Quit

Page 3 of 10

Let  $\mathcal{L}$  be a linear vector space of real-valued functions,  $p$  and  $q$  be two real numbers such that  $p \geq q \geq 1$ . Assume that  $f$  and  $g$  are two positive functions in  $\mathcal{L}$  and  $G$  is a positive linear form on  $\mathcal{L}$  such that  $G(g) > 0$ ,  $fg \in \mathcal{L}$ , and  $gf^p \in \mathcal{L}$ . If

$$(1.3) \quad [G(gf)]^{p-q} \geq [G(g)]^{p-1},$$

then


$$(1.4) \quad G(gf^p) \geq [G(gf)]^q.$$

Very recently, Csiszár and Móri [1] interpreted inequality (1.2) in terms of moments as

$$(1.5) \quad E(X^\alpha) \geq C(EX)^\beta,$$

where  $C = (b - a)^{\beta-1}$  and  $X = f$  is a random variable. To demonstrate the power of the convexity method in probability theory, among other things, they found sharp conditions on the range of  $X$ , under which (1.5) or the converse inequality holds for fixed  $0 < \beta < \alpha$ . Hence, the results by Pogány [5] were improved slightly by a factor of at least  $\left(\frac{3}{2}\right)^{\frac{\alpha}{\alpha-\beta}}$ .

In this short note, by utilizing the reversed Hölder inequality in [2] and a reversed convolution inequality in [7], we establish some new Qi type integral inequalities which extend related results in references.



## Notes on Qi Type Integral Inequalities

Lazhar Bougoffa

Title Page

Contents



Go Back

Close

Quit

Page 4 of 10

## 2. Two Lemmas

In order to prove our main results, the following two lemmas are necessary.

**Lemma 2.1 ([2]).** For two positive functions  $f$  and  $g$  satisfying  $0 < m \leq \frac{f}{g} \leq M < \infty$  on the set  $X$ , and for  $p > 1$  and  $q > 1$  with  $\frac{1}{p} + \frac{1}{q} = 1$ , we have

$$(2.1) \quad \left( \int_X f d\mu \right)^{\frac{1}{p}} \left( \int_X g d\mu \right)^{\frac{1}{q}} \leq A_{p,q} \left( \frac{m}{M} \right) \int_X f^{\frac{1}{p}} g^{\frac{1}{q}} d\mu,$$

where

$$(2.2) \quad A_{p,q}(t) = \frac{1}{p^{\frac{1}{p}} q^{\frac{1}{q}}} \cdot \frac{1-t}{t^{\frac{1}{pq}} \left(1-t^{\frac{1}{p}}\right)^{\frac{1}{p}} \left(1-t^{\frac{1}{q}}\right)^{\frac{1}{q}}}.$$

Inequality (2.1) is called the reverse Hölder inequality.

**Lemma 2.2 ([7]).** For two positive functions  $f$  and  $g$  satisfying  $0 < m \leq \frac{f^p}{g^q} \leq M < \infty$  on the set  $X$ , and for  $p > 1$  and  $q > 1$  with  $\frac{1}{p} + \frac{1}{q} = 1$ , we have

$$(2.3) \quad \left( \int_X f^p d\mu \right)^{\frac{1}{p}} \left( \int_X g^q d\mu \right)^{\frac{1}{q}} \leq \left( \frac{M}{m} \right)^{\frac{1}{pq}} \int_X f g d\mu.$$



---

Notes on Qi Type Integral Inequalities

Lazhar Bougoffa

---

Title Page

Contents



Go Back

Close

Quit

Page 5 of 10

### 3. Main Results and Proofs

In this section, we will state our main results and give their proofs as follows.

Using Lemma 2.1 and an estimation due to Lars-Erik Persson, we obtain:

**Theorem 3.1.** *If  $0 < m \leq f \leq M < \infty$  on  $[a, b]$ , then*

$$(3.1) \quad \int_{[a,b]} f^{\frac{1}{p}} d\mu \geq B \left( \int_{[a,b]} f d\mu \right)^{\frac{1}{p}-1},$$

where  $B = m[\mu(b) - \mu(a)]^{1+\frac{1}{q}} \left(\frac{m}{M}\right)^{\frac{1}{pq}}$  and  $p > 1, q > 1$  with  $\frac{1}{p} + \frac{1}{q} = 1$ .

*Proof.* In Lemma 2.1, putting  $g \equiv 1$  yields

$$(3.2) \quad [\mu(b) - \mu(a)]^{\frac{1}{q}} \left( \int_{[a,b]} f d\mu \right)^{\frac{1}{p}} \leq A_{p,q} \left( \frac{m}{M} \right) \int_{[a,b]} f^{\frac{1}{p}} d\mu,$$


and so

$$(3.3) \quad \int_{[a,b]} f^{\frac{1}{p}} d\mu \geq \frac{[\mu(b) - \mu(a)]^{\frac{1}{q}}}{A_{p,q} \left(\frac{m}{M}\right)} \left( \int_{[a,b]} f d\mu \right)^{\frac{1}{p}-1} \left( \int_{[a,b]} f d\mu \right).$$

Since  $0 < m \leq f$ , we have

$$(3.4) \quad \int_{[a,b]} f^{\frac{1}{p}} d\mu \geq \frac{m [\mu(b) - \mu(a)]^{1+\frac{1}{q}}}{A_{p,q} \left(\frac{m}{M}\right)} \left( \int_{[a,b]} f d\mu \right)^{\frac{1}{p}-1}.$$

Now, using the estimation  $A_{p,q}(t) < t^{-\frac{1}{pq}}$  due to Lars-Erik Persson (see [7]), we obtain inequality (3.1).  $\square$



Notes on Qi Type Integral  
Inequalities

Lazhar Bougoffa

Title Page

Contents




Go Back

Close

Quit

Page 6 of 10



Title Page

Contents



Go Back

Close

Quit

Page 7 of 10

**Corollary 3.2.** Let  $p > 1$  and  $q > 1$  with  $\frac{1}{p} + \frac{1}{q} = 1$ . If

$$(3.5) \quad m \left( \frac{m}{M} \right)^{\frac{1}{pq}} = \frac{1}{[\mu(b) - \mu(a)]^{1+\frac{1}{q}}}$$

and  $0 < m \leq f \leq M < \infty$  on  $[a, b]$ , then

$$(3.6) \quad \int_{[a,b]} f^{\frac{1}{p}} d\mu \geq \left( \int_{[a,b]} f d\mu \right)^{\frac{1}{p}-1}.$$

Using Lemma 2.1, Lemma 2.2 and the estimation due to Lars-Erik Persson, we have the following:

**Theorem 3.3.** Let  $p > 1$  and  $q > 1$  with  $\frac{1}{p} + \frac{1}{q} = 1$ . If  $0 < m^{\frac{1}{p}} \leq f \leq M^{\frac{1}{p}} < \infty$  on  $[a, b]$ , then

$$(3.7) \quad \left( \int_{[a,b]} f^{\frac{1}{p}} d\mu \right)^p \geq [\mu(b) - \mu(a)]^{\frac{p+1}{q}} \left( \frac{m}{M} \right)^{\frac{2}{pq}} \left( \int_{[a,b]} f^p d\mu \right)^{\frac{1}{p}}.$$

*Proof.* Putting  $g \equiv 1$  into Lemma 2.2, we obtain

$$(3.8) \quad \left( \int_{[a,b]} f^p d\mu \right)^{\frac{1}{p}} [\mu(b) - \mu(a)]^{\frac{1}{q}} \leq \left( \frac{m}{M} \right)^{-\frac{1}{pq}} \int_{[a,b]} f d\mu.$$

Therefore

$$(3.9) \quad \left( \int_{[a,b]} f^p d\mu \right)^{\frac{1}{p}} \leq \left( \frac{m}{M} \right)^{-\frac{1}{pq}} [\mu(b) - \mu(a)]^{-\frac{1}{q}} \int_{[a,b]} f d\mu.$$

Again, substituting  $g \equiv 1$  in the reverse Hölder inequality in Lemma 2.1 leads to

$$\left( \int_{[a,b]} f d\mu \right)^{\frac{1}{p}} \leq [\mu(b) - \mu(a)]^{-\frac{1}{q}} A_{p,q} \left( \frac{m^{\frac{1}{p}}}{M^{\frac{1}{p}}} \right) \int_{[a,b]} f^{\frac{1}{p}} d\mu,$$

and so,

$$(3.10) \quad \int_{[a,b]} f d\mu \leq [\mu(b) - \mu(a)]^{-\frac{p}{q}} A_{p,q}^p \left( \frac{m^{\frac{1}{p}}}{M^{\frac{1}{p}}} \right) \left( \int_{[a,b]} f^{\frac{1}{p}} d\mu \right)^p.$$


Combining (3.9) with (3.10), we obtain

$$(3.11) \quad \left( \int_{[a,b]} f^{\frac{1}{p}} d\mu \right)^p \geq [\mu(b) - \mu(a)]^{\frac{p+1}{q}} \left( \frac{m}{M} \right)^{\frac{1}{pq}} A_{p,q}^{-p} \left( \frac{m^{\frac{1}{p}}}{M^{\frac{1}{p}}} \right) \left( \int_{[a,b]} f^p d\mu \right)^{\frac{1}{p}}.$$

Then, by using the estimation  $A_{p,q}(t) < t^{-\frac{1}{pq}}$  due to Lars-Erik Persson (see [7]), we obtain inequality (3.7).  $\square$

**Corollary 3.4.** *If  $0 < m^{\frac{1}{p}} \leq f \leq M^{\frac{1}{p}} < \infty$  on  $[a, b]$  and  $\frac{m}{M} = [\mu(b) - \mu(a)]^{-\frac{p(p+1)}{2}}$  for  $p > 1$ , then*

$$(3.12) \quad \left( \int_{[a,b]} f^{\frac{1}{p}} d\mu \right)^p \geq \left( \int_{[a,b]} f^p d\mu \right)^{\frac{1}{p}}.$$



## Notes on Qi Type Integral Inequalities

Lazhar Bougoffa

Title Page

Contents



Go Back

Close


Quit

Page 8 of 10



## References

- [1] V. CSISZÁR AND T.F. MÓRI, The convexity method of proving moment-type inequalities, *Statist. Probab. Lett.*, (2003), accepted.
- [2] X.-H. LIU, On the inverses of Hölder inequalities, *Math. Practice Theory*, **20**(1) (1990), 84–88. (Chinese)
- [3] S. MAZOUZI AND F. QI, On an open problem by Feng Qi regarding an integral inequality, *RGMIA Res. Rep. Coll.*, **6**(1) (2003), Art. 6. Available online at <http://rgmia.vu.edu.au/v6n1.html>.
- [4] S. MAZOUZI AND F. QI, On an open problem regarding an integral inequality, *J. Inequal. Pure Appl. Math.*, **4**(2) (2003), Art. 31. Available online at [http://jipam.vu.edu.au/v4n2/029\\_03.html](http://jipam.vu.edu.au/v4n2/029_03.html).
- [5] T.K. POGÁNY, On an open problem of F. Qi, *J. Inequal. Pure Appl. Math.*, **3**(4) (2002), Art. 54. Available online at [http://jipam.vu.edu.au/v3n4/016\\_01.html](http://jipam.vu.edu.au/v3n4/016_01.html).
- [6] F. QI, Several integral inequalities, *J. Inequal. Pure Appl. Math.*, **1**(2) (2000), Art. 19. Available online at [http://jipam.vu.edu.au/v1n2.html/001\\_00.html](http://jipam.vu.edu.au/v1n2.html/001_00.html). *RGMIA Res. Rep. Coll.*, **2**(7) (1999), Art. 9, 1039–1042. Available online at <http://rgmia.vu.edu.au/v2n7.html>.
- [7] S. SAITOH, V.K. TUAN AND M. YAMAMOTO, Reverse convolution inequalities and applications to inverse heat source problems, *J. Inequal. Pure Appl. Math.*, **3**(5) (2003), Art. 80. Available online at [http://jipam.vu.edu.au/v3n5/029\\_02.html](http://jipam.vu.edu.au/v3n5/029_02.html).



---

### Notes on Qi Type Integral Inequalities

Lazhar Bougoffa

---

Title Page

Contents




Go Back

Close

Quit

Page 9 of 10

- [8] N. TOWGHI, Notes on integral inequalities, *RGMI*A Res. Rep. Coll., 4(2) (2001), Art. 12, 277–278. Available online at <http://rgmia.vu.edu.au/v4n2.html>.
- [9] K.-W. YU AND F. QI, A short note on an integral inequality, *RGMI*A Res. Rep. Coll., 4(1) (2001), Art. 4, 23–25. Available online at [http://rgmia.vu.edu.edu.au/v4n1.html](http://rgmia.vu.edu.au/v4n1.html).



---

### Notes on Qi Type Integral Inequalities

Lazhar Bougoffa

---

Title Page

Contents



Go Back

Close

Quit

Page 10 of 10