


GENERALIZATION OF A RESULT FOR COSINE SERIES ON THE L^1 NORM

JÓZSEF NÉMETH

Bolyai Institute
University of Szeged
Aradi vértanúk tere 1.
H-6720 Hungary.
EMail: nemethj@math.u-szeged.hu



volume 4, issue 5, article 99,
2003.

*Received 13 November, 2002;
accepted 03 July, 2003.*

Communicated by: A. Babenko

Abstract

Contents



Home Page

Go Back

Close

Quit

Abstract

A L^1 -estimate will be established for cosine series, considering the generalized Fomin-class \mathcal{F}_φ , where φ is a function more general than the power function

2000 Mathematics Subject Classification: 26D15; 42A20.

Key words: Trigonometric series, Fomin-class, L^1 -estimate.

This research was supported by the Hungarian National Foundation for Scientific Research under Grant No. 19/75/1K703.

Contents

1	Introduction	3
2	Result	6
3	Lemmas	7
4	Proof	10
	References	



**Generalization of a Result for
Cosine Series on the L^1 Norm**

József Németh

Title Page

Contents



Go Back

Close

Quit

Page 2 of 14

1. Introduction

Let

$$(1.1) \quad f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos nx.$$

Many authors ([1], [2], [9] – [15]) have investigated coefficient conditions guaranteeing that (1.1) is a Fourier series of some function $f \in L^1$ and they have given estimates for $\int_0^\pi |f(x)|dx$ via the sequence $\{a_n\}$.

Recently Z. Tomovski [15] proved a theorem of this type by using the class of coefficients defined by Fomin [1] as follows: a sequence $\{a_n\}$ belongs to \mathcal{F}_p ($p > 1$) if $a_k \rightarrow 0$ and


$$(1.2) \quad \sum_{k=1}^{\infty} \left\{ \frac{1}{k} \sum_{i=k}^{\infty} |\Delta a_i|^p \right\}^{\frac{1}{p}} < \infty.$$

Now we can formulate Z. Tomovski's result [15]:

Theorem 1.1. *Let $\{a_n\} \in \mathcal{F}_p$, $1 < p \leq 2$, then the series (1.1) is a Fourier series and the following inequality holds:*

$$(1.3) \quad \int_0^\pi |f(x)|dx \leq C_p \sum_{n=1}^{\infty} \left\{ \frac{1}{n} \sum_{k=n}^{\infty} |\Delta a_k|^p \right\}^{\frac{1}{p}},$$

where C_p depends only on p .



Generalization of a Result for
Cosine Series on the L^1 Norm

József Németh

Title Page

Contents



Go Back

Close

Quit

Page 3 of 14

Recently we ([7]) investigated the properties of classes of numerical sequences obtained by using functions more general than the power functions. Such functions were used first of all in the works of H.P. Mulholland [5] and M. Mateljevič and M. Pavlovič [8]. The following definition is due to Mateljevič and Pavlovič.


$\Delta(q, p)$ ($q \geq p > 0$) denotes the family of the nonnegative real functions $\varphi(x)$ defined on $[0; \infty)$ with the following properties: $\varphi(0) = 0$, $\frac{\varphi(t)}{t^q}$ is non-increasing and $\frac{\varphi(t)}{t^p}$ is nondecreasing on $(0; \infty)$. Δ will denote the set of the functions $\varphi(x) \in \Delta(q, p)$ for some $q \geq p > 0$.

Using this notion in [7] we defined the following class: a nullsequence $\{a_n\}$ belongs to the class \mathcal{F}_φ for some $\varphi \in \Delta$ if

$$(1.4) \quad \sum_{n=1}^{\infty} \bar{\varphi} \left(\frac{1}{n} \sum_{k=n}^{\infty} \varphi(|\Delta a_k|) \right) < \infty,$$

where $\bar{\varphi}$ is the inverse of the function φ .

The aim of the present paper is to generalize Theorem 1.1 using \mathcal{F}_φ instead of \mathcal{F}_p , where $\varphi \in \Delta$. Since our goal is to get a result concerning such functions as $\varphi(x) = x \log^\alpha(1+x)$ and not only for functions which are generalizations of x^p ($p > 1$), we therefore need to define two subclasses of Δ . Namely we use the following definitions: $\Delta^{(1)}$ denotes the family of functions $\varphi(x)$ belonging to $\Delta(q, p)$ for some $q \geq p > 1$ and $\Delta^{(2)}$ is the collection of functions $\varphi(x)$ from $\Delta(q, 1)$ for some $q > 1$ such that for all $A > 0$ there exists $p := p(A) > 1$ satisfying the condition that $\frac{\varphi(x)}{x^p}$ is nondecreasing on $(0; A)$. It is obvious that $\Delta^{(1)} \subset \Delta^{(2)}$.



Title Page

Contents




Go Back

Close

Quit

Page 4 of 14

After giving these definitions we can formulate our result which generalizes Theorem 1.1 (if $\varphi \in \Delta^{(1)}$). Furthermore, it contains the case like $\varphi(x) = x \log^\alpha(1+x)$ ($\alpha > 0$).



**Generalization of a Result for
Cosine Series on the L^1 Norm**

József Németh

Title Page

Contents



Go Back

Close

Quit

Page 5 of 14


2. Result

Theorem 2.1. Let $\{a_n\} \in \mathcal{F}_\varphi$ for $\varphi \in \Delta^{(2)}$. Then the series (1.1) is a Fourier series and the following estimate holds

$$(2.1) \quad \int_0^\pi |f(x)| dx \leq C_\varphi \sum_{n=1}^{\infty} \bar{\varphi} \left(\frac{1}{n} \sum_{k=n}^{\infty} \varphi(|\Delta a_k|) \right),$$

where C_φ is a constant depending only on φ .

Remark 2.1. In [3] and [4] L. Leindler investigated a relation among classes of numerical sequences other than \mathcal{F}_p (see the classes denoted by S_p , \mathcal{F}_p^* , $S_p(\delta)$, $S_p(A)$) and he proved that all these classes coincide. Later in [7] we defined the classes of sequences S_φ , \mathcal{F}_φ^* , $S_\varphi(\delta)$, $S_\varphi(A)$ exchanging the functions x^p to $\varphi(x)$ and showed that all these classes also coincide. Therefore in Theorem 2.1 the class \mathcal{F}_φ can be replaced by any of the above mentioned classes, if $\varphi \in \Delta^{(2)}$.



Generalization of a Result for
Cosine Series on the L^1 Norm

József Németh

Title Page

Contents



Go Back

Close

Quit

Page 6 of 14

3. Lemmas

Lemma 3.1. ([11]). Let the nullsequence $\{a_n\}$ be of bounded variation and

$$\sum_{i=2}^{\infty} \sum_{k=1}^{[i/2]} \left| \frac{\Delta a_{i-k} - \Delta a_{i+k}}{k} \right| < \infty$$

then (1.1) is a Fourier series and the following estimate holds:

$$(3.1) \quad \int_0^{\pi} |f(x)| dx \leq C \left(\sum_{k=0}^{\infty} |\Delta a_k| + \sum_{i=2}^{\infty} \left| \sum_{k=1}^{[i/2]} \frac{\Delta a_{i-k} - \Delta a_{i+k}}{k} \right| \right),$$

where C is some absolute constant.


Lemma 3.2. ([1]). Let $\{a_n\}$ be a nullsequence. Then the following estimate holds:

$$(3.2) \quad \sum_{i=2}^{\infty} \sum_{k=1}^{[i/2]} \left| \frac{\Delta a_{i-k} - \Delta a_{i+k}}{k} \right| \leq C_p \sum_{s=1}^{\infty} \Delta_s^{(p)},$$

where

$$\Delta_s^{(p)} = \left\{ \frac{1}{2^{s-1}} \sum_{k=2^{s-1}+1}^{2^s} |\Delta a_k|^p \right\}^{\frac{1}{p}}, \quad 1 < p \leq 2,$$

and C_p is a constant depending only on p .



Generalization of a Result for
Cosine Series on the L^1 Norm

József Németh

Title Page

Contents



Go Back

Close

Quit

Page 7 of 14

Lemma 3.3. ([5]). If $\frac{\Psi(x)}{x}$ is increasing then for all sequences $\{a_n\}$ and $\{b_n\}$ of nonnegative numbers

$$\Psi \left(\frac{\sum_{i=1}^n a_i b_i}{\sum_{i=1}^n a_i} \right) \leq \frac{\sum_{i=1}^n a_i \Psi(2b_i)}{\sum_{i=1}^n a_i}$$

holds.

Lemma 3.4. ([6]). Let $\rho(x)$ denote a nonnegative function increasing to infinity such that $\frac{\rho(x)}{x}$ is decreasing to zero when x is increasing from zero to infinity. Furthermore, if $a_n \geq 0$, $\lambda_n > 0$ for all n , then

$$(3.3) \quad \sum_{n=1}^{\infty} \lambda_n \rho \left(\frac{a_n}{\lambda_n} \sum_{k=1}^n \lambda_k \right) \leq C_\rho \sum_{n=1}^{\infty} \lambda_n \rho \left(\sum_{k=n}^{\infty} a_k \right),$$

where C_ρ depends only on the function $\rho(x)$.


Lemma 3.5. Let $b_n \geq 0$ for all n and let $\bar{\varphi}$ be the inverse of the function $\varphi(x) \in \Delta^{(2)}$. Then

$$(3.4) \quad \sum_{n=1}^{\infty} \bar{\varphi}(b_n) \leq K_{\bar{\varphi}} \cdot \sum_{n=1}^{\infty} \bar{\varphi} \left(\frac{\sum_{k=n}^{\infty} b_k}{n} \right),$$

where $K_{\bar{\varphi}}$ is a constant depending only on $\bar{\varphi}$.

Proof. Since $\frac{\bar{\varphi}(x)}{x}$ is decreasing to zero if $x \rightarrow \infty$, therefore using Lemma 3.4 and taking $\rho(x) = \bar{\varphi}(x)$, $b_n = n \cdot a_n$, $\lambda_n = 1$, we get

$$\sum_{n=1}^{\infty} \bar{\varphi}(b_n) \leq K_{\bar{\varphi}} \cdot \sum_{n=1}^{\infty} \bar{\varphi} \left(\sum_{k=n}^{\infty} \frac{b_k}{k} \right),$$



Generalization of a Result for
Cosine Series on the L^1 Norm

József Németh

Title Page

Contents



Go Back


Close

Quit

Page 8 of 14

whence the statement of Lemma 3.5 is obtained. \square

\square



**Generalization of a Result for
Cosine Series on the L^1 Norm**

József Németh

Title Page

Contents



Go Back

Close

Quit

Page 9 of 14

4. Proof

Proof of Theorem 2.1. Let $\varphi \in \Delta^{(2)}$ and $b_n = \varphi(|\Delta a_n|)$. Using Lemma 3.5 and that $\{a_n\} \in \mathcal{F}_\varphi$ we have that

$$(4.1) \quad \sum_{n=1}^{\infty} |\Delta a_n| \leq K \cdot \sum_{n=1}^{\infty} \bar{\varphi} \left(\frac{1}{n} \sum_{k=n}^{\infty} \varphi(|\Delta a_k|) \right) < \infty,$$

where K depends only on φ . It now follows that the sequence $\{a_n\}$ is of bounded variation.

Further on, we use the following notations:

$$\Delta_s^{(\varphi)} := \bar{\varphi} \left\{ \frac{1}{2^{s-1}} \sum_{k=2^{s-1}+1}^{2^s} \varphi(|\Delta a_k|) \right\}, \quad p' > 1$$

denotes a number for which $\frac{\varphi(x)}{x^{p'}} \uparrow$ on $(0; A)$, where $A = \sup_k |\Delta a_k|$, and by q

we denote a number satisfying $\frac{\varphi(x)}{x^q} \downarrow$ (see the definition of $\varphi \in \Delta^{(2)}$).


Now we will prove that for any $1 < p < p'$,

$$(4.2) \quad \sum_{s=1}^{\infty} 2^s \Delta_s^{(p)} \leq 2^{q/p} \sum_{s=1}^{\infty} 2^s \Delta_s^{(\varphi)}$$

holds.

Since for $\psi(t) = \varphi(t^{1/p})$ the function $\frac{\psi(t)}{t}$ is increasing thus using Lemma 3.3 and that $\varphi(2^{1/p}x) \leq 2^{\frac{q}{p}}\varphi(x)$ we get

$$(4.3) \quad \psi \left(\frac{\sum_{k=2^{s-1}+1}^{2^s} |\Delta a_k|^p}{2^{s-1}} \right) \leq 2^{\frac{q}{p}} \frac{\sum_{k=2^{s-1}+1}^{2^s} \varphi(|\Delta a_k|)}{2^{s-1}}.$$



Generalization of a Result for
Cosine Series on the L^1 Norm

József Németh

Title Page

Contents



Go Back

Close

Quit

Page 10 of 14

From (4.3), taking into account that $\bar{\varphi}(cx) \leq c\bar{\varphi}(x)$ if $c > 1$, we obtain

$$(4.4) \quad \bar{\varphi} \left[\psi \left(\frac{\sum_{k=2^{s-1}+1}^{2^s} |\Delta a_k|^p}{2^{s-1}} \right) \right] \leq 2^{\frac{q}{p}} \bar{\varphi} \left(\frac{\sum_{k=2^{s-1}+1}^{2^s} \varphi(|\Delta a_k|)}{2^{s-1}} \right).$$

Since $\bar{\varphi}(\psi(t)) = t^{1/p}$ so from (4.4) we have

$$(4.5) \quad \Delta_s^{(p)} = \left\{ \frac{\sum_{k=2^{s-1}+1}^{2^s} |\Delta a_k|^p}{2^{s-1}} \right\}^{\frac{1}{p}} \\ \leq 2^{q/p} \bar{\varphi} \left(\frac{\sum_{k=2^{s-1}+1}^{2^s} \varphi(|\Delta a_k|)}{2^{s-1}} \right) = 2^{q/p} \Delta_s^{(\varphi)}.$$


From (4.5), (4.2) immediately follows.

Now we show that

$$(4.6) \quad \sum_{s=1}^{\infty} 2^s \Delta_s^{(\varphi)} \leq 4 \cdot \sum_{s=1}^{\infty} \bar{\varphi} \left(\frac{\sum_{k=s}^{\infty} \varphi(|\Delta a_k|)}{s} \right).$$

Since the sequence $U_s = \frac{1}{s} \sum_{k=s}^{\infty} \varphi(|\Delta a_k|)$ is monotone decreasing, we obtain

$$(4.7) \quad \sum_{s=1}^n 2^s \Delta_s^{(\varphi)} = 2 \cdot \sum_{s=1}^n 2^{s-1} \bar{\varphi} \left(\frac{1}{2^{s-1}} \sum_{k=2^{s-1}+1}^{2^s} \varphi(|\Delta a_k|) \right) \\ \leq 4 \cdot \sum_{s=1}^{2^{n-1}} \bar{\varphi}(U_s) = 4 \cdot \sum_{s=1}^{2^{n-1}} \bar{\varphi} \left(\frac{1}{s} \sum_{k=s}^{\infty} \varphi(|\Delta a_k|) \right).$$



Generalization of a Result for
Cosine Series on the L^1 Norm

József Németh

Title Page

Contents



Go Back


Close

Quit

Page 11 of 14

Setting $n \rightarrow \infty$ we have from (4.7) the inequality (4.6).

Collecting (4.1), (4.2), (4.6), using Lemma 3.1 and Lemma 3.2, we get that (1.1) is a Fourier series and (2.1) is true. Thus Theorem 2.1 is proved. \square



**Generalization of a Result for
Cosine Series on the L^1 Norm**

József Németh

Title Page

Contents



Go Back


Close

Quit

Page 12 of 14

References

- [1] G.A. FOMIN, A class of trigonometric series, *Math. Zametki*, **23** (1978), 117–124 (Russian).
- [2] A.N. KOLMOGOROV, Sur l'ordre de grandeur des coefficients de la serie de Fourier-Lebesgue, *Bull. Acad. Polon. Ser. A, Sci. Math.* (1923), 83–86.
- [3] L. LEINDLER, On the equivalence of classes of Fourier coefficients, *Math. Inequal. Appl.*, **3** (2000), 45–50.
- [4] L. LEINDLER, A note on some classes of real sequences, *Math. Inequal. Appl.*, **4** (2001), 53–58.
- [5] H.P. MULHOLLAND, The generalization of certain inequality theorems involving powers, *Proc. London Math. Soc.*, **33** (1932), 481–516.
- [6] J. NÉMETH, Generalization of the Hardy-Littlewood inequality. II, *Acta Sci. Math.*, **35** (1973), 127–134.
- [7] J. NÉMETH, A note on two theorems of Leindler, *Math. Inequal. Appl.*, **5** (2002), 225–233.
- [8] M. MATELJEVIČ AND M. PAVLOVIČ, L^p -behavior of power series with positive coefficients and Hardy spaces, *Proc. Amer. Math. Soc.*, **87** (1983), 309–316.
- [9] S. SIDON, Hinreichende Bedingungen für den Fourier-charakter einer trigonometrischen Reihe, *J. London Math. Soc.*, **14** (1939), 158–160.



Generalization of a Result for
Cosine Series on the L^1 Norm

József Németh

Title Page

Contents




Go Back

Close

Quit

Page 13 of 14

- [10] S. SIDON, Reihen theoretische Sätze und ihre Anwendungen in der Theorie der Fourierschen Reihen, *Math. Z.*, **10** (1921), 121–127.
- [11] S.A. TELYAKOVSKĪI, Integrability conditions for trigonometric series and their application to the study of linear summation methods of Fourier series, *Izv. Akad. Nauk SSSR Ser. Mat.*, **28** (1964), 1209–1238.
- [12] S.A. TELYAKOVSKĪI, Some estimates for trigonometric series with quasi-convex coefficients, *Mat. Sb.*, **63**(105) (1964), 426–444. (Russian)
- [13] S.A. TELYAKOVSKĪI, An asymptotic estimate of the integral of the absolute value of a function given by sine series, *Sibirsk. Mat. Ž.*, **8** (1967), 1416–1422. (Russian)
- [14] S.A. TELYAKOVSKĪI, An estimate, useful in problems of approximation theory of the norm of a function by means of its Fourier coefficients, *Trudy Mat. Inst. Steklov*, **109** (1971), 73–109. (Russian)
- [15] Ž. TOMOVSKI, Two new L^1 -estimates for trigonometric series with Fomin's coefficient condition, *RGMIA Research Report Collection*, **2**(6) (1999), <http://rgmia.vu.edu.au/>.



Generalization of a Result for
Cosine Series on the L^1 Norm

József Németh

Title Page

Contents



Go Back

Close

Quit

Page 14 of 14