

BOUNDARY VALUE PROBLEM FOR SECOND-ORDER DIFFERENTIAL OPERATORS WITH MIXED NONLOCAL BOUNDARY CONDITIONS

M. DENCHE AND A. KOURTA

Laboratoire Equations Differentielles,
Departement de Mathematiques,
Faculte des Sciences, Universite Mentouri,
25000 Constantine, Algeria.

EMail: denech@wissal.dz



volume 5, issue 2, article 38,
2004.

*Received 15 October, 2003;
accepted 16 April, 2004.*

Communicated by: A.G. Ramm

Abstract

Contents



Home Page

Go Back

Close

Quit

Abstract

In this paper, we study a second order differential operator with mixed nonlocal boundary conditions combined weighting integral boundary condition with another two point boundary condition. Under certain conditions on the weighting functions and on the coefficients in the boundary conditions, called non regular boundary conditions, we prove that the resolvent decreases with respect to the spectral parameter in $L^p(0, 1)$, but there is no maximal decreasing at infinity for $p \geq 1$. Furthermore, the studied operator generates in $L^p(0, 1)$ an analytic semi group with singularities for $p \geq 1$. The obtained results are then used to show the correct solvability of a mixed problem for a parabolic partial differential equation with non regular boundary conditions.

2000 Mathematics Subject Classification: 47E05, 35K20.


Key words: Green's Function, Non Regular Boundary Conditions, Regular Boundary Conditions, Semi group with Singularities.

The authors are grateful to the editor and the referees for their valuable comments and helpful suggestions which have much improved the presentation of the paper.

Contents

1	Introduction	3
2	Green's Function	5
3	Bounds on the Resolvent	10
	3.1 Estimation of $N(x, s; \lambda)$	10
	3.2 Estimation of the Characteristic Determinant	17
	3.3 Application	26

References



Boundary Value Problem for Second-Order Differential Operators with Mixed Nonlocal Boundary Conditions

M. Denche and A. Kourta

Title Page

Contents



Go Back

Close

Quit

Page 2 of 36

1. Introduction

In the space $L^p(0, 1)$ we consider the boundary value problem


$$(1.1) \quad \begin{cases} L(u) = u'' = f(x), \\ B_1(u) = a_0 u(0) + b_0 u'(0) + c_0 u(1) + d_0 u'(1) = 0, \\ B_2(u) = \int_0^1 R(t) u(t) dt + \int_0^1 S(t) u'(t) dt = 0, \end{cases}$$

where the functions $R, S \in C([0, 1], \mathbb{C})$. We associate to problem (1.1) in space $L^p(0, 1)$ the operator:

$$L_p(u) = u'',$$

with domain $D(L_p) = \{u \in W^{2,p}(0, 1) : B_i(u) = 0, i = 1, 2\}$.

Many papers and books give the full spectral theory of Birkhoff regular differential operators with two point linearly independent boundary conditions, in terms of coefficients of boundary conditions. The reader should refer to [7, 10, 20, 21, 22, 28, 31, 33] and references therein. Few works have been devoted to the study of a non regular situation. The case of separated non regular boundary conditions was studied by W. Eberhard, J.W. Hopkins, D. Jackson, M.V. Keldysh, A.P. Khromov, G. Seifert, M.H. Stone, L.E. Ward (see S. Yakubov and Y. Yakubov [33] for exact references). A situation of non regular non-separated boundary conditions was considered by H. E. Benzinger [2], M. Denche [4], W. Eberhard and G. Freiling [8], M.G. Gasumov and A.M. Mageramov [12, 13], A.P. Khromov [18], Yu. A. Mamedov [19], A.A. Shkalikov [24], Yu. T. Silchenko [26], C. Tretter [29], A.I. Vagabov [30], S. Yakubov [32] and Y. Yakubov [34].



**Boundary Value Problem for
Second-Order Differential
Operators with Mixed Nonlocal
Boundary Conditions**

M. Denche and A. Kourta

Title Page

Contents



Go Back

Close

Quit

Page 3 of 36

A mathematical model with integral boundary conditions was derived by [9, 23] in the context of optical physics. The importance of this kind of problem has been also pointed out by Samarskii [27].

In this paper, we study a problem for second order ordinary differential equations with mixed nonlocal boundary conditions combined with weighted integral boundary conditions and another two point boundary condition. Following the technique in [11, 20, 21, 22], we should bound the resolvent in the space $L^p(0, 1)$ by means of a suitable formula for Green's function. Under certain conditions on the weighting functions and on the coefficients in the boundary conditions, called non regular boundary conditions, we prove that the resolvent decreases with respect to the spectral parameter in $L^p(0, 1)$, but there is no maximal decreasing at infinity for $p \geq 1$. In contrast to the regular case this decreasing is maximal for $p = 1$ as shown in [11]. Furthermore, the studied operator generates in $L^p(0, 1)$ an analytic semi group with singularities for $p \geq 1$. The obtained results are then used to show the correct solvability of a mixed problem for a parabolic partial differential equation with non regular non local boundary conditions.



**Boundary Value Problem for
Second-Order Differential
Operators with Mixed Nonlocal
Boundary Conditions**

M. Denche and A. Kourta

Title Page

Contents



Go Back

Close

Quit

Page 4 of 36

2. Green's Function

Let $\lambda \in \mathbb{C}$, $u_1(x) = u_1(x, \lambda)$ and $u_2(x) = u_2(x, \lambda)$ be a fundamental system of solutions to the equation

$$L(u) - \lambda u = 0.$$

Following [20], the Green's function of problem (1.1) is given by

$$(2.1) \quad G(x, s; \lambda) = \frac{N(x, s; \lambda)}{\Delta(\lambda)},$$

where $\Delta(\lambda)$ is the characteristic determinant of the considered problem defined by

$$(2.2) \quad \Delta(\lambda) = \begin{vmatrix} B_1(u_1) & B_1(u_2) \\ B_2(u_1) & B_2(u_2) \end{vmatrix}$$


and

$$(2.3) \quad N(x, s; \lambda) = \begin{vmatrix} u_1(x) & u_2(x) & g(x, s, \lambda) \\ B_1(u_1) & B_1(u_2) & B_1(g)_x \\ B_2(u_1) & B_2(u_2) & B_2(g)_x \end{vmatrix}$$

for $x, s \in [0, 1]$. The function $g(x, s, \lambda)$ is defined as follows

$$(2.4) \quad g(x, s; \lambda) = \pm \frac{1}{2} \frac{u_1(x)u_2(s) - u_1(s)u_2(x)}{u_1'(s)u_2(s) - u_1(s)u_2'(s)},$$

where it takes the plus sign for $x > s$ and the minus sign for $x < s$.



**Boundary Value Problem for
Second-Order Differential
Operators with Mixed Nonlocal
Boundary Conditions**

M. Denche and A. Kourta

Title Page

Contents



Go Back

Close

Quit

Page 5 of 36

Given an arbitrary $\delta \in (\frac{\pi}{2}, \pi)$, we consider the sector

$$\Sigma_\delta = \{\lambda \in \mathbb{C}, |\arg \lambda| \leq \delta, \lambda \neq 0\}.$$

For $\lambda \in \Sigma_\delta$, define ρ as the square root of λ with positive real part. For $\lambda \neq 0$, we can consider a fundamental system of solutions of the equation $u'' = \lambda u = \rho^2 u$ given by $u_1(t) = e^{-\rho t}$ and $u_2(t) = e^{\rho t}$.

In the following we are going to deduce an adequate formulae for $\Delta(\lambda)$ and $G(x, s; \lambda)$. First of all, for $j = 1, 2$, we have

$$B_1(u_j) = a_0 + (-1)^j b_0 \rho + c_0 e^{(-1)^j \rho} + (-1)^j d_0 \rho e^{(-1)^j \rho},$$

$$B_2(u_j) = \int_0^1 R(t) e^{(-1)^j \rho t} dt + (-1)^j \rho \int_0^1 S(t) e^{(-1)^j \rho t} dt.$$

so we obtain from (2.2)

$$(2.5) \quad \Delta(\lambda) = (a_0 - b_0 \rho + c_0 e^{-\rho} - d_0 \rho e^{-\rho}) \left(\int_0^1 (R(t) + \rho S(t)) e^{\rho t} dt \right) \\ - (a_0 + b_0 \rho + c_0 e^{\rho} + d_0 \rho e^{\rho}) \left(\int_0^1 (R(t) - \rho S(t)) e^{-\rho t} dt \right),$$

and $g(x, s; \lambda)$ has the form

$$g(x, s; \lambda) = \begin{cases} \frac{1}{4\rho} (e^{\rho(x-s)} - e^{\rho(s-x)}) & \text{if } x > s, \\ \frac{1}{4\rho} (e^{\rho(s-x)} - e^{\rho(x-s)}) & \text{if } x < s. \end{cases}$$



**Boundary Value Problem for
Second-Order Differential
Operators with Mixed Nonlocal
Boundary Conditions**

M. Denche and A. Kourta

Title Page

Contents



Go Back

Close

Quit

Page 6 of 36

Thus we have


$$\begin{aligned}
 B_1(g) &= (a_0 - b_0\rho - c_0e^{-\rho} + d_0\rho e^{-\rho}) \frac{e^{\rho s}}{4\rho} \\
 &\quad + (-a_0 - b_0\rho + c_0e^{\rho} + d_0\rho e^{\rho}) \frac{e^{-\rho s}}{4\rho}, \\
 B_2(g) &= \frac{e^{\rho s}}{4\rho} \left(\int_0^s (R(t) - \rho S(t)) e^{-\rho t} dt + \int_s^1 (-R(t) + \rho S(t)) e^{-\rho t} dt \right) \\
 &\quad + \frac{e^{-\rho s}}{4\rho} \left(-\int_0^s (R(t) + \rho S(t)) e^{\rho t} dt + \int_s^1 (R(t) + \rho S(t)) e^{\rho t} dt \right).
 \end{aligned}$$

After a long calculation, formula (2.3) can be written as

$$(2.6) \quad N(x, s; \lambda) = \varphi(x, s; \lambda) + \varphi_i(x, s; \lambda),$$

where

$$\begin{aligned}
 (2.7) \quad \varphi(x, s; \lambda) &= \frac{1}{2\rho} \left[\left(\int_0^1 (a_0 + \rho b_0) (R(t) + \rho S(t)) e^{\rho t} dt \right. \right. \\
 &\quad \left. \left. - e^{\rho} (c_0 + \rho d_0) \int_0^s (R(t) + \rho S(t)) e^{\rho t} dt \right) e^{-\rho(x+s)} \right. \\
 &\quad \left. + \left(\int_s^1 (a_0 - \rho b_0) (R(t) - \rho S(t)) e^{-\rho t} dt \right. \right. \\
 &\quad \left. \left. - e^{\rho} (c_0 - \rho d_0) \int_0^s (R(t) - \rho S(t)) e^{-\rho t} dt \right) e^{\rho(x+s)} \right]
 \end{aligned}$$



**Boundary Value Problem for
Second-Order Differential
Operators with Mixed Nonlocal
Boundary Conditions**

M. Denche and A. Kourta

Title Page

Contents



Go Back

Close

Quit

Page 7 of 36

and the function $\varphi_i(x, s; \lambda)$ is given by


$$(2.8) \quad \varphi_i(x, s; \lambda) = \begin{cases} \varphi_1(x, s; \lambda) & \text{if } x > s, \\ \varphi_2(x, s; \lambda) & \text{if } x < s, \end{cases}$$

with

$$(2.9) \quad \begin{aligned} \varphi_1(x, s; \lambda) &= \frac{e^{\rho(s-x)}}{2\rho} \left(\int_0^s (a_0 + \rho b_0 + c_0 e^\rho + \rho e^\rho d_0) (R(t) - \rho S(t)) e^{-\rho t} dt \right. \\ &\quad \left. - \int_s^1 (a_0 - \rho b_0) (R(t) + \rho S(t)) e^{\rho t} dt \right) \\ &\quad + \frac{e^{\rho(x-s)}}{2\rho} \left(\int_0^s (a_0 - \rho b_0 + c_0 e^{-\rho} - \rho e^{-\rho} d_0) (R(t) + \rho S(t)) e^{\rho t} dt \right. \\ &\quad \left. - \int_0^1 (a_0 + \rho b_0) (R(t) - \rho S(t)) e^{-\rho t} dt \right), \end{aligned}$$

and

$$(2.10) \quad \begin{aligned} \varphi_2(x, s; \lambda) &= \frac{1}{2\rho} \left[\left(- \int_s^1 (a_0 + \rho b_0 + c_0 e^\rho + \rho e^\rho d_0) (R(t) - \rho S(t)) e^{-\rho t} dt \right. \right. \\ &\quad \left. \left. + \int_0^1 (c_0 - \rho d_0) e^{-\rho} (R(t) + \rho S(t)) e^{\rho t} dt \right) e^{\rho(s-x)} \right] \end{aligned}$$



**Boundary Value Problem for
Second-Order Differential
Operators with Mixed Nonlocal
Boundary Conditions**

M. Denche and A. Kourta

Title Page

Contents




Go Back

Close

Quit

Page 8 of 36

$$- \left(\int_s^1 (a_0 - \rho b_0 + c_0 e^{-\rho} - \rho e^{-\rho} d_0) (R(t) + \rho S(t)) e^{\rho t} dt - \int_0^1 (c_0 + \rho d_0) e^{\rho} (R(t) - \rho S(t)) e^{-\rho t} dt \right) e^{\rho(x-s)} \Big].$$



**Boundary Value Problem for
Second-Order Differential
Operators with Mixed Nonlocal
Boundary Conditions**

M. Denche and A. Kourta

Title Page

Contents



Go Back

Close

Quit

Page 9 of 36

3. Bounds on the Resolvent

Every $\lambda \in \mathbb{C}$ such that $\Delta(\lambda) \neq 0$ belongs to $\rho(L_p)$, and the associated resolvent operator $R(\lambda, L_p)$ can be expressed as a Hilbert Schmidt operator

$$(3.1) \quad (\lambda I - L_p)^{-1} f = R(\lambda, L_p) f = - \int_0^1 G(\cdot, s; \lambda) f(s) ds, \quad f \in L^p(0, 1).$$

Then, for every $f \in L^p(0, 1)$ we estimate (3.1)

$$\|R(\lambda; L_p) f\|_{L^p(0,1)} \leq \left(\sup_{0 \leq s \leq 1} \int_0^1 |G(x, s; \lambda)|^p dx \right)^{\frac{1}{p}} \|f\|_{L^p(0,1)},$$

and so we need to bound

$$\left(\sup_{0 \leq s \leq 1} \int_0^1 |G(x, s; \lambda)|^p dx \right)^{\frac{1}{p}} = \frac{1}{|\Delta(\lambda)|} \left(\sup_{0 \leq s \leq 1} \int_0^1 |N(x, s; \lambda)|^p dx \right)^{\frac{1}{p}}.$$

3.1. Estimation of $N(x, s; \lambda)$

We will denote by $\|\cdot\|$ the supremum norm which is defined by $\|R\| = \sup_{0 \leq s \leq 1} |R(s)|$.

Since

$$N(x, s; \lambda) = \begin{cases} \varphi(x, s; \lambda) + \varphi_1(x, s; \lambda) & \text{if } x > s \\ \varphi(x, s; \lambda) + \varphi_2(x, s; \lambda) & \text{if } x < s \end{cases};$$

then

$$(3.2) \quad \|N(x, s; \lambda)\|_{L^p} \leq \|\varphi(x, s; \lambda)\|_{L^p} + \|\varphi_i(x, s; \lambda)\|_{L^p},$$



Boundary Value Problem for
Second-Order Differential
Operators with Mixed Nonlocal
Boundary Conditions

M. Denche and A. Kourta

Title Page

Contents



Go Back

Close

Quit

Page 10 of 36

from (2.8), we have


$$(3.3) \quad \|\varphi_i(x, s; \lambda)\|_{L^p} \leq 2^{1/p} \left\{ \left[\int_0^s |\varphi_2(x, s; \lambda)|^p dx \right]^{\frac{1}{p}} + \left[\int_s^1 |\varphi_1(x, s; \lambda)|^p dx \right]^{\frac{1}{p}} \right\}.$$

From (2.7) we have

$$\begin{aligned} |\varphi(x, s; \lambda)| \leq & \frac{e^{-(x+s) \operatorname{Re} \rho}}{2|\rho|} \left[(\|R\| + |\rho| \|S\|) (|a_0| + |\rho| |b_0|) \int_s^1 e^{t \operatorname{Re} \rho} dt \right. \\ & \left. + (\|R\| + |\rho| \|S\|) (|c_0| + |\rho| |d_0|) e^{\operatorname{Re} \rho} \int_0^s e^{t \operatorname{Re} \rho} dt \right] \\ & + \frac{e^{(x+s) \operatorname{Re} \rho}}{2|\rho|} \left[(\|R\| + |\rho| \|S\|) (|a_0| + |\rho| |b_0|) \int_s^1 e^{-t \operatorname{Re} \rho} dt \right. \\ & \left. + e^{-\operatorname{Re} \rho} (\|R\| + |\rho| \|S\|) (|c_0| + |\rho| |d_0|) \int_0^s e^{-t \operatorname{Re} \rho} dt \right] \end{aligned}$$

then

$$\begin{aligned} |\varphi(x, s; \lambda)| \leq & \frac{e^{-(x+s) \operatorname{Re} \rho}}{2|\rho| \operatorname{Re} \rho} [(\|R\| + |\rho| \|S\|) (|a_0| + |\rho| |b_0|) (e^{\operatorname{Re} \rho} - e^{s \operatorname{Re} \rho}) \\ & + (\|R\| + |\rho| \|S\|) (|c_0| + |\rho| |d_0|) (e^{(s+1) \operatorname{Re} \rho} - e^{\operatorname{Re} \rho})] \\ & + \frac{e^{(x+s) \operatorname{Re} \rho}}{2|\rho| \operatorname{Re} \rho} [(\|R\| + |\rho| \|S\|) (|a_0| + |\rho| |b_0|) (e^{-s \operatorname{Re} \rho} - e^{-\operatorname{Re} \rho}) \\ & + (\|R\| + |\rho| \|S\|) (|c_0| + |\rho| |d_0|) (e^{-\operatorname{Re} \rho} - e^{-(s+1) \operatorname{Re} \rho})] \end{aligned}$$



**Boundary Value Problem for
Second-Order Differential
Operators with Mixed Nonlocal
Boundary Conditions**

M. Denche and A. Kourta

Title Page

Contents



Go Back

Close

Quit


Page 11 of 36

and

$$\begin{aligned} & \int_0^1 |\varphi(x, s; \lambda)|^p dx \\ & \leq \frac{2^p \int_0^1 e^{-px \operatorname{Re} \rho} dx}{(|\rho| \operatorname{Re} \rho)^p} \left[(\|R\| + |\rho| \|S\|)^p (|a_0| + |\rho| |b_0|)^p (e^{(1-s) \operatorname{Re} \rho} - 1)^p \right. \\ & \quad \left. + (\|R\| + |\rho| \|S\|)^p \times (|c_0| + |\rho| |d_0|)^p (e^{\operatorname{Re} \rho} - e^{(1-s) \operatorname{Re} \rho})^p \right] \\ & \quad + \frac{2^p \int_0^1 e^{px \operatorname{Re} \rho} dx}{(|\rho| \operatorname{Re} \rho)^p} \left[(\|R\| + |\rho| \|S\|)^p (|a_0| + |\rho| |b_0|)^p (1 - e^{(s-1) \operatorname{Re} \rho})^p \right. \\ & \quad \left. + (\|R\| + |\rho| \|S\|)^p (|c_0| + |\rho| |d_0|)^p (e^{(s-1) \operatorname{Re} \rho} - e^{-1 \operatorname{Re} \rho})^p \right], \end{aligned}$$

after calculation we obtain

$$\begin{aligned} & \left(\int_0^1 |\varphi(x, s; \lambda)|^p dx \right)^{\frac{1}{p}} \\ & \leq \frac{2^{1+\frac{2}{p}} e^{\operatorname{Re} \rho}}{|\rho| (\operatorname{Re} \rho)^{1+\frac{1}{p}} p^{1/p}} \left[(\|R\| + |\rho| \|S\|) (|a_0| + |\rho| |b_0|) [(e^{-s \operatorname{Re} \rho} - e^{-\operatorname{Re} \rho}) \right. \\ & \quad \times (1 - e^{-p \operatorname{Re} \rho})^{\frac{1}{p}} + (1 - e^{-p \operatorname{Re} \rho})^{\frac{1}{p}} (1 - e^{(s-1) \operatorname{Re} \rho})] + (\|R\| + |\rho| \|S\|) \\ & \quad \times (|c_0| + |\rho| |d_0|) \left[(1 - e^{-p \operatorname{Re} \rho})^{\frac{1}{p}} (1 - e^{-s \operatorname{Re} \rho}) \right. \\ & \quad \left. \left. + (1 - e^{-p \operatorname{Re} \rho})^{\frac{1}{p}} (e^{(s-1) \operatorname{Re} \rho} - e^{-\operatorname{Re} \rho}) \right] \right] \end{aligned}$$



**Boundary Value Problem for
Second-Order Differential
Operators with Mixed Nonlocal
Boundary Conditions**

M. Denche and A. Kourta

Title Page

Contents



Go Back

Close

Quit


Page 12 of 36

as $\operatorname{Re} \rho > 0$ so

$$(3.4) \quad \sup_{0 \leq s \leq 1} \left(\int_0^1 |\varphi(x, s; \lambda)|^p dx \right)^{\frac{1}{p}} \leq \frac{2^{1+\frac{2}{p}} e^{\operatorname{Re} \rho} (\|R\| + |\rho| \|S\|) [(|a_0| + |c_0|) + |\rho| (|b_0| + |d_0|)]}{|\rho| (\operatorname{Re} \rho)^{1+\frac{1}{p}} p^{1/p}}.$$

From (2.9) we have

$$\begin{aligned} |\varphi_1(x, s; \lambda)| \leq & \frac{e^{(s-x)\operatorname{Re} \rho}}{2|\rho|} \left[[(|a_0| + |\rho| |b_0|) + e^{\operatorname{Re} \rho} (|c_0| + |\rho| |d_0|)] \right. \\ & \times (\|R\| + |\rho| \|S\|) \int_0^s e^{-t\operatorname{Re} \rho} dt \\ & \left. + (|a_0| + |\rho| |b_0|) (\|R\| + |\rho| \|S\|) \int_0^1 e^{t\operatorname{Re} \rho} dt \right] \\ & + \frac{e^{(x-s)\operatorname{Re} \rho}}{2|\rho|} \left[(\|R\| + |\rho| \|S\|) \left[(|a_0| + |\rho| |b_0|) \right. \right. \\ & \left. \left. + e^{\operatorname{Re} \rho} (|c_0| + |\rho| |d_0|) \right] \int_0^s e^{t\operatorname{Re} \rho} dt \right. \\ & \left. + (|a_0| + |\rho| |b_0|) (\|R\| + |\rho| \|S\|) \int_0^1 e^{-t\operatorname{Re} \rho} dt \right] \end{aligned}$$



**Boundary Value Problem for
Second-Order Differential
Operators with Mixed Nonlocal
Boundary Conditions**

M. Denche and A. Kourta

Title Page

Contents



Go Back

Close

Quit

Page 13 of 36

then


$$\begin{aligned}
 |\varphi_1(x, s; \lambda)| \leq & \frac{e^{(s-x)\operatorname{Re} \rho}}{2|\rho|\operatorname{Re} \rho} [(|a_0| + |\rho| |b_0|) (\|R\| + |\rho| \|S\|) (e^{\operatorname{Re} \rho} - e^{-s\operatorname{Re} \rho}) \\
 & + (|c_0| + |\rho| |d_0|) (\|R\| + |\rho| \|S\|) (e^{\operatorname{Re} \rho} - e^{(1-s)\operatorname{Re} \rho})] \\
 & + \frac{e^{(x-s)\operatorname{Re} \rho}}{2|\rho|\operatorname{Re} \rho} [(\|R\| + |\rho| \|S\|) (|a_0| + |\rho| |b_0|) (e^{s\operatorname{Re} \rho} - e^{-\operatorname{Re} \rho}) \\
 & + (|c_0| + |\rho| |d_0|) (\|R\| + |\rho| \|S\|) (e^{(s-1)\operatorname{Re} \rho} - e^{-\operatorname{Re} \rho})]
 \end{aligned}$$

and

$$\begin{aligned}
 & \int_s^1 |\varphi_1(x, s; \lambda)|^p dx \\
 & \leq \frac{2^p \int_s^1 e^{-px\operatorname{Re} \rho} dx}{|\rho|^p (\operatorname{Re} \rho)^p} [(|a_0| + |\rho| |b_0|)^p (\|R\| + |\rho| \|S\|)^p (e^{(1+s)\operatorname{Re} \rho} - 1)^p \\
 & \quad + (|c_0| + |\rho| |d_0|)^p (\|R\| + |\rho| \|S\|)^p (e^{(1+s)\operatorname{Re} \rho} - e^{\operatorname{Re} \rho})^p] \\
 & \quad + \frac{2^p \int_s^1 e^{px\operatorname{Re} \rho} dx}{|\rho|^p (\operatorname{Re} \rho)^p} [(|a_0| + |\rho| |b_0|)^p (\|R\| + |\rho| \|S\|)^p (1 - e^{-(1+s)\operatorname{Re} \rho})^p \\
 & \quad + (|c_0| + |\rho| |d_0|)^p (\|R\| + |\rho| \|S\|)^p (e^{-\operatorname{Re} \rho} - e^{-(1+s)\operatorname{Re} \rho})^p],
 \end{aligned}$$

this yields

$$\begin{aligned}
 \int_s^1 |\varphi_1(x, s; \lambda)|^p dx \leq & \frac{2^{p+1} e^{p\operatorname{Re} \rho}}{|\rho|^p (\operatorname{Re} \rho)^{p+1}} [(|a_0| + |\rho| |b_0|)^p (\|R\| + |\rho| \|S\|)^p \\
 & \times (1 - e^{-(1+s)\operatorname{Re} \rho})^p (1 - e^{p(s-1)\operatorname{Re} \rho})
 \end{aligned}$$



**Boundary Value Problem for
Second-Order Differential
Operators with Mixed Nonlocal
Boundary Conditions**

M. Denche and A. Kourta

Title Page

Contents



Go Back

Close

Quit

Page 14 of 36

$$+ (|c_0| + |\rho| |d_0|)^p (\|R\| + |\rho| \|S\|)^p (1 - e^{-s \operatorname{Re} \rho})^p (1 - e^{p(s-1) \operatorname{Re} \rho})]$$

as $\operatorname{Re} \rho > 0$ we obtain


$$(3.5) \quad \sup_{0 \leq s \leq 1} \left(\int_s^1 |\varphi_1(x, s; \lambda)|^p dx \right)^{\frac{1}{p}} \leq \frac{2^{1+\frac{2}{p}} e^{\operatorname{Re} \rho} (\|R\| + |\rho| \|S\|) [(|a_0| + |c_0|) + |\rho| (|b_0| + |d_0|)]}{|\rho| (\operatorname{Re} \rho)^{1+\frac{1}{p}} p^{1/p}}.$$

From (2.10) we have

$$\begin{aligned} & |\varphi_2(x, s; \lambda)| \\ & \leq \frac{e^{(x-s) \operatorname{Re} \rho}}{2 |\rho|} \left[(\|R\| + |\rho| \|S\|) (|c_0| + |\rho| |d_0|) \int_0^1 e^{(1-t) \operatorname{Re} \rho} dt \right. \\ & \quad \left. + (\|R\| + |\rho| \|S\|) [(|a_0| + |\rho| |b_0|) + e^{-\operatorname{Re} \rho} (|c_0| + |\rho| |d_0|)] \int_0^1 e^{t \operatorname{Re} \rho} dt \right] \\ & \quad + \frac{e^{(s-x) \operatorname{Re} \rho}}{2 |\rho|} [(|c_0| + |\rho| |d_0|) (\|R\| + |\rho| \|S\|) \int_0^1 e^{(t-1) \operatorname{Re} \rho} dt \\ & \quad + [(|a_0| + |\rho| |b_0|) + e^{\operatorname{Re} \rho} (|c_0| + |\rho| |d_0|)] (\|R\| + |\rho| \|S\|) \int_s^1 e^{-t \operatorname{Re} \rho} dt] \end{aligned}$$

then

$$\begin{aligned} |\varphi_2(x, s; \lambda)| & \leq \frac{e^{x \operatorname{Re} \rho}}{2 |\rho| \operatorname{Re} \rho} [(\|R\| + |\rho| \|S\|) (|c_0| + |\rho| |d_0|) (e^{(1-s) \operatorname{Re} \rho} - e^{-\operatorname{Re} \rho}) \\ & \quad + (|a_0| + |\rho| |b_0|) (\|R\| + |\rho| \|S\|) (e^{(1-s) \operatorname{Re} \rho} - 1)] \end{aligned}$$



**Boundary Value Problem for
Second-Order Differential
Operators with Mixed Nonlocal
Boundary Conditions**

M. Denche and A. Kourta

Title Page

Contents




Go Back

Close

Quit

Page 15 of 36



Title Page

Contents



Go Back

Close

Quit

Page 16 of 36

$$+ \frac{e^{-x \operatorname{Re} \rho}}{2|\rho| \operatorname{Re} \rho} [(\|R\| + |\rho| \|S\|) (|c_0| + |\rho| |d_0|) (e^{\operatorname{Re} \rho} - e^{(s-1) \operatorname{Re} \rho}) + (|a_0| + |\rho| |b_0|) (\|R\| + |\rho| \|S\|) (1 - e^{(s-1) \operatorname{Re} \rho})].$$

So

$$\begin{aligned} & \int_0^s |\varphi_2(x, s; \lambda)|^p dx \\ & \leq \frac{2^{p+1} e^{p \operatorname{Re} \rho}}{p |\rho|^p (\operatorname{Re} \rho)^{p+1}} [(\|R\| + |\rho| \|S\|)^p (|c_0| + |\rho| |d_0|)^p (1 - e^{(s-2) \operatorname{Re} \rho})^p \\ & \quad \times (1 - e^{-ps \operatorname{Re} \rho}) + (|a_0| + |\rho| |b_0|)^p \\ & \quad \times (\|R\| + |\rho| \|S\|)^p (1 - e^{-ps \operatorname{Re} \rho}) (1 - e^{(s-1) \operatorname{Re} \rho})^p] \end{aligned}$$

as $\operatorname{Re} \rho > 0$ we obtain

$$(3.6) \quad \sup_{0 \leq s \leq 1} \left(\int_0^s |\varphi_2(x, s; \lambda)|^p dx \right)^{\frac{1}{p}} \leq \frac{2^{1+\frac{2}{p}} e^{\operatorname{Re} \rho} (\|R\| + |\rho| \|S\|) [(|a_0| + |c_0|) + |\rho| (|b_0| + |d_0|)]}{|\rho| (\operatorname{Re} \rho)^{1+\frac{1}{p}} p^{1/p}}.$$

From (3.4), (3.5) and (3.6), we obtain

$$\begin{aligned} & \|R(\lambda, L_p)\| \\ & \leq \frac{2^{1+\frac{2}{p}} e^{\operatorname{Re} \rho} (\|R\| + |\rho| \|S\|) [(|a_0| + |c_0|) + |\rho| (|b_0| + |d_0|)]}{|\Delta(\lambda)| |\rho| (\operatorname{Re} \rho)^{1+\frac{1}{p}} p^{1/p}} \left(2^{1+\frac{2}{p}} + 1 \right), \end{aligned}$$

for $\rho \in \Sigma_{\frac{\delta}{2}} = \{\rho \in \mathbb{C} : |\arg \rho| \leq \frac{\delta}{2}, \rho \neq 0\}$, we have $(\operatorname{Re} \rho)^{-1} < \frac{1}{|\rho| \cos(\frac{\delta}{2})}$, then

$$\begin{aligned} & \|R(\lambda, L_p)\| \\ & \leq \frac{2^{1+\frac{2}{p}} \left(2^{1+\frac{2}{p}} + 1\right) e^{\operatorname{Re} \rho} (\|R\| + |\rho| \|S\|) [(|a_0| + |c_0|) + |\rho| (|b_0| + |d_0|)]}{|\Delta(\lambda)| |\rho| \left(|\rho|^{1+\frac{1}{p}}\right) \left(\cos \frac{\delta}{2}\right)^{1+\frac{1}{p}} p^{1/p}}. \end{aligned}$$

Finally, we obtain for $\lambda = \rho^2 \in \Sigma_\delta$


$$(3.7) \quad \begin{aligned} & \|R(\lambda, L_p)\|_{L^p} \\ & \leq \frac{c e^{\operatorname{Re} \rho}}{|\Delta(\lambda)| |\rho|^{1+\frac{1}{p}}} \left(\frac{\|R\|}{|\rho|} + \|S\|\right) [(|a_0| + |c_0|) + |\rho| (|b_0| + |d_0|)], \end{aligned}$$

where

$$c = \frac{2^{1+\frac{2}{p}} \left(2^{1+\frac{2}{p}} + 1\right)}{p^{1/p} \left(\cos \frac{\delta}{2}\right)^{1+\frac{1}{p}}}.$$

3.2. Estimation of the Characteristic Determinant

The next step is to determine the cases for which $|\Delta(\lambda)|$ remains bounded below. It will then be necessary to bound $|\Delta(\lambda)|$ appropriately. However, formula (2.5) is not useful for this purpose. It will be then necessary to make some additional regularity hypotheses on the functions R and S , and so we assume that the functions R and S are in $C^2([0, 1], \mathbb{C})$.



**Boundary Value Problem for
Second-Order Differential
Operators with Mixed Nonlocal
Boundary Conditions**

M. Denche and A. Kourta

Title Page

Contents



Go Back

Close

Quit

Page 17 of 36


Integrating twice by parts in (2.5), we obtain

$$(3.8) \quad \Delta(\lambda) = e^\rho \left[\rho(d_0S(0) - b_0S(1)) + (d_0S'(0) + b_0S'(1) + a_0S(1) - b_0R(1) - d_0R(0) + c_0S(0)) + \frac{1}{\rho}(a_0R(1) - c_0R(0) + b_0R'(1) - a_0S'(1) - d_0R'(0) + c_0S'(0)) + \frac{1}{\rho^2}(-a_0R'(1) - c_0R'(0)) + \Phi(\rho) \right],$$

where

$$(3.9) \quad \Phi(\rho) = \left[(a_0 - \rho b_0)e^{-\rho} + (c_0 - \rho d_0)e^{-2\rho} \int_0^1 \left(\frac{R''(t)}{\rho^2} - \frac{S''(t)}{\rho} \right) e^{\rho t} dt + [(a_0 + \rho b_0)e^{-\rho} + (c_0 + \rho d_0)] \int_0^1 \left(\frac{R''(t)}{\rho^2} - \frac{S''(t)}{\rho} \right) e^{-\rho t} dt + e^{-\rho} \left[\frac{1}{\rho} \left(-b_0R'(0) + d_0R'(1) - c_0S'(1) + a_0S'(0) + c_0R(1) - a_0R(0) \right) + \rho(b_0S(0) - d_0S(1)) \right] + 2e^{-2\rho} [(a_0 + \rho b_0)(R(1) - \rho S(1)) - (c_0 - \rho d_0)(R(0) + \rho S(0)) + \frac{1}{\rho^2}(a_0 + \rho b_0)(R'(1) - \rho S'(1)) + \frac{1}{\rho^2}(c_0 - \rho d_0)(R'(0) + \rho S'(0))] \right].$$

After a straightforward calculation, we obtain the following inequality valid for



Boundary Value Problem for Second-Order Differential Operators with Mixed Nonlocal Boundary Conditions

M. Denche and A. Kourta

Title Page

Contents



Go Back

Close

Quit


Page 18 of 36

$\rho \in \Sigma_{\frac{\delta}{2}}$, with $|\rho|$ sufficiently large,

$$\begin{aligned}
 & |\Phi(\rho)| \\
 & \leq [(|a_0| + |\rho| |b_0|) e^{-\operatorname{Re} \rho} + (|c_0| + |\rho| |d_0|) e^{-2\operatorname{Re} \rho}] \\
 & \quad \times \left(\frac{\|R''\|}{|\rho|^2} + \frac{\|S''\|}{|\rho|} \right) \left(\frac{e^{\operatorname{Re} \rho} - 1}{\operatorname{Re} \rho} \right) \\
 & + [(|a_0| + |\rho| |b_0|) e^{-\operatorname{Re} \rho} + (|c_0| + |\rho| |d_0|)] \\
 & \quad \times \left(\frac{\|R''\|}{|\rho|^2} + \frac{\|S''\|}{|\rho|} \right) \left(\frac{1 - e^{-\operatorname{Re} \rho}}{\operatorname{Re} \rho} \right) \\
 & + 2e^{-\operatorname{Re} \rho} \left[\frac{1}{|\rho|} \left| -b_0 R'(0) + d_0 R'(1) - c_0 S'(1) + a_0 S'(0) + c_0 R(1) - a_0 R(0) \right| \right. \\
 & + |\rho| |b_0 S(0) - d_0 S(1)| \left. + e^{-2\operatorname{Re} \rho} \left[\frac{1}{|\rho|} (|a_0| + |\rho| |b_0|) (\|R\| + |\rho| \|S\|) \right. \right. \\
 & + \frac{1}{|\rho|} (|c_0| + |\rho| |d_0|) (\|R\| + |\rho| \|S\|) + (|a_0| + |\rho| |b_0|) \left. \left. \left(\frac{\|R'\| + |\rho| \|S'\|}{|\rho|^2} \right) \right. \right. \\
 & \left. \left. + (|c_0| + |\rho| |d_0|) \left(\frac{\|R'\| + |\rho| \|S'\|}{|\rho|^2} \right) \right] \right].
 \end{aligned}$$

Then

$$|\Phi(\rho)| \leq \frac{2}{|\rho| \cos \frac{\delta}{2}} \left[\frac{(|a_0| + |c_0|)}{|\rho|} + (|b_0| + |d_0|) \left(\frac{\|R''\|}{|\rho|} + \|S''\| \right) \right]$$



**Boundary Value Problem for
Second-Order Differential
Operators with Mixed Nonlocal
Boundary Conditions**

M. Denche and A. Kourta

Title Page

Contents




Go Back

Close

Quit

Page 19 of 36



**Boundary Value Problem for
Second-Order Differential
Operators with Mixed Nonlocal
Boundary Conditions**

M. Denche and A. Kourta

Title Page

Contents



Go Back

Close

Quit

Page 20 of 36

$$\begin{aligned}
 & + \frac{1}{|\rho| \cos \frac{\delta}{2}} \left[\frac{1}{|\rho|^2} \left| -b_0 R'(0) + d_0 R'(1) - c_0 S'(1) \right. \right. \\
 & + \left. \left. + a_0 S'(0) + c_0 R(1) - a_0 R(0) \right| |b_0 S(0) - d_0 S(1)| \right] \\
 & + \frac{1}{|\rho| (\cos \frac{\delta}{2})^2} \left[\left(\frac{|a_0|}{|\rho|} + |b_0| \right) \left(\frac{\|R\|}{|\rho|} + \|S\| \right) \right. \\
 & + \left(\frac{|c_0|}{|\rho|} + |d_0| \right) \left(\frac{\|R\|}{|\rho|} + \|S\| \right) \\
 & + \left(\frac{|a_0|}{|\rho|} + |b_0| \right) \left(\frac{\|R'\|}{|\rho|^2} + \frac{\|S'\|}{|\rho|} \right) \\
 (3.10) \quad & \left. \left. + \left(\frac{|c_0|}{|\rho|} + |d_0| \right) \left(\frac{\|R'\|}{|\rho|^2} + \frac{\|S'\|}{|\rho|} \right) \right] .
 \end{aligned}$$

Where we have used that $Re(\rho) \geq |\rho| \cos(\frac{\delta}{2})$, $(1 - e^{-Re \rho}) < 1$,

$$e^{-Re \rho} \leq \frac{2}{|\rho|^2 (\cos \frac{\delta}{2})^2} \text{ and } e^{-2Re \rho} \leq \frac{1}{2 |\rho|^2 (\cos \frac{\delta}{2})^2}, e^{-Re \rho} < 1.$$

There are several cases to analyze depending on the functions R and S .

• **Case 1.**

Suppose that $S \neq 0$, $d_0 \neq 0$, $b_0 \neq 0$, $d_0 S(0) - b_0 S(1) = 0$ and

$$k_1 = d_0 S'(0) + b_0 S'(1) + a_0 S(1) - b_0 R(1) - d_0 R(0) + c_0 S(0) \neq 0.$$

From (3.8), we have for $|\rho|$ sufficiently large


$$\begin{aligned}
 |\Delta(\lambda)| &\geq e^{\operatorname{Re} \rho} \left[\left| d_0 S'(0) + b_0 S'(1) + a_0 S(1) - b_0 R(1) - d_0 R(0) + c_0 S(0) \right| \right. \\
 &\quad \left. - \frac{1}{|\rho|} \left| a_0 R(1) - c_0 R(0) + b_0 R'(1) - a_0 S'(1) - d_0 R'(0) + c_0 S'(0) \right| \right. \\
 &\quad \left. - \frac{1}{|\rho|^2} \left| a_0 R'(1) + c_0 R'(0) \right| - |\Phi(\rho)| \right].
 \end{aligned}$$

From (3.10) we deduce for $\rho \in \Sigma_{\frac{\delta}{2}}$, $|\rho| \geq r_0 > 0$.

$$|\Phi(\rho)| \leq \frac{c(r_0)}{|\rho|}.$$

Then, we have

$$\begin{aligned}
 |\Delta(\lambda)| &\geq e^{\operatorname{Re} \rho} \left[\left| d_0 S'(0) + b_0 S'(1) + a_0 S(1) - b_0 R(1) - d_0 R(0) + c_0 S(0) \right| \right. \\
 &\quad \left. - \frac{1}{|\rho|} \left| a_0 R(1) - c_0 R(0) + b_0 R'(1) - a_0 S'(1) - d_0 R'(0) + c_0 S'(0) \right| \right. \\
 &\quad \left. - \frac{1}{|\rho|^2} \left| a_0 R'(1) + c_0 R'(0) \right| - \frac{c(r_0)}{|\rho|} \right].
 \end{aligned}$$



**Boundary Value Problem for
Second-Order Differential
Operators with Mixed Nonlocal
Boundary Conditions**

M. Denche and A. Kourta

Title Page

Contents



Go Back

Close

Quit

Page 21 of 36

we can now choose $r_0 > 0$, such that

$$\begin{aligned} & \frac{1}{r_0} \left| a_0 R(1) - c_0 R(0) + b_0 R'(1) - a_0 S'(1) - d_0 R'(0) + c_0 S'(0) \right| \\ & + \frac{1}{r_0^2} \left| a_0 R'(1) + c_0 R'(0) \right| + \frac{c(r_0)}{|\rho|} \\ & \leq \frac{1}{2} \left| d_0 S'(0) + b_0 S'(1) + a_0 S(1) - b_0 R(1) - d_0 R(0) + c_0 S(0) \right|, \end{aligned}$$

then, for $\rho \in \Sigma_{\frac{\delta}{2}}$, $|\rho| \geq r_0 > 0$, we get

$$|\Delta(\lambda)| \geq \frac{e^{\operatorname{Re} \rho}}{2} |k_1|.$$

From (3.7), we deduce the following bound, valid for every $|\arg \rho| \leq \frac{\delta}{2}$, $\rho \neq 0$

$$\|R(\lambda, L_p)\| \leq \frac{2c}{|\rho|^{\frac{1}{p}} |k_1|} \left(\left(\frac{|a_0| + |c_0|}{|\rho|} \right) + (|b_0| + |d_0|) \right) \left(\frac{\|R\|}{|\rho|} + \|S\| \right),$$


then

$$\|R(\lambda, L_p)\| \leq \frac{c_1}{|\lambda|^{\frac{1}{2p}}},$$

as $|\lambda| \rightarrow +\infty$, where

$$c_1 = \frac{2c \|S\| (|b_0| + |d_0|)}{|k_1|}.$$

- **Case 2.**



**Boundary Value Problem for
Second-Order Differential
Operators with Mixed Nonlocal
Boundary Conditions**

M. Denche and A. Kourta

Title Page

Contents



Go Back

Close

Quit

Page 22 of 36

If $b_0 = d_0 = 0$, $S \neq 0$ and $a_0S(1) + c_0S(0) = 0$, with

$$k_2 = a_0R(1) - c_0R(0) - a_0S'(1) + c_0S'(0) \neq 0,$$

we have the following bound, valid for $\lambda \in \Sigma_\delta$ and $|\lambda|$ big enough,

$$\|R(\lambda, L_p)\| \leq \frac{2c(|a_0| + |c_0|)}{|\rho|^{\frac{1}{p}} |k_2|} \left(\frac{\|R\|}{|\rho|} + \|S\| \right),$$

then

$$\|R(\lambda, L_p)\| \leq \frac{c_1}{|\lambda|^{\frac{1}{2p}}},$$

as $|\lambda| \rightarrow +\infty$, where

$$c_1 = \frac{2c(|a_0| + |c_0|) \|S\|}{|k_2|}.$$


• **Case 3.**

If $S = 0$, $b_0 \neq 0$, $d_0 \neq 0$ and $b_0R(1) + d_0R(0) = 0$, with

$$k_3 = a_0R(1) - c_0R(0) + b_0R'(1) - d_0R'(0) \neq 0.$$

Similarly, we get

$$\|R(\lambda, L_p)\| \leq \frac{2c\|R\|}{|\rho|^{1+\frac{1}{p}} |k_3|} \left(\frac{(|a_0| + |c_0|)}{|\rho|} + (|b_0| + |d_0|) \right),$$



**Boundary Value Problem for
Second-Order Differential
Operators with Mixed Nonlocal
Boundary Conditions**

M. Denche and A. Kourta

Title Page

Contents



Go Back

Close

Quit

Page 23 of 36

then, we have

$$\|R(\lambda, L_p)\| \leq \frac{c_1}{|\lambda|^{\frac{1}{2p}}},$$

as $|\lambda| \rightarrow +\infty$, where

$$c_1 = \frac{2c \|R\| (|b_0| + |d_0|)}{|k_3|}.$$

• **Case 4.**

If $b_0 = d_0 = 0$, $S = 0$ and $a_0R(1) - c_0R(0) = 0$ with

$$k_4 = a_0R'(1) - c_0R'(0) \neq 0.$$

Again in this case, we have


$$\|R(\lambda, L_p)\| \leq \frac{2c(|a_0| + |c_0|) \|R\|}{|\rho|^{\frac{1}{p}} |k_4|},$$

then

$$\|R(\lambda, L_p)\| \leq \frac{c_1}{|\lambda|^{\frac{1}{2p}}},$$

as $|\lambda| \rightarrow +\infty$, where

$$c_1 = \frac{c \|R\| (|a_0| + |c_0|)}{|k_4|}.$$



**Boundary Value Problem for
Second-Order Differential
Operators with Mixed Nonlocal
Boundary Conditions**

M. Denche and A. Kourta

Title Page

Contents



Go Back

Close

Quit

Page 24 of 36

Definition 3.1. The boundary value conditions in (1.1) are called non regular if the functions $R, S \in C^2([0, 1], \mathbb{C})$ and if and only if one of the following conditions holds

1- $d_0 S(0) - b_0(1) = 0, b_0 \neq 0, d_0 \neq 0, S \neq 0$ and

$$d_0 S'(0) + b_0 S'(1) + a_0 S(1) - b_0 R(1) - d_0 R(0) + c_0 S(0) \neq 0$$

2- $b_0 = d_0 = 0, \|S\| \neq 0, a_0 S(1) + c_0 S(0) = 0$ with

$$a_0 R(1) - c_0 R(0) - a_0 S'(1) + c_0 S'(0) \neq 0$$

3- $S = 0, b_0 \neq 0, d_0 \neq 0, b_0 R(1) + d_0 R(0) \neq 0$ with

$$a_0 R(1) - c_0 R(0) + b_0 R'(1) - d_0 R'(0) \neq 0$$

4- $b_0 = d_0 = 0, S = 0, a_0 R(1) - c_0 R(0) = 0$ with


$$a_0 R'(1) - c_0 R'(0) \neq 0.$$

This proves the following theorem

Theorem 3.1. If the boundary value conditions in (1.1) are non regular, then $\Sigma_\delta \subset \rho(L_p)$ for sufficiently large $|\lambda|$ and there exists $c > 0$ such that

$$\|R(\lambda, L_p)\| \leq \frac{c}{|\lambda|^{\frac{1}{2p}}}.$$

Remark 3.1. From Theorem 3.1 it follows that the operator L_p , for $p \neq \infty$, generates an analytic semigroup with singularities [25] of type $A(2p - 1, 4p - 1)$.



Boundary Value Problem for
Second-Order Differential
Operators with Mixed Nonlocal
Boundary Conditions

M. Denche and A. Kourta

Title Page

Contents



Go Back

Close

Quit

Page 25 of 36


3.3. Application

In the following, we apply the above obtained results to the study of a class of a mixed problem for a parabolic equation with an weighted integral boundary condition combined with another two point boundary condition of the form

$$(3.11) \quad \begin{cases} \frac{\partial u(t, x)}{\partial t} - a \frac{\partial^2 u(t, x)}{\partial x^2} = f(t, x) \\ L_1(u) = a_0 u(0, t) + b_0 u'(0, t) + c_0 u(1, t) + d_0 u'(1, t) = 0, \\ L_2(u) = \int_0^1 R(\xi) u(t, \xi) d\xi + \int_0^1 S(\xi) u'(t, \xi) d\xi = 0, \\ u(0, x) = u_0(x), \end{cases}$$

where $(t, \xi) \in [0, T] \times [0, 1]$. Boundary value problems for parabolic equations with integral boundary conditions are studied by [1, 3, 5, 6, 14, 15, 16, 17, 35] using various methods. For instance, the potential method in [3] and [17], Fourier method in [1, 14, 15, 16] and the energy inequalities method has been used in [5, 6, 35]. In our case, we apply the method of operator differential equation. The study of the problem is then reduced to a cauchy problem for a parabolic abstract differential equation, where the operator coefficients has been previously studied. For this purpose, let E , E_1 , and E_2 be Banach spaces. Introduce two Banach spaces

$$C_\mu((0, T], E) = \left\{ f/f \in C((0, T], E), \|f\| = \sup_{t \in (0, T]} \|t^\mu f(t)\| < \infty \right\}, \mu \geq 0,$$



**Boundary Value Problem for
Second-Order Differential
Operators with Mixed Nonlocal
Boundary Conditions**

M. Denche and A. Kourta

Title Page

Contents



Go Back

Close

Quit

Page 26 of 36

$$C_{\mu}^{\gamma}((0, T], E) = \left\{ f/f \in C((0, T], E), \|f\| = \sup_{t \in (0, T]} \|t^{\mu} f(t)\| \right. \\ \left. + \sup_{0 < t < t+h \leq T} \|f(t+h) - f(t)\| h^{-\gamma} t^{\mu} < \infty \right\}, \\ \mu \geq 0, \gamma \in (0, 1],$$

and a linear space

$$C^1((0, T], E_1, E_2) = \{f/f \in C((0, T], E_1) \cap C^1((0, T], E_2)\}, E_1 \subset E_2,$$

where $C((0, T], E)$ and $C^1((0, T], E)$ are spaces of continuous and continuously differentiable, respectively, vector-function from $(0, T]$ into E . We denote, for a linear operator A in a Banach space E , by


$$E(A) = \left\{ u/u \in D(A), \|u\|_{E(A)} = (\|Au\|^2 + \|u\|^2)^{\frac{1}{2}} \right\},$$

and

$$C^1((0, T], E(A), E) = \left\{ f/f \in C((0, T], E(A)), f' \in C((0, T], E) \right\}.$$

Let us derive a theorem which was proved by various methods in [25] and [31, 33]. Consider, in a Banach space E , the Cauchy problem

$$(3.12) \quad \begin{cases} u'(t) = Au(t) + f(t), & t \in [0, T], \\ u(0) = u_0, \end{cases}$$



**Boundary Value Problem for
Second-Order Differential
Operators with Mixed Nonlocal
Boundary Conditions**

M. Denche and A. Kourta

Title Page

Contents



Go Back

Close

Quit

Page 27 of 36

where A is, generally speaking, unbounded linear operator in E , u_0 is a given element of E , $f(t)$ is a given vector-function and $u(t)$ is an unknown vector-function in E .

Theorem 3.2. *Let the following conditions be satisfied:*

1. A is a closed linear operator in a Banach space E and for some $\beta \in (0, 1]$, $\alpha > 0$

$$\|R(\lambda, A)\| \leq C |\lambda|^{-\beta}, \quad |\arg \lambda| \leq \frac{\pi}{2} + \alpha, \quad |\lambda| \rightarrow \infty;$$

2. $f \in C_\mu^\gamma((0, T], E)$ for some $\gamma \in (1 - \beta, 1]$, $\mu \in [0, \beta)$;

3. $u_0 \in D(A)$.


Then the Cauchy problem (3.12) has a unique solution

$$u \in C([0, T], E) \cap C^1((0, T], E(A), E)$$

and for the solution u the following estimates hold

$$\|u(t)\| \leq C \left(\|Au_0\| + \|u_0\| + \|f\|_{C_\mu^\gamma((0,t],E)} \right), \quad t \in (0, T],$$

$$\begin{aligned} & \left\| u'(t) \right\| + \|Au(t)\| \\ & \leq C \left(t^{\beta-1} (\|Au_0\| + \|u_0\|) + t^{\beta-\mu-1} \|f\|_{C_\mu^\gamma((0,t],E)} \right), \quad t \in (0, T]. \end{aligned}$$



**Boundary Value Problem for
Second-Order Differential
Operators with Mixed Nonlocal
Boundary Conditions**

M. Denche and A. Kourta

Title Page

Contents



Go Back

Close

Quit

Page 28 of 36

As a result of this we get the following theorem

Theorem 3.3. *Let the following conditions be satisfied*

1. $a \neq 0, |\arg a| < \frac{\pi}{2},$
2. *the functions $R(t), S(t) \in C^2([0, 1], \mathbb{C})$ and one of the following conditions is satisfied*
 $d_0 S(0) - b_0 S(1) = 0, b_0 \neq 0, d_0 \neq 0,$ and $S \neq 0$

$$d_0 S'(0) + b_0 S'(1) + a_0 S(1) - b_0 R(1) - d_0 R(0) + c_0 S(0) \neq 0,$$

or $b_0 = 0, d_0 = 0, S \neq 0, a_0 S(1) + c_0 S(0) = 0$ with

$$a_0 R(1) - c_0 R(0) - a_0 S'(1) + c_0 S'(0) \neq 0,$$

or $b_0 \neq 0, d_0 \neq 0, S = 0, b_0 R(1) + d_0 R(0) = 0$ with


$$a_0 R(1) - c_0 R(0) + b_0 R'(1) - d_0 R'(0) \neq 0,$$

or $b_0 = d_0 = 0, S = 0, a_0 R(1) - c_0 R(0) = 0$ with

$$a_0 R'(1) - c_0 R'(0) \neq 0.$$

3. $f \in C_{\mu}^{\gamma}((0, T], L^q(0, 1))$ for some $\gamma \in \left(1 - \frac{1}{2q}, 1\right]$ and some $\mu \in \left[0, \frac{1}{2q}\right),$

4. $u_0 \in W_q^2((0, 1), L_i u = 0, i = \overline{1, 2}).$



**Boundary Value Problem for
Second-Order Differential
Operators with Mixed Nonlocal
Boundary Conditions**

M. Denche and A. Kourta

Title Page

Contents



Go Back

Close

Quit

Page 29 of 36

Then the problem (3.11) has a unique solution

$$u \in C((0, T], L^q(0, 1)) \cap C^1((0, T], W_q^2(0, 1), L^q(0, 1))$$

and for this solution we have the estimates:

$$(3.13) \quad \|u(t, \cdot)\|_{L^q(0,1)} \leq c \left(\|u_0\|_{W_q^2(0,1)} + \|f\|_{C_\mu((0,t], L^q(0,1))} \right), \quad t \in (0, T],$$

$$(3.14) \quad \|u''(t, \cdot)\|_{L^q(0,1)} + \|u'(t, \cdot)\|_{L^q(0,1)} \\ \leq c \left(t^{\frac{1}{2q}-1} \|u_0\|_{W_q^2(0,1)} + t^{\frac{1}{2q}-1-\mu} \|f\|_{C_\mu^\gamma((0,t], L^q(0,1))} \right), \quad t \in (0, T].$$

Proof. We consider in the space $L^q(0, 1)$ $1 \leq q < \infty$, the operator A defined by

$$A(u) = au''(x), D(A) = \{u \in W_q^2(0, 1), L_i(u) = 0, i = \overline{1, 2}\}.$$


Then problem (3.11) can be written as

$$\begin{cases} u'(t) = Au(t) + f(t), \\ u(0) = u_0, \end{cases}$$

where $u(t) = u(t, \cdot)$, $f(t) = f(t, \cdot)$, and $u_0 = u_0(\cdot)$ are functions with values in the Banach space $L^q(0, 1)$. From Theorem 3.1 we conclude that $\|R(\lambda, A)\| \leq c |\lambda|^{-\frac{1}{2q}}$, for $|\arg \lambda| \leq \frac{\pi}{2} + \alpha$, as $|\lambda| \rightarrow \infty$.

Then, from Theorem 3.2 the problem (3.11) has a unique solution

$$u \in C((0, T], L^q(0, 1)) \cap C^1((0, T], W_q^2(0, 1), L^q(0, 1))$$



**Boundary Value Problem for
Second-Order Differential
Operators with Mixed Nonlocal
Boundary Conditions**

M. Denche and A. Kourta

Title Page

Contents



Go Back

Close

Quit

Page 30 of 36

and we have the following estimates

$$(3.15) \quad \|u(t, \cdot)\|_{L^q(0,1)} \leq c \left(\|Au_0\|_{L^q(0,1)} + \|u_0\|_{L^q(0,1)} + \|f\|_{C_\mu((0,t],L^q(0,1))} \right),$$

$$(3.16) \quad \left\| u'(t) \right\|_{L^q(0,1)} + \|Au(t, \cdot)\|_{L^q(0,1)} \\ \leq c \left(t^{\frac{1}{2q}-1} \left(\|Au_0\|_{L^q(0,1)} + \|u_0\|_{L^q(0,1)} \right) + t^{\frac{1}{2q}-1-\mu} \|f\|_{C_\mu^\gamma((0,t],L^q(0,1))} \right),$$


where $t \in [0, T]$, from (3.15) we get

$$\|u(t, \cdot)\|_{L^q(0,1)} \leq c \left(\|u_0''\|_{L^q(0,1)} + \|u_0\|_{L^q(0,1)} + \|f\|_{C_\mu((0,t],L^q(0,1))} \right) \\ \leq c \left(\|u_0\|_{W_q^2(0,1)} + \|f\|_{C_\mu((0,t],L^q(0,1))} \right), \quad t \in [0, T].$$

and from (3.16) we get

$$\left\| u'(t) \right\|_{L^q(0,1)} + \|u''(t, \cdot)\|_{L^q(0,1)} \\ \leq c \left(t^{\frac{1}{2q}-1} \left(\|Au_0\|_{L^q(0,1)} + \|u_0\|_{L^q(0,1)} \right) + t^{\frac{1}{2q}-1-\mu} \|f\|_{C_\mu^\gamma((0,t],L^q(0,1))} \right) \\ \leq c \left(t^{\frac{1}{2q}-1} \|u_0\|_{W_q^2(0,1)} + t^{\frac{1}{2q}-1-\mu} \|f\|_{C_\mu^\gamma((0,t],L^q(0,1))} \right), \quad t \in [0, T].$$

which gives the desired result. □



**Boundary Value Problem for
Second-Order Differential
Operators with Mixed Nonlocal
Boundary Conditions**

M. Denche and A. Kourta

Title Page

Contents



Go Back


Close

Quit

Page 31 of 36

References

- [1] S.A. BEILIN, Existence of solutions for one-dimensional wave equations with nonlocal conditions, *Electronic Journal of Differential Equations*, **2001** (2001), Art. 76, 1–8.
- [2] H.E. BENZINGER, Green's function for ordinary differential operators, *J. Differen. Equations*, **7** (1970), 478–496.
- [3] J.R. CANNON, Y. LIN AND J. van der HOEK, A quasi-linear parabolic equation with nonlocal boundary conditions, *Rend. Mat. Appl.*, **9**(7) (1989), 239–264.
- [4] M. DENCHE, Defect-Coerciveness for non regular transmission problem, *Advances in Mathematical Sciences and Applications*, **9**(1) (1999), 229–241.
- [5] M. DENCHE AND A.L. MARHOUNE, Mixed problem with nonlocal boundary conditions for a third-order partial differential equation of mixed type. *IJMMS*, **26**(7) (2001), 417–426.
- [6] M. DENCHE AND A.L. MARHOUNE, Mixed problem with integral boundary condition for a high order mixed type partial differential equation, *J. Appl. Math. Stochastic Anal.*, **16**(1) (2003), 69–79.
- [7] N. DUNFORD AND J.T. SCHWARTZ, *Linear Operators. Part III. Spectral operators*, Interscience, New York, 1971.
- [8] W. EBERHARD AND G. FREILING, Ston reguläre Eigenwert problem, *Math.Zeit.*, **160**(2) (1978), 139–161.



Boundary Value Problem for
Second-Order Differential
Operators with Mixed Nonlocal
Boundary Conditions

M. Denche and A. Kourta

Title Page

Contents




Go Back

Close

Quit

Page 32 of 36

- [9] J. FARJAS, J.I. ROSELL, R. HERRERO, R. PONS, F. PI AND G. ORRIOLS, Equivalent low-order model for a nonlinear diffusion equation, *Phys. D.*, **95** (1996), 107–127.
- [10] J.M. GALLARDO, Generation of analytic semi -groups by second-order differential operators with nonseparated boundary conditions, *Rocky Mountain J. Math.*, **30**(2) (2000), 869–899.
- [11] J.M. GALLARDO, Second-order differential operators with integral boundary conditions and generation of analytic semigroups, *Rocky Mountain J. Math.*, **30**(4) (2000), 1265–1291.
- [12] M.G. GASUMOV AND A.M. MAGERRAMOV, On fold-completeness of a system of eigenfunctions and associated functions of a class of differential operators, *DAN Azerb. SSR*, **30**(3) (1974), 9–12.
- [13] M.G. GASUMOV AND A.M. MAGERRAMOV, Investigation of a class of differential operator pencils of even order, *DAN SSR*, **265**(2) (1982), 277–280.
- [14] N.I. IONKIN, Solution of boundary value problem in heat conduction theory with nonlocal boundary conditions, *Differ. Uravn.*, **13** (1977), 294–304.
- [15] N.I. IONKIN, Stability of a problem in heat conduction theory with non-local boundary conditions, *Differ. Uravn.*, **15**(7) (1979), 1279–1283.
- [16] N.I. IONKIN AND E.I. MOISEEV, A problem for the heat conduction equation with two-point boundary condition, *Differ. Uravn.*, **15**(7) (1979), 1284–1295.



**Boundary Value Problem for
Second-Order Differential
Operators with Mixed Nonlocal
Boundary Conditions**

M. Denche and A. Kourta

Title Page

Contents




Go Back

Close

Quit

Page 33 of 36

- [17] N.I. KAMYNNIN, A boundary value problem in the theory of heat conduction with non classical boundary condition, *Theoret. Vychisl. Mat. Fiz.*, **4**(6) (1964), 1006–1024.
- [18] A.P. KHROMOV, Eigenfunction expansions of ordinary linear differential operators in a finite interval, *Soviet Math. Dokl.*, **3** (1962), 1510–1514.
- [19] Yu. A. MAMEDOV, On functions expansion to the series of solutions residue of irregular boundary value problems, in “*Investigations on Differential Equations*”, AGU named after S.M. Kirov (1984), 94–98.
- [20] M.A. NAIMARK, *Linear Differential Operators*, Vols. 1-2, Ungar Publishing, (1967).
- [21] M.L. RASULOV, *Methods of Contour Integration*, North-Holland, Amsterdam, 1967.
- [22] M.L. RASULOV, *Applications of the contour Integral Method*, Nauka, Moscow, 1975.
- [23] J.I. ROSELL, J. FARJAS, R. HERRERO, F. PI AND G. ORRIOLS, Homoclinic phenomena in opto-thermal bistability with localized absorption, *Phys. D*, **85** (1995), 509–547.
- [24] A.A. SHKALIKOV, On completeness of eigenfunctions and associated function of an ordinary differential operator with separated irregular boundary conditions, *Func. Anal. i evo Prilozh*, **10**(4) (1976), 69–80.



**Boundary Value Problem for
Second-Order Differential
Operators with Mixed Nonlocal
Boundary Conditions**

M. Denche and A. Kourta

Title Page

Contents




Go Back

Close

Quit

Page 34 of 36

- [25] J.T. SILCHENKO, Differential equations with non-densely defined operator coefficients, generating semigroups with singularities, *Nonlinear Anal., Ser. A: Theory Methods*, **36**(3) (1999), 345–352.
- [26] Yu. T. SILCHENKO, An ordinary differential operator with irregular boundary conditions, (Russian) *Sibirsk. Mat. Zh.*, **40**(1) (1999), 183–190, iv; translation in *Siberian Math. J.*, **40**(1) (1999), 158–164.
- [27] A.A. SAMARSKII, Some problems in differential equations theory, *Differ. Uravn.*, **16**(11) (1980), 1925–1935.
- [28] J.D. TAMARKIN, *About Certain General Problems of Theory of Ordinary Linear Differential Equations and about Expansions of Derivative Functions into Series*, Petrograd, 1917.
- [29] C. TRETTER, *On λ -Nonlinear Boundary Eigenvalue Problems*, Academic Verlag, Berlin, 1993.
- [30] A.I. VAGABOV, On eigenvectors of irregular differential pencils with non-separated boundary conditions, *DAN SSSR*, **261**(2) (1981), 268–271.
- [31] S. YAKUBOV, *Completeness of Root Functions of Regular Differential Operators*, Longman Scientific Technical, New York, (1994).
- [32] S. YAKUBOV, On a new method for solving irregular problems, *J. Math. Anal. Appl.*, **220**(1) (1998), 224–249.
- [33] S. YAKUBOV AND Y. YAKUBOV, Differential-operator equations, *Ordinary and Partial Differential Equations*, Chapman & Hall/CRC Monographs



**Boundary Value Problem for
Second-Order Differential
Operators with Mixed Nonlocal
Boundary Conditions**

M. Denche and A. Kourta

Title Page

Contents



Go Back


Close

Quit

Page 35 of 36

and Surveys in Pure and Applied Mathematics, 103. Chapman & Hall/CRC, Boca Raton, FL, (2000).

- [34] Y. YAKUBOV, Irregular boundary value problems for ordinary differential equations, *Analysis (Munich)* **18**(4) (1998), 359–402.
- [35] N.I. YURCHUK, Mixed problem with an integral condition for certain parabolic equations, *Differ. Uravn.*, **22**(12) (1986), 2117–2126.



**Boundary Value Problem for
Second-Order Differential
Operators with Mixed Nonlocal
Boundary Conditions**

M. Denche and A. Kourta

Title Page

Contents



Go Back

Close

Quit

Page 36 of 36