



ON CERTAIN INEQUALITIES FOR MEANS IN TWO VARIABLES

JÓZSEF SÁNDOR

BABEȘ-BOLYAI UNIVERSITY
DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCES
STR. KOGĂLNICEANU NR. 1
400084 CLUJ-NAPOCA, ROMANIA
jsandor@math.ubbcluj.ro

Received 26 May, 2008; accepted 10 April, 2009

Communicated by S.S. Dragomir

ABSTRACT. We show that certain inequalities for the means A , G , L , I proved by H.-J. Seiffert [12] as well as by H. Alzer and S.-L. Qiu [3] are consequences of some results of the author [5], [7], [9].

Key words and phrases: Means and their inequalities.

2000 Mathematics Subject Classification. 26D07, 27E60.

1. INTRODUCTION

The logarithmic and identric means of two positive real numbers a and b with $a \neq b$ are defined by

$$L = L(a, b) = \frac{b - a}{\log b - \log a} \quad \text{and} \quad I = I(a, b) = \frac{1}{e} \left(\frac{a^a}{b^b} \right)^{\frac{1}{a-b}},$$

respectively. These means have been the subject of much intensive research, partly because they are related to many other important means and partly because these means have applications in physics, economics, meteorology, statistics, etc. For a survey of results, with an extended literature, see [3], [6]. For identities involving these, and other means, see e.g. [8], [10]. In particular, the identity

$$\frac{I(a^2, b^2)}{I(a, b)} = (a^a \cdot b^b)^{\frac{1}{a+b}} = S = S(a, b)$$

leads to the weighted geometric mean of a and b , denoted by $S(a, b)$ in [6], [8], [9].

In paper [12], the following two inequalities are proved

$$(1.1) \quad G(a, b) \exp \left(\frac{1}{3} \left(\frac{b-a}{b+a} \right)^2 \right) < I(a, b) < A(a, b) \exp \left(-\frac{1}{6} \left(\frac{b-a}{b+a} \right)^2 \right),$$

where $a \neq b$, $a, b > 0$.

In [3], the following relation appears among many other ones:

$$(1.2) \quad G(a, b) \exp\left(\frac{1}{6} \left(\frac{b-a}{b+a}\right)^2\right) < L(a, b) < A(a, b) \exp\left(-\frac{1}{3} \left(\frac{b-a}{b+a}\right)^2\right),$$

$a \neq b, a, b > 0$.

We note that, the right hand side inequality of (1.1) was first proved by the author in 1989 [5]. In that paper the following inequality was also shown:

$$(1.3) \quad \frac{A^2(a, b)}{I(a^2, b^2)} < \exp\left(-\frac{1}{3} \left(\frac{b-a}{b+a}\right)^2\right).$$

The aim of this note is to prove that the above inequalities are connected to each other by a chain of relations, and that, in fact, all are consequences of (1.3).

2. MAIN RESULTS

To begin with, we write all the inequalities in another form. The left and right sides of (1.1) may be written respectively as

$$(2.1) \quad \exp\left(\frac{1}{3} \left(\frac{b-a}{b+a}\right)^2\right) < \frac{I(a, b)}{G(a, b)},$$

$$(2.2) \quad \exp\left(\frac{1}{3} \left(\frac{b-a}{b+a}\right)^2\right) < \frac{A^2(a, b)}{I^2(a, b)}$$

and the inequalities of (1.2) as

$$(2.3) \quad \exp\left(\frac{1}{3} \left(\frac{b-a}{b+a}\right)^2\right) < \frac{L^2(a, b)}{G^2(a, b)},$$

$$(2.4) \quad \exp\left(\frac{1}{3} \left(\frac{b-a}{b+a}\right)^2\right) < \frac{A(a, b)}{L(a, b)}.$$

Finally note that, (1.3) may be written as

$$(2.5) \quad \exp\left(\frac{1}{3} \left(\frac{b-a}{b+a}\right)^2\right) < \frac{I(a^2, b^2)}{A^2(a, b)} = \frac{I(a, b)S(a, b)}{A^2(a, b)}.$$

Theorem 2.1. *The following chain of implications holds true:*

$$(2.5) \Rightarrow (2.2) \Rightarrow (2.4) \Rightarrow (2.1) \Rightarrow (2.3).$$

Proof. (2.5) \Rightarrow (2.2) means that $\frac{I \cdot S}{A^2} < \frac{A^2}{I^2}$, or $S < \frac{A^4}{I^3}$. This inequality is proved in [9, Theorem 1].

(2.2) \Rightarrow (2.4) by $\frac{A^2}{I^2} < \frac{A}{L}$, i.e. $I^2 > A \cdot L$. For this inequality, see [7, Relation (9)].

(2.4) \Rightarrow (2.1) by $\frac{A}{L} < \frac{I}{G}$, i.e. $A \cdot G < L \cdot I$, see [1].

(2.1) \Rightarrow (2.3) by $\frac{I}{G} < \frac{L^2}{G^2}$, i.e. $\sqrt{GI} < L$, see [2].

Therefore all implications are valid. \square

We note that inequality (2.5) was a consequence of an integral inequality due to the author [4], (discovered in 1982), to the effect that:

Theorem 2.2. Let $f : [a, b] \rightarrow \mathbb{R}$ be a $2k$ -times ($k \geq 1$) differentiable function such that $f^{(2k)}(x) > 0$. Then

$$(2.6) \quad \int_a^b f(x)dx > \sum_{j=0}^{k-1} \frac{(b-a)^{2j+1}}{2^{2j}(2j+1)!} f^{(2j)}\left(\frac{a+b}{2}\right).$$

For $k = 2$ we obtain that if f is 4-times differentiable, then

$$(2.7) \quad \frac{1}{b-a} \int_a^b f(x)dx > f\left(\frac{a+b}{2}\right) + \frac{(b-a)^2}{24} f''\left(\frac{a+b}{2}\right).$$

Clearly, (2.6) and (2.7) are extensions of the classical Hadamard inequality, which says that, if f is convex on $[a, b]$ then

$$(2.8) \quad \frac{1}{b-a} \int_a^b f(x)dx > f\left(\frac{a+b}{2}\right).$$

Applying (2.7) for $f(x) = x \log x$, and using the identity

$$(2.9) \quad \int_a^b x \log x dx = \frac{1}{4}(b^2 - a^2) \log I(a^2, b^2)$$

(see [6]), we get (2.5). Applying (2.7) to $f(x) = -\log x$, we get (2.2), i.e. the right side of (1.1) (see [5]). For another proof, see [11].

REFERENCES

- [1] H. ALZER, Ungleichungen für Mittelwerte, *Arch. Math.*, **47** (1986), 422–426.
- [2] H. ALZER, Two inequalities for means, *C.R. Math. Rep. Acad. Sci. Canada*, **9** (1987), 11–16.
- [3] H. ALZER AND S.-L. QIU, Inequalities for means in two variables, *Arch. Math.*, **80** (2003), 201–215.
- [4] J. SÁNDOR, Some integral inequalities, *El. Math.*, **43** (1988), 177–180.
- [5] J. SÁNDOR, Inequalities for means, in: *Proc. 3rd Symposium of Math. and its Appl.*, 3–4 Nov. 1989, 87–90, Timișoara (Romania).
- [6] J. SÁNDOR, On the identric and logarithmic means, *Aequat. Math.*, **40** (1990), 261–270.
- [7] J. SÁNDOR, A note on some inequalities for means, *Arch. Math.*, **56** (1991), 471–473.
- [8] J. SÁNDOR, On certain identities for means, *Studia Univ. Babeș-Bolyai Math.*, **38** (1993), 7–14.
- [9] J. SÁNDOR AND I. RAȘA, Inequalities for certain means in two arguments, *Nieuw Arch. Wisk.*, **15** (1997), 51–55.
- [10] J. SÁNDOR AND W. WANG, On certain identities for means (Chinese), *J. Chengdu Univ.*, **20** (2001), 6–8.
- [11] J. SÁNDOR AND T. TRIF, Some new inequalities for means of two arguments, *Internat. J. Math. Math. Sci.*, **25** (2001), 525–532.
- [12] H.-J. SEIFFERT, Ungleichungen für Elementare Mittelwerte, *Arch. Math.*, **64** (1995), 129–131.