

Journal of Inequalities in Pure and Applied Mathematics


AN INTEGRAL INEQUALITY SIMILAR TO QI'S INEQUALITY

LAZHAR BOUGOFFA

Department of Mathematics
Faculty of Science
King Khalid University
P.O. Box 9004, Abha, Saudi Arabia

EMail: abogafah@kku.edu.sa

©2000 Victoria University
ISSN (electronic): 1443-5756
161-04



volume 6, issue 1, article 27,
2005.

*Received 15 August, 2004;
accepted 18 February, 2005.*

Communicated by: F. Qi

[Abstract](#)

[Contents](#)



[Home Page](#)

[Go Back](#)

[Close](#)

[Quit](#)

Abstract

In this note, as a complement of an open problem by F. Qi in the paper [*Several integral inequalities*, J. Inequal. Pure Appl. Math. **1** (2002), no. 2, Art. 54. <http://jipam.vu.edu.au/article.php?sid=113>. RGMIA Res. Rep. Coll. **2** (1999), no. 7, Art. 9, 1039–1042. <http://rgmia.vu.edu.au/v2n7.html>], a similar problem is posed and an affirmative answer to it is established.

2000 Mathematics Subject Classification: Primary: 26D15.

Key words: Hölder's inequality, Qi's inequality, Integral inequality.

The following problem was posed by F. Qi in his paper [6]:


Problem 1. *Under what conditions does the inequality*

$$(1) \quad \int_a^b [f(x)]^t dx \geq \left(\int_a^b f(x) dx \right)^{t-1}$$

hold for $t > 1$?

This problem has attracted much attention from some mathematicians [5]. Its meanings of probability and statistics is found in [2]. See also [1, 3, 4] and the references therein.

Similar to Problem 1, we propose the following



An Integral Inequality Similar to Qi's Inequality

Lazhar Bougoffa

Title Page

Contents




Go Back

Close

Quit

Page 2 of 6



An Integral Inequality Similar to Qi's Inequality

Lazhar Bougoffa

Title Page

Contents



Go Back

Close

Quit

Page 3 of 6

Problem 2. Under what conditions does the inequality

$$(2) \quad \int_a^b [f(x)]^t dx \leq \left(\int_a^b f(x) dx \right)^{1-t}$$

hold for $t < 1$?

Before giving an affirmative answer to Problem 2, we establish the following

Proposition 1. Let f and g be nonnegative functions with $0 < m \leq f(x)/g(x) \leq M < \infty$ on $[a, b]$. Then for $p > 1$ and $q > 1$ with $\frac{1}{p} + \frac{1}{q} = 1$ we have

$$(3) \quad \int_a^b [f(x)]^{\frac{1}{p}} [g(x)]^{\frac{1}{q}} dx \leq M^{\frac{1}{p^2}} m^{-\frac{1}{q^2}} \int_a^b [f(x)]^{\frac{1}{q}} [g(x)]^{\frac{1}{p}} dx,$$

and then

$$(4) \quad \int_a^b [f(x)]^{\frac{1}{p}} [g(x)]^{\frac{1}{q}} dx \leq M^{\frac{1}{p^2}} m^{-\frac{1}{q^2}} \left(\int_a^b f(x) dx \right)^{\frac{1}{q}} \left(\int_a^b g(x) dx \right)^{\frac{1}{p}}.$$

Proof. From Hölder's inequality, we obtain

$$(5) \quad \int_a^b [f(x)]^{\frac{1}{p}} [g(x)]^{\frac{1}{q}} dx \leq \left(\int_a^b f(x) dx \right)^{\frac{1}{p}} \left(\int_a^b g(x) dx \right)^{\frac{1}{q}},$$

that is,

$$(6) \quad \int_a^b [f(x)]^{\frac{1}{p}} [g(x)]^{\frac{1}{q}} dx \leq \left(\int_a^b [f(x)]^{\frac{1}{p}} [f(x)]^{\frac{1}{q}} dx \right)^{\frac{1}{p}} \left(\int_a^b [g(x)]^{\frac{1}{p}} [g(x)]^{\frac{1}{q}} dx \right)^{\frac{1}{q}}.$$

Since $[f(x)]^{\frac{1}{p}} \leq M^{\frac{1}{p}}[g(x)]^{\frac{1}{p}}$ and $[g(x)]^{\frac{1}{q}} \leq m^{-\frac{1}{q}}[f(x)]^{\frac{1}{q}}$, from the above inequality it follows that

$$(7) \quad \int_a^b [f(x)]^{\frac{1}{p}} [g(x)]^{\frac{1}{q}} dx \leq M^{\frac{1}{p^2}} m^{-\frac{1}{q^2}} \left(\int_a^b [f(x)]^{\frac{1}{q}} [g(x)]^{\frac{1}{p}} dx \right)^{\frac{1}{p}} \left(\int_a^b [f(x)]^{\frac{1}{q}} [g(x)]^{\frac{1}{p}} dx \right)^{\frac{1}{q}},$$

that is

$$(8) \quad \int_a^b [f(x)]^{\frac{1}{p}} [g(x)]^{\frac{1}{q}} dx \leq M^{\frac{1}{p^2}} m^{-\frac{1}{q^2}} \int_a^b [f(x)]^{\frac{1}{q}} [g(x)]^{\frac{1}{p}} dx.$$

Hence, the inequality (3) is proved.

The inequality (4) follows from substituting the following


$$(9) \quad \int_a^b [f(x)]^{\frac{1}{q}} [g(x)]^{\frac{1}{p}} dx \leq \left(\int_a^b f(x) dx \right)^{\frac{1}{q}} \left(\int_a^b g(x) dx \right)^{\frac{1}{p}}$$

into (8), which can be obtained by Hölder's inequality. \square

Now we are in a position to give an affirmative answer to Problem 2 as follows.

Proposition 2. For a given positive integer $p \geq 2$, if $0 < m \leq f(x) \leq M$ on $[a, b]$ with $M \leq m^{(p-1)^2} / (b-a)^p$, then

$$(10) \quad \int_a^b [f(x)]^{\frac{1}{p}} dx \leq \left(\int_a^b f(x) dx \right)^{1-\frac{1}{p}}.$$



An Integral Inequality Similar to Qi's Inequality

Lazhar Bougoffa

Title Page

Contents



Go Back

Close

Quit

Page 4 of 6

Proof. Putting $g(x) \equiv 1$ into (4) yields

$$(11) \quad \int_a^b [f(x)]^{\frac{1}{p}} dx \leq K \left(\int_a^b f(x) dx \right)^{1-\frac{1}{p}},$$

where $K = M^{\frac{1}{p^2}}(b-a)^{\frac{1}{p}} / m^{(1-\frac{1}{p})^2}$.

From $M \leq m^{(p-1)^2} / (b-a)^p$, we conclude that $K \leq 1$. Thus the inequality (10) is proved. \square

Remark 1. Now we discuss a simple case of "equality" in Proposition 2. If we make the substitution $f(x) = M = m$ and $b - a = 1$ with $p = 2$, then the equality in (10) holds.


In order to illustrate a possible practical use of Proposition 2, we shall give in the following two simple examples in which we can apply inequality (10).

Example 1. Let $f(x) = 8x^2$ on $[1/2, 1]$ with $M = 8$ and $m = 2$. Taking $p = 2$, we see that the conditions of Proposition 2 are fulfilled and straightforward computation yields

$$\int_{1/2}^1 (8x^2)^{1/2} dx = \frac{3}{4}\sqrt{2} < \left(\int_{1/2}^1 8x^2 dx \right)^{\frac{1}{2}} = \frac{\sqrt{7}}{\sqrt{3}}.$$

Example 2. Let $f(x) = e^x$ on $[1, 2]$ with $M = e^2$ and $m = e$. Taking $p = 3$, all the conditions of Proposition 2 are satisfied and direct calculation produces

$$\int_1^2 (e^x)^{1/3} dx = 3(e^{2/3} - e^{1/3}) \approx 1.65 < \left(\int_1^2 e^x dx \right)^{\frac{2}{3}} = (e^2 - e)^{2/3} \approx 2.78.$$



An Integral Inequality Similar to Qi's Inequality

Lazhar Bougoffa

Title Page

Contents



Go Back


Close

Quit

Page 5 of 6

References

- [1] L. BOUGOFFA, Notes on Qi type inequalities, *J. Inequal. Pure and Appl. Math.*, **44**(4) (2003), Art. 77. Available online at <http://jipam.vu.edu.au/article.php?sid=318>.
- [2] V. CSISZAR AND T.F. MÓRI, The convexity method of proving moment-type inequalities, *Statist. Probab. Lett.*, **66** (2004), 303–313.
- [3] S. MAZOUZI AND F. QI, On an open problem regarding an integral inequality, *J. Inequal. Pure Appl. Math.*, **4**(2) (2003), Art. 31. Available online at <http://jipam.vu.edu.au/article.php?sid=269>.
- [4] J. PEČARIĆ AND T. PEJKOVIĆ, Note on Feng Qi's integral inequality, *J. Inequal. Pure Appl. Math.*, **5**(3) (2004), Art. 51. Available online at <http://jipam.vu.edu.au/article.php?sid=418>.
- [5] T.K. POGÁNY, On an open problem of F. Qi, *J. Inequal. Pure Appl. Math.*, **3**(4) (2002), Art. 54. Available online at <http://jipam.vu.edu.au/article.php?sid=206>.
- [6] F. QI, Several integral inequalities, *J. Inequal. Pure Appl. Math.*, **1**(2) (2002), Art. 54. Available online at <http://jipam.vu.edu.au/article.php?sid=113>. *RGMIA Res. Rep. Coll.*, **2**(7) (1999), Art. 9, 1039–1042. Available online at <http://rgmia.vu.edu.au/v2n7.html>.



An Integral Inequality Similar to Qi's Inequality

Lazhar Bougoffa

Title Page

Contents



Go Back

Close

Quit

Page 6 of 6