

Journal of Inequalities in Pure and Applied Mathematics

DIRECT RESULTS FOR CERTAIN FAMILY OF INTEGRAL TYPE OPERATORS

VIJAY GUPTA AND OGÜN DOĞRU


School of Applied Sciences
Netaji Subhas Institute of Technology
Sector 3 Dwarka, New Delhi-110045, India.

E-Mail: vijay@nsit.ac.in

Ankara University
Faculty of Science
Department of Mathematics
Tandoğan, Ankara-06100, Turkey

E-Mail: dogru@science.ankara.edu.tr

©2000 Victoria University
ISSN (electronic): 1443-5756
165-04



volume 6, issue 1, article 21,
2005.

*Received 08 September, 2004;
accepted 22 January, 2005.*

Communicated by: D. Hinton

Abstract

Contents



Home Page

Go Back

Close

Quit



Abstract

In the present paper we introduce a certain family of linear positive operators and study some direct results which include a pointwise convergence, asymptotic formula and an estimation of error in simultaneous approximation.


2000 Mathematics Subject Classification: 41A25, 41A30.

Key words: Linear positive operators, Simultaneous approximation, Steklov mean, Modulus of continuity.

The authors are thankful to the referee for his/her useful recommendations and valuable remarks.

Contents

1	Introduction	3
2	Auxiliary Results	4
3	Simultaneous Approximation	10
References		



Direct Results for Certain Family of Integral Type Operators

Vijay Gupta and Ogün Dođru

Title Page

Contents



Go Back

Close

Quit

Page 2 of 21

1. Introduction

We consider a certain family of integral type operators, which are defined as

$$(1.1) \quad B_n(f, x) = \frac{1}{n} \sum_{\nu=1}^{\infty} p_{n,\nu}(x) \int_0^{\infty} p_{n,\nu-1}(t) f(t) dt + (1+x)^{-n-1} f(0), \quad x \in [0, \infty),$$

where

$$p_{n,\nu}(t) = \frac{1}{B(n, \nu + 1)} t^\nu (1+t)^{-n-\nu-1}$$

with $B(n, \nu + 1) = \nu!(n-1)/(n+\nu)!$ the Beta function.


Alternatively the operators (1.1) may be written as

$$B_n(f, x) = \int_0^{\infty} W_n(x, t) f(t) dt,$$

where

$$W_n(x, t) = \frac{1}{n} \sum_{\nu=1}^{\infty} p_{n,\nu}(x) p_{n,\nu-1}(t) + (1+x)^{-n-1} \delta(t),$$

$\delta(t)$ being the Dirac delta function. The operators B_n are discretely defined linear positive operators. It is easily verified that these operators reproduce only the constant functions. As far as the degree of approximation is concerned these operators are very similar to the operators considered by Srivastava and Gupta [5], but the approximation properties of these operators are different. In this paper, we study some direct theorems in simultaneous approximation for the operators (1.1).



Direct Results for Certain
Family of Integral Type
Operators

Vijay Gupta and Ogün Doğru

Title Page

Contents



Go Back

Close

Quit

Page 3 of 21

2. Auxiliary Results

In this section we mention some lemmas which are necessary to prove the main theorems.

Lemma 2.1 ([3]). For $m \in \mathbb{N}^0$, if the m -th order moment is defined as

$$U_{n,m}(x) = \frac{1}{n} \sum_{\nu=0}^{\infty} p_{n,\nu}(x) \left(\frac{\nu}{n+1} - x \right)^m,$$

then

$$(n+1)U_{n,m+1}(x) = x(1+x) [U'_{n,m}(x) + mU_{n,m-1}(x)].$$

Consequently

$$U_{m,n}(x) = \mathcal{O}(n^{-(m+1)/2}).$$

Lemma 2.2. Let the function $\mu_{n,m}(x)$, $n > m$ and $m \in \mathbb{N}^0$, be defined as


$$\mu_{n,m}(x) = \frac{1}{n} \sum_{\nu=1}^{\infty} p_{n,\nu}(x) \int_0^{\infty} p_{n,\nu-1}(t)(t-x)^m dt + (-x)^m(1+x)^{-n-1}.$$

Then

$$\mu_{n,0}(x) = 1, \mu_{n,1}(x) = \frac{2x}{(n-1)}, \mu_{n,2}(x) = \frac{2nx(1+x) + 2x(1+4x)}{(n-1)(n-2)}$$

and there holds the recurrence relation

$$(n-m-1)\mu_{n,m+1}(x) = x(1+x) [\mu'_{n,m}(x) + 2m\mu_{n,m-1}(x)] + [m(1+2x) + 2x]\mu_{n,m}(x).$$



Direct Results for Certain
Family of Integral Type
Operators

Vijay Gupta and Ogün Dođru

Title Page

Contents



Go Back

Close

Quit

Page 4 of 21

Consequently for each $x \in [0, \infty)$, we have from this recurrence relation that


$$\mu_{n,m}(x) = \mathcal{O}(n^{-[(m+1)/2]}).$$

Proof. The values of $\mu_{n,0}(x)$, $\mu_{n,1}(x)$ and $\mu_{n,2}(x)$ easily follow from the definition. We prove the recurrence relation as follows

$$\begin{aligned} & x(1+x)\mu'_{n,m}(x) \\ &= \frac{1}{n} \sum_{\nu=1}^{\infty} x(1+x)p'_{n,\nu}(x) \int_0^{\infty} p_{n,\nu-1}(t)(t-x)^m dt \\ &\quad - m \frac{1}{n} \sum_{\nu=1}^{\infty} x(1+x)p_{n,\nu}(x) \int_0^{\infty} p_{n,\nu-1}(t)(t-x)^{m-1} dt \\ &\quad - [(n+1)(-x)^m(1+x)^{-n-2} + m(-x)^{m-1}(1+x)^{-n-1}] x(1+x). \end{aligned}$$

Now using the identities $x(1+x)p'_{n,\nu}(x) = [\nu - (n+1)x]p_{n,\nu}(x)$, we obtain

$$\begin{aligned} & x(1+x) [\mu'_{n,m}(x) + m\mu_{n,m-1}(x)] \\ &= \frac{1}{n} \sum_{\nu=1}^{\infty} [\nu - (n+1)x] p_{n,\nu}(x) \\ &\quad \times \int_0^{\infty} p_{n,\nu-1}(t)(t-x)^m dt + (n+1)(-x)^{m+1}(1+x)^{-n-1} \\ &= \frac{1}{n} \sum_{\nu=1}^{\infty} p_{n,\nu}(x) \int_0^{\infty} [\{(\nu-1) - (n+1)t\} + (n+1)(t-x) + 1] \\ &\quad \times p_{n,\nu-1}(t)(t-x)^m dt + (n+1)(-x)^{m+1}(1+x)^{-n-1} \end{aligned}$$



Direct Results for Certain
Family of Integral Type
Operators

Vijay Gupta and Ogün Dođru

Title Page

Contents



Go Back


Close

Quit

Page 5 of 21

$$\begin{aligned}
&= \frac{1}{n} \sum_{\nu=1}^{\infty} p_{n,\nu}(x) \int_0^{\infty} t(1+t)p'_{n,\nu-1}(t)(t-x)^m dt \\
&\quad + (n+1) \frac{1}{n} \sum_{\nu=1}^{\infty} p_{n,\nu}(x) \int_0^{\infty} p_{n,\nu-1}(t)(t-x)^{m+1} dt \\
&\quad + \frac{1}{n} \sum_{\nu=1}^{\infty} p_{n,\nu}(x) \int_0^{\infty} p_{n,\nu-1}(t)(t-x)^m dt + (n+1)(-x)^{m+1}(1+x)^{-n-1} \\
&= \frac{1}{n} \sum_{\nu=1}^{\infty} p_{n,\nu}(x) \int_0^{\infty} [(1+2x)(t-x) \\
&\quad + (t-x)^2 + x(1+x)] p'_{n,\nu-1}(t)(t-x)^m dt \\
&\quad + (n+1) \frac{1}{n} \sum_{\nu=1}^{\infty} p_{n,\nu}(x) \int_0^{\infty} p_{n,\nu-1}(t)(t-x)^{m+1} dt \\
&\quad + \frac{1}{n} \sum_{\nu=1}^{\infty} p_{n,\nu}(x) \int_0^{\infty} p_{n,\nu-1}(t)(t-x)^m dt + (n+1)(-x)^{m+1}(1+x)^{-n-1} \\
&= -(m+1)(1+2x) [\mu_{n,m}(x) - (-x)^m(1+x)^{-n-1}] \\
&\quad - (m+2) [\mu_{n,m+1}(x) - (-x)^{m+1}(1+x)^{-n-1}] \\
&\quad - mx(1+x) [\mu_{n,m-1}(x) - (-x)^{m-1}(1+x)^{-n-1}] \\
&\quad + (n+1) [\mu_{n,m+1}(x) - (-x)^{m+1}(1+x)^{-n-1}] \\
&\quad + [\mu_{n,m}(x) - (-x)^m(1+x)^{-n-1}] + (n+1)(-x)^{m+1}(1+x)^{-n-1} \\
&= -[m(1+2x) + 2x] \mu_{n,m}(x) + (n-m-1) \mu_{n,m+1}(x) - mx(1+x) \mu_{n,m-1}(x).
\end{aligned}$$

This completes the proof of recurrence relation. \square



**Direct Results for Certain
Family of Integral Type
Operators**

Vijay Gupta and Ogün Doğru

Title Page

Contents



Go Back

Close

Quit

Page 6 of 21

Remark 1. It is easily verified from Lemma 2.2 and by the principle of mathematical induction, that for $n > i$ and each $x \in (0, \infty)$

$$B_n(t^i, x) = \frac{(n+i)!(n-i-1)!}{n!(n-1)!} x^i + i(i-1) \frac{(n+i-1)!(n-i-1)!}{n!(n-1)!} x^{i-1} + i(i-1)(i-2) \mathcal{O}(n^{-2}).$$

Corollary 2.3. Let δ be a positive number. Then for every $n > \gamma > 0$, $x \in (0, \infty)$, there exists a constant $M(s, x)$ independent of n and depending on s and x such that


$$\frac{1}{n} \sum_{\nu=1}^{\infty} p_{n,\nu}(x) \int_{|t-x|>\delta} p_{n,\nu-1}(t) t^\gamma dt \leq M(s, x) n^{-s}, \quad s = 1, 2, 3, \dots$$

Lemma 2.4 ([3]). There exist the polynomials $Q_{i,j,r}(x)$ independent of n and ν such that

$$\{x(1+x)\}^r D^r [p_{n,\nu}(x)] = \sum_{\substack{2i+j \leq r \\ i,j \geq 0}} (n+1)^i [\nu - (n+1)x]^j Q_{i,j,r}(x) p_{n,\nu}(x),$$

where $D \equiv \frac{d}{dx}$.

Lemma 2.5. Let f be r times differentiable on $[0, \infty)$ such that $f^{(r-1)}$ is absolutely continuous with $f^{(r-1)}(t) = \mathcal{O}(t^\gamma)$ for some $\gamma > 0$ as $t \rightarrow \infty$. Then for



Direct Results for Certain
Family of Integral Type
Operators

Vijay Gupta and Ogün Doğru

Title Page

Contents



Go Back

Close

Quit

Page 7 of 21

$r = 1, 2, 3, \dots$ and $n > \gamma + r$ we have

$$B_n^{(r)}(f, x) = \frac{(n+r-1)!(n-r-1)!}{n!(n-1)!} \sum_{\nu=0}^{\infty} p_{n+r,\nu}(x) \int_0^{\infty} p_{n-r,\nu+r-1}(t) f^{(r)}(t) dt.$$


Proof. It follows by simple computation the following relations:

$$(2.1) \quad p'_{n,\nu}(t) = n [p_{n+1,\nu-1}(t) - p_{n+1,\nu}(t)],$$

where $t \in [0, \infty)$.

Furthermore, we prove our lemma by mathematical induction. Using the above identity (2.1), we have

$$\begin{aligned} B'_n(f, x) &= \frac{1}{n} \sum_{\nu=1}^{\infty} p'_{n,\nu}(x) \int_0^{\infty} p_{n,\nu-1}(t) f(t) dt - (n+1)(1+x)^{-n-2} f(0) \\ &= \sum_{\nu=1}^{\infty} [p_{n+1,\nu-1}(x) - p_{n+1,\nu}(x)] \int_0^{\infty} p_{n,\nu-1}(t) f(t) dt \\ &\quad - (n+1)(1+x)^{-n-2} f(0) \\ &= np_{n+1,0}(x) \int_0^{\infty} p_{n,0}(t) f(t) dt - (n+1)(1+x)^{-n-2} f(0) \\ &\quad + \sum_{\nu=1}^{\infty} p_{n+1,\nu}(x) \int_0^{\infty} [p_{n,\nu}(t) - p_{n,\nu-1}(t)] f(t) dt \end{aligned}$$



Direct Results for Certain
Family of Integral Type
Operators

Vijay Gupta and Ogün Doğru

Title Page

Contents



Go Back

Close

Quit

Page 8 of 21

$$\begin{aligned}
&= (n+1)(1+x)^{-n-2} \int_0^\infty n(1+t)^{-n-1} f(t) dt \\
&\quad + \sum_{\nu=1}^\infty p_{n+1,\nu}(x) \int_0^\infty \left(\frac{-1}{n-1}\right) p'_{n-1,\nu}(t) f(t) dt \\
&\quad - (n+1)(1+x)^{-n-2} f(0).
\end{aligned}$$

Applying the integration by parts, we get


$$\begin{aligned}
&B'_n(f, x) \\
&= (n+1)(1+x)^{-n-2} f(0) + (n+1)(1+x)^{-n-2} \int_0^\infty (1+t)^{-n} f'(t) dt \\
&\quad + \frac{1}{n-1} \sum_{\nu=1}^\infty p_{n+1,\nu}(x) \int_0^\infty p_{n-1,\nu}(t) f'(t) dt - (n+1)(1+x)^{-n-2} f(0) \\
&= \frac{1}{n-1} \sum_{\nu=0}^\infty p_{n+1,\nu}(x) \int_0^\infty p_{n-1,\nu}(t) f'(t) dt,
\end{aligned}$$

which was to be proved.

If we suppose that

$$B_n^{(i)}(f, x) = \frac{(n+i-1)!(n-i-1)!}{n!(n-1)!} \sum_{\nu=0}^\infty p_{n+i,\nu}(x) \int_0^\infty p_{n-i,\nu+i-1}(t) f^{(i)}(t) dt$$

then by (2.1), and using a similar method to the one above it is easily verified that the result is true for $r = i + 1$. Therefore by the principle of mathematical induction the result follows. \square



Direct Results for Certain
Family of Integral Type
Operators

Vijay Gupta and Ogün Doğru

Title Page

Contents



Go Back

Close

Quit

Page 9 of 21

3. Simultaneous Approximation

In this section we study the rate of pointwise convergence of an asymptotic formula and an error estimation in terms of a higher order modulus of continuity in simultaneous approximation for the operators defined by (1.1). Throughout the section, we have $C_\gamma[0, \infty) := \{f \in C[0, \infty) : |f(t)| \leq Mt^\gamma \text{ for some } M > 0, \gamma > 0\}$.

Theorem 3.1. *Let $f \in C_\gamma[0, \infty)$, $\gamma > 0$ and $f^{(r)}$ exists at a point $x \in (0, \infty)$, then*

$$B_n^{(r)}(f, x) = f^{(r)}(x) + o(1) \text{ as } n \rightarrow \infty.$$


Proof. By Taylor's expansion of f , we have

$$f(t) = \sum_{i=0}^r \frac{f^{(i)}(x)}{i!} (t-x)^i + \varepsilon(t, x)(t-x)^r,$$

where $\varepsilon(t, x) \rightarrow 0$ as $t \rightarrow x$.

Hence

$$\begin{aligned} B_n^{(r)}(f, x) &= \int_0^\infty W_n^{(r)}(t, x) f(t) dt \\ &= \sum_{i=0}^r \frac{f^{(i)}(x)}{i!} \int_0^\infty W_n^{(r)}(t, x) (t-x)^i dt \\ &\quad + \int_0^\infty W_n^{(r)}(t, x) \varepsilon(t, x) (t-x)^r dt \\ &=: R_1 + R_2. \end{aligned}$$



Direct Results for Certain
Family of Integral Type
Operators

Vijay Gupta and Ogün Dođru

Title Page

Contents



Go Back

Close

Quit

Page 10 of 21


First to estimate R_1 , using the binomial expansion of $(t - x)^m$, Lemma 2.2 and Remark 1, we have

$$\begin{aligned}
 R_1 &= \sum_{i=0}^r \frac{f^{(i)}(x)}{i!} \sum_{\nu=0}^i \binom{i}{\nu} (-x)^{i-\nu} \frac{\partial^r}{\partial x^r} \int_0^\infty W_n(t, x) t^\nu dt \\
 &= \frac{f^{(r)}(x)}{r!} \frac{\partial^r}{\partial x^r} \int_0^\infty W_n(t, x) t^r dt \\
 &= \frac{f^{(r)}(x)}{r!} \left\{ \frac{(n+r)!(n-r-1)!}{n!(n-1)!} r! + \text{terms containing lower powers of } x \right\} \\
 &= f^{(r)}(x) + o(1), \quad n \rightarrow \infty.
 \end{aligned}$$

Using Lemma 2.4, we obtain

$$\begin{aligned}
 R_2 &= \int_0^\infty W_n^{(r)}(t, x) \varepsilon(t, x) (t - x)^r dt \\
 &= \sum_{\substack{2i+j \leq r \\ i, j \geq 0}} n^i \frac{Q_{i,j,r}(x)}{\{x(1+x)\}^r} \sum_{\nu=1}^\infty [\nu - (n+1)x]^j \frac{p_{n,\nu}(x)}{n} \\
 &\quad \times \int_0^\infty p_{n,\nu-1}(t) \varepsilon(t, x) (t - x)^r dt \\
 &\quad + (-1)^r \frac{(n+r)!}{(n+1)!} (1+x)^{-n-r-1} \varepsilon(0, x) (-x)^r \\
 &=: R_3 + R_4.
 \end{aligned}$$

Since $\varepsilon(t, x) \rightarrow 0$ as $t \rightarrow x$ for a given $\varepsilon > 0$ there exists a $\delta > 0$ such that



Direct Results for Certain
Family of Integral Type
Operators

Vijay Gupta and Ogün Dođru

Title Page

Contents



Go Back

Close

Quit

Page 11 of 21

$|\varepsilon(t, x)| < \varepsilon$ whenever $0 < |t - x| < \delta$. Thus for some $M_1 > 0$, we can write

$$|R_3| \leq M_1 \sum_{\substack{2i+j \leq r \\ i, j \geq 0}} n^{i-1} \sum_{\nu=1}^{\infty} p_{n,\nu}(x) |\nu - (n+1)x|^j \left\{ \varepsilon \int_{|t-x| < \delta} p_{n,\nu-1}(t) |t-x|^r dt + \int_{|t-x| \geq \delta} p_{n,\nu-1}(t) M_2 t^r dt \right\} \\ =: R_5 + R_6,$$


where

$$M_1 = \sup_{\substack{2i+j \leq r \\ i, j \geq 0}} \frac{|Q_{i,j,r}(x)|}{\{x(1+x)\}^r}$$

and M_2 is independent of t .

Applying Schwarz's inequality for integration and summation respectively, we obtain

$$R_5 \leq \varepsilon M_1 \sum_{\substack{2i+j \leq r \\ i, j \geq 0}} n^{i-1} \sum_{\nu=1}^{\infty} p_{n,\nu}(x) |\nu - (n+1)x|^j \left(\int_0^{\infty} p_{n,\nu-1}(t) dt \right)^{\frac{1}{2}} \\ \times \left(\int_0^{\infty} p_{n,\nu-1}(t) (t-x)^{2r} dt \right)^{\frac{1}{2}} \\ \leq \varepsilon M_1 \sum_{\substack{2i+j \leq r \\ i, j \geq 0}} n^i \sum_{\nu=1}^{\infty} p_{n,\nu}(x) \left(\frac{1}{n} \sum_{\nu=1}^{\infty} p_{n,\nu}(x) [\nu - (n+1)x]^{2j} \right)^{\frac{1}{2}}$$



Direct Results for Certain
Family of Integral Type
Operators

Vijay Gupta and Ogün Dođru

Title Page

Contents



Go Back

Close

Quit

Page 12 of 21


$$\times \left(\frac{1}{n} \sum_{\nu=1}^{\infty} p_{n,\nu}(x) \int_0^{\infty} p_{n,\nu-1}(t)(t-x)^{2r} dt \right)^{\frac{1}{2}}.$$

Using Lemma 2.1 and Lemma 2.2, we get

$$R_5 \leq \varepsilon M_1 \mathcal{O}(n^{j/2}) \mathcal{O}(n^{-r/2}) = \varepsilon \mathcal{O}(1).$$

Again using the Schwarz inequality, Lemma 2.1 and Corollary 2.3, we obtain

$$\begin{aligned} R_6 &\leq M_2 \sum_{\substack{2i+j \leq r \\ i,j \geq 0}} n^{i-1} \sum_{\nu=1}^{\infty} p_{n,\nu}(x) |\nu - (n+1)x|^j \int_{|t-x| \geq \delta} p_{n,\nu-1}(t) t^\gamma dt \\ &\leq M_2 \sum_{\substack{2i+j \leq r \\ i,j \geq 0}} n^{i-1} \sum_{\nu=1}^{\infty} p_{n,\nu}(x) |\nu - (n+1)x|^j \left(\int_{|t-x| \geq \delta} p_{n,\nu-1}(t) dt \right)^{\frac{1}{2}} \\ &\quad \times \left(\int_{|t-x| \geq \delta} p_{n,\nu-1}(t) t^{2\gamma} dt \right)^{\frac{1}{2}} \\ &\leq M_2 \sum_{\substack{2i+j \leq r \\ i,j \geq 0}} n^i \left(\frac{1}{n} \sum_{\nu=1}^{\infty} p_{n,\nu}(x) [\nu - (n+1)x]^{2j} \right)^{\frac{1}{2}} \\ &\quad \times \left(\frac{1}{n} \sum_{\nu=1}^{\infty} p_{n,\nu}(x) \int_0^{\infty} p_{n,\nu-1}(t) t^{2\gamma} dt \right)^{\frac{1}{2}} \\ &= \sum_{\substack{2i+j \leq r \\ i,j \geq 0}} n^i \mathcal{O}(n^{j/2}) \mathcal{O}(n^{-s/2}) \end{aligned}$$



Direct Results for Certain
Family of Integral Type
Operators

Vijay Gupta and Ogün Dođru

Title Page

Contents



Go Back

Close

Quit

Page 13 of 21

for any $s > 0$.

Choosing $s > r$ we get $R_6 = o(1)$. Thus, due to arbitrariness of $\varepsilon > 0$, it follows that $R_3 = o(1)$. Also $R_4 \rightarrow 0$ as $n \rightarrow \infty$ and hence $R_2 = o(1)$. Collecting the estimates of R_1 and R_2 , we get the required result. \square

The following result holds.

Theorem 3.2. *Let $f \in C_\gamma[0, \infty)$, $\gamma > 0$. If $f^{(r+2)}$ exists at a point $x \in (0, \infty)$, then*

$$\begin{aligned} \lim_{n \rightarrow \infty} n[B_n^{(r)}(f, x) - f^{(r)}(x)] \\ = r(r+1)f^{(r)}(x) + [2x(1+r) + r]f^{(r+1)}(x) + x(1+x)f^{(r+2)}(x). \end{aligned}$$

Proof. Using Taylor's expansion of f , we have

$$f(t) = \sum_{i=0}^{r+2} \frac{f^{(i)}(x)}{i!} (t-x)^i + \varepsilon(t, x)(t-x)^{r+2},$$

where $\varepsilon(t, x) \rightarrow 0$ as $t \rightarrow x$ and $\varepsilon(t, x) = \mathcal{O}((t-x)^\beta)$, $t \rightarrow \infty$ for some $\beta > 0$. Applying Lemma 2.2, we have

$$\begin{aligned} n[B_n^{(r)}(f, x) - f^{(r)}(x)] &= n \left[\sum_{i=0}^{r+2} \frac{f^{(i)}(x)}{i!} \int_0^\infty W_n^{(r)}(x, t)(t-x)^i dt - f^{(r)}(x) \right] \\ &\quad + \left[n \int_0^\infty W_n^{(r)}(x, t)\varepsilon(t, x)(t-x)^{r+2} dt \right] \\ &=: E_1 + E_2. \end{aligned}$$



Direct Results for Certain
Family of Integral Type
Operators

Vijay Gupta and Ogün Doğru

Title Page

Contents



Go Back

Close


Quit

Page 14 of 21

$$\begin{aligned}
E_1 &= n \sum_{i=0}^{r+2} \frac{f^{(i)}(x)}{i!} \sum_{j=0}^i \binom{i}{j} (-x)^{i-j} \int_0^\infty W_n^{(r)}(x, t) t^j dt - n f^{(r)}(x) \\
&= \frac{f^{(r)}(x)}{r!} n [B_n^{(r)}(t^r, x) - r!] + \frac{f^{(r+1)}(x)}{(r+1)!} n [(r+1)(-x)B_n^{(r)}(t^r, x) \\
&\quad + B_n^{(r)}(t^{r+1}, x)] + \frac{f^{(r+2)}(x)}{(r+2)!} n \left[\frac{(r+2)(r+1)}{2} x^2 B_n^{(r)}(t^r, x) \right. \\
&\quad \left. + (r+2)(-x)B_n^{(r)}(t^{r+1}, x) + B_n^{(r)}(t^{r+2}, x) \right].
\end{aligned}$$

Therefore by applying Remark 1, we get

$$\begin{aligned}
E_1 &= n f^{(r)}(x) \left[\frac{(n+r)!(n-r-1)!}{n!(n-1)!} - 1 \right] \\
&+ n \frac{f^{(r+1)}(x)}{(r+1)!} \left[(r+1)(-x)r! \left\{ \frac{(n+r)!(n-r-1)!}{n!(n-1)!} \right\} \right. \\
&+ \left. \left\{ \frac{(n+r+1)!(n-r-2)!}{n!(n-1)!} (r+1)!x + r(r+1) \frac{(n+r)!(n-r-2)!}{n!(n-1)!} r! \right\} \right] \\
&+ n \frac{f^{(r+2)}(x)}{(r+2)!} \left[\frac{(r+2)(r+1)x^2}{2} \cdot r! \frac{(n+r)!(n-r-1)!}{n!(n-1)!} \right. \\
&+ (r+2)(-x) \left\{ \frac{(n+r+1)!(n-r-2)!}{n!(n-1)!} (r+1)!x \right. \\
&\quad \left. \left. + r(r+1) \frac{(n+r)!(n-r-2)!}{n!(n-1)!} r! \right\} \right]
\end{aligned}$$



**Direct Results for Certain
Family of Integral Type
Operators**

Vijay Gupta and Ogün Doğru

Title Page

Contents



Go Back

Close

Quit

Page 15 of 21

$$+ \left\{ \frac{(n+r+2)!(n-r-3)!}{n!(n-1)!} \frac{(r+2)!}{2} x^2 + (r+1)(r+2) \frac{(n+r+1)!(n-r-3)!}{n!(n-1)!} (r+1)! x \right\} + \mathcal{O}(n^{-2}).$$

In order to complete the proof of the theorem it is sufficient to show that $E_2 \rightarrow 0$ as $n \rightarrow \infty$, which can be easily proved along the lines of the proof of Theorem 3.1 and by using Lemma 2.1, Lemma 2.2 and Lemma 2.4. \square


Let us assume that $0 < a < a_1 < b_1 < b < \infty$, for sufficiently small $\delta > 0$, the m -th order Steklov mean $f_{m,\delta}(t)$ corresponding to $f \in C_\gamma[0, \infty)$ is defined by

$$f_{m,\delta}(t) = \delta^{-m} \int_{-\frac{\delta}{2}}^{\frac{\delta}{2}} \int_{-\frac{\delta}{2}}^{\frac{\delta}{2}} \dots \int_{-\frac{\delta}{2}}^{\frac{\delta}{2}} [f(t) - \Delta_\eta^m f(t)] \prod_{i=1}^m dt_i$$

where $\eta = \frac{1}{m} \sum_{i=1}^m t_i$, $t \in [a, b]$ and $\Delta_\eta^m f(t)$ is the m -th forward difference with step length η .

It is easily checked (see e. g. [1], [4]) that

- (i) $f_{m,\delta}$ has continuous derivatives up to order m on $[a, b]$;
- (ii) $\left\| f_{m,\delta}^{(r)} \right\|_{C[a_1, b_1]} \leq M_1 \delta^{-r} \omega_r(f, \delta, a_1, b_1)$, $r = 1, 2, 3, \dots, m$;
- (iii) $\|f - f_{m,\delta}\|_{C[a_1, b_1]} \leq M_2 \omega_m(f, \delta, a, b)$;
- (iv) $\|f_{m,\delta}\|_{C[a_1, b_1]} \leq M_3 \|f\|_\gamma$,



Direct Results for Certain Family of Integral Type Operators

Vijay Gupta and Ogün Dođru

Title Page

Contents



Go Back

Close

Quit

Page 16 of 21

where M_i , for $i = 1, 2, 3$ are certain unrelated constants independent of f and δ . The r -th order modulus of continuity $\omega_r(f, \delta, a, b)$ for a function f continuous on the interval $[a, b]$ is defined by:

$$\omega_r(f, \delta, a, b) = \sup \{ |\Delta_h^r f(x)| : |h| \leq \delta; x, x+h \in [a, b] \}.$$

For $r = 1$, $\omega_1(f, \delta)$ is written simply $\omega_f(\delta)$ or $\omega(f, \delta)$.

The following error estimation is in terms of higher order modulus of continuity:

Theorem 3.3. *Let $f \in C_\gamma[0, \infty)$, $\gamma > 0$ and $0 < a < a_1 < b_1 < b < \infty$. Then for all n sufficiently large*

$$\|B_n^{(r)}(f, *) - f^{(r)}(x)\|_{C[a_1, b_1]} \leq \max \left\{ M_3 \omega_2(f^{(r)}, n^{-1/2}, a, b), M_4 n^{-1} \|f\|_\gamma \right\}$$


where $M_3 = M_3(r)$, $M_4 = M_4(r, f)$.

Proof. First by the linearity property, we have

$$\begin{aligned} \|B_n^{(r)}(f, *) - f^{(r)}\|_{C[a_1, b_1]} &\leq \|B_n^{(r)}((f - f_{2,\delta}), *)\|_{C[a_1, b_1]} \\ &\quad + \|B_n^{(r)}(f_{2,\delta}, *) - f_{2,\delta}^{(r)}\|_{C[a_1, b_1]} + \|f^{(r)} - f_{2,\delta}^{(r)}\|_{C[a_1, b_1]} \\ &=: A_1 + A_2 + A_3. \end{aligned}$$

By property (iii) of the Steklov mean, we have

$$A_3 \leq C_1 \omega_2(f^{(r)}, \delta, a, b).$$



**Direct Results for Certain
Family of Integral Type
Operators**

Vijay Gupta and Ogün Dođru

Title Page

Contents



Go Back

Close

Quit

Page 17 of 21

Next using Theorem 3.2, we have

$$A_2 \leq C_2 n^{-(k+1)} \sum_{j=r}^{r+2} \left\| f_{2,\delta}^{(j)} \right\|_{C[a,b]}.$$

By applying the interpolation property due to Goldberg and Meir [2] for each $j = r, r + 1, r + 2$, we have

$$\left\| f_{2,\delta}^{(j)} \right\|_{C[a,b]} \leq C_3 \left\{ \left\| f_{2,\delta} \right\|_{C[a,b]} + \left\| f_{2,\delta}^{(r+2)} \right\|_{C[a,b]} \right\}.$$

Therefore by applying properties (ii) and (iv) of the Steklov mean, we obtain

$$A_2 \leq C_4 n^{-1} \left\{ \left\| f \right\|_{\gamma} + \delta^{-2} \omega_2(f^{(r)}, \delta) \right\}.$$

Finally we estimate A_1 , choosing a^*, b^* satisfying the condition $0 < a < a^* < a_1 < b_1 < b^* < b < \infty$. Also let $\psi(t)$ denote the characteristic function of the interval $[a^*, b^*]$, then

$$\begin{aligned} A_1 &\leq \left\| B_n^{(r)}(\psi(t)(f(t) - f_{2,\delta}(t)), *) \right\|_{C[a_1, b_1]} \\ &\quad + \left\| B_n^{(r)}((1 - \psi(t))(f(t) - f_{2,\delta}(t)), *) \right\|_{C[a_1, b_1]} \\ &=: A_4 + A_5. \end{aligned}$$

We may note here that to estimate A_4 and A_5 , it is enough to consider their expressions without the linear combinations. By Lemma 2.5, we have

$$\begin{aligned} &B_n^{(r)}(\psi(t)(f(t) - f_{2,\delta}(t)), x) \\ &= \frac{(n - r - 1)!(n + r - 1)!}{n!(n - 1)!} \sum_{\nu=0}^{\infty} p_{n+r,\nu}(x) \int_0^{\infty} p_{n-1,\nu+r-1}(t) f^{(r)}(t) dt. \end{aligned}$$



Direct Results for Certain
Family of Integral Type
Operators

Vijay Gupta and Ogün Dođru

Title Page

Contents



Go Back

Close

Quit


Page 18 of 21

Hence

$$\|B_n^{(r)}(\psi(t)(f(t) - f_{2,\delta}(t)), *)\|_{C[a,b]} \leq C_5 \|f^{(r)} - f_{2,\delta}^{(r)}\|_{C[a^*,b^*]}.$$

Now for $x \in [a_1, b_1]$ and $t \in [0, \infty) \setminus [a^*, b^*]$, we choose a $\delta_1 > 0$ satisfying $|t - x| \geq \delta_1$. Therefore by Lemma 2.4 and the Schwarz inequality, we have

$$\begin{aligned} I &= |B_n^{(r)}((1 - \psi(t))(f(t) - f_{2,\delta}(t)), x)| \\ &\leq \sum_{\substack{2i+j \leq r \\ i,j \geq 0}} n^i \frac{|Q_{i,j,r}(x)|}{x^r} \\ &\quad \times \frac{1}{n} \sum_{\nu=1}^{\infty} p_{n,\nu}(x) |\nu - (n+1)x|^j \int_0^{\infty} p_{n,\nu-1}(t) (1 - \psi(t)) |f(t) - f_{2,\delta}(t)| dt \\ &\quad + (1+x)^{-n-1} |(-n-1)(-n) \cdots (-n-r)| (1 - \psi(0)) |f(0) - f_{2,\delta}(0)| \\ &\leq C_6 \|f\|_{\gamma} \left\{ \sum_{\substack{2i+j \leq r \\ i,j \geq 0}} n^{i-1} \sum_{\nu=1}^{\infty} p_{n,\nu}(x) |\nu - (n+1)x|^j \right. \\ &\quad \left. \times \int_{|t-x| \geq \delta_1} p_{n,\nu-1}(t) dt + (1+x)^{-n-1} |(-n-1)(-n) \cdots (-n-r)| \right\} \\ &\leq C_6 \|f\|_{\gamma} \left\{ \delta_1^{-2s} \sum_{\substack{2i+j \leq r \\ i,j \geq 0}} n^{i-1} \sum_{\nu=1}^{\infty} p_{n,\nu}(x) |\nu - (n+1)x|^j \left(\int_0^{\infty} p_{n,\nu-1}(t) dt \right)^{\frac{1}{2}} \right. \end{aligned}$$



Direct Results for Certain
Family of Integral Type
Operators

Vijay Gupta and Ogün Dođru

Title Page

Contents



Go Back

Close

Quit

Page 19 of 21

$$\begin{aligned}
& \times \left(\int_0^\infty p_{n,\nu-1}(t)(t-x)^{4s} dt \right)^{\frac{1}{2}} + (1+x)^{-n-1} |(-n-1)(-n)\cdots(-n-r)| \Big\} \\
& \leq C_6 \|f\|_\gamma \delta_1^{-2s} \\
& \times \sum_{\substack{2i+j \leq r \\ i,j \geq 0}} n^i \left\{ \frac{1}{n} \sum_{\nu=0}^\infty p_{n,\nu}(x) [\nu - (n+1)x]^{2j} - (1+x)^{-n-1} - \{-(n+1)x\}^{2j} \right\}^{\frac{1}{2}} \\
& \times \left\{ \frac{1}{n} \sum_{\nu=0}^\infty p_{n,\nu}(x) \int_0^\infty p_{n,\nu-1}(t)(t-x)^{4s} dt - (1+x)^{-n-1}(-x)^{4s} \right. \\
& \left. - (1+x)^{-n-1}(-x)^{4s} \right\}^{\frac{1}{2}} + C_6 \|f\|_\gamma (1+x)^{-n-1} |(-n-1)(-n)\cdots(-n-r)|.
\end{aligned}$$

Hence by Lemma 2.1 and Lemma 2.2, we have

$$I \leq C_7 \|f\|_\gamma \delta_1^{-2s} \mathcal{O} \left(n^{(i+\frac{j}{2}-s)} \right) \leq C_7 n^{-q} \|f\|_\gamma, \quad q = s - \frac{r}{2},$$

where the last term vanishes as $n \rightarrow \infty$. Now choosing q satisfying $q \geq 1$, we obtain

$$I \leq C_7 n^{-1} \|f\|_\gamma.$$

Therefore by property (iii) of the Steklov mean, we get

$$\begin{aligned}
A_1 & \leq C_8 \left\| f^{(r)} - f_{2,\delta}^{(r)} \right\|_{C[a^*,b^*]} + C_7 n^{-1} \|f\|_\gamma \\
& \leq C_9 \omega_2(f^{(r)}, \delta, a, b) + C_7 n^{-1} \|f\|_\gamma.
\end{aligned}$$

Choosing $\delta = n^{-1/2}$, the theorem follows. \square



**Direct Results for Certain
Family of Integral Type
Operators**

Vijay Gupta and Ogün Doğru

Title Page

Contents



Go Back


Close

Quit

Page 20 of 21

References

- [1] G. FREUD AND V. POPOV, On approximation by spline functions, *Proc. Conf. on Constructive Theory Functions*, Budapest (1969), 163–172.
- [2] S. GOLDBERG AND V. MEIR, Minimum moduli of ordinary differential operators, *Proc. London Math. Soc.*, **23** (1971), 1–15.
- [3] V. GUPTA AND G.S. SRIVASTAVA, Convergence of derivatives by summation-integral type operators, *Revista Colombiana de Matematicas*, **29** (1995), 1–11.
- [4] E. HEWITT AND K. STROMBERG, *Real and Abstract Analysis*, McGraw Hill, New York, 1956.
- [5] H.M. SRIVASTAVA AND V. GUPTA, A certain family of summation integral type operators, *Mathematical and Computer Modelling*, **37** (2003), 1307–1315.



Direct Results for Certain
Family of Integral Type
Operators

Vijay Gupta and Ogün Dođru

Title Page

Contents



Go Back

Close

Quit

Page 21 of 21