

## ON ČEBYŠEV TYPE INEQUALITIES INVOLVING FUNCTIONS WHOSE DERIVATIVES BELONG TO $L_p$ SPACES

B.G. PACHPATTE

57 Shri Niketan Colony  
Near Abhinay Talkies  
Aurangabad 431 001 (Maharashtra)  
India.

*EMail:* [bgpachpatte@gmail.com](mailto:bgpachpatte@gmail.com)



---

volume 7, issue 2, article 58,  
2006.

*Received 10 November, 2005;  
accepted 16 November, 2005.*

*Communicated by: I. Gavrea*

---

Abstract

Contents



Home Page

Go Back

Close

Quit

## Abstract


In this note we establish new Čebyšev type integral inequalities involving functions whose derivatives belong to  $L_p$  spaces via certain integral identities.

*2000 Mathematics Subject Classification:* 26D15, 26D20.

*Key words:* Čebyšev type inequalities,  $L_p$  spaces, Integral identities, Absolutely continuous functions, Hölder's integral inequality.

## Contents

|   |                            |    |
|---|----------------------------|----|
| 1 | Introduction .....         | 3  |
| 2 | Statement of Results ..... | 4  |
| 3 | Proof of Theorem 2.1 ..... | 7  |
| 4 | Proof of Theorem 2.2 ..... | 11 |
|   | References                 |    |



---

### On Čebyšev Type Inequalities Involving Functions Whose Derivatives Belong to $L_p$ Spaces

B.G. Pachpatte

---

Title Page

Contents



Go Back

Close

Quit

Page 2 of 16

# 1. Introduction

One of the classical and important inequalities discovered by P.L.Čebyšev [1] is the following integral inequality (see also [10, p. 207]):


$$(1.1) \quad |T(f, g)| \leq \frac{1}{12} (b - a)^2 \|f'\|_\infty \|g'\|_\infty,$$

where  $f, g : [a, b] \rightarrow \mathbb{R}$  are absolutely continuous functions whose derivatives  $f', g' \in L_\infty [a, b]$  and

$$(1.2) \quad T(f, g) = \frac{1}{b - a} \int_a^b f(x) g(x) dx - \left( \frac{1}{b - a} \int_a^b f(x) dx \right) \left( \frac{1}{b - a} \int_a^b g(x) dx \right),$$

which is called the Čebyšev functional, provided the integrals in (1.2) exist.

Because of fundamental importance of (1.1) in analysis and applications, many researchers have given considerable attention to it and a number of extensions, generalizations and variants have appeared in the literature, see [5], [6], [8] – [10] and the references given therein. The main purpose of the present note is to establish new inequalities similar to the inequality (1.1) involving functions whose derivatives belong to  $L_p$  spaces. The analysis used in the proofs is based on the integral identities proved in [3] and [2] and our results provide new estimates on these types of inequalities.



---

On Čebyšev Type Inequalities  
Involving Functions Whose  
Derivatives Belong to  $L_p$   
Spaces

B.G. Pachpatte

---

Title Page

Contents



Go Back

Close

Quit

Page 3 of 16

## 2. Statement of Results


In what follows  $\mathbb{R}$  and  $'$  denote the set of real numbers and the derivative of a function. Let  $[a, b] \subset \mathbb{R}$ ,  $a < b$ ; and as usual for any function  $h \in L_p[a, b]$ ,  $p > 1$  we define  $\|h\|_p = \left( \int_a^b |h(t)|^p dt \right)^{\frac{1}{p}}$ . We use the following notations to simplify the details of presentation. For suitable functions  $f, g : [a, b] \rightarrow \mathbb{R}$  we set

$$F = \frac{1}{3} \left[ \frac{f(a) + f(b)}{2} + 2f\left(\frac{a+b}{2}\right) \right],$$
$$G = \frac{1}{3} \left[ \frac{g(a) + g(b)}{2} + 2g\left(\frac{a+b}{2}\right) \right],$$

$$S(f, g) = FG - \frac{1}{b-a} \left[ F \int_a^b g(x) dx + G \int_a^b f(x) dx \right]$$
$$+ \left( \frac{1}{b-a} \int_a^b f(x) dx \right) \left( \frac{1}{b-a} \int_a^b g(x) dx \right),$$

$$H(f, g) = \frac{1}{b-a} \int_a^b [Fg(x) + Gf(x)] dx$$
$$- 2 \left( \frac{1}{b-a} \int_a^b f(x) dx \right) \left( \frac{1}{b-a} \int_a^b g(x) dx \right).$$

Now, we state our main results as follows.



On Čebyšev Type Inequalities  
Involving Functions Whose  
Derivatives Belong to  $L_p$   
Spaces

B.G. Pachpatte

Title Page

Contents



Go Back

Close

Quit

Page 4 of 16

**Theorem 2.1.** Let  $f, g : [a, b] \rightarrow \mathbb{R}$  be absolutely continuous functions whose derivatives  $f', g' \in L_p[a, b]$ ,  $p > 1$ . Then we have the inequalities

$$(2.1) \quad |T(f, g)| \leq \frac{1}{(b-a)^3} \|f'\|_p \|g'\|_p \int_a^b (B(x))^{\frac{2}{q}} dx,$$

$$(2.2) \quad |T(f, g)| \leq \frac{1}{2(b-a)^2} \int_a^b \left[ |g(x)| \|f'\|_p + |f(x)| \|g'\|_p \right] (B(x))^{\frac{1}{q}} dx,$$

where

$$(2.3) \quad B(x) = \frac{(x-a)^{q+1} + (b-x)^{q+1}}{q+1},$$


for  $x \in [a, b]$  and  $\frac{1}{p} + \frac{1}{q} = 1$ .

The following variants of the inequalities in (2.1) and (2.2) hold.

**Theorem 2.2.** Let  $f, g : [a, b] \rightarrow \mathbb{R}$  be absolutely continuous functions whose derivatives  $f', g' \in L_p[a, b]$ ,  $p > 1$ . Then we have the inequalities

$$(2.4) \quad |S(f, g)| \leq \frac{1}{(b-a)^2} M^{\frac{2}{q}} \|f'\|_p \|g'\|_p,$$

$$(2.5) \quad |H(f, g)| \leq \frac{1}{(b-a)^2} M^{\frac{1}{q}} \int_a^b \left[ |g(x)| \|f'\|_p + |f(x)| \|g'\|_p \right] dx,$$



On Čebyšev Type Inequalities  
Involving Functions Whose  
Derivatives Belong to  $L_p$   
Spaces

B.G. Pachpatte

Title Page

Contents



Go Back

Close


Quit

Page 5 of 16

where

$$(2.6) \quad M = \frac{(2^{q+1} + 1)(b - a)^{q+1}}{3(q + 1)6^q},$$

and  $\frac{1}{p} + \frac{1}{q} = 1$ .



---

**On Čebyšev Type Inequalities  
Involving Functions Whose  
Derivatives Belong to  $L_p$   
Spaces**

B.G. Pachpatte

---

Title Page

Contents



Go Back

Close

Quit

Page 6 of 16

### 3. Proof of Theorem 2.1

From the hypotheses we have the following identities (see [3, 7]):

$$(3.1) \quad f(x) - \frac{1}{b-a} \int_a^b f(t) dt = \frac{1}{b-a} \int_a^b k(x,t) f'(t) dt,$$


$$(3.2) \quad g(x) - \frac{1}{b-a} \int_a^b g(t) dt = \frac{1}{b-a} \int_a^b k(x,t) g'(t) dt,$$

for  $x \in [a, b]$ , where

$$(3.3) \quad k(x,t) = \begin{cases} t-a & \text{if } t \in [a, x] \\ t-b & \text{if } t \in (x, b] \end{cases}.$$

Multiplying the left sides and right sides of (3.1) and (3.2) we have

$$(3.4) \quad f(x)g(x) - f(x) \left( \frac{1}{b-a} \int_a^b g(t) dt \right) - g(x) \left( \frac{1}{b-a} \int_a^b f(t) dt \right) \\ + \left( \frac{1}{b-a} \int_a^b f(t) dt \right) \left( \frac{1}{b-a} \int_a^b g(t) dt \right) \\ = \frac{1}{(b-a)^2} \left( \int_a^b k(x,t) f'(t) dt \right) \left( \int_a^b k(x,t) g'(t) dt \right).$$



---

On Čebyšev Type Inequalities  
Involving Functions Whose  
Derivatives Belong to  $L_p$   
Spaces

B.G. Pachpatte

---

Title Page

Contents



Go Back

Close

Quit

Page 7 of 16

Integrating both sides of (3.4) with respect to  $x$  over  $[a, b]$  and dividing both sides of the resulting identity by  $(b - a)$  we get


$$(3.5) \quad T(f, g) = \frac{1}{(b-a)^3} \int_a^b \left( \int_a^b k(x, t) f'(t) dt \right) \left( \int_a^b k(x, t) g'(t) dt \right) dx.$$

From (3.5) and using the properties of modulus and Hölder's integral inequality we have

$$(3.6) \quad |T(f, g)| \leq \frac{1}{(b-a)^3} \int_a^b \left( \int_a^b |k(x, t)| |f'(t)| dt \right) \left( \int_a^b |k(x, t)| |g'(t)| dt \right) dx \\ \leq \frac{1}{(b-a)^3} \int_a^b \left( \left\{ \int_a^b |k(x, t)|^q dt \right\}^{\frac{1}{q}} \left\{ \int_a^b |f'(t)|^p dt \right\}^{\frac{1}{p}} \right) \\ \times \left( \left\{ \int_a^b |k(x, t)|^q dt \right\}^{\frac{1}{q}} \left\{ \int_a^b |g'(t)|^p dt \right\}^{\frac{1}{p}} \right) dx \\ = \frac{1}{(b-a)^3} \|f'\|_p \|g'\|_p \int_a^b \left( \left\{ \int_a^b |k(x, t)|^q dt \right\}^{\frac{1}{q}} \right)^2 dx.$$

A simple calculation shows that (see [4])

$$\int_a^b |k(x, t)|^q dt = \int_a^x |t-a|^q dt + \int_x^b |t-b|^q dt \\ = \int_a^x (t-a)^q dt + \int_x^b (b-t)^q dt$$



On Čebyšev Type Inequalities  
Involving Functions Whose  
Derivatives Belong to  $L_p$   
Spaces

B.G. Pachpatte

Title Page

Contents



Go Back

Close

Quit

Page 8 of 16



$$(3.7) \quad = \frac{(x-a)^{q+1} + (b-x)^{q+1}}{q+1} = B(x).$$

Using (3.7) in (3.6) we get (2.1).

Multiplying both sides of (3.1) and (3.2) by  $g(x)$  and  $f(x)$  respectively and adding the resulting identities we get


$$(3.8) \quad 2f(x)g(x) - \left[ g(x) \left( \frac{1}{b-a} \int_a^b f(t) dt \right) + f(x) \left( \frac{1}{b-a} \int_a^b g(t) dt \right) \right] = g(x) \left( \frac{1}{b-a} \int_a^b k(x,t) f'(t) dt \right) + f(x) \left( \frac{1}{b-a} \int_a^b k(x,t) g'(t) dt \right).$$

Integrating both sides of (3.8) with respect to  $x$  over  $[a, b]$  and rewriting we obtain

$$(3.9) \quad T(f, g) = \frac{1}{2(b-a)^2} \int_a^b \left[ g(x) \int_a^b k(x,t) f'(t) dt + f(x) \int_a^b k(x,t) g'(t) dt \right] dx.$$

From (3.9) and using the properties of modulus, Hölder's integral inequality and (3.7) we have

$$|T(f, g)| \leq \frac{1}{2(b-a)^2} \int_a^b \left[ |g(x)| \int_a^b |k(x,t)| |f'(t)| dt + |f(x)| \int_a^b |k(x,t)| |g'(t)| dt \right] dx$$



On Čebyšev Type Inequalities  
Involving Functions Whose  
Derivatives Belong to  $L_p$   
Spaces

B.G. Pachpatte

Title Page

Contents



Go Back


Close

Quit

Page 9 of 16

$$\begin{aligned}
&\leq \frac{1}{2(b-a)^2} \int_a^b \left[ |g(x)| \left\{ \int_a^b |k(x,t)|^q dt \right\}^{\frac{1}{q}} \left\{ \int_a^b |f'(t)|^p dt \right\}^{\frac{1}{p}} \right. \\
&\quad \left. + |f(x)| \left\{ \int_a^b |k(x,t)|^q dt \right\}^{\frac{1}{q}} \left\{ \int_a^b |g'(t)|^p dt \right\}^{\frac{1}{p}} \right] dx \\
&= \frac{1}{2(b-a)^2} \int_a^b \left[ |g(x)| \|f'\|_p + |f(x)| \|g'\|_p \right] \left\{ \int_a^b |k(x,t)|^q dt \right\}^{\frac{1}{q}} dx \\
&= \frac{1}{2(b-a)^2} \int_a^b \left[ |g(x)| \|f'\|_p + |f(x)| \|g'\|_p \right] (B(x))^{\frac{1}{q}} dx.
\end{aligned}$$

This is the required inequality in (2.2). The proof is complete.



**On Čebyšev Type Inequalities  
Involving Functions Whose  
Derivatives Belong to  $L_p$   
Spaces**

B.G. Pachpatte

Title Page

Contents



Go Back

Close

Quit

Page 10 of 16

## 4. Proof of Theorem 2.2

From the hypotheses we have the following identities (see [2]):

$$(4.1) \quad F - \frac{1}{b-a} \int_a^b f(x) dx = \frac{1}{b-a} \int_a^b m(x) f'(x) dx,$$

$$(4.2) \quad G - \frac{1}{b-a} \int_a^b g(x) dx = \frac{1}{b-a} \int_a^b m(x) g'(x) dx,$$

where


$$(4.3) \quad m(x) = \begin{cases} x - \frac{5a+b}{6} & \text{if } x \in [a, \frac{a+b}{2}] \\ x - \frac{a+5b}{6} & \text{if } x \in [\frac{a+b}{2}, b] \end{cases}.$$

Multiplying the left sides and right sides of (4.1) and (4.2) we get

$$(4.4) \quad S(f, g) = \frac{1}{(b-a)^2} \left( \int_a^b m(x) f'(x) dx \right) \left( \int_a^b m(x) g'(x) dx \right).$$

From (4.4) and using the properties of modulus and Hölder's integral inequality, we have

$$|S(f, g)| \leq \frac{1}{(b-a)^2} \left( \int_a^b |m(x)| |f'(x)| dx \right) \left( \int_a^b |m(x)| |g'(x)| dx \right)$$



On Čebyšev Type Inequalities  
Involving Functions Whose  
Derivatives Belong to  $L_p$   
Spaces

B.G. Pachpatte

Title Page

Contents



Go Back

Close

Quit

Page 11 of 16



Title Page

Contents



Go Back

Close

Quit

Page 12 of 16

$$\begin{aligned}
 &\leq \frac{1}{(b-a)^2} \left( \left\{ \int_a^b |m(x)|^q dx \right\}^{\frac{1}{q}} \left\{ \int_a^b |f'(x)|^p dx \right\}^{\frac{1}{p}} \right) \\
 &\quad \times \left( \left\{ \int_a^b |m(x)|^q dx \right\}^{\frac{1}{q}} \left\{ \int_a^b |g'(x)|^p dx \right\}^{\frac{1}{p}} \right) \\
 (4.5) \quad &= \frac{1}{(b-a)^2} \left( \left\{ \int_a^b |m(x)|^q dx \right\}^{\frac{1}{q}} \right)^2 \|f'\|_p \|g'\|_p.
 \end{aligned}$$

A simple computation gives (see [2])

$$\begin{aligned}
 \int_a^b |m(x)|^q dx &= \int_a^{\frac{a+b}{2}} \left| x - \frac{5a+b}{6} \right|^q dx + \int_{\frac{a+b}{2}}^b \left| x - \frac{a+5b}{6} \right|^q dx \\
 &= \int_a^{\frac{5a+b}{6}} \left( \frac{5a+b}{6} - x \right)^q dx + \int_{\frac{5a+b}{6}}^{\frac{a+b}{2}} \left( x - \frac{5a+b}{6} \right)^q dx \\
 &\quad + \int_{\frac{a+b}{2}}^{\frac{a+5b}{6}} \left( \frac{a+5b}{6} - x \right)^q dx + \int_{\frac{a+5b}{6}}^b \left( x - \frac{a+5b}{6} \right)^q dx \\
 &= \frac{1}{q+1} \left[ \left( \frac{5a+b}{6} - a \right)^{q+1} + \left( \frac{a+b}{2} - \frac{5a+b}{6} \right)^{q+1} \right. \\
 &\quad \left. + \left( \frac{a+5b}{6} - \frac{a+b}{2} \right)^{q+1} + \left( b - \frac{a+5b}{6} \right)^{q+1} \right] \\
 (4.6) \quad &= \frac{(2^{q+1} + 1)(b-a)^{q+1}}{3(q+1)6^q} = M.
 \end{aligned}$$

Using (4.6) in (4.5) we get the required inequality in (2.4).

Multiplying both sides of (4.1) and (4.2) by  $g(x)$  and  $f(x)$  respectively and adding the resulting identities we get


$$(4.7) \quad Fg(x) + Gf(x) - \left[ g(x) \left( \frac{1}{b-a} \int_a^b f(x) dx \right) + f(x) \left( \frac{1}{b-a} \int_a^b g(x) dx \right) \right] = g(x) \left( \frac{1}{b-a} \int_a^b m(x) f'(x) dx \right) + f(x) \left( \frac{1}{b-a} \int_a^b m(x) g'(x) dx \right).$$

Integrating both sides of (4.7) with respect to  $x$  over  $[a, b]$  and dividing both sides of the resulting identity by  $(b-a)$  we get

$$(4.8) \quad H(f, g) = \frac{1}{(b-a)^2} \int_a^b \left[ g(x) \int_a^b m(x) f'(x) dx + f(x) \int_a^b m(x) g'(x) dx \right] dx.$$

From (4.8) and using the properties of modulus, Hölder's integral inequality and (4.6) we have

$$|H(f, g)| \leq \frac{1}{(b-a)^2} \int_a^b \left[ |g(x)| \int_a^b |m(x)| |f'(x)| dx + |f(x)| \int_a^b |m(x)| |g'(x)| dx \right] dx$$



On Čebyšev Type Inequalities  
Involving Functions Whose  
Derivatives Belong to  $L_p$   
Spaces

B.G. Pachpatte

Title Page

Contents



Go Back


Close

Quit

Page 13 of 16

$$\begin{aligned}
&\leq \frac{1}{(b-a)^2} \int_a^b \left[ |g(x)| \left\{ \int_a^b |m(x)|^q dx \right\}^{\frac{1}{q}} \left\{ \int_a^b |f'(x)|^p dx \right\}^{\frac{1}{p}} \right. \\
&\quad \left. + |f(x)| \left\{ \int_a^b |m(x)|^q dx \right\}^{\frac{1}{q}} \left\{ \int_a^b |g'(x)|^p dx \right\}^{\frac{1}{p}} \right] dx \\
&= \frac{1}{(b-a)^2} \int_a^b \left[ |g(x)| \|f'\|_p + |f(x)| \|g'\|_p \right] \left( \int_a^b |m(x)|^q dx \right)^{\frac{1}{q}} \\
&= \frac{1}{(b-a)^2} M^{\frac{1}{q}} \int_a^b \left[ |g(x)| \|f'\|_p + |f(x)| \|g'\|_p \right] dx.
\end{aligned}$$

This is the desired inequality in (2.5). The proof is complete.



**On Čebyšev Type Inequalities  
Involving Functions Whose  
Derivatives Belong to  $L_p$   
Spaces**

B.G. Pachpatte

Title Page

Contents



Go Back


Close

Quit

Page 14 of 16

## References

- [1] P.L. ČEBYŠEV, Sur les expressions approximatives des intégrales par les autres prises entre les mêmes limites, *Proc.Math.Soc. Charkov*, **2** (1882), 93–98.
- [2] S.S. DRAGOMIR, On Simpson's quadrature formula for differentiable mappings whose derivatives belong to  $L_p$  spaces and applications, *RGMIA Res. Rep. Coll.*, **1**(2) (1998), 89–96.
- [3] S.S. DRAGOMIR AND N.S. BARNETT, An Ostrowski type inequality for mappings whose second derivatives are bounded and applications, *RGMIA Res. Rep. Coll.*, **1**(2) (1998), 69–77.
- [4] S.S. DRAGOMIR AND S. WANG, A new inequality of Ostrowski's type in  $L_p$  norm, *Indian J. Math.*, **40**(3) (1998), 299–304.
- [5] H.P. HEINIG AND L. MALIGRANDA, Chebyshev inequality in function spaces, *Real Analysis Exchange*, **17** (1991-92), 211–247.
- [6] D.S. MITRINOVIĆ, J.E. PEČARIĆ AND A.M. FINK, *Classical and New Inequalities in Analysis*, Kluwer Academic Publishers, Dordrecht, 1993.
- [7] D.S. MITRINOVIĆ, J.E. PEČARIĆ AND A.M. FINK, *Inequalities Involving Functions and Their Integrals and Derivatives*, Kluwer Academic Publishers, Dordrecht, 1994.
- [8] B.G. PACHPATTE, On Trapezoid and Grüss like integral inequalities, *Tamkang J. Math.*, **34**(4) (2003), 365–369.



---

On Čebyšev Type Inequalities  
Involving Functions Whose  
Derivatives Belong to  $L_p$   
Spaces

B.G. Pachpatte

---

Title Page

Contents




Go Back

Close

Quit

Page 15 of 16

- [9] B.G. PACHPATTE, On Ostrowski-Grüss-Čebyšev type inequalities for functions whose modulus of derivatives are convex, *J. Inequal. Pure and Appl. Math.*, **6**(4) (2005), Art. 128. [ONLINE: <http://jipam.vu.edu.au/article.php?sid=602>].
- [10] J.E. PEČARIĆ, F. PORCHAN AND Y. TONG, *Convex Functions, Partial Orderings and Statistical Applications*, Academic Press, San Diego, 1992.



---

**On Čebyšev Type Inequalities  
Involving Functions Whose  
Derivatives Belong to  $L_p$   
Spaces**

B.G. Pachpatte

---

Title Page

Contents



Go Back

Close

Quit

Page 16 of 16