



Patterns in Inversion Sequences II: Inversion Sequences Avoiding Triples of Relations

Megan Martinez

Department of Mathematics

Ithaca College

Ithaca, NY 14850

USA

mmartinez@ithaca.edu

Carla Savage

Department of Computer Science

North Carolina State University

Raleigh, NC 27695

USA

savage@ncsu.edu

Abstract

Inversion sequences of length n , \mathbf{I}_n , are integer sequences (e_1, \dots, e_n) with $0 \leq e_i < i$ for each i . The study of patterns in inversion sequences was initiated recently by Mansour-Shattuck and Corteel-Martinez-Savage-Weselcouch through a systematic study of inversion sequences avoiding words of length 3. We continue this investigation by reframing the notion of a length-3 pattern from a “word of length 3,” $w_1w_2w_3$, to a “triple of binary relations,” (ρ_1, ρ_2, ρ_3) , and consider the set $\mathbf{I}_n(\rho_1, \rho_2, \rho_3)$ consisting of those $e \in \mathbf{I}_n$ with no $i < j < k$ such that $e_i\rho_1e_j$, $e_j\rho_2e_k$, and $e_i\rho_3e_k$. We show that “avoiding a triple of relations” can characterize inversion sequences with a variety of monotonicity or unimodality conditions, or with multiplicity constraints on the elements. We uncover several interesting enumeration results and relate pattern avoiding inversion sequences to familiar combinatorial families. We highlight open questions about the relationship between pattern avoiding inversion sequences and a variety of

classes of pattern avoiding permutations. For several combinatorial sequences, pattern avoiding inversion sequences provide a simpler interpretation than otherwise known.

1 Introduction

Pattern avoiding permutations have been studied extensively for their connections in computer science, biology, and other fields of mathematics. Within combinatorics they have proven their usefulness, providing an interpretation that relates a vast array of combinatorial structures. See the comprehensive survey of Kitaev [17].

The notion of pattern avoidance in inversion sequences was introduced by Corteel, Martinez, Savage, and Weselcouch [13] and Mansour and Shattuck [23]. An *inversion sequence* is an integer sequence (e_1, e_2, \dots, e_n) satisfying $0 \leq e_i < i$ for all $i \in [n] = \{1, 2, \dots, n\}$. There is a natural bijection $\Theta : \mathbf{S}_n \rightarrow \mathbf{I}_n$ from \mathbf{S}_n , the set of permutations of $[n]$, to \mathbf{I}_n , the set of inversion sequences of length n . Under this bijection, $e = \Theta(\pi)$ is obtained from a permutation $\pi = \pi_1\pi_2 \dots \pi_n \in \mathbf{S}_n$ by setting $e_i = |\{j \mid j < i \text{ and } \pi_j > \pi_i\}|$.

The encoding of permutations as inversion sequences suggests that it could be illuminating to study patterns in inversion sequences in the same way that patterns have been studied in permutations. The paper of Corteel, et al. [13] focused on the enumeration of inversion sequences that avoid words of length three and the paper of Mansour and Shattuck [23] targets permutations of length 3. For example, the inversion sequences $e \in \mathbf{I}_n$ that avoid the pattern 021 are those with no $i < j < k$ such that $e_i < e_j > e_k$ and $e_i < e_k$. We denote these by $\mathbf{I}_n(021)$. Similarly, $\mathbf{I}_n(010)$ is the set of $e \in \mathbf{I}_n$ with no $i < j < k$ such that $e_i < e_j > e_k$ and $e_i = e_k$. The results in each of these works [13, 23] related pattern avoidance in inversion sequences to a number of well-known combinatorial sequences including the Fibonacci numbers, Bell numbers, large Schröder numbers, and Euler up/down numbers. They also gave rise to natural sequences that previously had not appeared in the On-Line Encyclopedia of Integer Sequences (OEIS) [15].

In this paper we reframe the notion of a length-3 pattern from a word of length 3 to a triple of binary relations. For a fixed triple of binary relations (ρ_1, ρ_2, ρ_3) , we study the set $\mathbf{I}_n(\rho_1, \rho_2, \rho_3)$ consisting of those $e \in \mathbf{I}_n$ with no $i < j < k$ such that $e_i\rho_1e_j$, $e_j\rho_2e_k$, and $e_i\rho_3e_k$. For example, $\mathbf{I}_n(<, >, <) = \mathbf{I}_n(021)$ and $\mathbf{I}_n(<, >, =) = \mathbf{I}_n(010)$. As another example, $\mathbf{I}_n(\geq, \geq, \geq)$ is the set of inversion sequences in \mathbf{I}_n with no weakly decreasing subsequence of length 3, i.e., those inversion sequences avoiding all of the words in the set $\{000, 100, 110, 210\}$.

Table 1 illustrates that avoiding a single triple of relations can characterize inversion sequences with a variety of natural monotonicity or unimodality conditions, or with multiplicity constraints on the appearance of elements in the inversion sequence. Moreover, as will be seen, inversion sequences avoiding a single triple of relations can provide realizations of certain combinatorial sequences that are simpler than known realizations as pattern avoiding permutations or other combinatorial structures.

For this project, we considered all triples of relations in the set $\{<, >, \leq, \geq, =, \neq, -\}^3$. The relation “ $-$ ” on a set S is all of $S \times S$; that is, x “ $-$ ” y for all $x, y \in S$. There are 343 possi-

ble triples of relations (patterns). For each pattern, (ρ_1, ρ_2, ρ_3) , we can consider the avoidance set, $\mathbf{I}_n(\rho_1, \rho_2, \rho_3)$, or the avoidance sequence $|\mathbf{I}_1(\rho_1, \rho_2, \rho_3)|, |\mathbf{I}_2(\rho_1, \rho_2, \rho_3)|, |\mathbf{I}_3(\rho_1, \rho_2, \rho_3)|, \dots$. We say two patterns are *equivalent* if they give rise to the same avoidance sets and two patterns are *Wilf-equivalent* if they yield the same avoidance sequence. For example, patterns (\geq, \geq, \geq) and $(\geq, \geq, -)$ are equivalent, whereas patterns $(\geq, \geq, -)$ and $(-, \leq, \geq)$ are inequivalent, but Wilf equivalent (see Section 2.16).

The 343 patterns partition into 98 equivalence classes of patterns. Additionally, we conjecture that there are 63 Wilf-equivalence classes. In this paper, we enumerate a number of avoidance sets either directly or by relating them to familiar combinatorial structures. These relationships establish Wilf-equivalence between a number of inequivalent patterns. However, in many cases, even where enumeration is elusive, Wilf-equivalence can be proved via a bijection. This paper presents the results we have been able to prove, documents what has not yet been settled, and highlights the most intriguing open questions.

We uncovered several interesting enumeration results beyond those discovered in the initial introduction of patterns in inversion sequences [13, 23]. For example, the inversion sequences e with no $i < j < k$ such that $e_i = e_j \leq e_k$ are counted by the Fibonacci numbers (as are, e.g., permutations avoiding the pair (321, 3412)). Inversion sequences with no $i < j < k$ such that $e_i < e_j \leq e_k$ are counted by powers of two (as are, e.g., permutations avoiding (213, 312)). Inversion sequences avoiding $(-, \neq, =)$ are counted by the Bell numbers; inversion sequences avoiding $(\geq, -, >)$ are counted by the large Schröder numbers; and inversion sequences avoiding $(-, \geq, <)$ are counted by the Catalan numbers. There are the same number of inversion sequences in \mathbf{I}_n avoiding (\neq, \neq, \neq) as there are Grassmannian permutations in \mathbf{S}_n . $\mathbf{I}_n(\neq, <, \leq)$ is counted by the number of 321-avoiding separable permutations in \mathbf{S}_n . $\mathbf{I}_n(\neq, <, \neq)$ has the same number of elements as the set of permutations in \mathbf{S}_n avoiding both of the patterns 321 and 2143.

Since an earlier draft of this paper was posted [24], several conjectures we made have been proven. Inversion sequences avoiding $e_i > e_j \geq e_k$ were shown to be counted by the “semi-Baxter sequence” as defined by Bouvel, et al. [7]. It was also shown that the semi-Baxter sequence enumerates plane permutations, which are defined as those permutations avoiding the barred pattern 21 $\bar{3}$ 54 or, equivalently, the vincular pattern 21 $\bar{4}$ 3. Inversion sequences avoiding $e_i \geq e_j \geq e_k$ were shown to have the same counting sequence as set partitions avoiding enhanced 3-crossings. This was shown using the obstinate kernel method by Lin [21] and by a bijection that utilizes 0,1 fillings of Ferrers shapes by Yan [31]. Additionally, the set $\mathbf{I}_n(\geq, \geq, >)$ was shown to have the same counting sequence as the Baxter permutations through use of the obstinate kernel method by Kim and Lin [16]. A number of open questions concerning inversion sequences still remain, and these will be highlighted throughout this paper.

Of the 63 conjectured Wilf-equivalence classes, five classes are counted by sequences that are ultimately constant. In the remaining 58 classes, 30 have counting sequences that appear to match already existing sequences in the OEIS. The remaining 28 resulted in new entries in the OEIS.

In Sections 2.1 through 2.30 we present our results and conjectures for the 30 Wilf classes

of pattern-avoiding inversion sequences that (appear to) match sequences in the OEIS. Table 2 gives an overview. Even for patterns with a “no” in this table, we are able to prove some Wilf-equivalence results.

For the patterns whose counting sequence did not originally match a sequence in the OEIS we have some limited results on Wilf-equivalence and counting. Table 3 gives an overview of the patterns in these 28 Wilf classes in addition to their newly created OEIS number. Our results and conjectures for a few of these patterns are presented in Section 3.

In Tables 1, 2, and 3, each row represents an equivalence class of patterns whose identifier is given in the last column. A Wilf class of patterns is identified by the number a_7 , the number of inversion sequences of length 7 avoiding a pattern in the class. Within a Wilf class, equivalence classes are labeled A,B,C, etc. So, for example, there are three equivalence classes of patterns counted by the Catalan numbers and these classes are labeled 429A, 429B, and 429C.

This paper is intended to provide a comprehensive overview of the enumeration sequences for a large number of classes of pattern avoiding inversion sequences. Many classes can be enumerated through basic counting techniques; in these situations, we have omitted the relevant proofs for the sake of brevity.

In the remainder of this section we give some definitions that will be needed throughout this paper, keeping the number of definitions to a minimum so that subsections can be somewhat self-contained. Finally, for completeness, we list the triples of relations whose avoidance sequences are ultimately constant.

1.1 Encodings of permutations

We compare Θ with a few other common encodings of permutations mentioned later in this paper: Lehmer codes, and *inv*codes, which are reverse Lehmer codes.

For a sequence $t = (t_1, \dots, t_n)$, let $t^R = (t_n, t_{n-1}, \dots, t_1)$, and, for a set of sequences T , let $T^R = \{t^R \mid t \in T\}$. For a permutation $\pi = (\pi_1, \dots, \pi_n) \in \mathbf{S}_n$, let $\pi^C = (n + 1 - \pi_1, \dots, n + 1 - \pi_n)$, and, for a set of permutations P , let $P^C = \{\pi^C \mid \pi \in P\}$. We use the following encodings.

- Define $\Theta : \mathbf{S}_n \rightarrow \mathbf{I}_n$ such that $e = \Theta(\pi)$ if and only if $e_i = |\{j \mid j < i \text{ and } \pi_j > \pi_i\}|$.
- Define $L : \mathbf{S}_n \rightarrow \mathbf{I}_n^R$ such that $e = L(\pi)$ if and only if $e_i = |\{j \mid j > i \text{ and } \pi_j < \pi_i\}|$.
- Define *inv*code : $\mathbf{S}_n \rightarrow \mathbf{I}_n$ such that $e = \text{invcode}(\pi)$ if and only if $e^R = L(\pi)$.

Note that $\text{invcode}(\pi) = e$ if and only if $e = \Theta((\pi^C)^R)$. Additionally, notice that if $\Theta(\pi) = e$, then i is a *descent* of π (that is, $\pi_i > \pi_{i+1}$), if and only if i is an *ascent* of e (that is, $e_i < e_{i+1}$).

We will make use of another encoding, ϕ , in Sections 2.24 and 2.29.

1.2 Operations on inversion sequences

For $e = (e_1, e_2, \dots, e_n) \in \mathbf{I}_n$ and any integer t , define $\sigma_t(e) = (e'_1, e'_2, \dots, e'_n)$, where $e'_i = e_i$ if $e_i = 0$, and $e'_i = e_i + t$ otherwise. So, σ_t adds t to the nonzero elements of a sequence (notice that t could be negative).

Concatenation is used to add an element to the beginning or end of an inversion sequence. For $e = (e_1, e_2, \dots, e_n) \in \mathbf{I}_n$, we have $0 \cdot e = (0, e_1, e_2, \dots, e_n) \in \mathbf{I}_{n+1}$, and, if $0 \leq i \leq n$, we have $e \cdot i = (e_1, e_2, \dots, e_n, i) \in \mathbf{I}_{n+1}$. More generally, $x \cdot y$ denotes the concatenation of two sequences or two words x, y .

1.3 Statistics on inversion sequences

In several cases, statistics on inversion sequences helped to prove or refine the results in Table 2 and make connections with statistics on other combinatorial families. The statistics used in this paper are defined below for an inversion sequence $e \in \mathbf{I}_n$.

$$\begin{aligned}
 \text{asc}(e) &= |\{i \in [n-1] \mid e_i < e_{i+1}\}| \\
 \text{zeros}(e) &= |\{i \in [n] \mid e_i = 0\}| \\
 \text{dist}(e) &= |\{e_1, e_2, \dots, e_n\}| \\
 \text{repeats}(e) &= |\{i \in [n-1] \mid e_i \in \{e_{i+1}, \dots, e_n\}\}| = n - \text{dist}(e) \\
 \text{maxim}(e) &= |\{i \in [n] \mid e_i = i - 1\}| \\
 \text{maxx}(e) &= \max\{e_1, e_2, \dots, e_n\} \\
 \text{last}(e) &= e_n.
 \end{aligned}$$

These statistics are, respectively, the number of ascents, the number of zeros, the number of distinct elements, the number of repeats, the number of maximal elements, the maximum element, and the last element of e .

1.4 Ultimately constant avoidance sequences

It can be easily checked that the following are the equivalence classes whose avoidance sequences are eventually constant.

pattern	avoidance sequence
$(-, -, -)$	$1, 2, 0, 0, 0, 0, \dots$
$(\leq, \leq, -)$	$1, 2, 1, 0, 0, 0, \dots$
$(-, -, \neq)$	$1, 2, 2, 1, 1, 1, \dots$
$(-, -, <)$	$1, 2, 2, 2, 2, 2, \dots$
$(-, \neq, -)$	$1, 2, 2, 2, 2, 2, \dots$
$(-, \neq, \neq)$	$1, 2, 3, 3, 3, 3, \dots$

Inversion sequences e satisfying:	are those with no $i < j < k$ such that:	and are counted by:	a_7 , equiv class	Sect.
Monotonicity constraints:				
$e_1 = e_2 = \dots = e_{n-1}$	$e_i \neq e_j$	n	7,D	2.1
$\exists t: e_1 = \dots = e_t \leq e_{t+1} = e_{t+2} = \dots = e_n$	$e_i < e_j \neq e_k$	$1 + n(n-1)/2$	22,A	2.4
$\exists t: e_1 = \dots = e_t < e_{t+1} < e_{t+2} < \dots < e_n$	$e_i < e_j \geq e_k$	2^{n-1}	64,C	2.6
$e_1 \leq e_2 \leq \dots \leq e_{n-1}$	$e_i > e_j$	Binom($2n-2, n-1$)	924	2.18
$e_1 \leq e_2 \leq \dots \leq e_{n-1} \leq e_n$	$e_j > e_k$	Catalan number C_n	429,A	2.14
$e_1 \leq e_2 < e_3 < \dots < e_n$	$e_j \geq e_k$	n	7,B	2.1
$e_1 < e_2 < \dots < e_{n-1}$	$e_i = e_j$	n	7,C	2.1
$e_2 \geq e_3 \geq \dots \geq e_n$	$e_j < e_k$	n	7,A	2.1
Unimodality constraints:				
$\exists t: e_1 = \dots = e_t \leq e_{t+1} \geq 0 = \dots = 0$	$e_i < e_j, e_i < e_k$	$1 + n(n-1)/2$	22,B	2.4
$\exists t: e_1 = \dots = e_t \leq e_{t+1} \geq e_{t+2} \geq \dots \geq e_n$	$e_i \neq e_j < e_k$	Grassmannian perms	121,A	2.7
$\exists t: e_1 = \dots = e_t < e_{t+1} > e_{t+2} > \dots > e_n$	$e_i \neq e_j \leq e_k$	$F_{n+2} - 1$	33,A	2.5
$\exists t: e_1 \leq \dots \leq e_t > e_{t+1} \geq e_{t+2} \geq \dots \geq e_n$	$e_i > e_j < e_k$	A033321	1265	2.20
$\exists t: e_1 \leq \dots \leq e_t > e_{t+1} > e_{t+2} > \dots > e_n$	$e_i > e_j \leq e_k$	A071356	1064	2.19
$\exists t: e_1 \leq \dots \leq e_t \geq e_{t+1} = e_{t+2} = \dots = e_n$	$e_i > e_j \neq e_k$	See Section 3	1079,A	3.1.3
$\exists t: e_1 < \dots < e_t \geq e_{t+1} = e_{t+2} = \dots = e_n$	$e_i \geq e_j \neq e_k$	$1 + n(n-1)/2$	22,C	2.4
$\exists t: e_1 < \dots < e_t \geq e_{t+1} \geq e_{t+2} \geq \dots \geq e_n$	$e_i = e_j < e_k$	2^{n-1}	64,A	2.6
$\exists t: e_1 < \dots < e_t \geq e_{t+1} > e_{t+2} > \dots > e_n$	$e_i = e_j \leq e_k$	F_{n+1}	21	2.3
$\exists(t \leq s): e_1 < \dots < e_t = \dots = e_s > \dots > e_n$	$e_i \geq e_j \leq e_k, e_i \neq e_k$	$F_{n+2} - 1$	33,B	2.5
Positive elements monotone:				
positive entries are strictly decreasing	$e_i < e_j \leq e_k$	2^{n-1}	64,B	2.6
positive entries are weakly decreasing	$e_i < e_j < e_k$	F_{2n-1}	233	2.12
positive entries are strictly increasing	$e_j \geq e_k, e_i < e_k$	Catalan number C_n	429,B	2.14
positive entries are weakly increasing	$e_j > e_k, e_i < e_k$	large Schröder number	1806,A	2.24
Multiplicity constraints:				
entries e_2, \dots, e_n are all distinct	$e_i \leq e_j = e_k$	2^{n-1}	64,D	2.6
$ \{e_1, e_2, \dots, e_n\} \leq 2$	$e_i \neq e_j \neq e_k, e_i \neq e_k$	Grassmannian perms	121,B	2.7
positive entries are distinct	$e_i < e_j = e_k$	Bell numbers	877,A	2.17
no three entries equal	$e_i = e_j = e_k$	Euler up/down nos.	1385	2.22
only adjacent entries can be equal	$e_i \neq e_j \neq e_k, e_i \neq e_k$	Bell numbers	877,C	2.17
$e_s = e_t \implies s - t \leq 1$	$e_i = e_k$	A229046	304	2.13

Table 1: Characterizations of inversion sequences avoiding triples of relations.

Inversion sequences with no $i < j < k$ such that:	appear to be counted by OEIS seq:	proven?	notes/OEIS description	a_7 , equiv class	Sect.
$e_i \neq e_j, e_i \neq e_k$	A004275	yes	$2(n-1)$ for $n > 1$	12,A	2.2
$e_i \geq e_j, e_i \neq e_k$	A004275	yes	$2(n-1)$ for $n > 1$	12,B	2.2
$e_i = e_j \leq e_k$	A000045	yes	Fibonacci numbers, F_{n+1}	21	2.3
$e_i < e_j \neq e_k$	A000124	yes	Lazy caterer sequence	22,A	2.4
$e_i < e_j, e_i < e_k$	A000124	yes	Lazy caterer sequence	22,B	2.4
$e_i \geq e_j \neq e_k$	A000124	yes	Lazy caterer sequence	22,C	2.4
$e_i \neq e_j \leq e_k$	A000071	yes	$F_{n+2} - 1$	33,A	2.5
$e_i \geq e_j \leq e_k, e_i \neq e_k$	A000071	yes	$F_{n+2} - 1$	33,B	2.5
$e_i = e_j < e_k$	A000079	yes	$\mathbf{I}_n(001)$, 2^{n-1} (see [13])	64,A	2.6
$e_i < e_j \leq e_k$	A000079	yes	2^{n-1}	64,B	2.6
$e_i < e_j \geq e_k$	A000079	yes	2^{n-1}	64,C	2.6
$e_i \leq e_j = e_k$	A000079	yes	2^{n-1}	64,D	2.6
$e_i \neq e_j < e_k$	A000325	yes	Grassmannian permutations	121,A	2.7
$e_i \neq e_j \neq e_k, e_i \neq e_k$	A000325	yes	Grassmannian permutations	121,B	2.7
$e_j \geq e_k, e_i \neq e_k$	A000325	yes	Grassmannian permutations	121,C	2.7
$e_i \neq e_j < e_k, e_i \leq e_k$	A034943	yes	321-avoiding separable perms	151	2.8
$e_i \neq e_j < e_k, e_i \neq e_k$	A088921	yes	$\mathbf{S}_n(321, 2143)$	185	2.9
$e_i \geq e_k$	A049125	no	ordered trees, internal nodes, adj. to ≤ 1 leaf	187	2.10
$e_i \leq e_j \geq e_k, e_i \neq e_k$	A005183	yes	$\mathbf{S}_n(132, 4312)$, $n2^{n-1} + 1$	193	2.11
$e_i < e_j < e_k$	A001519	yes	$\mathbf{I}_n(012)$, F_{2n-1} (see [13, 23])	233	2.12
$e_i = e_k$	A229046	no	recurrence \rightarrow gf?	304	2.13
$e_j > e_k$	A000108	yes	Catalan numbers	429,A	2.14
$e_j \geq e_k, e_i < e_k$	A000108	yes	Catalan numbers	429,B	2.14
$e_i \geq e_j, e_i \geq e_k$	A000108	yes	Catalan numbers [16]	429,C	2.14
$e_i \neq e_j = e_k$	A047970	yes	$\mathbf{S}_n(\overline{31542})$, nexus numbers	523	2.15
$e_j \leq e_k, e_i \geq e_k$	A108307	yes	set partitions avoiding enhanced 3-crossings	772,A	2.16
$e_i \geq e_j \geq e_k$	A108307	yes	set partitions avoiding enhanced 3-crossings [21, 31]	772,B	2.16
$e_i < e_j = e_k$	A000110	yes	$\mathbf{I}_n(011)$ (see [13]), Bell numbers B_n	877,A	2.17
$e_i = e_j \leq e_k$	A000110	no	$\mathbf{I}_n(000, 110)$, B_n	877,B	2.17
$e_j \neq e_k, e_i = e_k$	A000110	yes	$\mathbf{I}_n(010, 101)$, B_n	877,C	2.17
$e_i \geq e_j, e_i = e_k$	A000110	no	$\mathbf{I}_n(000, 101)$, B_n	877,D	2.17
$e_i > e_j$	A000984	yes	central binomial coefficients	924	2.18
$e_i > e_j \leq e_k$	A071356	no	certain underdiagonal lattice paths	1064	2.19
$e_i > e_j < e_k$	A033321	yes	$\mathbf{S}_n(2143, 3142, 4132)$ (see [8])	1265	2.20
$e_i > e_j, e_i \leq e_k$	A106228	no	$\mathbf{I}_n(101, 102)$, $\mathbf{S}_n(4123, 4132, 4213)$	1347	2.21
$e_i = e_j = e_k$	A000111	yes	$\mathbf{I}_n(000)$ (see [13]), Euler up/down numbers	1385	2.22
$e_i > e_j, e_i < e_k$	A200753	yes	$\mathbf{I}_n(102)$, [23]	1694	2.23
$e_j > e_k, e_i < e_k$	A006318	yes	$\mathbf{I}_n(021)$ [13, 23], large Schröder numbers R_{n-1}	1806,A	2.24
$e_i > e_j, e_i \geq e_k$	A006318	yes	$\mathbf{I}_n(210, 201, 101, 100)$, R_{n-1}	1806,B	2.24
$e_i \geq e_j, e_i > e_k$	A006318	yes	$\mathbf{I}_n(210, 201, 100, 110)$, R_{n-1}	1806,C	2.24
$e_i \geq e_j \neq e_k, e_i \geq e_k$	A006318	yes	$\mathbf{I}_n(210, 201, 101, 110)$, R_{n-1}	1806,D	2.24
$e_i \geq e_j \geq e_k, e_i > e_k$	A001181	yes	Baxter permutations [16]	2074	2.25
$e_i > e_j, e_i > e_k$	A098746	no	$\mathbf{I}_n(210, 201, 100)$, $\mathbf{S}_n(4231, 42513)$	2549,A	2.26
$e_i > e_j \neq e_k, e_i \geq e_k$	A098746	no	$\mathbf{I}_n(210, 201, 101)$, $\mathbf{S}_n(4231, 42513)$	2549,B	2.26
$e_i \geq e_j \neq e_k, e_i > e_k$	A098746	no	$\mathbf{I}_n(210, 201, 110)$, $\mathbf{S}_n(4231, 42513)$	2549,C	2.26
$e_j < e_k, e_i \geq e_k$	A117106	yes	$\mathbf{I}_n(201, 101)$, $\mathbf{S}_n(21\overline{354})$	2958,A	2.27
$e_i > e_j \geq e_k$	A117106	yes	$\mathbf{I}_n(210, 100)$, $\mathbf{S}_n(21\overline{354})$ [7]	2958,B	2.27
$e_i \geq e_j > e_k$	A117106	yes	$\mathbf{I}_n(210, 110)$, $\mathbf{S}_n(21\overline{354})$	2958,C	2.27
$e_j \leq e_k, e_i > e_k$	A117106	yes	$\mathbf{I}_n(201, 100)$, $\mathbf{S}_n(21\overline{354})$	2958,D	2.27
$e_j < e_k, e_i = e_k$	A113227	yes	$\mathbf{I}_n(101)$, $\mathbf{S}_n(1-23-4)$, (see [13])	3207,A	2.28
$e_i = e_j > e_k$	A113227	yes	$\mathbf{I}_n(110)$, $\mathbf{S}_n(1-23-4)$, (see [13])	3207,B	2.28
$e_i > e_j \neq e_k, e_i > e_k$	A212198	yes	$\mathbf{I}_n(201, 210)$, $MMP(0, 2, 0, 2)$ -avoiding perms	3720	2.28

Table 2: Patterns whose avoidance sequences appear to match sequences in the OEIS. Those marked as “yes” are cited, if known, and otherwise are proven in this paper.

Inversion sequences with no $i < j < k$ such that:	comments	initial terms a_1, \dots, a_9	a_7 , equiv class
$e_j \geq e_k, e_i \geq e_k$	$\mathbf{I}_n(000, 010, 011, 021)$ (A279544)	1, 2, 4, 10, 26, 73, 214, 651, 2040	214
$e_i \leq e_j, e_i \geq e_k$	$\mathbf{I}_n(000, 010, 110, 120)$ (A279551)	1, 2, 4, 10, 27, 79, 247, 816, 2822	247
$e_j \geq e_k, e_i = e_k$	$\mathbf{I}_n(000, 010)$ (A279552)	1, 2, 4, 10, 29, 95, 345, 1376, 5966	345
$e_j \neq e_k, e_i \geq e_k$	Wilf-eq. to 663B (Sec. 3.2, A279553)	1, 2, 5, 15, 50, 178, 663, 2552, 10071	663,A
$e_i \neq e_j, e_i \geq e_k$	Wilf-eq. to 663A (Sec. 3.2, A279553)	1, 2, 5, 15, 50, 178, 663, 2552, 10071	663,B
$e_i \neq e_j \neq e_k, e_i \geq e_k$	$\mathbf{I}_n(010, 101, 120, 201)$ (A279554)	1, 2, 5, 15, 51, 188, 733, 2979, 12495	733
$e_j > e_k, e_i \geq e_k$	Wilf-eq. to 746B (Sec. 3.2, A279555)	1, 2, 5, 15, 51, 189, 746, 3091, 13311	746,A
$e_i \neq e_j \geq e_k, e_i \geq e_k$	Wilf-eq. to 746A (Sec. 3.2, A279555)	1, 2, 5, 15, 51, 189, 746, 3091, 13311	746,B
$e_i \leq e_j \neq e_k, e_i \geq e_k$	$\mathbf{I}_n(010, 110, 120)$ (A279556)	1, 2, 5, 15, 51, 190, 759, 3206, 14180	759
$e_i \leq e_j > e_k, e_i \neq e_k$	counted - See Section 3.1 (A279557)	1, 2, 6, 20, 68, 233, 805, 2807, 9879	805
$e_i \neq e_j > e_k, e_i \geq e_k$	$\mathbf{I}_n(010, 120, 210)$ (A279558)	1, 2, 5, 15, 52, 200, 830, 3654, 16869	830
$e_i < e_j, e_i \geq e_k$	$\mathbf{I}_n(010, 120)$ (A279559)	1, 2, 5, 15, 52, 201, 845, 3801, 18089	845
$e_j > e_k, e_i = e_k$	$\mathbf{I}_n(010)$ (A263779)	1, 2, 5, 15, 53, 215, 979, 4922, 26992	979
$e_i > e_j, e_i \neq e_k$	counted - See Section 3.1 (A279560)	1, 2, 6, 21, 76, 277, 1016, 3756, 13998	1016
$e_i > e_j \neq e_k$	counted - See Section 3.1 (A279561)	1, 2, 6, 21, 77, 287, 1079, 4082, 15522	1079,A
$e_i < e_j > e_k, e_i \neq e_k$	$\mathbf{I}_n(021, 120)$ (A279561)	1, 2, 6, 21, 77, 287, 1079, 4082, 15522	1079,B
$e_i > e_j \leq e_k, e_i \neq e_k$	$\mathbf{I}_n(100, 102, 201)$ (A279562)	1, 2, 6, 21, 78, 299, 1176, 4729, 19378	1176
$e_i > e_j \neq e_k, e_i \neq e_k$	$\mathbf{I}_n(102, 201, 210)$ (A279563)	1, 2, 6, 22, 85, 328, 1253, 4754, 17994	1253
$e_i \geq e_j = e_k$	$\mathbf{I}_n(000, 100)$ (A279564)	1, 2, 5, 16, 60, 260, 1267, 6850, 40572	1267
$e_i > e_k$	$\mathbf{I}_n(100, 110, 120, 210, 201)$ (A279565)	1, 2, 6, 21, 81, 332, 1420, 6266, 28318	1420
$e_i > e_j < e_k, e_i \neq e_k$	$\mathbf{I}_n(102, 201)$ (A279566)	1, 2, 6, 22, 87, 354, 1465, 6154, 26223	1465
$e_j \geq e_k, e_i > e_k$	$\mathbf{I}_n(100, 110, 120, 210)$ (A279567)	1, 2, 6, 21, 82, 343, 1509, 6893, 32419	1509
$e_j \neq e_k, e_i > e_k$	Wilf-eq. to 1833B (Sec. 3.2, A279568)	1, 2, 6, 22, 90, 396, 1833, 8801, 43441	1833,A
$e_i \neq e_j, e_i > e_k$	Wilf-eq. to 1833A (Sec. 3.2, A279568)	1, 2, 6, 22, 90, 396, 1833, 8801, 43441	1833,B
$e_j > e_k, e_i > e_k$	Wilf-eq. to 1953B (Sec. 3.2, A279569)	1, 2, 6, 22, 91, 409, 1953, 9763, 50583	1953,A
$e_i \neq e_j \geq e_k, e_i > e_k$	Wilf-eq. to 1953A (Sec. 3.2, A279569)	1, 2, 6, 22, 91, 409, 1953, 9763, 50583	1953,B
$e_i \leq e_j > e_k, e_i > e_k$	$\mathbf{I}_n(110, 120)$ (A279570)	1, 2, 6, 22, 92, 423, 2091, 10950, 60120	2091
$e_i > e_j \leq e_k, e_i \geq e_k$	$\mathbf{I}_n(100, 101, 201)$ (A279571)	1, 2, 6, 22, 92, 424, 2106, 11102, 61436	2106
$e_i \neq e_j \neq e_k, e_i > e_k$	$\mathbf{I}_n(120, 210, 201)$ (A279572)	1, 2, 6, 23, 101, 484, 2468, 13166, 72630	2468
$e_i \neq e_j > e_k, e_i > e_k$	$\mathbf{I}_n(210, 120)$ (A279573)	1, 2, 6, 23, 102, 499, 2625, 14601, 84847	2625
$e_i < e_j, e_i > e_k$	$\mathbf{I}_n(120)$ (A263778)	1, 2, 6, 23, 103, 515, 2803, 16334, 100700	2803
$e_i > e_j = e_k$	$\mathbf{I}_n(100)$ (A263780)	1, 2, 6, 23, 106, 565, 3399, 22678, 165646	3399
$e_j < e_k, e_i > e_k$	$\mathbf{I}_n(201)$ (A263777)	1, 2, 6, 24, 118, 674, 4306, 29990, 223668	4306,A
$e_i > e_j > e_k$	$\mathbf{I}_n(210)$ Wilf-eq to 4306A [13]	1, 2, 6, 24, 118, 674, 4306, 29990, 223668	4306,B

Table 3: The patterns whose avoidance sequences did not match sequences in the OEIS. (OEIS numbers in parentheses were newly assigned.)

2 Patterns whose sequences appear in the OEIS

2.1 Classes 7(A,B,C,D): n

There are four equivalence classes of patterns whose avoidance sequences are counted by the positive integers. We characterize each, from which it is straightforward to prove that

$$|\mathbf{I}_n(-, <, -)| = |\mathbf{I}_n(-, \geq, -)| = |\mathbf{I}_n(=, -, -)| = |\mathbf{I}_n(\neq, -, -)| = n,$$

although these four patterns are not equivalent. The conditions on the entries of inversion sequences in each class are as follows.

- (Class **7A**: $e_j < e_k$) $\mathbf{I}_n(-, <, -)$ is the set of $e \in \mathbf{I}_n$ satisfying $e_2 \geq e_3 \geq \dots \geq e_n$.
- (Class **7B**: $e_j \geq e_k$) $\mathbf{I}_n(-, \geq, -)$ is the set of $e \in \mathbf{I}_n$ satisfying $e_1 \leq e_2 < e_3 < \dots < e_n$.
- (Class **7C**: $e_i = e_j$) $\mathbf{I}_n(=, -, -)$ is the set of $e \in \mathbf{I}_n$ satisfying $e_1 < e_2 < \dots < e_{n-1}$.
- (Class **7D**: $e_i \neq e_j$) $\mathbf{I}_n(\neq, -, -)$ is the set of $e \in \mathbf{I}_n$ satisfying $e_1 = e_2 = \dots = e_{n-1}$.

Note: Simion and Schmidt [27] showed that $|\mathbf{S}_n(123, 132, 231)| = n$. In fact, we get the following relationship between inversion sequences and permutations.

Theorem 1. For every n , $\Theta(\mathbf{S}_n(123, 132, 231)) = \mathbf{I}_n(=, -, -)$.

2.2 Classes 12(A,B): $2(n - 1)$ for $n > 1$

In the following theorem, we show that classes **12A** ($e_i \neq e_j$ and $e_i \neq e_k$) and **12B** ($e_i \geq e_j$ and $e_i \neq e_k$) are Wilf-equivalent, but not equivalent.

Theorem 2. $|\mathbf{I}_n(\neq, -, \neq)|$ and $|\mathbf{I}_n(\geq, -, \neq)|$ are both counted by 1 if $n = 1$ and by $2(n - 1)$ for $n > 1$. However, $\mathbf{I}_n(\neq, -, \neq) \neq \mathbf{I}_n(\geq, -, \neq)$ for $n > 2$.

Proof. For $n = 1$ this is clear. For $n > 1$ this follows by noting that any $e \in \mathbf{I}_n$ with no $i < j < k$ such that $e_i \neq e_j$ and $e_i \neq e_k$ can be $(0, 0, \dots, 0)$ or can be of the form $(0, 0, \dots, t, 0)$ or $(0, 0, \dots, 0, s)$ where $t \in [n - 2]$ and $s \in [n - 1]$. On the other hand, any $e \in \mathbf{I}_n$ with no $i < j < k$ such that $e_i \geq e_j$ and $e_i \neq e_k$ must have the form $(0, 1, 2, \dots, n - 2, t)$ for $t = 0, \dots, n - 1$ or the form $(0, 1, 2, \dots, t - 1, t, t, \dots, t)$ for $t = 0, \dots, n - 3$. \square

2.3 Class 21: F_{n+1}

Let F_n be the n -th Fibonacci number, where $F_0 = 0$, $F_1 = 1$, and $F_n = F_{n-1} + F_{n-2}$ for $n \geq 2$. The Fibonacci numbers count pattern-avoiding permutations such as $\mathbf{S}_n(123, 132, 213)$ [27]. Class **21** ($e_i = e_j \leq e_k$) is counted by the $(n + 1)$ -th Fibonacci number, as shown below.

Observation 3. The inversion sequences $e \in \mathbf{I}_n$ with no $i < j < k$ such that $e_i = e_j \leq e_k$ are those satisfying, for some $t \in [n]$,

$$e_1 < e_2 < \dots < e_t \geq e_{t+1} > \dots > e_n. \quad (1)$$

Theorem 4. $|\mathbf{I}_n(=, \leq, -)| = F_{n+1}$

Proof. This is clear for $n = 1, 2$. For $n \geq 3$, any $e \in \mathbf{I}_n(=, \leq, -)$ must have the form $(0, e_1 + 1, \dots, e_{n-1} + 1)$ for $(e_1, \dots, e_{n-1}) \in \mathbf{I}_{n-1}(=, \leq, -)$ or $(0, e_1 + 1, \dots, e_{n-2} + 1, 0)$ for $(e_1, \dots, e_{n-2}) \in \mathbf{I}_{n-2}(=, \leq, -)$. Conversely, strings of either of these forms are in $\mathbf{I}_n(=, \leq, -)$. \square

Among the 343 patterns checked, it can be shown that the six patterns whose avoidance sequence is counted by F_{n+1} are equivalent.

Observation 5. All of the following patterns are equivalent to $(=, \leq, -)$: $(=, -, \leq)$, $(=, \leq, \leq)$, $(\geq, -, \leq)$, $(\geq, \leq, -)$, (\geq, \leq, \geq) .

2.4 Classes 22(A,B,C): Lazy caterer sequence, $\binom{n}{2} + 1$

We show that there are three inequivalent patterns that are all counted by the sequence $\binom{n}{2} + 1$, which also counts $\mathbf{S}_n(132, 321)$ [27].

2.4.1 Class 22A: Avoiding $e_i < e_j \neq e_k$

It is not hard to see that inversion sequences avoiding $e_i < e_j \neq e_k$ are characterized by the following.

Observation 6. The inversion sequences $e \in \mathbf{I}_n$ with no $i < j < k$ such that $e_i < e_j \neq e_k$ are those satisfying, for some t where $1 \leq t \leq n$,

$$0 = e_1 = e_2 = \dots = e_{t-1} \leq e_t = e_{t+1} = \dots = e_n. \quad (2)$$

That is, either $e = (0, 0, \dots, 0)$ or, for some $t : 2 \leq t \leq n$ and $j : 1 \leq j \leq t - 1$, e consists of a string of $t - 1$ zeros followed by a string of $n - t + 1$ copies of j . This gives the following.

Theorem 7. $|\mathbf{I}_n(<, \neq, -)| = \binom{n}{2} + 1$.

The sequence whose n th entry is $\binom{n}{2} + 1$ is sequence A000124 in the OEIS, where it is called the *Lazy Caterer* sequence [15] because its n th entry is the maximum number of pieces that can be formed by making $n - 1$ straight cuts in a pizza. This is also the avoidance sequence for certain pairs of permutation patterns, as was shown by Simion and Schmidt [27].

Theorem 8 (Simion-Schmidt [27]). $|\mathbf{S}_n(\alpha, \beta)| = \binom{n}{2} + 1$ for any of the following pairs (α, β) of patterns:

$$(132, 321), (123, 231), (123, 312), (213, 321).$$

We can relate these permutations to the inversion sequences in $\mathbf{I}_n(<, \neq, -)$. Recall the bijection $\Theta(\pi) : \mathbf{S}_n \rightarrow \mathbf{I}_n$ for $\pi = \pi_1 \dots \pi_n \in \mathbf{S}_n$ defined by $\Theta(\pi) = (e_1, e_2, \dots, e_n)$, where $e_i = |\{j \mid j < i \text{ and } e_j > e_i\}|$.

Theorem 9. $\mathbf{I}_n(<, \neq, -) = \Theta(\mathbf{S}_n(213, 321))$.

Proof. Note that $e \in \mathbf{I}_n$ satisfies (2) if and only if $\pi = \Theta^{-1}(e)$ satisfies

$$\pi_1 < \pi_2 < \dots < \pi_t > \pi_{t+1} < \pi_{t+2} < \dots < \pi_n,$$

where $\pi_t, \pi_{t+1}, \dots, \pi_n$ are consecutive integers. Such permutations are precisely the ones that avoid both 213 and 321. \square

The patterns $(<, -, <)$ and $(\geq, \neq, -)$ are Wilf-equivalent to the pattern $(<, \neq, -)$ on inversion sequences, although the three patterns are pairwise inequivalent. This is clear from the following characterizations.

2.4.2 Class 22B: Avoiding $e_i < e_j$ and $e_i < e_k$

Observation 10. The inversion sequences $e \in \mathbf{I}_n$ with no $i < j < k$ such that $e_i < e_j$ and $e_i < e_k$ are those satisfying, for some t where $1 \leq t \leq n$,

$$0 = e_1 = e_2 = \dots = e_{t-1} \leq e_t \geq e_{t+1} = \dots = e_n = 0. \quad (3)$$

2.4.3 Class 22C: Avoiding $e_i \geq e_j \neq e_k$

Observation 11. The inversion sequences $e \in \mathbf{I}_n$ with no $i < j < k$ such that $e_i \geq e_j \neq e_k$ are those satisfying, for some t where $1 \leq t \leq n$,

$$e_1 < e_2 < \dots < e_{t-1} \geq e_t = e_{t+1} = \dots = e_n. \quad (4)$$

2.5 Classes 33(A,B): $F_{n+2} - 1$

We show that **33A**: $(\neq, \leq, -)$ and **33B**: (\geq, \leq, \neq) are inequivalent Wilf-equivalent patterns whose avoidance sequences are counted by $F_{n+2} - 1$.

2.5.1 Class 33A: Avoiding $e_i \neq e_j \leq e_k$

Theorem 12. $|\mathbf{I}_n(\neq, \leq, -)| = F_{n+2} - 1$.

Proof. Observe that the inversion sequences $e \in \mathbf{I}_n$ with no $i < j < k$ such that $e_i \neq e_j \leq e_k$ are those satisfying, for some t where $1 \leq t \leq n + 1$,

$$0 = e_1 = e_2 = \dots = e_{t-1} < e_t > e_{t+1} > \dots > e_n. \quad (5)$$

We can partition the inversion sequences in $\mathbf{I}_n(\neq, \leq, -)$ into three disjoint sets: $\{(0, 0, \dots, 0)\}$, $A = \{e \in \mathbf{I}_n(\neq, \leq, -) \mid e_n \neq 0\}$, and $B = \{e \in \mathbf{I}_n(\neq, \leq, -) \mid e \neq 0, e_n = 0\}$. Any inversion sequence in A can be constructed by taking any $e' = (e_1, e_2, \dots, e_{n-1}) \in \mathbf{I}_{n-1}(\neq, \leq, -)$ and letting t be the index of the first nonzero entry (if there is no nonzero entry, set $e_t = e_{n-1}$). Then we can use the characterization given by (5) to verify that $(0, e_1, e_2, \dots, e_{t-1}, e_t + 1, \dots, e_{n-1} + 1)$ is an element of A .

Any element of B can be constructed by taking some $e'' = (e_1, e_2, \dots, e_{n-2}) \in \mathbf{I}_{n-2}(\neq, \leq, -)$ and letting t be the index of the first nonzero entry (again, if no such entry exists, set $e_t = e_{n-2}$). Then $(0, e_1, \dots, e_{t-1}, e_t + 1, \dots, e_{n-2} + 1, 0)$ is an element of B .

Setting $a_n = |\mathbf{I}_n(\neq, \leq, -)|$, this gives $a_n = a_{n-1} + a_{n-2} + 1$, with initial conditions $a_1 = 1$, $a_2 = 2$. So $a_n = F_{n+2} - 1$. \square

2.5.2 Class 33B: Avoiding $e_i \geq e_j \leq e_k$ and $e_i \neq e_k$

Theorem 13. $|\mathbf{I}_n(\geq, \leq, \neq)| = F_{n+2} - 1$.

Proof. The inversion sequences $e \in \mathbf{I}_n$ with no $i < j < k$ such that $e_i \geq e_j \leq e_k$ and $e_i \neq e_k$ are those satisfying, for some t, s where $1 \leq t \leq s \leq n$,

$$e_1 < e_2 < \dots < e_{t-1} < e_t = e_{t+1} = \dots = e_s > e_{s+1} > \dots > e_n. \quad (6)$$

The following is a bijection mapping $\mathbf{I}_n(\geq, \leq, \neq)$ to $\mathbf{I}_n(\neq, \leq, -)$. For $e \in \mathbf{I}_n(\geq, \leq, \neq)$, let s be the first index, if any, such that $e_s > e_{s+1}$; if e is weakly increasing, set $s = n$. To obtain an element of $\mathbf{I}_n(\neq, \leq, -)$, set $e_i = 0$ for $i = 1, \dots, s - 1$. \square

2.6 Classes 64(A,B,C,D): 2^{n-1}

2.6.1 Class 64A: Avoiding $e_i = e_j < e_k$

Corteel, et al. [13] characterized $\mathbf{I}_n(=, <, -) = \mathbf{I}_n(001)$ as the set of $e \in \mathbf{I}_n$ satisfying, for some $t \in [n]$,

$$e_1 < e_2 < \dots < e_t \geq e_{t+1} \geq e_{t+2} \geq \dots \geq e_n.$$

They showed that $|\mathbf{I}_n(001)| = 2^{n-1}$ by showing that the bijection $\Theta : \mathbf{S}_n \rightarrow \mathbf{I}_n$ restricts to a bijection from $\mathbf{S}_n(132, 231)$ to $\mathbf{I}_n(001)$. Simion and Schmidt [27] showed that permutations avoiding both 132 and 231 are enumerated by 2^{n-1} .

We show that three other patterns are Wilf-equivalent, though inequivalent, to class 64A.

2.6.2 Class 64B: Avoiding $e_i < e_j \leq e_k$

Theorem 14. *The number of $e \in \mathbf{I}_n$ with no $i < j < k$ such that $e_i < e_j \leq e_k$ is 2^{n-1} .*

Proof. First observe that the inversion sequences e with no $i < j < k$ such that $e_i < e_j \leq e_k$ are those whose positive entries form a strictly decreasing sequence.

Let $B_n = \mathbf{I}_n(<, \leq, -)$. Notice that $|B_1| = |\{(0)\}| = 1$; we will show that for $n > 1$, $|B_n| = 2|B_{n-1}|$. Recall that $\sigma_1(e)$ adds 1 to each positive element in e .

An $e \in B_n$ has no “1” if and only if it has the form $0 \cdot \sigma_1(e')$ for some $e' \in B_{n-1}$, so there are $|B_{n-1}|$ such e . An $e \in B_n$ has a “1” if and only if it has the form $e' \cdot 0$ for some $e' \in B_{n-1}$ containing a “1” or the form $e' \cdot 1$ for some $e' \in B_{n-1}$ not containing a “1”, so there are also $|B_{n-1}|$ elements of B_n containing a “1”. \square

2.6.3 Class 64C: Avoiding $e_i < e_j \geq e_k$

Theorem 15. *The number of $e \in \mathbf{I}_n$ with no $i < j < k$ such that $e_i < e_j \geq e_k$ is 2^{n-1} .*

Proof. The inversion sequences avoiding the pattern $e_i < e_j \geq e_k$, where $i < j < k$, are those $e \in \mathbf{I}_n$ satisfying, for some $t \in [n]$,

$$0 = e_1 = e_2 = \cdots = e_t < e_{t+1} < e_{t+2} < \cdots < e_n.$$

Map $e \in \mathbf{I}_n(<, \geq, -)$ to the set consisting of its nonzero values. Clearly this is a bijection from $\mathbf{I}_n(<, \geq, -)$ to $2^{[n-1]}$. \square

In fact, we can show that $\mathbf{I}_n(<, \geq, -)$ is the image under Θ of $\mathbf{S}_n(213, 312)$.

Theorem 16. $\Theta(\mathbf{S}_n(213, 312)) = \mathbf{I}_n(<, \geq, -)$.

Proof. It is straightforward to prove that $\mathbf{S}_n(213, 312)$ consists of the unimodal permutations where

$$\pi_1 < \pi_2 < \cdots < \pi_t = n > \pi_{t+1} > \cdots > \pi_n.$$

The inversion sequences avoiding the pattern $e_i < e_j \geq e_k$, where $i < j < k$, are those $e \in \mathbf{I}_n$ satisfying, for some $t \in [n]$,

$$0 = e_1 = e_2 = \cdots = e_t < e_{t+1} < e_{t+2} < \cdots < e_n.$$

It immediately follows that $\Theta(\mathbf{S}_n(213, 312)) = \mathbf{I}_n(<, \geq, -)$. \square

2.6.4 Class 64D: Avoiding $e_i \leq e_j = e_k$

Theorem 17. *The number of $e \in \mathbf{I}_n$ with no $i < j < k$ such that $e_i \leq e_j = e_k$ is 2^{n-1} .*

Proof. The inversion sequences avoiding the pattern $e_i \leq e_j = e_k$, where $i < j < k$, are those $e \in \mathbf{I}_n$ in which all of the entries e_2, e_3, \dots, e_n are distinct.

Let $D_n = \mathbf{I}_n(\leq, =, -)$. Then $|D_1| = |\{(0)\}| = 1$. We show that for $n > 1$, $|D_n| = 2|D_{n-1}|$. An $e \in D_n$ ends in $n-1$ if and only if $e = e' \cdot (n-1)$ for some $e' \in D_{n-1}$, so there are $|D_{n-1}|$ such e . An $e \in D_n$ ends in $d \neq n-1$ if and only if $e = e' \cdot d$ where $e' \in D_{n-1}$ and d is the unique element in $\{0, 1, \dots, n-2\} \setminus \{e_2, \dots, e_{n-1}\}$, so there are again $|D_{n-1}|$ such e . \square

2.7 Classes 121(A,B,C): Grassmannian permutations, $2^n - n$

Permutations with at most one descent were called *Grassmannian* by Lascoux and Schützenberger [20], who also characterized them in terms of their Lehmer codes. Grassmannian permutations of length n are counted by $2^n - n$ and relate to three equivalence classes of patterns for inversion sequences.

2.7.1 Class 121A: Avoiding $e_i \neq e_j < e_k$

Theorem 18. $|\mathbf{I}_n(\neq, <, -)| = 2^n - n$.

Proof. First observe that those $e \in \mathbf{I}_n$ with no $i < j < k$ such that $e_i \neq e_j < e_k$ are exactly those with at most one ascent.

Using the mapping $\Theta : \mathbf{S}_n \rightarrow \mathbf{I}_n$, recall that π has a descent in a position i if and only if $\Theta(\pi)$ has an ascent in position i . Thus Θ restricts to a bijection from Grassmannian permutations of $[n]$ to $\mathbf{I}_n(\neq, <, -)$. \square

The inversion sequences in $\mathbf{I}_n(\neq, <, -)$ correspond to the Grassmannian Lehmer codes used by Lascoux and Schützenberger [20] via the natural bijection (reversal) between inversion sequences and Lehmer codes.

2.7.2 Class 121B: Avoiding $e_i \neq e_j \neq e_k$ and $e_i \neq e_k$

Theorem 19. $|\mathbf{I}_n(\neq, \neq, \neq)| = 2^n - n$.

Proof. Note that inversion sequences with no $i < j < k$ such that $e_i \neq e_j \neq e_k$ and $e_i \neq e_k$ are those with at most 2 distinct entries; precisely, $|\{e_1, \dots, e_n\}| \leq 2$.

The theorem is clear for $n = 1$. Now consider some $e \in \mathbf{I}_n(\neq, \neq, \neq)$ when $n > 1$. Note that $(e_1, \dots, e_{n-1}) \in \mathbf{I}_{n-1}(\neq, \neq, \neq)$. It follows that either (1) $|\{e_1, \dots, e_{n-1}\}| = 2$, and e_n is one of the two elements occurring in (e_1, \dots, e_{n-1}) ; or (2) $|\{e_1, \dots, e_{n-1}\}| = 1$, and $e_n \in \{0, 1, \dots, n-1\}$. Furthermore, the only inversion sequence in $\mathbf{I}_{n-1}(\neq, \neq, \neq)$ where $|\{e_1, \dots, e_{n-1}\}| = 1$ is the zero inversion sequence. This gives the recurrence $|\mathbf{I}_n(\neq, \neq, \neq)| = 2(|\mathbf{I}_{n-1}(\neq, \neq, \neq)| - 1) + n$ which has the claimed solution. \square

2.7.3 Class 121C: Avoiding $e_j \geq e_k$ and $e_i \neq e_k$

Theorem 20. $|\mathbf{I}_n(-, \geq, \neq)| = 2^n - n$.

Proof. Inversion sequences with no $i < j < k$ such that $e_j \geq e_k$ and $e_i \neq e_k$ are those satisfying

$$e_1 = \dots = e_{i-1} < e_i < \dots < e_n$$

or, if $e_{i+1} = 0$,

$$e_1 = \dots = e_{i-1} < e_i < e_{i+2} < \dots < e_n,$$

for some i with $2 \leq i \leq n+1$.

To count these, for each $t = 1, \dots, n-1$, and for any t -element subset $x_1 < x_2 < \dots < x_t$ of $[n-1]$, associate the length n inversion sequence $(0, 0, \dots, 0, x_1, x_2, \dots, x_t)$ and, unless $\{x_1, \dots, x_t\} = \{n-t, n-t+1, \dots, n-1\}$, also associate the length n inversion sequence $(0, 0, \dots, 0, x_1, 0, x_2, \dots, x_t)$, giving $2^{n-1} + (2^{n-1} - n) = 2^n - n$. \square

2.8 Class 151: 321-avoiding separable permutations

We show that the avoidance sequence for this pattern satisfies the recurrence $a_n = 3a_{n-1} - 2a_{n-2} + a_{n-3}$ with initial conditions $a_1 = 1$, $a_2 = 2$, and $a_3 = 5$. This coincides with sequence A034943 in the OEIS, where, among other things, it is said to count 321-avoiding separable permutations (OEIS entry by Vince Vatter) [15]. A *separable* permutation is one that avoids 2413 and 3142. Moreover, we show that ($\neq, <, \leq$)-avoiding inversion sequences have a simple characterization.

Theorem 21. *Let $A_n = \mathbf{I}_n(\neq, <, \leq)$ and $a_n = |A_n|$. Then $a_n = 3a_{n-1} - 2a_{n-2} + a_{n-3}$ with initial conditions $a_1 = 1$, $a_2 = 2$, and $a_3 = 5$.*

Proof. First, it can be shown that the set of $e \in \mathbf{I}_n$ such that there is no $i < j < k$ for which $e_i \neq e_j < e_k$ and $e_i \leq e_k$ is the set of $e \in \mathbf{I}_n$ where the nonzero elements are weakly decreasing and equal nonzero elements are consecutive. That is, (1) if $e_i < e_j$, then $e_i = 0$ and (2) if $0 < e_i = e_j$ for some $i < j$, then $e_i = e_{i+1} = \dots = e_j$.

Define X_n, Y_n, Z_n by

$$\begin{aligned} X_n &= \{e \in A_n \mid e_i \neq 1, \text{ for all } 1 \leq i \leq n\}, \\ Y_n &= \{e \in A_n \mid e_n = 1\}, \\ Z_n &= \{e \in A_n \mid e_n = 0 \text{ and } e_i = 1 \text{ for some } i < n\}. \end{aligned}$$

Then A_n is the disjoint union $A_n = X_n \cup Y_n \cup Z_n$. Recall that the operator σ_1 adds 1 to the positive elements of an inversion sequence. To get a recurrence, note that $|X_n| = |A_{n-1}| = a_{n-1}$ since $e \in A_{n-1}$ if and only if $0 \cdot \sigma_1(e) \in X_n$. Also, $(e_1, \dots, e_{n-1}, 0) \in Z_n$ if and only if $(e_1, \dots, e_{n-1}) \in Y_{n-1} \cup Z_{n-1} = A_{n-1} - X_{n-1}$; so $|Z_n| = |A_{n-1}| - |X_{n-1}| = a_{n-1} - a_{n-2}$. Finally, $(e_1, \dots, e_{n-1}, 1) \in Y_n$ if and only if $(e_1, \dots, e_{n-1}) \in A_{n-1} - Z_{n-1}$, so $|Y_n| = a_{n-1} - |Z_{n-1}| = a_{n-1} - (a_{n-2} - a_{n-3})$. Putting this together,

$$a_n = |A_n| = |X_n| + |Y_n| + |Z_n| = 3a_{n-1} - 2a_{n-2} + a_{n-3}$$

and the result follows by checking the initial conditions. \square

2.9 Class 185: 321-avoiding vexillary permutations, $2^{n+1} - \binom{n+1}{3} - 2n - 1$

Vexillary permutations, studied by Lascoux and Schützenberger [20], are 2143-avoiding permutations. The 321-avoiding vexillary permutations arose in work of Billey, Jockush and

Stanley [4] on the combinatorics of Schubert polynomials. It was shown that $|\mathbf{S}_n(321, 2143)| = 2^{n+1} - \binom{n+1}{3} - 2n - 1$ which is entry A088921 in the OEIS. In this entry, it is noted that the 321-avoiding vexillary permutations are exactly the Grassmannian permutations (see Section 2.7) and their inverses.

We show that the $(\neq, <, \neq)$ -avoiding inversion sequences are counted by the same function as the 321-avoiding vexillary permutations.

Lemma 22. $\mathbf{I}_n(\neq, <, \neq) = \mathbf{I}_n(\neq, \neq, \neq) \cup \mathbf{I}_n(\neq, <, -)$.

Proof. If $e \in \mathbf{I}_n(\neq, <, \neq)$, then either $e \in \mathbf{I}_n(\neq, <, -)$ or for any $i < j < k$ such that $e_i \neq e_j < e_k$, $e_i = e_k$ and therefore $e \in \mathbf{I}_n(\neq, \neq, \neq)$.

Conversely, if, for some $i < j < k$, $e_i \neq e_j < e_k$ and $e_i \neq e_k$, then e contains both $(\neq, <, -)$ and (\neq, \neq, \neq) . \square

Theorem 23. $|\mathbf{I}_n(\neq, <, \neq)| = 2^{n+1} - \binom{n+1}{3} - 2n - 1$.

Proof. By Theorem 19, $|\mathbf{I}_n(\neq, \neq, \neq)| = 2^n - n$ and by Theorem 18, $|\mathbf{I}_n(\neq, <, -)| = 2^n - n$. From the characterizations of these sets in the proof of Theorems 18 and 19, $\mathbf{I}_n(\neq, \neq, \neq) \cap \mathbf{I}_n(\neq, <, -)$ is the set of inversion sequences with at most one ascent and at most two distinct elements, that is, the set of $e \in \mathbf{I}_n$ satisfying, for some $1 \leq t < a < b \leq n + 1$,

$$0 = e_1 = \dots = e_{a-1}; \quad t = e_a = \dots = e_{b-1}; \quad 0 = e_b = \dots = e_n,$$

which is counted by $\binom{n+1}{3}$, together with $(0, 0, \dots, 0)$. Thus

$$|\mathbf{I}_n(\neq, \neq, \neq) \cap \mathbf{I}_n(\neq, <, -)| = \binom{n+1}{3} + 1$$

and the result follows. \square

2.10 Class 187: Conjectured to be counted by A049125

It appears that the number of $e \in \mathbf{I}_n$ avoiding this pattern is given by A049125 in the OEIS, where it is described by David Callan to be the number of ordered trees with n edges in which every non-leaf non-root vertex has at most one leaf child. However, we have not yet proven it. We can prove a characterization of the avoidance set and, from that, derive a 4-parameter recurrence that allows us to check against A049125 for several terms.

Observation 24. The sequences $e \in \mathbf{I}_n$ having no $i < j < k$ with $e_i \geq e_k$ are those for which $e_i > \max\{e_1, \dots, e_{i-2}\}$ for $i = 3, \dots, n$. For $e \in \mathbf{I}_n$, this is equivalent to the conditions $e_3 > e_1$ and, for $4 \leq i \leq n$, $e_i > \max\{e_{i-2}, e_{i-3}\}$.

2.11 Class 193: $\mathbf{S}_n(132, 4312)$, $(n-1)2^{n-2} + 1$

Sequence $(n-1)2^{n-2} + 1$ appears as A005183 in the OEIS, where Pudwell indicates that it counts $\mathbf{S}_n(132, 4312)$ [15]. We show it also counts $\mathbf{I}_n(\leq, \geq, \neq)$.

Theorem 25. $|\mathbf{I}_n(\leq, \geq, \neq)| = (n-1)2^{n-2}$

Proof. Observe that if $e \in \mathbf{I}_n$ has no $i < j < k$ such that $e_i \leq e_j \geq e_k$ and $e_i \neq e_k$ then e must have the form

$$e = (0, \dots, 0, e_a, 0, \dots, 0, e_{n-b+1}, e_{n-b+2}, \dots, e_n)$$

where $1 \leq a < n+1$ and $b < n-a+2$ and $1 \leq e_a < e_{n-b+1} < e_{n-b+2} < \dots < e_n < n$.

If $e_a > 1$ then $e = 0 \cdot \sigma_1(e')$ for some $e' \in \mathbf{I}_{n-1}(\leq, \geq, \neq)$. Otherwise, $e_a = 1$ and e can be obtained by first choosing a b -element subset of $\{2, \dots, n-1\}$ to place (sorted) in locations $n-b+1, \dots, n$, and then choosing one of the locations $2, \dots, n-b$ to be the location a such that $e_a = 1$. Thus the number of sequences containing a 1 is:

$$\sum_{b=0}^{n-2} \binom{n-2}{b} (n-1-b) = n2^{n-3}.$$

This gives the recurrence

$$|\mathbf{I}_n(\leq, \geq, \neq)| = |\mathbf{I}_{n-1}(\leq, \geq, \neq)| + n2^{n-3},$$

where $|\mathbf{I}_1(\leq, \geq, \neq)| = 1$, whose solution is as claimed in the theorem. \square

2.12 Class 233: $\mathbf{I}_n(012)$, F_{2n-1}

It was shown by Corteel, et al. [13] that the inversion sequences $e \in \mathbf{I}_n(<, <, -) = \mathbf{I}_n(012)$ are those in which the positive elements of e are weakly decreasing. From that characterization, it was proven that

$$|\mathbf{I}_n(<, <, -)| = |\mathbf{I}_n(012)| = F_{2n-1}.$$

The sequence F_{2n-1} also counts the *Boolean permutations*, given by $\mathbf{S}_n(321, 3412)$ [30, 25].

2.13 Class 304: Conjectured to be counted by A229046

We derive a recurrence to count the $(-, -, =)$ -avoiding inversion sequences. This sequence appears to be sequence A229046 in the OEIS. If true, this would give a combinatorial interpretation of A229046 which so far is defined only by a generating function and summation.

Note that $\mathbf{I}_n(-, -, =)$ is the set of $e \in \mathbf{I}_n$ with at most two copies of any entry and any equal entries must be adjacent.

Let $S_{n,k}$ be the set of $e \in \mathbf{I}_n(-, -, =)$ with k distinct elements; that is, $S_{n,k}$ consists of the inversion sequences $e = (e_1, e_2, \dots, e_n) \in \mathbf{I}_n(-, -, =)$ such that $|\{e_1, \dots, e_n\}| = k$. Let $s(n, k) = |S_{n,k}|$.

Theorem 26. for $1 \leq k \leq n$,

$$s(n, k) = (n - 1 + k)s(n - 1, k - 1) + (n - k)s(n - 2, k - 1),$$

with initial conditions $s(1, 1) = s(2, 1) = s(2, 2) = 1$ and otherwise $s(n, k) = 0$ for $k = 1$ or $n \leq 2$.

Proof. Let $A_{n,k}$ be the subset of $S_{n,k}$ consisting of those e in which e_n is unrepeated. Let $B_{n,k} = S_{n,k} \setminus A_{n,k}$. We can extend some $e \in S_{n,k}$ to strings in $S_{n+1,k}$ and $S_{n+1,k+1}$ in the following ways.

If $e \in B_{n,k}$, then $e \cdot n \in A_{n+1,k+1}$. Additionally, if x is one of the $n - k$ values in $\{0, 1, \dots, n - 1\}$ not used in e , then $e \cdot x \in A_{n+1,k+1}$.

If $e \in A_{n,k}$, then $e \cdot n \in A_{n+1,k+1}$. Furthermore, if x is one of the $n - k$ values in $\{0, 1, \dots, n - 1\}$ not used in e , then $e \cdot x \in A_{n+1,k+1}$. Finally, if $e_n = y$, then $e \cdot y \in B_{n+1,k}$. Letting $a(n, k) = |A_{n,k}|$ and $b(n, k) = |B_{n,k}|$, we have

$$\begin{aligned} s(n, k) &= a(n, k) + b(n, k); \\ b(n + 1, k) &= a(n, k); \\ a(n + 1, k + 1) &= (n - k + 1)b(n, k) + (n - k + 1)a(n, k) \\ &= (n - k + 1)s(n, k). \end{aligned}$$

So,

$$\begin{aligned} s(n, k) &= a(n, k) + b(n, k) \\ &= a(n, k) + a(n - 1, k) \\ &= (n - k + 1)s(n - 1, k - 1) + (n - k)s(n - 2, k - 1). \end{aligned}$$

□

Then $|\mathbf{I}_n(-, -, =)| = s(n, 1) + \dots + s(n, n)$. It is an open question to show that this theorem provides a refinement of A229046. Additionally, an interesting question is whether there is a natural description of the set $\Theta^{-1}(\mathbf{I}_n(-, -, =))$.

2.14 Classes 429(A,B,C): Catalan numbers

It is known that for any $\pi \in \mathbf{S}_3$, $|\mathbf{S}_n(\pi)|$ is the Catalan number $C_n = \binom{2n}{n}/(n + 1)$ [22, 27]. There are three inequivalent triples of relations $\rho = (\rho_1, \rho_2, \rho_3) \in \{\geq, \leq, <, >, =, \neq, -\}^3$ such that $|\mathbf{I}_n(\rho)| = C_n$. The first corresponds naturally under $\Theta : \mathbf{S}_n \rightarrow \mathbf{I}_n$ to a pattern $\pi \in \mathbf{S}_3$.

2.14.1 Class 429A: Avoiding $e_j > e_k$

Theorem 27. $\mathbf{I}_n(-, >, -) = \Theta(\mathbf{S}_n(213))$.

Proof. Observe that an $e \in \mathbf{I}_n$ has no $i < j < k$ with $e_j > e_k$ if and only if e is weakly increasing. Similarly, it can be checked that $\pi \in \mathbf{S}_n$ avoids 213 if and only if $\Theta(\pi)$ is weakly increasing. □

2.14.2 Class 429B: Avoiding $e_j \geq e_k$ and $e_i < e_k$

Theorem 28. $|\mathbf{I}_n(-, \geq, <)| = C_n$.

Proof. Observe that some $e \in \mathbf{I}_n$ has no $i < j < k$ with $e_i < e_k$ and $e_j \geq e_k$ if and only if the positive elements of e are strictly increasing.

Let $I(x) = \sum_{n \geq 0} \mathbf{I}_n(-, \geq, <)x^n$. We will show that

$$I(x) = 1 + xI^2(x), \tag{7}$$

which has the solution $\frac{1 - \sqrt{1 - 4x}}{2x}$; recall that this is the generating function for C_n .

Given any $e \in \mathbf{I}_n(-, \geq, <)$, consider the last maximal entry e_t ; this is the largest t such that $e_t = t - 1$. The string $(e_1, e_2, \dots, e_{t-1})$ is an element of $\mathbf{I}_{t-1}(-, \geq, <)$. Additionally, it is straightforward to show that the string $\sigma_{1-t}(e_{t+1}, e_{t+2}, \dots, e_n)$ (where $t - 1$ is subtracted from each positive value) is an element of $\mathbf{I}_{n-t}(-, \geq, <)$. Conversely, any element of $\mathbf{I}_n(-, \geq, <)$ with last maximal entry in position t is of the form $e' \cdot (t - 1) \cdot \sigma_{1-t}(e'')$ where $e' \in \mathbf{I}_{t-1}(-, \geq, <)$ and $e'' \in \mathbf{I}_{n-t}(-, \geq, <)$. This accounts for the “ $xI^2(x)$ ” term of equation 7. Since this construction doesn’t account for the length 0 inversion sequence, we must also add a “1.” \square

Alternatively, it can be checked that the following map from $\mathbf{I}_n(-, >, -)$ to $\mathbf{I}_n(-, \geq, <)$ is a bijection. Send $e \in \mathbf{I}_n(-, >, -)$ to e' , defined by $e'_i = 0$ if $e_i \in \{e_1, \dots, e_{i-1}\}$ and otherwise $e'_i = e_i$.

2.14.3 Class 429C: Avoiding $e_i \geq e_j$ and $e_i \geq e_k$

In [24] we conjectured that $|\mathbf{I}_n(\geq, -, \geq)| = C_n$. This has been proven by Kim and Lin [16] where they also prove the following conjectures from the first version of this paper [24]:

- The number of $e \in \mathbf{I}_n(\geq, -, \geq)$ with $\text{last}(e) = k$ is equal to the number of standard tableaux of shape $(n - 1, k)$ (ballot numbers A009766 in OEIS [15]).
- The number of $e \in \mathbf{I}_n(\geq, -, \geq)$ with $\text{dist}(e) = k$ is equal to the number of $\pi \in \mathbf{S}_n(123)$ with $k - 1$ descents (A166073 in the OEIS [15]). (The number of *distinct elements* is $\text{dist}(e) = |\{e_1, \dots, e_n\}|$.)

2.15 Class 523: $\mathbf{S}_n(\overline{3\bar{1}542})$ and the nexus numbers

In this section we show that inversion sequences avoiding the pattern $(\neq, =, -)$ are equinumerous with permutations avoiding $\overline{3\bar{1}542}$. Note that a permutation π avoids the pattern $\overline{3\bar{1}542}$ if any occurrence of 542 in π is contained in an occurrence of 31542. We do this by proving that the $(\neq, =, -)$ -avoiding inversion sequences with k distinct entries are counted by the nexus numbers, $(n + 1 - k)^k - (n - k)^k$.

Observe that the sequences $e \in \mathbf{I}_n$ with no $i < j < k$ satisfying $e_i \neq e_j = e_k$ are those in which the nonzero elements are distinct and once a nonzero element has occurred, at most

one more 0 can appear in e . We use this characterization to show that $\mathbf{I}_n(\neq, =, -)$ is counted by the sequence A047970 which counts diagonal sums of nexus numbers [15]. This sequence also counts permutations in \mathbf{S}_n avoiding the barred pattern $\bar{3}\bar{1}542$ (conjectured by Pudwell, and proved by Callan [11]).

Let $T_{n,k}$ be the set of $e \in \mathbf{I}_n(\neq, =, -)$ with k distinct elements. We prove the following refinement, which gives a new combinatorial interpretation of the *nexus numbers*, $(n+1-k)^k - (n-k)^k$ (see A047969 in the OEIS [15]).

Theorem 29. *For $1 \leq k \leq n$, $|T_{n,k}| = (n+1-k)^k - (n-k)^k$.*

Proof. We count $T_{n,k}$ directly. When $k = 1$, $|T_{n,k}| = 1$ and the result follows. When $k \geq 2$, any $e \in T_{n,k}$ will contain some e_t such that $0 = e_1 = e_2 = \dots = e_{t-1} < e_t$ and there are no repeated values among e_t, e_{t+1}, \dots, e_n . Therefore, if e has k distinct values, there are two cases: (1) e begins with $n-k+1$ zeros and contains no other zeros; or (2) e begins with $n-k$ zeros and contains one further zero after e_{n-k+1} (which is the first nonzero entry).

For Case (1), the values $e_{n-k+2}, e_{n-k+3}, \dots, e_n$ must all be distinct and nonzero. So, there are $n-k+1$ possibilities for each, giving $(n-k+1)^{k-1}$ inversion sequences.

For Case (2), e_{n-k+1} must be nonzero, so there are $n-k$ choices for this entry. Additionally, each of $e_{n-k+1}, e_{n-k+2}, e_{n-k+3}, \dots, e_n$ must be distinct, though zero could appear after e_{n-k+1} . In total, this gives $(n-k)(n-k+1)^{k-1}$ possible inversion sequences. Finally, we must remove any inversion sequence that does not include a zero among $e_{n-k+2}, e_{n-k+3}, \dots, e_n$; there are $(n-k)^k$ such sequences. As a result, there are $(n-k)(n-k+1)^{k-1} - (n-k)^k$ inversion sequences that are part of Case (2).

Adding Cases (1) and (2), we have $|T_{n,k}| = (n-k+1)^{k-1} + (n-k)(n-k+1)^{k-1} - (n-k)^k = (n-k+1)^k - (n-k)^k$, as desired. \square

2.16 Classes 772(A,B): Set partitions avoiding enhanced 3-crossings

In the first version of this paper [24], we conjectured that the avoidance sets for the patterns $(-, \leq, \geq)$ and $(\geq, \geq, -)$ are counted by A108307. It was shown by Bousquet-Mélou and Xin that A108307 gives the number of set partitions of $[n]$ avoiding enhanced 3-crossings [6].

The following theorem was proven by Lin through use of generating trees and the obstinate kernel method [21] and by Yan through construction of a bijection with 0,1 fillings of Ferrers shapes [31]. Lin's proof makes use of our characterization of $I_n(\geq, \geq, -)$ in Observation 31.

Theorem 30 (Lin [21], Yan [31]). *$|I_n(\geq, \geq, -)|$ is the number set partitions of $[n]$ that avoid enhanced 3-crossings (or 3-nestings).*

It follows that $(\geq, \geq, -)$ is indeed counted by A108307. We can additionally show that there is a bijection that not only proves Wilf-equivalence of the patterns 772A and 772B below, but also preserves a number of statistics. First consider the following characterizations of each class.

Observation 31. The inversion sequences with no $i < j < k$ such that $e_i \geq e_j \geq e_k$ are precisely those that can be partitioned into two increasing subsequences.

Proof. Suppose e has such a partition $e_{a_1} < e_{a_2} < \cdots < e_{a_t}$ and $e_{b_1} < e_{b_2} < \cdots < e_{b_{n-t}}$. If there exists $i < j < k$ such that $e_i \geq e_j \geq e_k$, then no two of i, j, k can both be in $\{a_1, \dots, a_t\}$ or both be in $\{b_1, \dots, b_{n-t}\}$, so e avoids $(\geq, \geq, -)$. Conversely, if e avoids $(\geq, \geq, -)$, let $a = (a_1, \dots, a_t)$ be the sequence of left-to-right maxima of e . Then $e_{a_1} < e_{a_2} < \cdots < e_{a_t}$. Consider $i, j \notin \{a_1, \dots, a_t\}$ where $i < j$. The fact that e_i is not a left-to-right maxima implies there exists some e_s such that $s < i$ and $e_s \geq e_i$. Thus to avoid $(\geq, \geq, -)$, we must have $e_i < e_j$. \square

Observation 32. Let $(e_1, e_2, \dots, e_n) \in \mathbf{I}_n$. Additionally, for any $i \in [n]$, let $M_i = \max(e_1, e_2, \dots, e_{i-1})$. Then $e \in \mathbf{I}_n(-, \leq, \geq)$ if and only if for every $i \in [n]$, the entry e_i is a left-to-right maximum, or for every j where $i < j$, we have $e_i > e_j$ or $M_i < e_j$.

Proof. Let $e \in \mathbf{I}_n$ satisfy the conditions of Observation 32 and, to obtain a contradiction, assume there exist $i < j < k$ such that $e_j \leq e_k$ and $e_i \geq e_k$ (that is $e_j \leq e_k \leq e_i$). Notice that $M_j = \max\{e_1, e_2, \dots, e_{j-1}\} \geq e_i$. It follows that $M_j \geq e_k \geq e_j$, which contradicts our assumption.

Conversely, if $(e_1, e_2, \dots, e_n) \in \mathbf{I}_n(-, \leq, \geq)$, consider any e_i . If e_i is not a left-to-right maximum, then there exists some maximum value $M_i = e_s$ such that $s < i$ and $e_s \geq e_i$. Therefore, in order to avoid a 201 pattern, any e_j where $j > i$ must have $e_i > e_j$ or $e_j > M_i = e_s$. \square

Theorem 33. For $n \geq 1$, $|\mathbf{I}_n(\geq, \geq, -)| = |\mathbf{I}_n(-, \leq, \geq)|$.

Proof. We exhibit a bijection based on the characterizations in Observations 31 and 32.

Given $e \in \mathbf{I}_n(\geq, \geq, -)$, define $f \in \mathbf{I}_n(-, \leq, \geq)$ as follows. Let $e_{a_1} < e_{a_2} < \cdots < e_{a_t}$ be the sequence of left-to-right maxima of e and let $e_{b_1} < e_{b_2} < \cdots < e_{b_{n-t}}$ be the subsequence of remaining elements of e .

For $i = 1, \dots, t$, set $f_{a_i} = e_{a_i}$. For each $j = 1, 2, \dots, n - t$, we extract an element of the multiset $B = \{e_{b_1}, e_{b_2}, \dots, e_{b_{n-t}}\}$ and assign it to $f_{b_1}, f_{b_2}, \dots, f_{b_{n-t}}$ as follows:

$$f_{b_j} = \max\{k \mid k \in B - \{f_{b_1}, f_{b_2}, \dots, f_{b_{j-1}}\} \text{ and } k < \max(e_1, \dots, e_{b_{j-1}})\}.$$

By definition, f satisfies the characterization property in Observation 32 of $\mathbf{I}_n(-, \leq, \geq)$. \square

The bijection in Theorem 33 preserves a number of statistics, as is shown below.

Corollary 34. All of the following statistics have the same distribution over $\mathbf{I}_n(-, \leq, \geq)$ and $\mathbf{I}_n(\geq, \geq, -)$.

- the number of locations i such that $e_i = i - 1$;
- the largest entry of e ;

- the number of zeros of e (there can be at most two in either class);
- the number of distinct elements of e (and therefore the number of repeats in e);
- the number of left-to-right maxima of e .

An interesting future direction would be to relate these statistics on the classes 772A and 772B of inversion sequences to corresponding statistics on set partitions avoiding enhanced 3-crossings.

2.17 Classes 877(A,B,C,D): Bell numbers and Stirling numbers

The Bell number B_n is the number of partitions of the set $[n]$ into nonempty blocks. The Stirling number of the second kind, $S_{n,k}$ is the number of partitions of $[n]$ into k blocks.

Among the triples of relations under consideration in this paper, four equivalence classes of patterns have avoidance sets that appear to be counted by the Bell numbers. We have shown this to be true for the classes 877A and 877C, whose inversion sequences have a similar character. We have not confirmed this for the classes 877B and 877D, nor have we confirmed that 877B and 877D are Wilf-equivalent, even though our experiments show that there is likely a bijection that preserves several statistics.

It is interesting to note that $B(n)$ counts permutations avoiding $4\bar{1}32$ and several other barred patterns of length 4, as shown by Callan [9]. Can any of these be related to one of the four patterns 877(A,B,C,D)?

2.17.1 Class 877A: Avoiding $e_i < e_j = e_k$

These are the 011-avoiding sequences. It was observed by Corteel, et al. [13] that these are the $e \in \mathbf{I}_n$ in which the positive elements of e are distinct, and that the 011-avoiding sequences in \mathbf{I}_n with k zeros are counted by the Stirling number of the second kind, $S_{n,k}$. (This also appears in Stanley's *Enumerative Combinatorics, Vol. 1* [28].) Thus $\mathbf{I}_n(011)$ is counted by the Bell numbers.

2.17.2 Class 877B: Avoiding $e_i = e_j \geq e_k$

The set $\mathbf{I}_n(=, \geq, -)$ consists of all $e \in \mathbf{I}_n$ such that no element appears more than twice and if an element, x , is repeated, all elements following the second occurrence of x must be larger than x .

From our calculations, it appears that $\mathbf{I}_n(=, \geq, -)$ is counted by the Bell numbers and, in fact, that the number of $e \in \mathbf{I}_n(=, \geq, -)$ with k repeats is given by A124323, the number of set partitions of $[n]$ with k blocks of size larger than 1, but we have not proven this.

2.17.3 Class 877C: Avoiding $e_i \neq e_j \neq e_k$ and $e_i = e_k$

Observe that these are the $e \in \mathbf{I}_n$ in which only adjacent elements of e can be equal.

Theorem 35. *The number of $e \in \mathbf{I}_n$ in which only adjacent elements of e can be equal is B_n , the n th Bell number.*

Proof. It can be checked that the following map from $\mathbf{I}_n(\neq, \neq, =)$ to $\mathbf{I}_n(011)$ is a bijection. Send $e \in \mathbf{I}_n(\neq, \neq, =)$ to e' , defined by $e'_i = 0$ if $e_i \in \{e_1, \dots, e_{i-1}\}$ and otherwise $e'_i = e_i$. \square

2.17.4 Class 877D: Avoiding $e_i \geq e_j$ and $e_i = e_k$

$\mathbf{I}_n(\geq, -, =)$ is the set of $e \in \mathbf{I}_n$ such that no element appears more than twice and if an element x is repeated, all elements between the two occurrences of x must be larger than x (Note the similarity to 877B).

From our calculations, it appears that $\mathbf{I}_n(\geq, -, =)$ is also counted by the Bell numbers. Moreover, it appears that all of the following statistics are equally distributed over the classes 877B and 877D:

- the number of locations i such that $e_i = i - 1$;
- the largest entry of e ;
- the number of zeros of e (there can be at most two in either class);
- the number of distinct elements of e (and therefore the number of repeats in e , which appears to be A124323).

2.18 Class 924: Central binomial coefficients

Theorem 36. $|\mathbf{I}_n(>, -, -)| = \binom{2n-2}{n-1}$.

Proof. $\mathbf{I}_n(>, -, -)$ is the set of $e \in \mathbf{I}_n$ with $e_1 \leq \dots \leq e_{n-1}$ (counted by the Catalan number C_{n-1} as shown in Section 2.14) and with e_n chosen arbitrarily from $\{0, \dots, n-1\}$. Thus

$$|\mathbf{I}_n(>, -, -)| = nC_{n-1} = n \left(\frac{1}{n} \binom{2n-2}{n-1} \right) = \binom{2n-2}{n-1}.$$

\square

2.19 Class 1064: Conjectured to be counted by A071356

Class 1064 are those inversion sequences with no $i < j < k$ such that $e_i > e_j \leq e_k$. These are the inversion sequences $e \in \mathbf{I}_n$ satisfying, for some t such that $1 < t \leq n$,

$$e_1 \leq \dots \leq e_t > e_{t+1} > \dots > e_n.$$

Our experiments suggest that these are counted by A071356 in the OEIS, which Emeric Deutsch notes counts the number of underdiagonal lattice paths from $(0, 0)$ to the line $x = n$ using only steps $R = (1, 0)$, $V = (0, 1)$, and $D = (1, 2)$ [15].

It also appears from our experiments that the distribution of the number of distinct elements of e is symmetric and unimodal on $\mathbf{I}_n(<, \leq, -)$. The number of $e \in \mathbf{I}_n(<, \leq, -)$ with $\text{dist}(e) = k$ is given in the table below for $n = 1, \dots, 7$.

1						
1	1					
1	4	1				
1	9	9	1			
1	16	38	16	1		
1	25	110	110	25	1	
1	36	255	480	255	36	1

If these observations are true in general, this provides a new simple combinatorial interpretation for A071356 with a natural refinement via a symmetric statistic.

2.20 Class 1265: $\mathbf{S}_n(2143, 3142, 4132)$

Observe that $\mathbf{I}_n(>, <, -)$ is the set of $e \in \mathbf{I}_n$ satisfying, for some t with $1 < t \leq n$,

$$e_1 \leq e_2 \leq \dots \leq e_t > e_{t+1} \geq \dots \geq e_n.$$

Our experiments suggested that $\mathbf{I}_n(>, <, -)$ is counted by A033321, which counts $\mathbf{S}_n(2143, 3142, 4132)$, as well as permutations avoiding several other triples of 4-permutations. Burstein and Stromquist confirmed this by recognizing a natural bijection between $\mathbf{S}_n(2143, 3142, 4132)$ and $\mathbf{I}_n(>, <, -)$ [8]. Their theorem is as follows. Recall from Section 1 that $\text{invcode} : \mathbf{S}_n \rightarrow \mathbf{I}_n$ is the reverse of the Lehmer code.

Theorem 37 (Burstein, Stromquist [8]). *For $n \geq 1$, $\text{invcode}(\mathbf{S}_n(2143, 3142, 4132)) = \mathbf{I}_n(>, <, -)$.*

From Section 1, $\text{invcode}(\pi) = e$ if and only if $e = \Theta((\pi^C)^R)$, giving the following.

Corollary 38. $\Theta(\mathbf{S}_n(2143, 3142, 3241)) = \mathbf{I}_n(>, <, -)$.

2.21 Class 1347: Conjectured to be counted by $\mathbf{S}_n(4123, 4132, 4213)$

Our calculations suggest that $\mathbf{I}_n(>, -, \leq)$ is counted by A106228 in the OEIS, which was recently shown to count $\mathbf{S}_n(4123, 4132, 4213)$ by Albert, Homberger, Pantone, Shar and Vatter [3]. We have not been able to confirm that our avoidance sequence is A106228.

2.22 Class 1385: $\mathbf{I}_n(000)$ and the Euler up/down numbers

$\mathbf{I}_n(=, =, -)$ is the set of inversion sequences avoiding the pattern “000”. It was shown by Corteel, et al. [13] that $|\mathbf{I}_n(000)| = E_{n+1}$, where E_n is the Euler up/down number which counts the number of $\pi \in \mathbf{S}_n$ such that $\pi_1 < \pi_2 > \pi_3 < \pi_4 > \dots$. The proof was via a bijection with n -vertex 0-1-2 increasing trees, which are also counted by E_{n+1} .

Another family of permutations counted by E_{n+1} is the number of simsun permutations of $[n]$, introduced by Simion and Sundaram [29]. A simsun permutation is one with no double descents, even after the removal of the elements $\{n, n-1, \dots, k\}$ for any k . It was shown that the number of $e \in \mathbf{I}_n(000)$ with $n-k$ distinct elements is the number of simsun permutations of n with k descents [13]. The method of proof was to show that they satisfy the same recurrence. Recently, Kim and Lin proved bijectively that the joint distribution of $\text{asc}(\pi)$ and $\text{last}(\pi)$ over simsun permutations π of $[n]$ is the same as the joint distribution of $\text{dist}(e)$ and $\text{last}(e) + 1$ for $e \in \mathbf{I}_n(000)$.

An interesting question is whether there is a natural bijection between $\mathbf{I}_n(000)$ and up-down (or down-up) permutations of $[n+1]$. For example, our calculations suggest that the number of $e \in \mathbf{I}_n(000)$ with $e_n = k-1$ is the number of down-up permutations π of $[n+1]$ with $\pi_1 = k+1$.

2.23 Class 1694: $\mathbf{I}_n(102)$

It was suggested by Corteel, et al. [13] that $\mathbf{I}_n(>, -, <)$ is counted by A200753 in the OEIS [15], a sequence defined by the generating function

$$A(x) = 1 + (x - x^2)(A(x))^3. \quad (8)$$

This was confirmed by Mansour and Shattuck [23] who derived an explicit formula for $|\mathbf{I}_n(102)|$.

Theorem 39 (Mansour-Shattuck [23]). *The generating function $\sum_{n \geq 0} |\mathbf{I}_n(102)|x^n$ satisfies (8).*

It would be interesting to find a direct combinatorial argument.

2.24 1806(A,B,C,D): large Schröder numbers

The *large Schröder number* R_n is the number of Schröder n -paths; that is, the number of paths in the plane from $(0, 0)$ to $(2n, 0)$ never venturing below the x -axis, and using only the steps $(1, 1)$ (up), $(1, -1)$ (down) and $(2, 0)$ (flat).

In the area of pattern avoiding permutations, R_{n-1} counts the separable permutations $\mathbf{S}_n(2413, 3142)$, as well as $\mathbf{S}_n(\alpha, \beta)$ for many other pairs (α, β) of patterns of length 4 [19]. We have four inequivalent triples of relations whose avoidance sets are counted by the large Schröder numbers, two of which (1806B and 1806D) correspond in natural ways to a pair of patterns of length 4.

2.24.1 Class 1806A: Avoiding $e_j > e_k$ and $e_i < e_k$

These are the sequences avoiding 021. It was shown by Corteel, et al. [13] that $e \in \mathbf{I}_n$ avoids 021 if and only if its positive entries are weakly increasing. It was also proven in [13] that $|\mathbf{I}_n(021)| = R_{n-1}$.

The following refinements were shown:

- The number of 021-avoiding inversion sequences e in \mathbf{I}_n with k positions i such that $e_i = i - 1$ is equal to the number of Schröder $(n - 1)$ -paths with $k - 1$ initial up steps.
- The number of 021-avoiding inversion sequences e in \mathbf{I}_n with k zeros is equal to the number of Schröder $(n - 1)$ -paths with $k - 1$ peaks (or $k - 1$ flat steps).

It was also shown that the ascent polynomial for $\mathbf{I}_n(021)$ is palindromic and corresponds to sequence A175124 in the OEIS.

2.24.2 Class 1806B: Avoiding $e_i > e_j$ and $e_i \geq e_k$

It is known that $\mathbf{S}_n(2134, 2143)$ is counted by R_{n-1} [19]. This is a member of “Class VI” in Kitaev’s book [17]; we use this fact to count $\mathbf{I}_n(>, -, \geq)$.

Theorem 40. $|\mathbf{I}_n(>, -, \geq)| = R_{n-1}$.

Proof. We show that $\Theta(\mathbf{S}_n(2134, 2143)) = \mathbf{I}_n(>, -, \geq)$.

Let $e \in \mathbf{I}_n$ satisfy $e_i > e_j$ and $e_i \geq e_k$ for some $i < j < k$. Let $\pi = \Theta^{-1}(e)$. Then $\min\{\pi_j, \pi_k\} > \pi_i$ and, since $e_i > e_j$, there must exist $a < i$ such that both $\pi_a > \pi_i$ and $\min\{\pi_j, \pi_k\} > \pi_a$. Thus $\pi_a \pi_i \pi_j \pi_k$ forms either a 2134 or a 2143.

Conversely, suppose, for some $\pi \in \mathbf{S}_n$, that $\pi_a \pi_i \pi_j \pi_k$ is one of the patterns 2134 or 2143 and let $e = \Theta(\pi)$. Let j' be the smallest index larger than i for which $\pi_{j'} > \pi_a$. Then $\pi_{i+1}, \dots, \pi_{j'-1}$ are all smaller than $\pi_a < \pi_{j'}$ and so $e_i > e_{j'}$. Let k' be the smallest index larger than j' such that $\pi_{k'} > \pi_a$. Then, with the possible exception of $\pi_{j'}$, all of $\pi_{i+1}, \dots, \pi_{k'-1}$ are smaller than $\pi_{k'}$ (since these entries are necessarily smaller than π_a). In addition, since $\pi_i < \pi_a < \pi_{k'}$, we have $e_i \geq e_{k'}$. Thus e has the pattern $(>, -, \geq)$. \square

From our calculations, it appears that the ascent polynomial for $\mathbf{I}_n(>, -, \geq)$ is the same as that for 1806A, which was palindromic. This has been confirmed by Kim and Lin [16]. Since Θ sends descents to ascents, this implies that the descent polynomial for $\mathbf{S}_n(2134, 2143)$ is palindromic. This is not true in general for permutations avoiding pairs of patterns of length 4, even those counted by the large Schröder numbers. For example, it is not true of $\mathbf{S}_n(1234, 2134)$ or $\mathbf{S}_n(1324, 2314)$. However Fu, Lin, and Zeng have recently shown that the descent polynomial for the separable permutations $\mathbf{S}_n(2413, 3142)$ is γ -positive and therefore palindromic [14].

2.24.3 Class 1806C: Avoiding $e_i \geq e_j$ and $e_i > e_k$

Theorem 41. $|\mathbf{I}_n(\geq, -, >)| = R_{n-1}$.

Proof. We will construct a generating function for $\mathbf{I}_n(\geq, -, >)$.

Let $E(x) = \sum_{i=1}^{\infty} |\mathbf{I}_n(\geq, -, >)| x^n$. We will show that $E(x)$ satisfies

$$E(x) = x + xE(x) + E^2(x),$$

whose solution is

$$E(x) = \frac{1 - x - \sqrt{x^2 - 6x + 1}}{2}.$$

This implies that $|\mathbf{I}_n(\geq, -, >)|$ is the $(n - 1)$ th large Schröder number.

Let $e = e_1 e_2 \dots e_n \in \mathbf{I}_n(\geq, -, >)$. Let e_t be the latest maximal entry of e ; that is, $\max\{i \mid e_i = i - 1\}$. If $t = 1$, then either $e = (0)$ or $e = (0, f_1, f_2, \dots, f_{n-1})$ for some $(f_1, f_2, \dots, f_{n-1}) \in \mathbf{I}_{n-1}(\geq, -, >)$.

Now consider the case where $t > 1$. Notice that $e_{t+1} \leq t - 1$ since e_{t+1} cannot be maximal. This implies that $(e_1, e_2, \dots, e_{t-1}, e_{t+1})$ is an inversion sequence of length t . Furthermore, it is straightforward to show that $(e_1, e_2, \dots, e_{t-1}, e_{t+1}) \in \mathbf{I}_n(\geq, -, >)$.

Additionally, for all e_j where $j > t + 1$, we must have $e_j \geq t - 1$; if this is not the case, and there exists some j where $e_j < t - 1$, then we have $t < t + 1 < j$ where $e_t \geq e_{t+1}$ and $e_t > e_j$. Therefore $(e_t - t + 1, e_{t+2} - t + 1, \dots, e_n - t + 1) \in \mathbf{I}_{n-t}(\geq, -, >)$.

Conversely, for any sequences $(e_1, e_2, \dots, e_t) \in \mathbf{I}_t(\geq, -, >)$ and $(f_1, f_2, \dots, f_{n-t}) \in \mathbf{I}_{n-t}(\geq, -, >)$, we can construct the inversion sequence

$$(e_1, e_2, \dots, e_{t-1}, f_1 + t - 1, e_t, f_2 + t - 1, f_3 + t - 1, \dots, f_{n-t} + t - 1).$$

It is straightforward to show that this inversion sequence is in $\mathbf{I}_n(\geq, -, >)$ and the last maximal entry is in the t -th position. \square

2.24.4 Class 1806D: Avoiding $e_i \geq e_j \neq e_k$ and $e_i \geq e_k$

Due to a result of Kremer [19], we know that $|\mathbf{S}_n(4321, 4312)| = R_{n-1}$. We can prove that $|\mathbf{I}_n(\geq, \neq, \geq)|$ is also enumerated by R_{n-1} by constructing a bijection between \mathbf{S}_n and \mathbf{I}_n that restricts to a bijection between $\mathbf{S}_n(4321, 4312)$ and $\mathbf{I}_n(\geq, \neq, \geq)$. This bijection is useful for another class of inversion sequences: it will be used later to prove results related to $\mathbf{I}_n(>, \neq, >)$ (classified as 3720).

Definition 42. Let $\pi \in S_n$ and define $\phi(\pi) = e_1 e_2 \dots e_n \in \mathbf{I}_n$ as follows, starting with e_n and defining entries in reverse order.

1. $e_n = \pi_n - 1$
2. For $1 \leq i < n$,

- (a) if $\pi_i \leq i$, then $e_i = \pi_i - 1$.
- (b) otherwise, if π_i is the k -th largest element of $\{\pi_1, \dots, \pi_i\}$ then e_i is the k -th smallest element of the set $\{e_{i+1}, \dots, e_n\}$.

Lemma 43. For $\pi \in \mathbf{S}_n$, $\phi(\pi) \in \mathbf{I}_n$.

Proof. To show that $\phi(\pi) \in \mathbf{I}_n$, we need to prove that $0 \leq e_i \leq i - 1$ for every $i \in [n]$. We will use an inductive argument, starting with e_n , to show this. We defined $e_n = \pi_n - 1$; since $1 \leq \pi_n \leq n$, it follows that $0 \leq e_n \leq n - 1$, as desired. Now consider e_i and assume that for all e_j among $e_{i+1}e_{i+2}\dots e_n$, $0 \leq e_j \leq j - 1$. If $\pi_i \leq i$, then $e_i = \pi_i - 1$ and it immediately follows that $0 \leq e_i \leq i - 1$.

If instead $\pi_i > i$, assume that π_i is in the k -th largest element of $\{\pi_1, \dots, \pi_i\}$. Notice that each value of $\{e_{i+1}, \dots, e_n\}$ corresponds to an entry π_j where $i < j$ and $\pi_j \leq j$ (any entry π_j where $\pi_j > j$ will repeat a value). So, there are $n - \pi_i - k + 1$ entries π_j such that $i < j$ and $\pi_j > \pi_i$; in turn, this implies that there are $(n - i) - (n - \pi_i - k + 1) = \pi_i - (i + 1) + k$ entries π_j such that $i < j$ and $\pi_j < \pi_i$. At a maximum, $\pi_i - (i + 1)$ of these entries are greater than i ; this leaves k entries occurring after π_i that are less than or equal to i . Each of these entries corresponds to a value in $\{e_{i+1}, \dots, e_n\}$ that is less than or equal to $i - 1$. It follows, that $e_i \leq i - 1$, as desired. \square

Lemma 44. $\phi : \mathbf{S}_n \rightarrow \mathbf{I}_n$ is a bijection.

Proof. Let $e = e_1e_2\dots e_n \in \mathbf{I}_n$. We can define the inverse image of e , $\phi^{-1}(e) = \pi_1\pi_2\dots\pi_n$ in reverse order, starting with π_n so that $\pi_n = e_n + 1$. For $1 \leq i < n$, if $e_i \neq e_j$ for all j where $i < j \leq n$, then $\pi_i = e_i + 1$; otherwise, if e_i is the k -th smallest value of $\{e_{i+1}, \dots, e_n\}$, π_i is the k -th largest value of $[n]$ that does not appear among π_{i+1}, \dots, π_n . \square

It is interesting to note that for any $\pi \in \mathbf{S}_n$, $\text{exc}(\pi) = \text{repeats}(\phi(\pi))$, where $\text{exc}(\pi)$ is the number of positions i such that $\pi_i > i$.

Now, we show that ϕ restricts to a bijection between $\mathbf{S}_n(4321, 4312)$ and $\mathbf{I}_n(\geq, \neq, \geq)$ by proving the following:

Theorem 45. $\phi(\mathbf{S}_n(4321, 4312)) = \mathbf{I}_n(\geq, \neq, \geq)$

Proof. Consider some $\pi \in \mathbf{S}_n$ that contains an occurrence of 4321 or 4312. So, there exists some $a < i < j < k$ such that $\pi_a\pi_i\pi_j\pi_k$ form a 4321 or 4312 pattern. We will show that there exists an occurrence of the pattern (\geq, \neq, \geq) in $\phi(\pi) = e$.

We must consider two cases. If $\pi_i \leq i$, then $j > \pi_i > \pi_j$ and $k > \pi_i > \pi_k$. Therefore, $e_i = \pi_i - 1$, $e_j = \pi_j - 1$ and $e_k = \pi_k - 1$. So, since $\pi_i > \max\{\pi_j, \pi_k\}$ and $\pi_j \neq \pi_k$, e_i, e_j, e_k forms an occurrence of (\geq, \neq, \geq) .

Now assume $\pi_i > i$. Recall that e_i is the t -th smallest element of $\{e_{i+1}, \dots, e_n\}$ if π_i is the t -th largest element of $\{\pi_1, \pi_2, \dots, \pi_i\}$. Since π_a is larger than and occurs before π_i , we know that t is at least 2. So, if $e_{j'}$ and $e_{k'}$, where $j' < k'$, are the two smallest distinct values

in the set $\{e_{i+1}, \dots, e_n\}$, we are guaranteed that $e_i \geq e_{j'}$ and $e_i \geq e_{k'}$; so, $e_i, e_{j'}, e_{k'}$ form the pattern (\geq, \neq, \geq) .

For the converse, let $\pi \in \mathbf{S}_n$ such that $e = \phi(\pi)$ has indices $i < j < k$ such $e_i \geq e_j$, $e_i \geq e_k$, and $e_i \neq e_k$. We show that π contains 4321 or 4312.

If $e_i \in \{e_{i+1}, \dots, e_n\}$ then, by definition of ϕ , $\pi_i > i$ and e_i is the k th smallest element of $\{e_{i+1}, \dots, e_n\}$ for some $k \geq 2$, noting that the distinct elements e_j and e_k are both at most e_i . Then, again by definition of ϕ , π_i is the k th largest element of $\{\pi_1, \dots, \pi_i\}$, so there is some $a < i$ with $\pi_a > \pi_i$. To show that π contains one of the patterns 4321 or 4312, it remains to show there are at least two elements of $\{\pi_{i+1}, \dots, \pi_n\}$ that are smaller than π_i . The number of elements in π that are larger than π_i is $n - \pi_i < n - i$. At least one of these is to the left of π_i in π . Thus at most $n - i - 2$ can be in the $n - i$ positions to the right of π_i .

If $e_i \notin \{e_{i+1}, \dots, e_n\}$ then, by definition of ϕ , $\pi_i \leq i$ and $e_i = \pi_i - 1$. Since $e_j \leq e_i \leq i < j$ and $e_k \leq e_i \leq i < k$, we have $e_j = \pi_j - 1$ and $e_k = \pi_k - 1$, so that $\pi_i \pi_j \pi_k$ has the pattern 321 or 312. To show that π contains 4321 or 4312, it remains to show there is an $a < i$ such that $\pi_a > \pi_i$. Note that there are $n - \pi_i \geq n - i$ elements of π larger than π_i . The $n - i$ positions to the right of π_i hold at least two elements smaller than π_i . So there must be a larger element to the left of π_i , that is, an $a < i$ such that $\pi_a > \pi_i$. □

Our calculations suggested that the ascent polynomial for the inversion sequences in $\mathbf{I}_n(\geq, \neq, \geq)$ is the same as the (symmetric) ascent polynomial for 1806A. This has been proven by Kim and Lin in [16]. It also appears that the number of these inversion sequences with k “repeats” is counted by A090981, the number of Schröder paths with k ascents.

2.25 Class 2074: Baxter numbers

In the earlier version of this paper [24], it was conjectured that the avoidance sequence for the pattern $(\geq, \geq, >)$ is A001181 in the OEIS. This sequence counts the Baxter permutations, which is a result of Chung, et al. in [12]. A *Baxter permutation* π is one that avoids the vincular patterns $\underline{3142}$ and $\underline{2413}$; that is, there is no $i < j < k$ such that $\pi_j < \pi_k < \pi_i < \pi_{j+1}$ or $\pi_j > \pi_k > \pi_i > \pi_{j+1}$.

Kin and Lin prove this via the so-called obstinate kernel method [16]:

Theorem 46 ([16]). $|I_n(\geq, \geq, >)|$ is the number of Baxter permutations of $[n]$.

Note that the Baxter permutations contain the separable permutations $\mathbf{S}_n(3142, 2413)$ which are counted by the large Schröder numbers. Similarly, $\mathbf{I}_n(\geq, \geq, >)$ contains the inversion sequences $\mathbf{I}_n(\geq, -, >)$ (which define class 1806C) which are also counted by the large Schröder numbers. It would be nice to find a bijection between $(\geq, \geq, >)$ -avoiding inversion sequences and Baxter permutations that restricts to a bijection between $(\geq, -, >)$ -avoiding inversion sequences and separable permutations.

2.26 Classes 2549(A,B,C): Conjectured to be counted by $S_n(4231, 42513)$

Albert, et al. [1] showed that

$$|S_n(4231, 42513)| = \sum_{i=0}^n \frac{n-i}{2i+n} \binom{2i+n}{i},$$

which is sequence A098746 in the OEIS [15]. It appears from our calculations that the inequivalent classes $\mathbf{I}_n(>, -, >)$, $\mathbf{I}_n(>, \neq, \geq)$, and $\mathbf{I}_n(\geq, \neq, >)$ have the same avoidance sequence A098746. We will show at least that the classes 2549A ($\mathbf{I}_n(>, -, >)$) and 2549C ($\mathbf{I}_n(\geq, \neq, >)$) are Wilf-equivalent.

Theorem 47. *The patterns $(>, -, >)$ and $(\geq, \neq, >)$, defining classes 2549A and 2549C respectively, are Wilf-equivalent.*

The proof relies on Theorem 53 in the next section and will be given there.

2.27 Classes 2958(A,B,C,D): Plane permutations and the semi-Baxter numbers

In an earlier version of this paper [24], we conjectured that all four pattern equivalence classes 2958(A,B,C,D) have avoidance sets equinumerous with the *plane permutations*, $S_n(21\bar{3}54)$. These are permutations in which every occurrence of the pattern 2154 is contained in an occurrence of 21354. We showed that the classes 2958(B,C,D) are Wilf-equivalent (Theorems 53 and 54 below); additionally, we provided a characterization of the inversion sequences avoiding 2958B (Observation 50 below) and proved a recurrence for the avoidance sequence of 2958B.

We can now make use of a recent paper of Bouvel, Guerrini, Rechnitzer, and Rinaldi [7] to confirm our conjecture.

Bouvel, et al. coined the term “semi-Baxter” for the sequence of integers that enumerates the class of permutations avoiding the vincular pattern $\underline{2413}$ (an occurrence of $\underline{2413}$ is an occurrence of 2413 where the “4” and “1” occur consecutively) [7]. They derived a functional equation and used it to get a closed-form formula for the semi-Baxter sequence.

Moreover, Bouvel, et al. showed that semi-Baxter permutations are equinumerous with plane permutations: they used generating trees and proved that both classes of permutations are generated by the same succession rule Ω_{semi} . Therefore, their formula for semi-Baxter numbers also enumerates plane permutations, which fulfills a challenge posed by Bousquet-Mélou and Butler [5].

Then, using our characterization of the inversion sequences avoiding 2958B, Bouvel, et al. showed that those inversion sequences are also generated by the succession rule Ω_{semi} and are therefore equinumerous with plane and semi-Baxter permutations. This result, together with

our proof below of the Wilf-equivalence of 2958(B,C,D), establishes that all three patterns give rise to avoidances sets that are equinumerous with plane permutations.

To complete the picture and confirm our original conjecture, we will show that inversion sequences avoiding the pattern 2958A also grow according to the succession rule Ω_{semi} .

2.27.1 Class 2958A: Avoiding $e_j < e_k$ and $e_i \geq e_k$

Bouvel, et al. utilize generating trees to relate distinct combinatorial objects to the semi-Baxter sequence, providing a *succession rule* for the sequence. To use this succession rule, each combinatorial object is identified with a label, (h, k) , which is based on the properties of the object. The rule then describes the labels of those members of the same class of objects that can be obtained by systematically adding one atom. For our purposes, adding one atom will correspond to adding a new last entry to an inversion sequence.

The semi-Baxter sequence follows the succession rule:

$$\Omega_{semi} = \begin{cases} (1, 1) \\ (h, k) \rightsquigarrow (1, k+1), \dots, (h, k+1) \\ (h+k, 1), \dots, (h+1, k) \end{cases}$$

The top line of the succession rule gives the label for the “root” object (which will be the inversion sequence (0) in our case) and the second line shows the labels of the inversion sequences obtained by systematically adding a new last entry.

We show how to define the labels (h, k) for inversion sequences avoiding $(-, <, \geq)$ in such a way that those sequences grow according to the succession rule Ω_{semi} .

Theorem 48. $\mathbf{I}_n(-, <, \geq) = \mathbf{I}_n(101, 201)$ grows according to the succession rule Ω_{semi} and is therefore enumerated by the semi-Baxter sequence.

Proof. For each $e \in \mathbf{I}_n(-, <, \geq)$, we will account for all $e' \in \mathbf{I}_{n+1}(-, <, \geq)$ obtainable by appending a new last entry e_{n+1} . Define an *active site* for e to be a value c such that $(e_1, e_2, \dots, e_n, c) \in \mathbf{I}_{n+1}(-, <, \geq)$. Now, for any inversion sequence e , let h denote the number of active sites less than or equal to $\maxx(e)$, and let k denote the number of active sites greater than $\maxx(e)$. First notice that for $e = (0) \in \mathbf{I}_1(-, <, \geq)$, since $\maxx(e) = 0$, the set of active sites is $\{0, 1\}$, with $0 \leq \maxx(e)$ and $1 > \maxx(e)$, so $(h, k) = (1, 1)$.

For $n \geq 2$, let $e \in \mathbf{I}_n(-, <, \geq)$. Let e_{n+1} be the value of one of the active sites for e and consider $e' = (e_1, \dots, e_{n+1}) \in \mathbf{I}_{n+1}(-, <, \geq)$.

If $e_{n+1} \leq \maxx(e)$, then since e' must only avoid 101 and 201, an active site for e' must either be less than or equal to e_{n+1} or greater than $\maxx(e) = \maxx(e')$. So, the labels of the inversion sequences we obtain by appending such an active site e_{n+1} to e are $(1, k+1), (2, k+1), \dots, (h, k+1)$.

If $e_{n+1} > \maxx(e)$, again since e' must only avoid 101 and 201, all sites active for e are also active for e' . Additionally, e' will have a new active site: $n+1$. Since $\maxx(e') > \maxx(e)$, some active sites for e that were counted with k will now be counted with h when finding

the label for e' . The labels of the inversion sequences we obtain by appending such an active site e_{n+1} to e are $(h+k, 1), (h+k-1, 2), \dots, (h+1, k)$.

It follows that $\mathbf{I}_n(-, <, \geq)$ grows by the succession rule Ω_{semi} and is therefore enumerated by the semi-Baxter sequence. \square

2.27.2 Class 2958B: Avoiding $e_i > e_j \geq e_k$

Note that $\mathbf{I}_n(>, \geq, -) = \mathbf{I}_n(210, 100)$. It was shown by Corteel, et al. [13] that the inversion sequences avoiding 210 (class 4306A) have the following useful characterization.

Define a *weak left-to-right maximum* in an inversion sequence e to be a position j such that $e_i \leq e_j$ for all $i \in [j-1]$.

For $e \in \mathbf{I}_n$, let $a_1 \leq a_2 \leq \dots \leq a_t$ be the sequence of weak left-to-right maxima of e . Let $b_1 < b_2 < \dots < b_{n-t}$ be the sequence of remaining indices in $[n]$. Let $e^{top} = (e_{a_1}, e_{a_2}, \dots, e_{a_t})$ and $\text{top}(e) = e_{a_t}$. Let $e^{bottom} = (e_{b_1}, e_{b_2}, \dots, e_{b_{n-t}})$ and $\text{bottom}(e) = e_{b_{n-t}}$. If every entry of e is a weak left-to-right maximum, then e^{bottom} is empty and we set $\text{bottom}(e) = -1$.

Observation 49 ([13]). The inversion sequence e avoids 210 if and only if e^{top} and e^{bottom} are weakly increasing sequences.

We can extend this to $\mathbf{I}_n(210, 100)$.

Observation 50. The inversion sequence e avoids both 210 and 100 if and only if e^{top} is weakly increasing and e^{bottom} is strictly increasing.

Using *Observation 50*, Bouvel, et al. showed the following (where SB_n denotes the n th semi-Baxter number):

Theorem 51 ([7]. Theorem 20). *There are as many inversion sequences of size n avoiding 210 and 100 as plane permutations of size n . In other words $|\mathbf{I}_n(210, 100)| = SB_n$.*

The characterization in *Observation 50* can be used to derive a recurrence.

Theorem 52. *Let $S_{n,a,b}$ be the number of $e \in \mathbf{I}_n(201, 100)$ with $\text{top}(e) = a$ and $\text{bottom}(e) = b$. Then*

$$S_{n,a,b} = \sum_{i=-1}^{b-1} S_{n-1,a,i} + \sum_{j=b+1}^a S_{n-1,j,b},$$

with initial conditions $S_{n,a,b} = 0$ if $a \geq n$ and $S_{n,a,-1} = \frac{n-a}{n} \binom{n-1+a}{a}$.

From *Theorem 52* we get:

$$|\mathbf{I}_n(210, 100)| = \sum_{a=0}^{n-1} \sum_{b=-1}^{a-1} S_{n,a,b} = \frac{1}{n+1} \binom{2n}{n} + \sum_{a=0}^{n-1} \sum_{b=0}^{a-1} S_{n,a,b}. \quad (9)$$

2.27.3 Class 2958C: Avoiding $e_i \geq e_j > e_k$

Now we will show that the class 2958B is Wilf-equivalent to 2958C.

Theorem 53. *The patterns $(-, <, \geq)$ and $(\geq, >, -)$, defining classes 2958B and 2958C respectively, are Wilf-equivalent.*

Proof. The avoidance sets for classes 2958B and 2958C are $\mathbf{I}_n(100, 210)$ and $\mathbf{I}_n(110, 210)$, respectively. We describe a bijection

$$\alpha : \mathbf{I}_n(110, 210) \rightarrow \mathbf{I}_n(100, 210).$$

For $e \in \mathbf{I}_n(110, 210)$, let $\alpha(e) = e' = (e'_1, \dots, e'_n)$ where, for $1 \leq j \leq n$,

$$e'_j = \begin{cases} \max\{e_1, \dots, e_j\}, & \text{if } e_j = e_k \text{ for some } k > j; \\ e_j, & \text{otherwise.} \end{cases} \quad (10)$$

Note that for $1 \leq j \leq n$,

$$e'_j \leq \max\{e_1, \dots, e_j\} = \max\{e'_1, \dots, e'_j\} \quad (11)$$

and if $e'_j \neq e_j$ then

$$e'_j = \max\{e_1, \dots, e_j\} \geq e'_i \text{ for } 1 \leq i < j. \quad (12)$$

To see that e' avoids 100, suppose $e'_i > e'_j = e'_k$ for some $i < j < k$. Then by (12), we must have $e'_j = e_j$ and $e'_k = e_k$. But from the definition of α (10), if $e_j = e_k$ where $j < k$ then $e'_j = \max\{e_1, \dots, e_j\} \geq e'_i$, which is a contradiction.

To see that e' avoids 210, suppose that $e'_i > e'_j > e'_k$ for some $i < j < k$. Again by (12), $e'_j = e_j$ and $e'_k = e_k$. Since e avoids 210, $e'_i \neq e_i$ so $e'_i = \max\{e_1, \dots, e_i\} = e_s$ for some $s \in [i]$. But then $e_s > e_j > e_k$ is a 210 in e .

Thus $e' = \alpha(e) \in \mathbf{I}_n(100, 210)$. To show that α is a bijection, we define its inverse β .

First we make an observation. Consider some $e \in \mathbf{I}_n(110, 210)$ and an entry e_j such that $e'_j \neq e_j$ in $\alpha(e) = e'$. This implies that there exists some index k with $j < k$ and $e_j = e_k = m$ for some value m . Additionally, it must be the case that $m < M = \max\{e_1, \dots, e_{j-1}\} = e_i$ where $i \in [j-1]$ (else, $e'_j = m$). Then, since e avoids 210, we must have $m = \min\{e_j, \dots, e_n\}$. Thus, for $e' = \alpha(e)$ we have

$$e'_i = e_i = M = e'_j > e_j = m = \min\{e_j, \dots, e_n\} = \min\{e'_j, \dots, e'_n\}.$$

So, we can reconstruct e from e' by defining $\beta : \mathbf{I}_n(100, 210) \rightarrow \mathbf{I}_n(110, 210)$ as follows.

For $e \in \mathbf{I}_n(100, 210)$, let $\beta(e) = e' = (e'_1, \dots, e'_n)$ where for $1 \leq j \leq n$

$$e'_j = \begin{cases} \min\{e_j, \dots, e_n\}, & \text{if } e_i = e_j \text{ for some } i < j; \\ e_j, & \text{otherwise.} \end{cases}$$

Then $\beta(\alpha(e)) = e$. We can check similarly that $\beta(e) \in \mathbf{I}_n(110, 210)$ and $\alpha(\beta(e)) = e$. \square

We now return to the proof of the Wilf-equivalence of the classes 2549A and 2549C from the previous section.

Proof of Theorem 47. Observe that $2549C \preceq 2958C$ and $2549A \preceq 2958B$ in the following sense:

$$2549C : \mathbf{I}_n(\geq, \neq, >) = \mathbf{I}_n(110, 210, 201) \subseteq \mathbf{I}_n(110, 210) = \mathbf{I}_n(\geq, >, -) : 2958C$$

$$2549A : \mathbf{I}_n(>, -, >) = \mathbf{I}_n(100, 210, 201) \subseteq \mathbf{I}_n(100, 210) = \mathbf{I}_n(>, \geq, -) : 2958B$$

We check that the mapping α (10) restricts to a bijection between inversion sequences in class 2549C and in class 2549A.

Let $e \in \mathbf{I}_n(110, 210)$ and let $e' = \alpha(e)$. Suppose e avoids the pattern 201, but that for some $i < j < k$, $e'_i > e'_j < e'_k$ and $e'_i > e'_k$. Then by (12), $e'_j = e_j$ and $e'_k = e_k$. But, since e avoids 201, $e'_i \neq e_i$. Then, by definition of α , there is an $s \in [i - 1]$ such that $e'_i = e_s$. But then $e_s e_j e_k$ forms a 201 pattern in e , a contradiction.

For the converse, let $e \in \mathbf{I}_n(100, 210)$ and let $e' = \beta(e)$. Suppose there is $i < j < k$ such that $e'_i e'_j e'_k$ form a 201. We show e must also contain 201. By definition of β , since $e'_i > e'_j$, $e_i = e'_i$. Also, there exists $s : j \leq s \leq n$ such that $e'_j = e_s$ and $t : k \leq t \leq n$ such that $e'_k = e_t$. Since e avoids 210, it must be that $s < t$. But then $e_i e_s e_t$ is a 201 in e . □

2.27.4 2958D: Avoiding $e_j \leq e_k$ and $e_i > e_k$

Theorem 54. *The patterns $(>, \geq, -)$ and $(-, \leq, >)$, defining classes 2958B and 2958D respectively, are Wilf-equivalent.*

Proof. The inversion sequences in classes 2958B and 2958D are defined by $\mathbf{I}_n(210, 100)$ and $\mathbf{I}_n(201, 100)$, respectively. It was shown in [13], Theorem 5, that the following gives a bijection from $\mathbf{I}_n(210)$ to $\mathbf{I}_n(201)$. It can be checked that this mapping preserves 100-avoidance and therefore restricts to a bijection from $\mathbf{I}_n(210, 100)$ to $\mathbf{I}_n(201, 100)$.

Given $e \in \mathbf{I}_n(210)$, define $f \in \mathbf{I}_n(201)$ as follows. Let $e_{a_1} \leq e_{a_2} \leq \dots \leq e_{a_t}$ be the sequence of weak left-to-right maxima of e and let $e_{b_1}, e_{b_2}, \dots, e_{b_{n-t}}$ be the subsequence of remaining elements of e . Since e avoids both 210 and 100, $e_{b_1} < e_{b_2} < \dots < e_{b_{n-t}}$.

For $i = 1, \dots, t$, set $f_{a_i} = e_{a_i}$. For each $j = 1, 2, \dots, n - t$, we extract an element of the multiset $B = \{e_{b_1}, e_{b_2}, \dots, e_{b_{n-t}}\}$ and assign it to $f_{b_1}, f_{b_2}, \dots, f_{b_{n-t}}$ as follows:

$$f_{b_j} = \max\{k \mid k \in B - \{f_{b_1}, f_{b_2}, \dots, f_{b_{j-1}}\} \text{ and } k < \max(e_1, \dots, e_{b_{j-1}})\}.$$

□

This is the same mapping that was used in Section 2.16 to show that the patterns 772A and 772B are Wilf-equivalent. Note that

$$772B : \mathbf{I}_n(210, 110, 100, 000) \subseteq \mathbf{I}_n(210, 100) : 2958B;$$

$$772A : \mathbf{I}_n(201, 101, 100, 000) \subseteq \mathbf{I}_n(201, 100) : 2958D.$$

By the Wilf-equivalencies between classes B, C, and D, we know that all of 2958(A,B,C,D) are enumerated by the semi-Baxter numbers.

2.28 Classes 3207(A,B): $\mathbf{I}_n(101)$, $\mathbf{I}_n(110)$

It was shown by Corteel, et al. [13] that both $\mathbf{I}_n(<, -, =) = \mathbf{I}_n(101)$ and $\mathbf{I}_n(=, >, -) = \mathbf{I}_n(110)$ are counted by the sequence A113227 in the OEIS [15], where it is said to count $\mathbf{S}_n(\underline{1234})$. $\mathbf{S}_n(\underline{1234})$ is the set of permutations with no $i < j < k < \ell$ such that $\pi_i < \pi_j < \pi_k < \pi_\ell$ and $k = j + 1$.

It was proven by David Callan in [10], that $\mathbf{S}_n(\underline{1234})$ is in bijection with increasing ordered trees with $n + 1$ vertices whose leaves, taken in preorder, are also increasing. He showed that if $u_{n,k}$ is the number of such trees with $n + 1$ vertices in which the root has k children then

$$u_{n,k} = u_{n-1,k-1} + k \sum_{j=k}^{n-1} u_{n-1,j} \quad (13)$$

with initial conditions $u_{0,0} = 1$ and $u_{n,k} = 0$ if $k > n$, or $n > 0$ and $k = 0$.

It was shown by Corteel, et al. [13] that the number of $e \in \mathbf{I}_n(101)$ with exactly k zeros is $u_{n,k}$, as is the number of $e \in \mathbf{I}_n(110)$ with exactly k zeros. As a consequence, 101 and 110 are Wilf-equivalent and both avoidance sets are counted by A113227.

2.29 Class 3720: Quadrant Marked Mesh Patterns

In this section we prove that $\mathbf{I}_n(>, \neq, >)$ is counted by the sequence A212198 in the OEIS [15] where it is said to count permutations avoiding a particular marked mesh pattern.

Kitaev and Remmel [18] introduced the idea of quadrant marked mesh patterns, a definition of which is given below.

Definition 55. Let $\pi = \pi_1\pi_2 \dots \pi_n \in S_n$. Consider the graph of π , $G(\pi)$, consisting of the points (i, π_i) for all $i \in [n]$. The entry π_i is said to *match* the quadrant marked mesh pattern $\text{MMP}(a, b, c, d)$, if in $G(\pi)$ there are at least a points to the northeast of (i, π_i) , at least b points to the northwest of (i, π_i) , at least c points to the southwest of (i, π_i) , and at least d points to the southeast of (i, π_i) . The order in which we consider quadrants proceeds counterclockwise, beginning with the top right quadrant.

Let $S_n(\text{MMP}(a, b, c, d))$ denote the set of permutations of length n where no π_i matches $\text{MMP}(a, b, c, d)$. We will prove that $|S_n(\text{MMP}(0, 2, 0, 2))| = |\mathbf{I}_n(>, \neq, >)|$ for all n . By symmetry established by Kitaev and Remmel [18], this implies that $|\mathbf{I}_n(>, \neq, >)| = |S_n(\text{MMP}(2, 0, 2, 0))|$. In our proof, we make use of the bijection ϕ from Section 2.24, whose definition was given in Definition 42. Specifically, we can prove the following:

Theorem 56. For all n , $\phi(S_n(\text{MMP}(0, 2, 0, 2))) = \mathbf{I}_n(>, \neq, >)$.

Proof. This proof is very similar to the proof of Theorem 45. Consider some $\pi \in \mathbf{S}_n$ such that there exists some π_i that matches $\text{MMP}(0, 2, 0, 2)$. This implies that there exist indices $a < b < i < j < k$ such that $\min\{\pi_a, \pi_b\} > \pi_i > \max\{\pi_j, \pi_k\}$. We will show that there exists an occurrence of the pattern $(>, \neq, >)$ in $\phi(\pi) = e$.

We must consider two cases. If $\pi_i \leq i$, then $j > \pi_i > \pi_j$ and $k > \pi_i > \pi_k$. Therefore, $e_i = \pi_i - 1$, $e_j = \pi_j - 1$ and $e_k = \pi_k - 1$. So, since $\pi_i > \max\{\pi_j, \pi_k\}$ and $\pi_j \neq \pi_k$, e_i, e_j, e_k forms an occurrence of $(>, \neq, >)$.

Now assume $\pi_i > i$. Recall that e_i is the t -th smallest element of $\{e_{i+1}, \dots, e_n\}$ if π_i is the t -th largest element of $\{\pi_1, \pi_2, \dots, \pi_i\}$. Since π_a, π_b are larger than and occur before π_i , we know that t is at least 3. If $e_{j'}$ and $e_{k'}$ are the two smallest distinct values in the set $\{e_{i+1}, \dots, e_n\}$, we are guaranteed that $e_i > e_{j'}$ and $e_i > e_{k'}$; so, $e_i, e_{j'}, e_{k'}$ form the pattern $(>, \neq, >)$.

For the converse, let $\pi \in \mathbf{S}_n$ such that $e = \phi(\pi)$ has indices $i < j < k$ such $e_i > e_j$, $e_i > e_k$, and $e_i \neq e_k$. We show that π contains an entry that matches $\text{MMP}(0, 2, 0, 2)$.

If $e_i \in \{e_{i+1}, \dots, e_n\}$ then, by definition of ϕ , $\pi_i > i$ and e_i is the k th smallest element of $\{e_{i+1}, \dots, e_n\}$ for some $k \geq 2$, noting that the distinct elements e_j and e_k are both less than e_i . Then, again by definition of ϕ , π_i is the k th largest element of $\{\pi_1, \dots, \pi_i\}$, so there is some $a, b < i$ with $\pi_a, \pi_b > \pi_i$. To show that π_i matches $\text{MMP}(0, 2, 0, 2)$ it remains to show there are at least two elements of $\{\pi_{i+1}, \dots, \pi_n\}$ that are smaller than π_i . The number of elements in π that are larger than π_i is $n - \pi_i < n - i$. At least two of these are to the left of π_i in π . Thus at most $n - i - 2$ can be in the $n - i$ positions to the right of π_i .

If $e_i \notin \{e_{i+1}, \dots, e_n\}$ then, by definition of ϕ , $\pi_i \leq i$ and $e_i = \pi_i - 1$. Since $e_j \leq e_i \leq i < j < k$ and $e_k \leq e_i \leq i < k$, we have $e_j = \pi_j - 1$ and $e_k = \pi_k - 1$, so that $\pi_i \pi_j \pi_k$ has the pattern 321 or 312. To show that π_i matches $\text{MMP}(0, 2, 0, 2)$, it remains to show there is an $a < i$ and $b < i$ such that $\pi_a, \pi_b > \pi_i$. Note that there are $n - \pi_i \geq n - i$ elements of π larger than π_i . The $n - i$ positions to the right of π_i hold at least two elements smaller than π_i . So there must be two larger element to the left of π_i , as desired. \square

The bijection ϕ turns out to be a versatile tool, giving interesting results when restricted to $\mathbf{S}_n(\text{MMP}(k, 0, k, 0))$ for any positive integer k . In this case, $\phi(\mathbf{S}_n(\text{MMP}(k, 0, k, 0)))$ maps to inversion sequences that avoid a particular set of length $k + 1$ patterns.

2.30 Class 5040: $n!$

The last equivalence class of patterns in this section is the set of those avoided by all inversion sequences. There are 41 such patterns among our 343, including the representative below. One such class is all inversion sequences with no $i > j > k$ such that $e_i = e_j = e_k$ and $e_i \neq e_k$.

3 Results about patterns whose sequences don't appear in the OEIS [15]

Table 3 lists all equivalence classes of the patterns $\rho \in \{<, >, \leq, \geq, =, \neq, -\}^3$ whose avoidance sequences did not appear in the OEIS (and have subsequently been entered). We were able to derive the avoidance sequences for a few of these patterns and prove Wilf-equivalence of some others. In this section we describe our results and leave identification of the avoidance sequences of the remaining patterns in Table 3 as questions for future study.

3.1 Counting results

3.1.1 Class 805: Avoiding $e_i \leq e_j > e_k$ and $e_i \neq e_k$

We were able to prove that $\mathbf{I}_n(\leq, >, \neq)$ (A279557) is counted by a sum of Catalan numbers. Recall that $C_n = \frac{1}{n+1} \binom{2n}{n}$ and $C(x) = \sum_{n \geq 0} C_n x^n = \frac{1 - \sqrt{1-4x}}{2x}$.

Observe that $\mathbf{I}_n(\leq, >, \neq)$ is the set of $e \in \mathbf{I}_n$ satisfying, for some t with $1 < t \leq n$ and k with $t < k < n$,

$$0 = e_1 = e_2 = \dots = e_{t-1} < e_t \leq e_k \leq e_{k+1} \leq \dots \leq e_n$$

with $0 = e_{t+1} = \dots = e_{k-1}$.

Lemma 57. *Let $A(x) = \sum_{n \geq 0} |\mathbf{I}_n(\leq, >, \neq)| x^n$. Then $A(x) = \frac{(1-2x)(1-2x-\sqrt{1-4x})}{2x^2(1-x)}$.*

Proof. For any $e = (e_1, e_2, \dots, e_n) \in \mathbf{I}_n(\leq, >, \neq)$, e satisfies one of the following three cases:

1. $e = \emptyset$ or e has no maximal entries,
2. $e = (0, 0, \dots, 0, e_t, e_{t+1}, \dots, e_n)$ where t is the index of the last occurrence of a maximal entry of e , or
3. $e = (0, 0, \dots, 0, e_t, \dots, e_j, \dots, e_n)$ where $e_t \neq 0$ and j is the index of the last occurrence of a maximal entry of e .

The generating function for case 1 is $(1 + xA(x))$, since any inversion sequence in $\mathbf{I}_n(\leq, >, \neq)$ with no maximal entries can be constructed by appending a 0 to the front of an inversion sequence in $\mathbf{I}_{n-1}(\leq, >, \neq)$.

Let $f = (f_1, f_2, \dots, f_k)$ be an inversion sequence that avoids $(-, >, -)$ (that is, f is an inversion sequence with weakly increasing entries, as seen in Section 2.14). Recall that $|\mathbf{I}_n(-, >, -)| = C_n$. Then any inversion sequence in case 2 is of the form $(0, 0, \dots, f_1, t-1, \sigma_{t-1}(f_2, f_3, \dots, f_k))$, where $t-1$ occurs in position t (and is therefore maximal). Notice that, in order for this construction to fall in case 2, f must have length greater than zero. It follows that the generating function for case 2 is $\frac{x}{1-x} \cdot (C(x) - 1)$.

Finally, case 3 consists of all inversion sequences of the form $e' \cdot (j-1) \cdot (f+j-1)$ where $e' \in (\mathbf{I}_{j-1}(\leq, >, \neq) \setminus \{(0, 0, \dots, 0)\})$ and $(f+j-1)$ denotes the sequence obtained by adding $j-1$ to each entry of some $f \in \mathbf{I}_{n-t}(-, >, -)$. The generating function for this case is $(A(x) - \frac{x}{1-x}) \cdot x \cdot C(x)$.

From these three disjoint cases, we have

$$A(x) = 1 + xA(x) + \frac{x}{1-x} \cdot (C(x) - 1) + (A(x) - \frac{x}{1-x}) \cdot x \cdot C(x).$$

Solving for $A(x)$ gives

$$A(x) = \frac{1 - 2x}{(1-x)(1-x-xC(x))}$$

By substituting the known expression for $C(x)$, and through conjugation and algebraic rearrangement, the result follows. \square

Theorem 58. $|\mathbf{I}_n(\leq, >, \neq)| = C_{n+1} - \sum_{i=1}^n C_i$, where $C_n = \frac{1}{n+1} \binom{2n}{n}$.

Proof. Let $B_n = C_{n+1} - \sum_{i=1}^n C_i$. Then $\sum_{n \geq 0} B_n x^n = \frac{C(x)-1}{x} - \frac{C(x)-1}{1-x}$. By substituting $C(x) = \frac{1-\sqrt{1-4x}}{2x}$ and simplifying, it follows that $A(x) = B(x)$ and therefore $B_n = |\mathbf{I}_n(\leq, >, \neq)|$. \square

3.1.2 Class 1016: Avoiding $e_i > e_j$ and $e_i \neq e_k$

We can show that the counting sequence for $|\mathbf{I}_n(>, -, \neq)|$ (A279560) is as follows. We omit the details since we hope to find a simpler formula and nicer explanation.

$$|\mathbf{I}_n(>, -, \neq)| = \binom{2(n-1)}{n-1} + \sum_{k=2}^{n-2} \sum_{i=1}^{k-1} \sum_{u=1}^i \sum_{d=0}^{u-1} \frac{i-d+1}{i+1} \binom{i+d}{d}.$$

3.1.3 1079(A,B): sum of binomial coefficients

The sequences in $\mathbf{I}_n(>, \neq, -)$ (Class 1079A, A279561) have a nice unimodality characterization. They are the inversion sequences $e \in \mathbf{I}_n$ satisfying for some t :

$$e_1 \leq e_2 \leq \dots \leq e_t \geq e_{t+1} = e_{t+2} = \dots = e_n.$$

From this characterization, we can show the following.

Theorem 59. $|\mathbf{I}_n(>, \neq, -)| = 1 + \sum_{i=1}^{n-1} \binom{2i}{i-1}$.

Additionally, we conjecture that $\mathbf{I}_n(<, >, \neq)$ (Class 1079B) is Wilf-equivalent, but have not proven this.

3.1.4 4306A,B: $\mathbf{I}_n(210)$, $\mathbf{I}_n(201)$

The sets $\mathbf{I}_n(-, <, >) = \mathbf{I}_n(201)$ and $\mathbf{I}_n(>, >, -) = \mathbf{I}_n(210)$ were shown to be Wilf-equivalent by Corteel, et al. [13] via a bijection, which we made use of in Section 2.27. An alternate proof of this fact was discovered by Mansour and Shattuck [23]. A recurrence to compute $|\mathbf{I}_n(201)|$ was also derived in [13].

3.2 Wilf-equivalence results

In the remainder of the section we show that the bijection α described in (10) of Section 2.27 proves Wilf-equivalence of all of the following pairs of patterns: 663A,B; 746A,B; 1833A,B; and 1953A,B.

3.2.1 Class 1953A: Avoiding $e_j > e_k$ and $e_i > e_k$, and Class 1953B: Avoiding $e_i \neq e_j \geq e_k$ and $e_i > e_k$

Theorem 60. *The patterns $(-, >, >)$ and $(\neq, \geq, >)$, defining classes 1953A and 1953B respectively, are Wilf-equivalent.*

Proof. The inversion sequences in classes 1953A and 1953B are those in $\mathbf{I}_n(110, 210, 120)$ and $\mathbf{I}_n(100, 210, 120)$, respectively. Notice that these are the inversion sequences in $\mathbf{I}_n(110, 210)$ (class 2958C) and $\mathbf{I}_n(100, 210)$ (class 2958A), respectively, that avoid 120. So it suffices to show that both $\alpha : \mathbf{I}_n(110, 210) \rightarrow \mathbf{I}_n(100, 210)$ and $\beta = \alpha^{-1}$ preserves 120-avoidance.

Suppose $e \in \mathbf{I}_n(110, 210)$ avoids 120, but for $e' = \alpha(e)$ there exist $i < j < k$ such that $e'_i < e'_j > e'_k$ and $e'_i > e'_k$. By (12), $e'_k = e_k$. Notice that we cannot have both $e'_i = e_i$ and $e'_j = e_j$, since this would create a 120 in e .

Suppose first that $e'_j = e_j$. Since e avoids 120, $e'_i \neq e_i$ so, by definition of α , there is an $s \in [i - 1]$ such that $e_s = e'_i$. But then $e_s e_j e_k$ forms a 120 in e .

So, assume that $e'_i = e_i$. Then, since e avoids 120, $e'_j \neq e_j$. So, there must be a $t \in [j - 1]$ such that $e_t = e'_j$. If $i < t < k$ then $e_i e_t e_k$ is a 120 in e . Otherwise, $t < i < k$ and $e_t e_i e_k$ is a 210 in e , which is impossible.

Finally, if both $e'_i \neq e_i$ and $e'_j \neq e_j$, then let s and t be as above. If $s < t$ then $e_s e_t e_k$ forms a 102 in e . Otherwise, $e_t e_s e_k$ forms a 210 in e . Both cases lead to a contradiction.

If both $e'_i \neq e_i$ and $e'_j \neq e_j$, then let $s \in [i - 1]$ and $j \in [j - 1]$ be indices such that $e'_i = e_s$ and $e'_j = e_t$. Since $e'_i = \max\{e_1, e_2, \dots, e_i\}$ and $e'_j = \max\{e_1, e_2, \dots, e_j\}$, and we have $e'_i < e'_j$, it must be the case the $e_s < e_t$ and $s < t$. Therefore $e_s e_t e_k$ is an occurrence of 120 in e , giving a contradiction.

It follows that α preserves 120-avoidance. Showing that β preserves 120-avoidance is similar. \square

3.2.2 Class 1833A: Avoiding $e_j \neq e_k$ and $e_i > e_k$, and Class 1833B: Avoiding $e_i \neq e_j$ and $e_i > e_k$

Theorem 61. *The patterns $(-, \neq, >)$ and $(\neq, -, >)$, defining classes 1833A and 1833B respectively, are Wilf-equivalent.*

Proof. The inversion sequences in classes 1833A and 1833B are those in $\mathbf{I}_n(110, 210, 120, 201)$ and $\mathbf{I}_n(100, 210, 120, 201)$, respectively. It was shown in Section 2.27 that the bijection $\alpha : \mathbf{I}_n(110, 210) \rightarrow \mathbf{I}_n(100, 210)$ preserves 201-avoidance, as does its inverse β . It follows from Theorem 60 that α and β preserves 120-avoidance as well. So $\alpha(\mathbf{I}_n(-, \neq, >)) = \mathbf{I}_n(\neq, -, >)$. \square

3.2.3 Class 746A: Avoiding $e_j > e_k$ and $e_i \geq e_k$, and Class 746B: Avoiding $e_i \neq e_j \geq e_k$ and $e_i \geq e_k$

Theorem 62. *The patterns $(-, >, \geq)$ and (\neq, \geq, \geq) , defining classes 746A and 746B respectively, are Wilf-equivalent.*

Proof. The inversion sequences in classes 746A and 746B are those in $\mathbf{I}_n(110, 210, 120, 010)$ and $\mathbf{I}_n(100, 210, 120, 010)$, respectively. These are, in turn, the inversion sequences in $\mathbf{I}_n(110, 210, 120)$ (class 1953A) and $\mathbf{I}_n(100, 210, 120)$ (class 1953B), respectively, that avoid 010. By Theorem 60, $\alpha : \mathbf{I}_n(110, 210) \rightarrow \mathbf{I}_n(100, 210)$ and $\alpha^{-1} = \beta$ preserve 120-avoidance. So it suffices to show that both $\alpha : \mathbf{I}_n(110, 210) \rightarrow \mathbf{I}_n(100, 210)$ and $\alpha^{-1} = \beta$ preserves 010-avoidance.

Suppose $e \in \mathbf{I}_n(110, 210)$ avoids 010, but for $e' = \alpha(e)$ there exist $i < j < k$ such that $e'_i < e'_j > e'_k$ and $e'_i = e'_k$. By (12), $e'_k = e_k$. Since e avoids 010, it follows that we cannot have both $e'_i = e_i$ and $e'_j = e_j$.

Suppose first that $e'_j = e_j$. Since e avoids 010, $e'_i \neq e_i$ so, by definition of α there is an $s \in [i - 1]$ such that $e_s = e'_i$. But then $e_s e_j e_k$ forms a 010 in e .

So, assume that $e'_i = e_i$. Then, since e avoids 010, $e'_j \neq e_j$. So, there must be a $t \in [j - 1]$ such that $e_t = e'_j$. If $i < t < k$ then $e_i e_t e_k$ is a 010 in e . Otherwise, $t < i < k$ and $e_t e_i e_k$ is a 100 in e , which is impossible.

If both $e'_i \neq e_i$ and $e'_j \neq e_j$, then let $s \in [i - 1]$ and $t \in [j - 1]$ be indices such that $e'_i = e_s$ and $e'_j = e_t$. Since $e'_i = \max\{e_1, e_2, \dots, e_i\}$ and $e'_j = \max\{e_1, e_2, \dots, e_j\}$, and we have $e'_i < e'_j$, it must be the case the $e_s < e_t$ and $s < t$. Additionally, $e_s = e'_i = e'_k = e_k$. Therefore $e_s e_t e_k$ is an occurrence of 010 in e , giving a contradiction.

It follows that α preserves 010-avoidance. Showing that β preserves 010-avoidance is similar. \square

3.2.4 Class 663A: Avoiding $e_j \neq e_k$ and $e_i \geq e_k$, and Class 663B: Avoiding $e_i \neq e_j$ and $e_i \geq e_k$

Theorem 63. *The patterns $(-, \neq, \geq)$ and $(\neq, -, \geq)$, defining classes 663A and 6633B respectively, are Wilf-equivalent.*

Proof. The inversion sequences in classes 663A and 663B are those in $\mathbf{I}_n(110, 210, 120, 201, 010)$ and $\mathbf{I}_n(100, 210, 120, 201, 010)$, respectively. It was shown in Section 2.27 that the bijection $\alpha : \mathbf{I}_n(110, 210) \rightarrow \mathbf{I}_n(100, 210)$ preserves 201-avoidance, as does its inverse. It follows from Theorem 60 that α, β preserve 120-avoidance and from Theorem 62 that they preserve 010-avoidance as well. So $\alpha(\mathbf{I}_n(-, \neq, \geq)) = \mathbf{I}_n(\neq, -, \geq)$. \square

4 Concluding remarks

The results in this work demonstrate that inversion sequences avoiding a triple of relations have connections to a vast array of pattern-avoiding classes and combinatorial sequences. Moreover, as a pattern to avoid, a triple of relations can be more expressive than a single word of length 3, even though it is always equivalent to an avoidance *set* of words of length 3.

On the other hand, $\mathbf{I}_n(001, 210)$ is not the avoidance set of any triple of relations considered in this paper. In ongoing work, we construct and examine the partially ordered set whose elements are the avoidance sets $\mathbf{I}_n(S)$, ordered by inclusion, where

$$S \subseteq \{000, 001, 010, 100, 011, 101, 110, 123, 132, 213, 231, 312, 321\}.$$

Several interesting questions remain for future work, such those highlighted in Table 2 with a “no” in column 3. For instance, is $\mathbf{I}_n(>, -, >)$ equinumerous with $\mathbf{S}_n(4231, 42513)$ and is $\mathbf{I}_n(>, -, \leq)$ equinumerous with $\mathbf{S}_n(4123, 4132, 4213)$? Are $\mathbf{I}_n(=, \geq, -)$ and $\mathbf{I}_n(\geq, =, -)$ counted by the Bell numbers?

Another fascinating open question is whether there is a bijective mapping to relate the Baxter permutations in \mathbf{S}_n to $\mathbf{I}_n(\geq, \geq, >)$. There are also a number of lingering open enumeration problems: can enumeration formulas be found for some of the avoidance sets in Table 3, such as $\mathbf{I}_n(010)$, $\mathbf{I}_n(100)$, $\mathbf{I}_n(120)$, or $\mathbf{I}_n(201) = \mathbf{I}_n(210)$?

Pattern avoidance can be studied in more general s -inversion sequences. For a given sequence $s = (s_1, s_2, \dots, s_n)$ of positive integers, the s -inversion sequences $\mathbf{I}_n^{(s)}$ are defined by

$$\mathbf{I}_n^{(s)} = \{(e_1, e_2, \dots, e_n) \in Z^n \mid 0 \leq e_i < s_i, 1 \leq i \leq n\}.$$

When $s = (1, 2, \dots, n)$, these are the usual inversion sequences. For $s = (2, 4, \dots, 2n)$, there is a statistics-preserving bijection between $\mathbf{I}_n^{(s)}$ and signed permutations of $[n]$. See [26] for other interesting families of s -inversion sequences, and for an overview of their combinatorial and geometric relationships to s -lecture hall partitions and cones.

5 Acknowledgments

Thanks to Sylvie Corteel for helpful discussions about several of the sequences herein. Thanks to Michael Weselcouch for independently verifying our computations of the avoidance sequences for the 343 patterns. Thanks to the referee of this paper for a careful reading

of the manuscript and many helpful suggestions. Thanks to Michael Albert for making his PermLab software freely available [2]. Thanks to the Simons Foundation for Collaboration Grant No. 244963 to the second author which supported the travel of the first author for collaboration. Some of this material is based upon work supported by the National Science Foundation under Grant No. 1440140, while the second author was in residence at the Mathematical Sciences Research Institute in Berkeley, California, during the fall semester of 2017.

We once again extend sincere thanks to Neil Sloane and the OEIS Foundation, Inc. The On-Line Encyclopedia of Integer Sequences [15] was an invaluable resource for this project.

References

- [1] M. H. Albert, R. E. L. Aldred, M. D. Atkinson, H. P. van Ditmarsch, C. C. Handley, and D. A. Holton, Restricted permutations and queue jumping, *Discrete Math.* **287** (2004), 129–133.
- [2] Michael Albert, Permlab: Software for permutation patterns, 2012.
<http://www.cs.otago.ac.nz/staffpriv/malbert/permlab.php>.
- [3] Michael H. Albert, Cheyne Hamburger, Jay Pantone, Nathaniel Shar, and Vincent Vatter, Generating permutations with restricted containers, (2015).
<http://arxiv.org/pdf/1510.00269.pdf>.
- [4] Sara C. Billey, William Jockusch, and Richard P. Stanley, Some combinatorial properties of Schubert polynomials, *J. Algebraic Combin.* **2** (1993), 345–374.
- [5] Mireille Bousquet-Mélou and Steve Butler, Forest-like permutations, *Ann. Comb.* **11** (2007), 335–354.
- [6] Mireille Bousquet-Mélou and Guoce Xin, On partitions avoiding 3-crossings, *Sém. Lothar. Combin.* **54** (2005/07), Art. B54e.
- [7] Mathilde Bouvel, Veronica Guerrini, Andrew Rechnitzer, and Simone Rinaldi, Semi-Baxter and strong-Baxter: two relatives of the Baxter sequence, (2017).
<https://arxiv.org/pdf/1702.04529.pdf>.
- [8] Alexander Burstein and Walter Stromquist, Unimodal inversion sequences and pattern-avoiding classes, *AMS Fall Central Sectional Meeting, Loyola University, Chicago*, 2015. Slides available at
<http://www.alexanderburstein.org/tiki-index.php?page=Papers+and+Talks/uis-chicago-t>
- [9] David Callan, A combinatorial interpretation of the eigensequence for composition, *J. Integer Sequences* **9** (2006), [Article 06.1.4](#).

- [10] David Callan, A bijection to count (1-23-4)-avoiding permutations, preprint, 2010. Available at <http://arxiv.org/abs/1008.2375>.
- [11] David Callan, The number of $\bar{2}\bar{1}542$ -avoiding permutations, preprint, 2011. Available at <http://arxiv.org/abs/1111.3088>.
- [12] F. R. K. Chung, R. L. Graham, V. E. Hoggatt, Jr., and M. Kleiman, The number of Baxter permutations, *J. Combin. Theory Ser. A* **24** (1978), 382–394.
- [13] Sylvie Corteel, Megan A. Martinez, Carla D. Savage, and Michael Weselcouch, Patterns in inversion sequences I, *Discrete Math. Theor. Comput. Sci.* **18** (2016).
- [14] Shishuo Fu, Zhicong Lin, and Jiang Zeng, Two new unimodal descent polynomials, 2015. <http://arxiv.org/abs/1507.05184>.
- [15] OEIS Foundation Inc., The On-Line Encyclopedia of Integer Sequences, 2011. Available at <http://oeis.org>.
- [16] Dongsu Kim and Zhicong Lin, Refined restricted inversion sequences, preprint, 2017. Available at <https://arxiv.org/pdf/1706.07208.pdf>.
- [17] Sergey Kitaev, *Patterns in Permutations and Words*, Monographs in Theoretical Computer Science, Springer, 2011.
- [18] Sergey Kitaev and Jeffrey Remmel, Quadrant marked mesh patterns, *J. Integer Sequences* **15** (2012), [Article 12.4.7](#).
- [19] Darla Kremer, Permutations with forbidden subsequences and a generalized Schröder number, *Discrete Math.* **218** (2000), 121–130.
- [20] Alain Lascoux and Marcel-Paul Schützenberger, Schubert polynomials and the Littlewood-Richardson rule, *Lett. Math. Phys.* **10** (1985), 111–124.
- [21] Zhicong Lin, Restricted inversion sequences and enhanced 3-noncrossing partitions, preprint, 2017. Available at <https://arxiv.org/pdf/1706.07213.pdf>.
- [22] Percy A. MacMahon, *Combinatory Analysis*, Chelsea Publishing Co., New York, 1960.
- [23] Toufik Mansour and Mark Shattuck, Pattern avoidance in inversion sequences, *Pure Math. Appl.* **25** (2015), 157–176.
- [24] Megan Martinez and Carla Savage, Patterns in inversion sequences II: Inversion sequences avoiding triples of relations [v1], preprint, 2016. Available at <https://arxiv.org/pdf/1609.08106v1.pdf>.
- [25] T. Kyle Petersen and Bridget Eileen Tenner, The depth of a permutation, *J. Comb.* **6** (2015), 145–178.

- [26] Carla D. Savage, The mathematics of lecture hall partitions, *J. Combin. Theory Ser. A* **144** (2016), 443–475.
- [27] Rodica Simion and Frank W. Schmidt, Restricted permutations, *European J. Combin.* **6** (1985), 383–406.
- [28] Richard P. Stanley, *Enumerative combinatorics. Vol. 1*, Vol. 49 of *Cambridge Studies in Advanced Mathematics*, Cambridge University Press, 1997.
- [29] Sheila Sundaram, The homology of partitions with an even number of blocks, *J. Algebraic Combin.* **4** (1995), 69–92.
- [30] Bridget Eileen Tenner, Pattern avoidance and the Bruhat order, *J. Combin. Theory Ser. A* **114** (2007), 888–905.
- [31] Sherry H. F. Yan, Bijections for inversion sequences, ascent sequences and 3-nonnesting set partitions, preprint, 2017. Available at <https://arxiv.org/pdf/1707.02408.pdf>.

2010 *Mathematics Subject Classification*: Primary 05A05; Secondary 05A19.

Keywords: inversion sequence, pattern avoidance, triple of relations, Catalan number, Schröder number, Baxter number.

(Concerned with sequences [A000045](#), [A000071](#), [A000079](#), [A000108](#), [A000110](#), [A000111](#), [A000124](#), [A000325](#), [A000984](#), [A001181](#), [A001519](#), [A004275](#), [A005183](#), [A006318](#), [A009766](#), [A033321](#), [A034943](#), [A047969](#), [A047970](#), [A049125](#), [A071356](#), [A088921](#), [A090981](#), [A091156](#), [A098746](#), [A106228](#), [A108307](#), [A108759](#), [A113227](#), [A117106](#), [A124323](#), [A166073](#), [A175124](#), [A200753](#), [A212198](#), [A229046](#), [A263777](#), [A263778](#), [A263779](#), [A263780](#), [A279544](#), [A279551](#), [A279552](#), [A279553](#), [A279554](#), [A279555](#), [A279556](#), [A279557](#), [A279558](#), [A279559](#), [A279560](#), [A279561](#), [A279562](#), [A279563](#), [A279564](#), [A279565](#), [A279566](#), [A279567](#), [A279568](#), [A279569](#), [A279570](#), [A279571](#), [A279572](#), and [A279573](#).)

Received May 2 2017; revised version received January 5 2018; January 21 2018. Published in *Journal of Integer Sequences*, February 23 2018.

Return to [Journal of Integer Sequences home page](#).