

CONNECTIONS INDUCED BY (1,1)-TENSOR FIELDS ON
COTANGENT BUNDLES

ANTON DEKRÉT, Zvolen

(Received May 23, 1997)

Abstract. On cotangent bundles the Liouville field, the Liouville 1-form ε and the canonical symplectic structure $d\varepsilon$ exist. In this paper interactions between these objects and (1,1)-tensor fields on cotangent bundles are studied. Properties of the connections induced by the above structures are investigated.

Keywords: connection, almost complex structure, tensor field

MSC 1991: 58F05, 70H35, 58A20

INTRODUCTION

We assume that all manifolds and maps in this paper are infinitely differentiable.

Let M be a manifold and (x^i) a local chart on M . Let (x^i, z_i) be the induced chart on the cotangent bundle T^*M of all 1-forms on M . Let us recall that the Liouville vector field $V = z_i \partial / \partial z_i$ the flow of which is formed by the homotheties on individual fibres of T^*M , the Liouville 1-form $\varepsilon = z_i dx^i$, and the symplectic 2-form $\omega = d\varepsilon = dz_i \wedge dx^i$ on T^*M exist. Let F be a (1,1)-tensor field on M . It is known, [4], [7] that there is no connection on M , i.e. a linear connection on TM , which could be constructed by natural operators from F only. In other words no linear connection on T^*M can be constructed from natural lifts of F on T^*M only. We deal with the connections on the vector fibre bundle $\pi: T^*M \rightarrow M$ which are induced by (1,1)-tensor fields α on T^*M that are very close to the natural lifts of F on T^*M . We favour almost complex structures α (ACS). First of all, two cases are

Supported by the VEGA SR No. 1/1466/94

investigated: α preserves the vertical subbundle VT^*M or not. In the former case we deal with connections Γ_α when α preserves the horizontal subbundle $H\Gamma$ and in the latter with connections Γ_α for which $\alpha(VT^*M) = H\Gamma_\alpha^2$ or $\alpha(H\Gamma_\alpha^1) = VT^*M$.

The main results are in the third part of the paper. We deal with symmetric (1,1)-tensor fields α when the form $d\varepsilon^\alpha(X, Y) = d\varepsilon(\alpha X, Y)$ is symmetric, and with symmetric connections Γ on T^*M when $d\varepsilon|_{H\Gamma} = 0$, where $H\Gamma$ is the horizontal bundle of a connection Γ .

When α does not preserve VT^*M our investigations are concentrated on the semi-linear case of α when $B = T\pi \cdot \alpha|_{VT^*M}$ is a base morphism and the map $T\pi \cdot \alpha \cdot X : T^*M \rightarrow TM$ is a vector bundle morphism for any projectable linear vector field $X : T^*M \rightarrow TT^*M$ on T^*M . Propositions 5–7 determine sufficient conditions for the connection Γ_α^1 to be linear, for the equality $\gamma_\alpha^1 = TB(\Gamma_\alpha^1)$ and for the connection γ_α^1 to be just the Levi-Civita connection determined by the pseudo-Riemannian structure B^{-1} on M , where γ_Γ^1 is the connection on TM induced by the linear connection Γ_α^1 . Propositions 8 and 9 describe some properties of the ACS $\alpha(\Gamma, B)$ which are determined by a linear symmetric connection Γ on T^*M and by a vector bundle morphism B .

In the case $\alpha(VT^*M) \subset VT^*M$ there are morphisms $A = T\pi \cdot \alpha : TT^*M \rightarrow TM$, $H = \alpha|_{VT^*M}$. Remember that the complete lift $\alpha = F^C$ of a (1, 1)-tensor field F on M preserves VT^*M , $A = H$ and it is a VB -field, i.e. for any linear projectable vector field X on T^*M the vector field $\alpha(X)$ is again linear and projectable. When α is symmetric then $A = -H$. Propositions 12 and 13 state sufficient conditions under which a symmetric (1,1)-tensor field α (especially an ACS) determines connections Γ_α on T^*M .

Our investigations are local.

CONNECTIONS INDUCED BY (1,1)-AND (0,2)-TENSOR FIELDS ON FIBRE BUNDLES

Let $\pi : E \rightarrow M$ be a fibre bundle. Let (x^i, y^α) be a local fibre chart on E . A connection Γ on E can be regarded as a (1,1)-tensor field h_Γ on E (called the horizontal form of Γ) such that $h_\Gamma(VE) = 0$, $T\pi \cdot h_\Gamma = T\pi$, where VE is the vector fibre bundle of the vertical vectors on E and Tf denotes the tangent prolongation of a map f . In coordinates,

$$(1) \quad h_\Gamma = dx^i \otimes \partial/\partial x^i + \Gamma_j^\alpha(x, y) dx^j \otimes \partial/\partial y^\alpha.$$

Denote $H\Gamma := h_\Gamma(TE) \subset TE$ the vector fibre bundle of the Γ -horizontal vectors on E , i.e. such vectors $(x^i, y^\alpha, dx^i, dy^\alpha)$ on E which satisfy the equation

$$dy^\alpha = \Gamma_j^\alpha dx^j.$$

The functions Γ_j^α are called the local components of the connection Γ . Readers are referred to [6] for more details on connections on fibre bundles.

1. Let

$$\alpha = (a_j^i(x, y)dx^j + b_\alpha^i(x, y)dy^\alpha) \otimes \partial/\partial x^i + (c_j^\alpha(x, y)dx^j + h_\beta^\alpha(x, y)dy^\beta) \otimes \partial/\partial y^\alpha$$

be a (1,1)-tensor field on E . We will briefly denote by B the vector bundle morphism $T\pi \cdot \alpha|_{VE}: VE \rightarrow TM$ over $\pi: E \rightarrow M$. It can be interpreted as a section $E \rightarrow V^*E \otimes TM$, $B = b_\alpha^i dy^\alpha \otimes \partial/\partial x^i$.

We will say shortly that α is vertical if $B = 0$, i.e. if $\alpha(VE) \subset VE$. We will recall some properties of connections connected with a (1,1)-tensor field α , see [2].

Let h_Γ be the horizontal form of a connection Γ on E given by (1). Let $Y = \eta^\alpha \partial/\partial y^\alpha \in VE$ be an arbitrary vertical vector on E and let $X = \xi^i \partial/\partial x^i + \Gamma_i^\alpha \xi^i \partial/\partial y^\alpha$ be a Γ -horizontal vector. Then $\alpha(Y) = b_\beta^i \eta^\beta \partial/\partial x^i + h_\beta^\alpha \eta^\beta \partial/\partial y^\alpha$ or $\alpha(X) = (a_j^i + b_\beta^i \Gamma_j^\beta) \xi^j \partial/\partial x^i + (c_j^\alpha + h_\beta^\alpha \Gamma_j^\beta) \xi^j \partial/\partial y^\alpha$ is Γ -horizontal for any vertical vector Y or vertical for any Γ -horizontal vector X iff

$$(2) \quad \Gamma_k^\alpha b_\beta^k = h_\beta^\alpha \quad \text{or} \quad a_j^i + b_\beta^i \Gamma_j^\beta = 0.$$

We have

Lemma 1. *Let $\dim M$ be the dimension of the fibres on E . Then there is a unique connection Γ_α^1 and a unique connection Γ_α^2 on E such that $\alpha(H\Gamma_\alpha^1) = VE$, $\alpha(VE) = H\Gamma_\alpha^2$ if and only if the vector bundle morphism B is regular. Then $-\tilde{b}_k^\alpha a_j^k$ and $h_\beta^\alpha \tilde{b}_j^\beta$ are respectively the local components of Γ_α^1 or Γ_α^2 , and $\Gamma_\alpha^1 = \Gamma_\alpha^2$ if and only if α^2 is vertical.*

We will suppose that $\dim M$ is the dimension of fibres on E .

The coordinate conditions for α to be an almost complex structure on E , i.e. $\alpha^2 = -\text{Id}_{TE}$ are

$$(3) \quad a_s^i a_j^s + b_\beta^i c_j^\beta = -\delta_j^i, \quad a_s^i b_\beta^s + b_\gamma^i h_\beta^\gamma = 0, \quad c_s^\alpha a_j^s + h_\gamma^\alpha c_j^\gamma = 0, \quad c_s^\alpha b_\beta^s - h_\gamma^\alpha h_\beta^\gamma = -\delta_\beta^\alpha.$$

If B is regular then the third and fourth equations of (3) are consequences of the first and the second ones.

By the equalities (2) and (3) it is easy to prove

Lemma 2. *Let Γ be a connection on E . Let $B: E \rightarrow V^*E \otimes TM$ be a vector bundle isomorphism $VE \rightarrow TM$ over π . Then there exists a unique almost complex structure $\alpha(\Gamma, B)$ on E such that $T\pi \cdot \alpha|_{VE} = B$ and $\Gamma_\alpha^1 = \Gamma_\alpha^2 = \Gamma_\alpha$.*

In the case of $E = TM$, if we choose $B = \text{Id}|_{V_{TM}}$ then the almost complex structure $\alpha(\Gamma, B)$ is the canonical almost complex structure determined by the connection Γ on TM , see [5].

Let $Q = Q_j^\alpha dx^i \otimes \partial/\partial y^\alpha$ be a section $E \rightarrow T^*M \otimes VE$. Denote

$$\begin{aligned} \alpha^+ : Q &\rightarrow Q\alpha = (Q_k^\alpha a_j^k dx^j + Q_k^\alpha b_\beta^k dy^\beta) \otimes \partial/\partial y^\alpha, \quad \alpha^+ : T^*M \otimes VE \rightarrow T^*E \otimes VE \\ \alpha^- : Q &\rightarrow \alpha Q = b_\beta^i Q_j^\beta dx^j \otimes \partial/\partial x^i + h_\beta^\alpha Q_j^\beta dx^j \otimes \partial/\partial y^\alpha, \quad \alpha^- : T^*M \otimes VE \rightarrow T^*M \otimes TE. \end{aligned}$$

We say that two (1,1)-tensor fields α_1, α_2 on E are (+)-equivalent or (-)-equivalent if $\alpha_1^+ = \alpha_2^+$ or $\alpha_1^- = \alpha_2^-$ respectively for any $Q : E \rightarrow T^*M \otimes VE$.

If B is regular then using (3) we get ([2] Proposition 6)

Lemma 3. *In every class of all (+)-equivalent (1,1)-tensor fields on E there is a unique almost complex structure on E . The same is true for the class of (-)-equivalent (1,1)-tensor fields.*

In the case when α is vertical, i.e. when $B = 0$, denote

$$\begin{aligned} A &:= T\pi \cdot \alpha = a_j^i dx^j \otimes \partial/\partial x^i, \quad A : E \rightarrow T^*M \otimes_E TM, \\ H &: \alpha|_{V_Y} = h_\beta^\alpha dy^\beta \otimes \partial/\partial y^\alpha, \quad H : E \rightarrow V^*E \otimes VE. \end{aligned}$$

Let $\Gamma, \bar{\Gamma}$ be two connections on E and α a vertical (1,1)-tensor field. Then

$$(4) \quad \bar{\Gamma}_k^\alpha a_j^k = c_j^\alpha + h_\beta^\alpha \Gamma_j^\beta$$

is the coordinate condition for the inclusion $\alpha(H\Gamma) \subset H\bar{\Gamma}$.

Using (3) and (4) we get (see [2], Proposition 10)

Lemma 4. *Let $A : E \rightarrow T^*M \otimes_E TM$, $H : E \rightarrow V^*E \otimes VE$ be sections. Let Γ be a connection on E . Then there is a unique vertical (1,1)-tensor field $\alpha(A, H, \Gamma)$ such that $\alpha(H\Gamma) \subset H\Gamma$, $T\pi \cdot \alpha = A$, $\alpha|_{V_E} = H$. Moreover, if $A^2(u) = -\text{Id}_{T_{\pi u}M}$ for any $u \in E$ and $H^2 = -\text{Id}_{V_{TM}}$ then $\alpha(A, H, \Gamma)$ is an ACS on E .*

Remark. If A is regular then for any connection Γ there exists a unique connection $\bar{\Gamma}$ on E such that $\alpha(H\Gamma) \subset H\bar{\Gamma}$.

2. Let $\omega = f_{ij} dx^i \otimes dx^j + f_{i\alpha} dx^i \otimes dy^\alpha + f_{\alpha i} dy^\alpha \otimes dx^i + f_{\alpha\beta} dy^\alpha \otimes dy^\beta$ be a (0,2)-tensor field on E . Recall some well known facts.

Denote by ω^t the (0,2)-field transposed to ω , $\omega^t(X, Y) = \omega(Y, X)$. Then ω is symmetric or skew-symmetric if $\omega^t = \omega$ or $\omega^t = -\omega$, respectively.

Let α be a (1,1)-tensor field on E . Denote by $\omega^\alpha, \omega_\alpha, \omega\alpha$ the following (0,2)-tensor fields:

$$\omega^\alpha(X, Y) := \omega(\alpha X, Y), \quad \omega_\alpha(X, Y) := \omega(X, \alpha Y), \quad \omega\alpha(X, Y) := \omega(\alpha X, \alpha Y).$$

It is evident that

$$(5) \quad (\omega^\alpha)^t = (\omega^t)_\alpha, \quad (\omega_\alpha)^t = (\omega^t)^\alpha, \quad (\omega\alpha)^t = \omega^t\alpha$$

in the general case and

$$(6) \quad \omega^\alpha = \pm \omega_\alpha \Leftrightarrow \omega\alpha = \mp \omega, \quad \omega^\alpha\alpha = -\omega_\alpha, \quad \omega_\alpha\alpha = -\omega^\alpha$$

$$(7) \quad (\omega^\alpha)^t = \omega^\alpha \Leftrightarrow \omega\alpha = -\omega^t, \quad (\omega_\alpha)^t = -\omega^\alpha \Leftrightarrow \omega\alpha = \omega^t$$

in the case of an ACS α on E .

We say that a tangent subbundle $V_2 \subset TE$ is ω -orthogonal to a subbundle $V_1 \subset TE$ if $\omega(X, Y) = 0$ for any $X \in V_1(u), Y \in V_2(u)$ at any $u \in E$. A tangent subbundle $V \subset TE$ is said to be ω -zero if $\omega|_V = 0$.

Let Γ_1, Γ_2 be two connections on E . We say that Γ_2 is ω -orthogonal to Γ_1 if the Γ_2 -horizontal subbundle $H\Gamma_2$ is ω -orthogonal to $H\Gamma_1$.

If $X = \xi^i \partial / \partial x^i + \Gamma_j^\alpha \xi^j \partial / \partial y^\alpha$ is Γ_1 -horizontal and $\bar{X} = \bar{\xi}^i \partial / \partial x^i + \bar{\Gamma}_j^\alpha \bar{\xi}^j$ is Γ_2 -horizontal then $\omega(X, \bar{X}) = (f_{ij} + f_{i\alpha} \bar{\Gamma}_j^\alpha + f_{\alpha j} \Gamma_i^\alpha + f_{\alpha\beta} \Gamma_i^\alpha \bar{\Gamma}_j^\beta) \xi^i \bar{\xi}^j$ and so the connection Γ_2 is ω -orthogonal to Γ_1 if and only if

$$(8) \quad f_{ij} + f_{i\alpha} \bar{\Gamma}_j^\alpha + f_{\alpha j} \Gamma_i^\alpha + f_{\alpha\beta} \Gamma_i^\alpha \bar{\Gamma}_j^\beta = 0.$$

Consider the following restrictions of a (0,2)-tensor field ω :

$$\omega_1 := \omega|_{TE \times_E VE}, \quad \omega_1 = f_{i\alpha} dx^i \otimes dy^\alpha + f_{\alpha\beta} dy^\alpha \otimes dy^\beta,$$

$$\omega_2 := \omega|_{VE \times_E TE}, \quad \omega_2 = f_{\alpha i} dy^\alpha \otimes dx^i + f_{\alpha\beta} dy^\alpha \otimes dy^\beta,$$

$$\omega_v := \omega|_{VE \times_E VE}, \quad \omega_v = f_{\alpha\beta} dy^\alpha \otimes dy^\beta.$$

The equality (8) immediately gives

Lemma 5. *Let Γ_1, Γ_2 be two connections on a fibre bundle $\pi: E \rightarrow M$. Let $\varphi_1 = f_{i\alpha} dx^i \otimes dy^\alpha + f_{\alpha\beta} dy^\alpha \otimes dy^\beta$, $\varphi_1: E \rightarrow T^*E \otimes_E V^*E$, $\varphi_2 = f_{\alpha i} dy^\alpha \otimes dx^i + f_{\alpha\beta} dy^\alpha \otimes dy^\beta$, $\varphi_2: E \rightarrow V^*E \otimes_E T^*E$ be two bilinear forms such that $\varphi_1|_{VE \times_E VE} = \varphi_2|_{VE \times_E VE}$. Then there is a unique (0,2)-tensor field $\omega(\varphi_1, \varphi_2, \Gamma_1, \Gamma_2)$ on E such that $\omega_1 = \varphi_1$, $\omega_2 = \varphi_2$ and Γ_2 is ω -orthogonal to Γ_1 .*

If $\Gamma_2 = \Gamma_1 = \Gamma$ then the tensor field $\omega(\varphi_1, \varphi_2, \Gamma)$ from Lemma 5 is such that the connection Γ is ω -zero, i.e. $\omega|_{H\Gamma} = 0$.

We can find subbundles $\mathcal{H}, \mathcal{H}' \subset TE$ such that \mathcal{H} is ω -orthogonal to VE and VE is ω -orthogonal to \mathcal{H}' . Let $Y = \eta^\alpha \partial / \partial y^\alpha$ be vertical and let $X = dx^i \partial / \partial x^i + dy^\alpha \partial / \partial y^\alpha$ be an arbitrary tangent vector on E . Then the equation $\omega(Y, X) = 0$ or $\omega(X, Y) = 0$ is satisfied for any vertical vector Y iff

$$(8') \quad \begin{aligned} f_{\alpha i} dx^i + f_{\alpha \beta} dy^\beta &= 0 \quad \text{or} \\ f_{i\alpha} dx^i + f_{\beta\alpha} dy^\beta &= 0. \end{aligned}$$

This immediately gives

Lemma 6. *There exist unique connections $\Gamma_\omega, \Gamma'_\omega$ such that Γ_ω is ω -orthogonal to VE and VE is ω -orthogonal to Γ'_ω if and only if the form $\omega_v = \omega|_{VE}$ is regular. If ω is symmetric or skew-symmetric then $\Gamma_\omega = \Gamma'_\omega$. The vertical subbundle VE is ω -zero if and only if $\omega_v = 0$.*

In the following lemma we suppose that $\dim M$ is the dimension of fibres on E and that $B = T\pi \cdot \alpha|_{VE}$ is regular (so there exist connections $\Gamma_\alpha^1, \Gamma_\alpha^2, \alpha(H\Gamma_\alpha^1) = VE, \alpha(VE) = H\Gamma_\alpha^2$ on E).

Lemma 7. *Let ω be a $(0, 2)$ -tensor field and α a $(1, 1)$ -tensor field on E . Then*

1. VE is ω -zero iff VE is ω^α -orthogonal to Γ_α^1 or Γ_α^1 is ω_α -orthogonal to V or $H\Gamma_\alpha^1$ is $\omega\alpha$ -zero.
2. The connection Γ_α^1 is ω -orthogonal to VE iff Γ_α^1 is ω^α -zero.
3. VE is ω -orthogonal to the connection Γ_α^1 iff Γ_α^1 is ω_α -zero.
4. VE is ω^α -zero (ω_α -zero) iff Γ_α^1 is $\omega^\alpha\alpha$ -zero ($\omega_\alpha\alpha$ -zero).
5. VE is $\omega\alpha$ -zero iff the connection Γ_α^2 is ω -zero or $H\Gamma_\alpha^2$ is ω^α -orthogonal to VE or VE is ω_α -orthogonal to $H\Gamma_\alpha^2$.
6. VE is ω^α -zero iff VE is ω -orthogonal to Γ_α^2 .
7. VE is ω_α -zero iff Γ_α^2 is ω -orthogonal to VE .
8. The connection Γ_α^2 is ω^α -zero iff VE is $\omega\alpha$ -orthogonal to Γ_α^2 .
9. The connection Γ_α^2 is ω_α -zero iff Γ_α^2 is $\omega\alpha$ -orthogonal to VE .

In the induced chart (x^i, z_i) on T^*M a (1,1)-tensor field α on the fibre bundle $\pi: T^*M \rightarrow M$ is of the form

$$\alpha = (a_j^i(x, z)dx^j + b^{ij}dz_j) \otimes \partial/\partial x^i + (c_{ij}dx^j + h_i^j dz_j) \otimes \partial/\partial z_i.$$

According to the identification $VT^*M = T^*M \times_M T^*M$ the vector bundle morphism $B = T\pi \cdot \alpha|_{VT^*M}: VT^*M \rightarrow TM$, $B = b^{ij}dz_j \otimes \partial/\partial x^i$, can be interpreted as a vector bundle morphism $B: T^*M \times_M T^*M \rightarrow T^*M \times_M TM$, i.e. as a section $B: T^*M \rightarrow TM \otimes_{T^*M} T^*M$, i.e. as a bilinear form in VT^*M .

Definition 1. A (1,1)-tensor field α on T^*M is called v -symmetric or v -skew symmetric or v -basic if the section B is symmetric or skew symmetric or if B is the π -pullback of a section $M \rightarrow TM \otimes TM$.

Let us introduce the coordinate expression of some forms and tensor fields constructed from the Liouville form $\varepsilon = z_i dx^i$, $\omega \equiv d\varepsilon = dz_i \wedge dx^i$ and α :

$$\begin{aligned} i_\alpha \varepsilon &= z_k (a_j^k dx^j + b^{kj} dz_j), \\ i_\alpha d\varepsilon &= c_{ij} dx^j \wedge dx^i + (h_i^j + a_i^j) dx_j \wedge dx^i + b^{ij} dz_i \wedge dz_j, \\ i_\alpha d\varepsilon|_{VT^*M} &= b^{ij} dz_i \wedge dz_j, \end{aligned}$$

where i_α denotes the algebraic graded derivation determined by α ,

$$\begin{aligned} d\varepsilon^\alpha &= c_{ij} dx^i \otimes dx^j + h_i^j dz_j \otimes dx^i - a_i^j dx^i \otimes dz_j - b^{ij} dz_j \otimes dz_i, \\ d\varepsilon_\alpha &= -c_{ij} dx^i \otimes dx^j + a_i^j dx_j \otimes dx^i - h_i^j dx^i \otimes dz_j + b^{ji} dz_j \otimes dz_i, \\ d\varepsilon\alpha &= c_{ti} a_t^j dx^i \wedge dx^j + (c_{ti} b^{tj} - a_t^i h_t^j) dx^i \wedge dz_j + h_t^i b^{tj} dz_i \wedge dz_j. \end{aligned}$$

It is evident that

- (9) $d\varepsilon^\alpha + d\varepsilon_\alpha = i_\alpha d\varepsilon$,
- (10) $d\varepsilon^\alpha - d\varepsilon_\alpha = (c_{ji} + c_{ij})dx^i \otimes dx^j + (h_i^j - a_i^j)(dx^i \otimes dz_j + dz_j \otimes dx^i) - (b^{ij} + b^{ji})dz_i \otimes dx_j$ is symmetric,
- (11) $i_\alpha d\varepsilon$ is the antisymmetrization of $d\varepsilon^\alpha$,
- (12) $(d\varepsilon^\alpha)^t = -d\varepsilon_\alpha$, i.e. $(d\varepsilon)^\alpha$ is symmetric or exterior
iff $d\varepsilon^\alpha = -d\varepsilon_\alpha$ or $d\varepsilon^\alpha = d\varepsilon_\alpha$.

In coordinates $(d\varepsilon)^\alpha$ is symmetric or exterior if

$$\begin{aligned} c_{ij} = c_{ji}, \quad h_i^j = -a_i^j, \quad b^{ij} = b^{ji} \quad \text{or} \\ c_{ij} = -c_{ji}, \quad h_i^j = a_i^j, \quad b^{ij} = -b^{ji}. \end{aligned}$$

Proposition 1. *Let α be an ACS on T^*M . Then*

1. $(i_\alpha d\varepsilon)\alpha = -i_\alpha d\varepsilon$,
2. $(d\varepsilon^\alpha - d\varepsilon_\alpha)\alpha = d\varepsilon^\alpha - d\varepsilon_\alpha$,
3. $d\varepsilon^\alpha$ is symmetric or skew symmetric iff $d\varepsilon\alpha = d\varepsilon$ or $d\varepsilon\alpha = -d\varepsilon$, respectively,
4. $d\varepsilon^\alpha\alpha = d\varepsilon^\alpha$ if $d\varepsilon^\alpha$ is moreover symmetric.

Proof. 1 and 2 follow from (6) and (9). Assertion 3 is a consequence of (7). The equalities (6) and (12) imply 4. \square

Corollary. *Let α be an ACS on T^*M . Let $d\varepsilon^\alpha$ be symmetric, i.e. let $d\varepsilon$ be invariant under α . Then $d\varepsilon^\alpha$ is a pseudo-Hermite metric on T^*M and $(T^*M, \alpha, d\varepsilon^\alpha)$ is a pseudo-almost Kähler space, [7].*

Proof. By Proposition 1 $d\varepsilon^\alpha\alpha = d\varepsilon^\alpha$ so $d\varepsilon^\alpha$ is a pseudo-Hermite metric. As $(d\varepsilon^\alpha)^\alpha = -d\varepsilon$ is exact so $(T^*M, \alpha, d\varepsilon^\alpha)$ is pseudo-almost Kähler. \square

Definition 2. A (1,1)-tensor field α on T^*M is called symmetric or skew symmetric if $d\varepsilon^\alpha$ is symmetric or skew symmetric.

In the induced local chart (x^i, z_i) on T^*M a connection Γ on T^*M is given by the equations

$$dz_i = \Gamma_{ij}(x, z)dx^j.$$

As the form $d\varepsilon$ is symplectic, there exists a unique connection Γ^t which is $d\varepsilon$ -orthogonal to Γ . By (8) its local components are $\bar{\Gamma}_{ij} = \Gamma_{ji}$. So Γ is $d\varepsilon$ -zero iff $\Gamma^t = \Gamma$. In this case we will say that Γ is symmetric on T^*M .

Analogously in the case when the form $i_\alpha d\varepsilon$ is regular, i.e. almost-symplectic. Remember that if $d\varepsilon^\alpha$ is symmetric then $i_\alpha d\varepsilon = 0$.

In our further consideration we will deal with two cases of the (1,1)-tensor field α on T^*M .

I. Let α be such that $B = T\pi \cdot \alpha|_{VT^*M} = b^{ij}dz_j \otimes \partial/\partial x^i$ is regular. Then by Lemma 1 there are connections Γ_α^1 and Γ_α^2 such that $\alpha(H\Gamma_\alpha^1) = VT^*M$ and $\alpha(VT^*M) = H\Gamma_\alpha^2$. Then

$$(14) \quad \Gamma_{ij}^1 = -b_{ik}a_j^k, \quad \Gamma_{ij}^2 = h_i^k b_{kj}, \quad b_{ik}b^{kj} = \delta_i^j$$

are their local components.

By Lemma 6 there are two connections $\Gamma_{d\varepsilon^\alpha}, \Gamma'_{d\varepsilon^\alpha}$ such that $\Gamma_{d\varepsilon^\alpha}$ is $d\varepsilon^\alpha$ orthogonal to VT^*M and VT^*M is $d\varepsilon^\alpha$ -orthogonal to $\Gamma'_{d\varepsilon^\alpha}$. According to (8') their local components are respectively

$$\Gamma_{ij} = h_j^t b_{ti}, \quad \Gamma'_{ij} = -b_{it} a_j^t.$$

Comparing the local components of the connections $\Gamma_\alpha^1, \Gamma_\alpha^2, \Gamma_{d\varepsilon^\alpha}, \Gamma'_{d\varepsilon^\alpha}$ we get

Proposition 2. *If α is such a $(1, 1)$ -tensor field that B is regular then $d\varepsilon^\alpha|_{VT^*M}$ is also regular and*

- a) $\Gamma_\alpha^1 = \Gamma'_{d\varepsilon^\alpha}$ and so VT^*M is $d\varepsilon^\alpha$ -orthogonal to Γ_α^1 ,
- b) $\Gamma_{d\varepsilon^\alpha} = (\Gamma_\alpha^2)^t$, i.e. the connections $\Gamma_{d\varepsilon^\alpha}$ and Γ_α^2 are $d\varepsilon^\alpha$ -orthogonal and thus the connection $(\Gamma_\alpha^2)^t$ is $d\varepsilon^\alpha$ -orthogonal to VT^*M .

Remark. As $(d\varepsilon^\alpha)^t = -d\varepsilon_\alpha$ therefore $\Gamma_{d\varepsilon_\alpha} = \Gamma'_{d\varepsilon^\alpha}$ and $\Gamma'_{d\varepsilon_\alpha} = \Gamma_{d\varepsilon^\alpha}$. So the connection Γ_α^1 is $d\varepsilon_\alpha$ -orthogonal to VT^*M and VT^*M is $d\varepsilon_\alpha$ -orthogonal to the connection $(\Gamma_\alpha^2)^t$. By Lemma 7/1, Γ_α^1 is $d\varepsilon$ -zero because VT^*M is $d\varepsilon$ -zero.

Proposition 3. *If α is such a $(1, 1)$ -tensor field that B is regular, α^2 is vertical and $d\varepsilon^\alpha$ is symmetric or skew-symmetric then the connection $\Gamma_\alpha = \Gamma_\alpha^1 = \Gamma_\alpha^2$ is $d\varepsilon$ -zero, i.e. $(\Gamma_\alpha)^t = \Gamma_\alpha$.*

Proof. If $d\varepsilon^\alpha$ is symmetric then $b_{kj} = b_{jk}, -a_j^k = h_j^k$. When α^2 is vertical then $\Gamma_\alpha^1 = \Gamma_\alpha^2$, i.e. $-b_{ik}a_j^k = h_i^k b_{kj}$, i.e. $b_{ki}h_j^k = b_{kj}h_i^k$, i.e. $\Gamma_{ij} = \Gamma_{ji}$. Analogously when $-b_{kj} = b_{jk}, a_j^k = h_j^k$, i.e. when $d\varepsilon^\alpha$ is skew symmetric. \square

Let us remark that if α is an ACS then α^2 is vertical.

The inverse $B^{-1} = b_{ij}(x, z)dx^i \otimes \partial/\partial z_j: T^*M \times_M TM \rightarrow T^*M \times_M T^*M$ can be interpreted as a semibasic bilinear form $b_{ij}dx^i \otimes dx^j$ on T^*M , i.e. as a section $T^*M \rightarrow T^*M \otimes_{T^*M} T^*M$.

Proposition 4. *Let α be such an ACS on T^*M that B is regular and let h_{Γ_α} be the horizontal form of the connection $\Gamma_\alpha = \Gamma_\alpha^1 = \Gamma_\alpha^2$. Then $d\varepsilon^\alpha h_{\Gamma_\alpha} = -B^{-1}$ and $d\varepsilon_\alpha h_{\Gamma_\alpha} = (B^{-1})^t$.*

Proof. $h_{\Gamma_\alpha}(X_q) = \xi_q^i \partial/\partial x^i + h_i^s \tilde{b}_{si} \xi_q^j \partial/\partial z_j$, $q = 1, 2$. Then using (3) we get $d\varepsilon^\alpha(h_{\Gamma_\alpha} X_1, h_{\Gamma_\alpha} X_2) = (c_{ij} + h_i^t h_t^s b_{sj} - a_j^t h_t^s b_{si} - b^{ut} h_t^s b_{sj} h_u^r b_{ri}) \xi_1^j \xi_2^i = -b_{ij} \xi_1^j \xi_2^i$. This proves the first part. The second is a consequence of the equality $(d\omega^\alpha)^t = -d\omega_\alpha$. \square

Remark. As $B^{-1} = b_{ij}dx^i \otimes \partial/\partial z_j$ is a semibasic 1-form with values in VT^*M then if h_Γ is the horizontal form of a connection Γ on T^*M then $h_\Gamma + B^{-1}$ is the horizontal form of the other connection on T^*M .

Recall that a projectable linear vector field X on T^*M is such a vector field that $T\pi X$ is a vector field on M and its flow is formed by linear maps of fibres on T^*M , i.e. in coordinates $X = \xi^i(x) \partial/\partial x^i + \eta_i^j(x) z_j \partial/\partial z_i$.

Definition 3. A non-vertical (1,1)-tensor field α on T^*M is said to be semi-linear if it is v -basic and for any projectable linear vector field $X: T^*M \rightarrow TT^*M$ on T^*M the map $T\pi \cdot \alpha X: T^*M \rightarrow TM$ is a vector bundle morphism.

In a local chart it is easy to see that α is semi-linear iff $a_j^i(x, z) = a_j^{ik}(x)z_k$ and $b^{ij}(x, z) = b^{ij}(x)$.

Proposition 5. If a (1, 1)-tensor field α on T^*M is semi-linear and such that B is regular then the connection Γ_α^1 is linear.

Proof. The local components of Γ_α^1 are $\Gamma_{ij}^1 = -b_{is}a_j^s = -b_{is}(x)a_j^{sk}(x)z_k$, i.e. Γ_α^1 is linear. \square

Let us recall that every linear connection $\Gamma, \Gamma_{ij}^k = \Gamma_{ij}^k z_k$, on T^*M is induced by the linear connection γ on the tangent bundle TM with the local components $\gamma_i^k = -\Gamma_{ji}^k x_1^j$. The connection Γ is symmetric if and only if γ is symmetric.

So, if α is semi-linear and B is regular then the connection Γ_α^1 is induced by the connection γ_α^1 on TM with the local components $\gamma_j^i = \gamma_{jk}^i x_1^k$, $\gamma_{jk}^i = b_{ks}a_j^{si}$. As $B: T^*M \rightarrow TM$, $\bar{x}^i = x^i$, $\bar{x}_1^i = b^{ij}(x)z_j$, is a vector bundle isomorphism therefore $TB(\Gamma_\alpha^1)$ is a connection on TM . We find its functions.

$$\begin{aligned} TB: \bar{x}^i &= x^i, & x_1^i &= b^{ij}(x)z_j, & d\bar{x}^i &= dx^i, & dx_1^i &= b_k^{ij}z_j dx^k + b^{ij}dz_j, \\ h_{\Gamma_\alpha^1} &= dx^i \otimes \partial/\partial x^i - b_{is}a_j^{sk}z_k dx^j \otimes \partial/\partial z_i, \\ TB \cdot h_{\Gamma_\alpha^1} &= dx^i \otimes \partial/\partial x^i + (b_k^{ij} - b^{it}b_{ts}a_k^{sj})z_j dx^k \otimes \partial/\partial x^i. \end{aligned}$$

Then $\Gamma_j^i = (b_j^{is} - a_j^{is})b_{sk}x_1^k$ establish the components of the connection $TB(\Gamma_\alpha^1)$. Therefore $\gamma_\alpha^1 = TB(\Gamma_\alpha^1)$ if and only if

$$(15) \quad b_{ks}a_j^{si} = (b_j^{is} - a_j^{is})b_{sk}, \quad \text{i.e.} \quad b_{ku}a_j^{ui}b^{ks} = b_j^{is} - a_j^{is}.$$

Proposition 6. Let α be such a semilinear (1,1)-tensor field on T^*M that B is regular and symmetric, i.e. α is v -symmetric. Then $\gamma_\alpha^1 = TB(\Gamma_\alpha^1)$ if and only if $i_V d(i_\alpha \varepsilon) = 0$.

Proof.

$$\begin{aligned} \varepsilon &= z_k dx^k, & i_\alpha \varepsilon &= a_j^{kt}z_k z_t dx^j + b^{kj}z_k dz_j \\ di_\alpha \varepsilon &= a_{ji}^{kt}z_k z_t dx^i \wedge dx^j + a_j^{kt}(z_t dz_k - z_k dz_t) \wedge dx^j + b_i^{kj}z_k dx^i \wedge dz_j + b^{kj}dz_k \wedge dz_j \end{aligned}$$

where we use the notation $f_i := \frac{\partial f}{\partial x^i}$, $f^i := \frac{\partial f}{\partial z_i}$. Then

$$\begin{aligned} i_V di_\alpha \varepsilon &= (a_j^{kt} + a_j^{tk} - b_j^{kt})z_k z_t dx^j + (b^{kj} - b^{jk})z_k dz_j \\ &= (a_j^{kt} + a_j^{tk} - b_j^{kt})z_k z_t dx^j = 0 \end{aligned}$$

if and only if $b_j^{kt} = a_j^{kt} + a_j^{tk}$. When B is symmetric then (15) reads $a_j^{si} = b_j^{is} - a_j^{is}$. This completes our proof. \square

If $B: T^*M \rightarrow TM$ is regular and symmetric then $B^{-1} = b_{ij}(x)dx^i \otimes dx^j$ determines a pseudo-Riemannian structure on M the Levi-Civita connection γ_b of which is given by the local components $C_{jk}^i = -\frac{1}{2}b^{is}(b_{skj} + b_{sjk} - b_{jks})$.

Proposition 7. *Let α be such a semi-linear (1, 1)-tensor field α on T^*M that B is regular, $d\varepsilon^\alpha$ is symmetric, α^2 is vertical and $i_V d(i_\alpha \varepsilon) = 0$. Then the connection γ_α determined on TM by Γ_α is just the Levi-Civita connection γ_b of the pseudo-Riemannian structure on M induced by B^{-1} .*

Proof. By supposition $b^{ij} = b^{ji}$, $h_j^i = -a_j^{ik}z_k$, $a_s^{ij}b^{sk} = b^{is}a_s^{kj}$ or $b_{is}a_j^{st} = a_i^{st}b_{sj}$ and $b_j^{kt} = a_j^{kt} + a_j^{tk}$. Then

$$\begin{aligned} C_{jk}^i &= -\frac{1}{2}b^{is}(b_{skj} + b_{sjk} - b_{jks}) = \frac{1}{2}b_j^{is}b_{sk} + \frac{1}{2}b_k^{is}b_{sj} - \frac{1}{2}b^{is}b_{jt}b_s^{tu}b_{uk} \\ &= \frac{1}{2}(a_j^{is} + a_j^{si})b_{sk} + \frac{1}{2}(a_k^{is} + a_k^{si})b_{sj} - \frac{1}{2}b^{is}b_{jt}(a_s^{tu} + a_s^{ut})b_{uk} \\ &= \frac{1}{2}a_j^{is}b_{sk} + \frac{1}{2}b_{js}a_k^{si} + \frac{1}{2}a_k^{is}b_{sj} + \frac{1}{2}b_{ks}a_j^{si} - \frac{1}{2}a_j^{iu}b_{uk} - \frac{1}{2}a_k^{it}b_{jt} \\ &= \frac{1}{2}b_{js}a_k^{si} + \frac{1}{2}b_{ks}a_j^{si} = \frac{1}{2}a_j^{si}b_{sk} + \frac{1}{2}b_{ks}a_j^{si} = b_{ks}a_j^{si} = \gamma_{jk}^i. \end{aligned}$$

This completes our proof. \square

Recall in the sense of Lemma 2 that by $\alpha(\Gamma, B)$ we denote the almost complex structure α on T^*M determined uniquely by a connection Γ on T^*M and by a vector bundle isomorphism $B: VT^*M \rightarrow TM$ over $\pi: T^*M \rightarrow M$.

Proposition 8. *Let Γ be a symmetric connection on T^*M , i.e. $\Gamma^t = \Gamma$. Let $B: VT^*M \rightarrow TM$ be a symmetric vector bundle isomorphism. Then the almost complex structure $\alpha(\Gamma, B)$ is symmetric.*

Proof. Let $B = b^{ij}dz_j \otimes \partial/\partial x_1^i$, $b^{ij} = b^{ji}$, and $\Gamma_{ij} = \Gamma_{ji}$ be the components of Γ . Let α be such a ACS on T^*M that $T\pi\alpha|_{VT^*M} = B$ and $\Gamma_\alpha^1 = \Gamma$. Using (14) we get $a_j^i = -b^{is}\Gamma_{sj}$. Then by the second equality of (3)

$$h_j^i = -b_{jt}a_s^tb^{si} = b_{jt}b^{tu}\Gamma_{us}b^{si} = b^{is}\Gamma_{sj} = -a_j^i.$$

The first equality of (3) reads $c_{ij} = -b_{ij} - b_{it}a_s^ta_j^s = -b_{ij} - b^{sr}\Gamma_{si}\Gamma_{rj}$. So $c_{ij} = c_{ji}$. This completes our proof. \square

Proposition 9. *Let $b = b_{ij}dx^i \otimes dx^j$ be a symmetric and regular bilinear form on TM . Let Γ_b be the connection induced on T^*M by the Levi-Civita connection γ_b*

on TM established by the pseudo-Riemannian structure b . Let $B = b^{ij} dz_j \otimes \partial/\partial x^i$ be the inverse of b . Then the almost complex structure $\alpha(\Gamma_b, B)$ is symmetric and $i_V d(i_\alpha \varepsilon) = 0$.

Proof. The symmetry of $\alpha(\Gamma_b, B)$ is a consequence of Proposition 7 and the equality $i_V d(i_\alpha \varepsilon) = 0$ follows, according to Proposition 5, from the well known fact that $Tb(\gamma_b) = \Gamma_b$, where b is interpreted as a map $b: TM \rightarrow T^*M$. \square

It is easy to prove

Corollary. Let $J = dx^i \otimes \partial/\partial x^i$ be the almost tangent structure on TM (which can be identified with Id_{VTM}). Then the vector bundle isomorphism $b: TM \rightarrow T^*M$ is an almost complex map of the almost complex structures $\alpha(\gamma_b, J)$ and $\alpha(\Gamma_b, B)$ where we use the notation from Proposition 9.

We turn to the second case when B vanishes.

II. Let α be such a (1,1)-tensor field that $B = T\pi \cdot \alpha|_{VT^*M} = 0$.

Now,

$$\begin{aligned} A &:= T\pi \cdot \alpha = a_j^i dx^j \otimes \partial/\partial x^i: TT^*M \rightarrow TM, \\ H &:= \alpha|_{VT^*M} = h_i^j dz_j \otimes \partial/\partial z_i: VT^*M \rightarrow VT^*M. \end{aligned}$$

So A and H can be interpreted as sections $A: T^*M \rightarrow TM^* \otimes_{T^*M} TM$, $H: T^*M \rightarrow TM \otimes_{T^*M} T^*M$. If α is a VB -(1,1)-tensor field on T^*M , i.e. $\alpha(X)$ is a linear and projectable vector field on T^*M for any projectable and linear vector field X on T^*M , then A and H are the π -pull-backs of the sections $\bar{A}: M \rightarrow TM^* \otimes TM$, $\bar{A} = a_j^i(x) dx^j \otimes \partial/\partial x^i: TM \rightarrow TM$ and $\bar{H}: M \rightarrow TM \otimes T^*M$, $\bar{H} = h_i^j(x) \partial/\partial x^j \otimes dx^i: T^*M \rightarrow T^*M$. It is easy to see that in the VB -case $c_{ij}(x, z) = c_{ij}^k(x) z_k$, see [1].

Let $\bar{A}^*: T^*M \rightarrow T^*M$, $\bar{z}_i = a_i^j z_j$, denote the transposed vector bundle morphism to a vector bundle morphism $\bar{A}: TM \rightarrow TM$ over Id_M , $\bar{A}^*(\varepsilon)(X) = \varepsilon(\bar{A}X)$. If a VB -(1,1)-tensor field on T^*M is symmetric or skew symmetric then $\bar{H} = -\bar{A}^*$ or $\bar{H} = \bar{A}^*$, respectively.

Let $\Gamma, dz_i = \Gamma_{ij}(x, z) dx^j$, be a connection on T^*M . It is $d\varepsilon^\alpha$ -zero, i.e. $d\varepsilon^\alpha|_{H\Gamma} = 0$ if and only if

$$(16) \quad c_{ij} = \Gamma_{ti} a_j^t - \Gamma_{tj} h_i^t.$$

Proposition 10. Let Γ be a symmetric connection on T^*M . Then Γ is $d\varepsilon^\alpha$ -zero if and only if $\alpha(H\Gamma) \subset H\Gamma$.

P r o o f. The equalities (4) read

$$(4') \quad c_{ij} = \Gamma_{it}a_j^t - h_i^t\Gamma_{ij}.$$

Comparing (16) with (4') we get our assertion. \square

Proposition 11. *Let α be such a vertical (1,1)-tensor field that A is regular. Let a connection Γ be $d\varepsilon^\alpha$ -zero and $\alpha(H\Gamma) \subset H\Gamma$. Then Γ is symmetric.*

P r o o f. It follows from (16) and (4) that $(\Gamma_{it} - \Gamma_{ti})a_j^t = 0$, which completes our proof. \square

In the case of a vertical almost complex structure α on T^*M the formulas (3) read

$$(3') \quad a_s^i a_j^s = -\delta_j^i, \quad c_{it}a_j^t + h_i^t c_{tj} = 0, \quad h_s^i h_j^s = -\delta_j^i.$$

We suppose that A is the π -pull-back of a (1,1)-tensor \bar{A} on M . Denote by ω the exterior derivative of the form $i_\alpha \varepsilon = z_k a_j^k(x) dx^j$, i.e. $\omega := di_\alpha \varepsilon = a_j^k dz_k \wedge dx^j + z_k a_{jt}^k dx^i \wedge dx^j$. Then

$$\omega^\alpha = (a_j^t c_{ti} + z_k a_{jt}^k a_i^t - z_k a_{tj}^k a_i^t) dx^i \otimes dx^j + h_i^t a_j^t dz_i \otimes dx^j - a_i^t a_t^j dx^i \otimes dx_j.$$

Let Γ , $dz_i = \Gamma_{ij} dx^j$ be a connection on T^*M . Then

$$\omega^\alpha|_{H\Gamma} = (a_j^t c_{ti} + (a_{jt}^k - a_{tj}^k) z_k a_i^t + h_i^s a_j^t \Gamma_{si} - a_i^t a_t^s \Gamma_{sj}) dx^i \otimes dx^j.$$

If α is symmetric, A is an almost complex structure on M and Γ is symmetric, then $\omega^\alpha|_{H\Gamma} = 0$ if and only if

$$(17) \quad a_j^t c_{ti} + (a_{jt}^k - a_{tj}^k) z_k a_i^t + 2\Gamma_{ij} = 0.$$

Let us recall the Nijenhuis-Frölicher bracket, see for example [7],

$$[A, A] = (a_{jt}^k - a_{tj}^k) a_i^t dx^j \wedge dx^i \otimes \partial / \partial x^k.$$

We conclude

Proposition 12. *Let α be such a symmetric almost complex structure that A is an integrable almost complex structure on M , i.e. $[A, A] = 0$. Then there is a unique symmetric connection Γ_1 on T^*M such that $\omega^\alpha|_{H\Gamma_1} = 0$. If α moreover is a VB-tensor field then Γ_1 is linear.*

Let $\overline{H}: T^*M \rightarrow T^*M$ be a vector bundle isomorphism and Γ a linear connection on T^*M , $\overline{H} = h_i^j(x)\partial/\partial x^i \otimes dx_j$, $dz_i = \Gamma_{ij}^k(x)z_k dx^j$. Then

$$\begin{aligned} T\overline{H}: TT^*M &\rightarrow TT^*M; \quad \overline{x}^i = x^i, \quad \overline{z}_i = h_i^j z_j, \quad d\overline{x}^i = dx^i, \quad d\overline{z}_i = h_{ij}^t z_t dx^j + h_i^j dz_j, \\ T\overline{H} \cdot h_\Gamma &= dx^i \otimes \partial/\partial x^i + (h_{ij}^t z_t + h_i^t \Gamma_{tj}^k z_k) dx^j \otimes \partial/\partial z_i. \end{aligned}$$

Therefore $T\overline{H}(H\Gamma) \subset H\Gamma$ at any $(x^i, \overline{z}_i = h_i^t z_t)$ if and only if

$$(17') \quad \Gamma_{ij}^t h_t^k = h_{ij}^k + \Gamma_{tj}^k h_i^t.$$

Let α be such a symmetric (1,1)-tensor field on T^*M that $A = T\pi \cdot \alpha$ is the π -pull-back of a regular (1,1)-tensor field \overline{A} on M . Then $\overline{H} = -\overline{A}^*$. Let Γ be an arbitrary connection on T^*M . Denote $\beta := \overline{H}^* d\varepsilon = d\varepsilon T\overline{H} = h_{ij}^t z_t dx^j \wedge dx^i + h_i^j dz_j \wedge dx^i$. Then $\beta|_{H\Gamma} = (h_{ij}^t z_t + h_i^t \Gamma_{tj}^k) dx^j \wedge dx^i$ and $\beta|_{H\Gamma} = 0$ if and only if

$$(18) \quad (h_{ij}^t - h_{ji}^t) z_t + h_i^t \Gamma_{tj}^k - h_j^t \Gamma_{ti}^k = 0.$$

The equalities (16) and (18), using $a_j^i = -h_j^i$, give

$$(19) \quad c_{ij} + (h_{ij}^t - h_{ji}^t) z_t + 2h_i^t \Gamma_{tj}^k = 0, \quad \text{i.e.} \quad \Gamma_{ij}^k = -\frac{1}{2} \tilde{h}_i^t (c_{tj} + (h_{tj}^k - h_{jt}^k) z_k).$$

These functions Γ_{ij} satisfy (16) and (18).

We conclude

Proposition 13. *Let α be such a symmetric (1,1)-tensor field on T^*M that $A = T\pi\alpha$ is the π -pull-back of a regular (1,1)-tensor field \overline{A} on M . Then there exists a unique connection Γ_α such that $(d\varepsilon TH)|_{H\Gamma_\alpha} = 0$ and $d\varepsilon^\alpha|_{H\Gamma} = 0$. If moreover*

1. α is a VB-tensor field then Γ_α is linear,
2. α is such a VB-almost complex structure that the almost complex structure \overline{A} on M is integrable then $\Gamma_\alpha = \Gamma_1$.

Proof. The first part of Proposition 13 is evident. The equality of the functions of the connections Γ_1, Γ_α follows from the equalities (17), (19), (3') and $[\overline{A}, \overline{A}] = 0$. \square

Corollary. *If α is such a symmetric VB-almost complex structure that $[\overline{A}, \overline{A}] = 0$ then by Proposition 10, $\alpha(H\Gamma_\alpha) = H\Gamma_\alpha$.*

Remark. The connections Γ_α, Γ_1 cannot be constructed when $\overline{H} = \overline{A}^*$, for example when α is skew symmetric. Let us recall that if α is the so-called complete lift of a (1,1)-tensor field F on M then it is skew symmetric, see [8]. In a more general case when α is the first order natural lift of F then $\overline{H} = \overline{A}^*$, see [3], and so the connections Γ_α, Γ_1 do not exist.

Propositon 14. *Let α be such a symmetric VB-almost complex structure that $[\overline{A}, \overline{A}] = 0$. Then $T\overline{H}(H\Gamma_\alpha) \subset H\Gamma_\alpha$ if and only if the Nijenhuis tensor $[\alpha, \alpha]$ is semibasic with values in VT^*M .*

Proof. By virtue of $[\overline{A}, \overline{A}] = 0$ we get $\frac{1}{2}[\alpha, \alpha] = (B_{ij}^k dz_k \wedge dx^j + D_{kj}^i dx^k \wedge dx^j) \otimes \partial/\partial z_i$, where $B_{ij}^k = c_{ij}^u h_u^k + h_{ij}^u h_{uj}^k + h_{iu}^k h_j^u - h_i^u c_{uj}^k$. Using (3') we obtain $B_{ij}^k = (c_{ij}^u - h_{ij}^u)h_u^k + (h_{iu}^k - c_{iu}^k)h_j^u$. The relation $T\overline{H}(H\Gamma_\alpha) \subset H\Gamma_\alpha$ is true iff the functions Γ_{ij}^k established by (19) satisfy the equality (17'). Putting Γ_{ij}^k in (17') and using (3') we get $(c_{ij}^u - h_{ij}^u)h_u^k = h_j^u(c_{iu}^k - h_{iu}^k)$. So $T\overline{H}(H\Gamma_\alpha) \subset H\Gamma_\alpha$ if and only if $B_{ij}^k = 0$, i.e. iff $[\alpha, \alpha]$ is semibasic with values in VT^*M . The proof is complete. \square

Remark. It is easy to show that the condition $T\overline{H}(H\Gamma_\alpha) \subset H\Gamma_\alpha$ is equivalent to the one that $\nabla_\gamma A = 0$, i.e. the (1,1)-tensor field A on M is constant with respect to the covariant derivative established by the linear connection γ_α on TM which induces the connection Γ_α on T^*M .

References

- [1] Cabras, A., Kolář, I.: Special tangent valued forms and the Frölicher-Nijenhuis bracket. Arch. Math. (Brno) 29 (1993), 71–82.
- [2] Dekrét, A.: On almost complex structures on fibre bundles. Acta Univ. M. Belii Math. 3 (1995), 3–8.
- [3] Doupovec, M., Kurek, J.: Liftings of tensor fields to the cotangent bundle. Proc. Conf. Diff. Geometry Brno 1995. Masaryk University, Brno, 1996, p. 141–150.
- [4] Janyška, J.: Remarks on the Nijenhuis tensor and almost complex connections. Arch. Math. (Brno) 26 (1990), 229–240.
- [5] Klein, J.: On variational second order differential equations Polynomial case. Proc. Conf. Diff. Geometry and its Applications 1992. Silesian University, Opava, 1993, p. 449–459.
- [6] Kolář, I., Michor, P.W., Slovák, J.: Natural Operations in Differential Geometry. Springer-Verlag, Berlin, 1993.
- [7] Yano, K.: Differential Geometry on Complex and Almost Complex Spaces. Pergamon Press, New York, 1965.
- [8] Yano, K., Ishihara, S.: Tangent and Cotangent Bundles. M. Dekker Inc., New York, 1973.

Author's address: Anton Dekrét, Department of Mathematics, TU Zvolen, Masarykova 24, 960 53 Zvolen, Slovakia, e-mail: dekrret@vsld.tuzvo.sk.