

CARDINAL AND ORDINAL ARITHMETICS OF n -ARY
RELATIONAL SYSTEMS AND n -ARY ORDERED SETS

JIŘÍ KARÁSEK, Brno

(Received March 3, 1997)

Abstract. The aim of this paper is to define and study cardinal (direct) and ordinal operations of addition, multiplication, and exponentiation for n -ary relational systems. n -ary ordered sets are defined as special n -ary relational systems by means of properties that seem to suitably generalize reflexivity, antisymmetry, and transitivity from the case of $n = 2$ or 3 . The class of n -ary ordered sets is then closed under the cardinal and ordinal operations.

Keywords: n -ary relational system, n -ary ordered set, cardinal sum, cardinal product, cardinal power, ordinal sum, ordinal product, ordinal power

MSC 1991: 04A05, 06A99, 08A02

In what follows n always denotes an integer not less than 2; $(n]$ denotes the set $\{1, \dots, n\}$, P_n , O_n , E_n denotes the set of all, of all odd, of all even permutations of the set $(n]$, respectively. By an n -ary relational system we understand a set G together with a subset of G^n . More precisely, let G be a nonempty set. Then a subset $R \subseteq G^n$ is called an n -ary relation on G , and the ordered pair $\mathbb{G} = (G, R)$ is said to be an n -ary relational system. The set G is called the carrier of \mathbb{G} .

Birkhoff's arithmetic of ordered sets discussed in [1], [2] and [3] has been generalized by several authors—see e.g. [4], [5], [6], [7], [8]. Especially, in [5] V. Novák deals with direct (cardinal) operations of addition, multiplication, and exponentiation for n -ary relational structures. In [8], J. Šlapal generalizes Novák's results to relational systems of any (possibly infinite) arity. In this paper we return to finite arity, but introduce and study ordinal operations and n -ary ordered sets as well.

This research was supported by Grant Agency of the Czech Republic, grant no. 201/95/0468.

1. Definition. ([5], [7], [8]) Let $\mathbb{G} = (G, R)$, $\mathbb{H} = (H, S)$ be two n -ary relational systems. Let $\varphi: G \rightarrow H$ be a mapping such that, for any $(a_1, \dots, a_n) \in G^n$, $(a_1, \dots, a_n) \in R$ implies $(\varphi(a_1), \dots, \varphi(a_n)) \in S$. Then φ is called a homomorphism of \mathbb{G} into \mathbb{H} . The set of all homomorphisms of \mathbb{G} into \mathbb{H} is denoted by $\text{Hom}(\mathbb{G}, \mathbb{H})$. A homomorphism φ of \mathbb{G} into \mathbb{H} is called strong if, for any $(a_1, \dots, a_n) \in G^n$, $(a_1, \dots, a_n) \in R$ is equivalent to $(\varphi(a_1), \dots, \varphi(a_n)) \in S$. An injective strong homomorphism of \mathbb{G} into \mathbb{H} is called an embedding of \mathbb{G} into \mathbb{H} . A bijective strong homomorphism of \mathbb{G} onto \mathbb{H} is called an isomorphism of \mathbb{G} onto \mathbb{H} . We write $\mathbb{G} \sim \mathbb{H}$ and say that the n -ary relational systems \mathbb{G} and \mathbb{H} are isomorphic if there exists an isomorphism of \mathbb{G} onto \mathbb{H} .

Clearly, the class of all n -ary relational systems together with homomorphisms as morphisms forms a category.

2. Definition. Let $\mathbb{G} = (G, R)$ be an n -ary relational system. The n -ary relation R (and \mathbb{G} itself as well) is called:

- (1) discrete if $(a_1, \dots, a_n) \in R$ is equivalent to $a_1 = \dots = a_n$;
- (2) reflexive if $(a, a, \dots, a) \in R$ for any $a \in G$;
- (3) semisymmetric if $(a_1, \dots, a_n) \in R$, $\varepsilon \in E_n$ imply $(a_{\varepsilon(1)}, \dots, a_{\varepsilon(n)}) \in R$;
- (4) antisymmetric if $(a_1, \dots, a_n) \in R$, $(a_{\omega(1)}, \dots, a_{\omega(n)}) \in R$, $\omega \in O_n$ imply $a_1 = \dots = a_n$;
- (5) transitive if $(a_1, \dots, a_n) \in R$, $(a_1, \dots, a_{n-2}, a_n, a_{n+1}) \in R$ imply $(a_1, \dots, a_{n-1}, a_{n+1}) \in R$;
- (6) complete if for any pairwise different elements $a_1, \dots, a_n \in G$ there exists a $\pi \in P_n$ such that $(a_{\pi(1)}, \dots, a_{\pi(n)}) \in R$;
- (7) universal if $R = G^n$;
- (8) diagonal if $(a_{1j}, \dots, a_{nj}) \in R$ for any $j \in (n)$, $(a_{i1}, \dots, a_{in}) \in R$ for any $i \in (n)$ imply $(a_{11}, \dots, a_{nn}) \in R$.

3. Remark. Reflexivity is defined as in [7], transitivity, however, is weaker than n -transitivity in [7].

4. Definition. Let $\mathbb{G} = (G, R)$ be an n -ary relational system. The n -ary relation R is called an n -ary order if it is reflexive, semisymmetric, antisymmetric, and transitive; \mathbb{G} itself is then called an n -ary ordered set.

5. Remark. A binary order is clearly an order. A ternary order is a cyclic order in the sense of [6] enriched by the set of all triads consisting of identical elements.

6. Definition. ([5], [8]) Let $\mathbb{A} = (A, R)$, $\mathbb{B} = (B, S)$ be n -ary relational systems such that $A \cap B = \emptyset$. The cardinal (direct) sum $\mathbb{A} + \mathbb{B}$ of \mathbb{A} and \mathbb{B} is the n -ary relational system $\mathbb{C} = (C, T)$ where $C = A \cup B$ and $T = R \cup S$.

7. Theorem. *The cardinal sum of discrete, reflexive, semisymmetric, antisymmetric, transitive, or diagonal n -ary relational systems has the same property.*

Proof. It follows directly from the definition. Let us show, for example, the antisymmetry. Under the notation of 6, assume that $(a_1, \dots, a_n) \in T$, at least two of the elements a_1, \dots, a_n are different, and $\omega \in O_n$. We have either $(a_1, \dots, a_n) \in R$, or $(a_1, \dots, a_n) \in S$. In the first case $(a_{\omega(1)}, \dots, a_{\omega(n)}) \notin T$, for R is antisymmetric and $a_1, \dots, a_n \notin B$. Analogously in the second case. \square

8. Corollary. *The cardinal sum of n -ary ordered sets is an n -ary ordered set.*

9. Definition. ([5], [8]) Let $\mathbb{A} = (A, R)$, $\mathbb{B} = (B, S)$ be n -ary relational systems. The cardinal (direct) product $\mathbb{A}\mathbb{B}$ of \mathbb{A} and \mathbb{B} is the n -ary relational system $\mathbb{C} = (C, T)$ where $C = A \times B$ and T is the n -ary relation defined as follows: For any $(a_1, \dots, a_n) \in A^n$, $(b_1, \dots, b_n) \in B^n$, $((a_1, b_1), \dots, (a_n, b_n)) \in T \Leftrightarrow (a_1, \dots, a_n) \in R$, $(b_1, \dots, b_n) \in S$.

10. Theorem. *The cardinal product of discrete, reflexive, semisymmetric, antisymmetric, transitive, universal, or diagonal n -ary relational systems has the same property.*

Proof. We show the transitivity only, the other cases being similar. Under the notation of 9, assume that $((a_1, b_1), \dots, (a_n, b_n)) \in T$, $((a_1, b_1), \dots, (a_{n-2}, b_{n-2}), (a_n, b_n), (a_{n+1}, b_{n+1})) \in T$. Then $(a_1, \dots, a_n) \in R$, $(a_1, \dots, a_{n-2}, a_n, a_{n+1}) \in R$, $(b_1, \dots, b_n) \in S$, $(b_1, \dots, b_{n-2}, b_n, b_{n+1}) \in S$, so that $(a_1, \dots, a_{n-1}, a_{n+1}) \in R$, $(b_1, \dots, b_{n-1}, b_{n+1}) \in S$. This implies that $((a_1, b_1), \dots, (a_{n-1}, b_{n-1}), (a_{n+1}, b_{n+1}))$ belongs to T . \square

11. Corollary. *The cardinal product of n -ary ordered sets is an n -ary ordered set as well.*

12. Definition. ([5], [8]) Let $\mathbb{A} = (A, R)$, $\mathbb{B} = (B, S)$ be n -ary relational systems. The cardinal (direct) power $\mathbb{A}^{\mathbb{B}}$ with the base \mathbb{A} and exponent \mathbb{B} is the n -ary relational system $\mathbb{C} = (C, T)$ where $C = \text{Hom}(\mathbb{B}, \mathbb{A})$ and T is the n -ary relation defined as follows: For any $f_1, \dots, f_n \in C$,

$$(f_1, \dots, f_n) \in T \Leftrightarrow (f_1(x), \dots, f_n(x)) \in R \quad \text{for each } x \in B.$$

13. Theorem. *The cardinal power whose base is discrete, reflexive, semisymmetric, antisymmetric, transitive, universal, or diagonal has the same property.*

Proof. As before, we show one of the assertions only, namely the semisymmetry, under the notation of 12. Let $(f_1, \dots, f_n) \in T$ and let $\varepsilon \in E_n$. We have

$(f_1(x), \dots, f_n(x)) \in R$ for each $x \in B$, so that $(f_{\varepsilon(1)}(x), \dots, f_{\varepsilon(n)}(x)) \in R$ for each $x \in B$, and $(f_{\varepsilon(1)}, \dots, f_{\varepsilon(n)}) \in T$. \square

14. Corollary. *The cardinal power whose base is an n -ary ordered set is an n -ary ordered set as well.*

15. Definition. Let $\mathbb{A}_j = (A_j, R_j)$ be n -ary relational systems for each $j \in (n]$ such that $A_{j_1} \cap A_{j_2} = \emptyset$ whenever $j_1, j_2 \in (n], j_1 \neq j_2$. The ordinal sum $\mathbb{A}_1 \oplus \dots \oplus \mathbb{A}_n$ of $\mathbb{A}_1, \dots, \mathbb{A}_n$ is the n -ary relational system $\mathbb{B} = (B, S)$ where $B = \bigcup_{j=1}^n A_j$ and

$$S = \bigcup_{j=1}^n R_j \cup \bigcup_{\varepsilon \in E_n} A_{\varepsilon(1)} \times \dots \times A_{\varepsilon(n)}.$$

16. Theorem. *The ordinal sum of reflexive, semisymmetric, antisymmetric, or transitive n -ary relational systems has the same property.*

Proof. We limit ourselves to the case of transitivity under the notation of 15. Let $(a_1, \dots, a_n) \in S$ and $(a_1, \dots, a_{n-2}, a_n, a_{n+1}) \in S$.

(i) Let there exist a j_0 such that $a_1, \dots, a_n \in A_{j_0}$. If $n = 2$, then either $a_3 \in A_{j_0}$ and then $(a_1, a_3) \in R_{j_0} \subseteq S$, or $a_1, a_2 \in A_1, a_3 \in A_2$, for the identity is the only even permutation of $(2]$, and then $(a_1, a_3) \in A_1 \times A_2 \subseteq S$. Let $n \geq 3$. Then $a_{n+1} \in A_{j_0}$, for otherwise $(a_1, \dots, a_{n-2}, a_n, a_{n+1}) \notin S$, and $(a_1, \dots, a_n) \in R_{j_0}$, $(a_1, \dots, a_{n-2}, a_n, a_{n+1}) \in R_{j_0}$, so that $(a_1, \dots, a_{n-1}, a_{n+1}) \in R_{j_0} \subseteq S$.

(ii) Let there exist no j_0 such that $a_1, \dots, a_n \in A_{j_0}$. If $n = 2$, then $a_1 \in A_1, a_2 \in A_2$, hence $a_3 \in A_2$ and $(a_1, a_3) \in A_1 \times A_2 \subseteq S$. Admit that $n \geq 3$. We have $(a_1, \dots, a_n) \in \bigcup_{\varepsilon \in E_n} A_{\varepsilon(1)} \times \dots \times A_{\varepsilon(n)}$, therefore $a_j \in A_{\varepsilon(j)}$ for each $j \in (n]$ where $\varepsilon \in E_n$. There exists no j_0 such that $a_1, \dots, a_{n-2}, a_n, a_{n+1} \in A_{j_0}$, so that $(a_1, \dots, a_{n-2}, a_n, a_{n+1}) \in A_{\xi(1)} \times \dots \times A_{\xi(n)}$ for some $\xi \in E_n$. Thus $a_i \in A_{\xi(i)}$ for each $i \in (n-2]$, $a_n \in A_{\xi(n-1)}, a_{n+1} \in A_{\xi(n)}$. This implies that $\varepsilon(i) = \xi(i)$ for each $i \in (n-2]$, $\varepsilon(n) = \xi(n-1), \varepsilon(n-1) = \xi(n)$, which is a contradiction, for both ε and ξ are even. \square

17. Remark. In contrast to the binary case, the ordinal sum of complete n -ary relational systems need not be complete if $n \geq 3$. Let, for example, $A_1 = \{a, b, c\}, A_2 = \{d, e, f\}, A_3 = \{g, h, i\}, R_1 = \{(a, b, c)\}, R_2 = \{(d, e, f)\}, R_3 = \{(g, h, i)\}, \mathbb{A}_j = (A_j, R_j)$ for $j = 1, 2, 3$. Then each of the ternary relational systems $\mathbb{A}_1, \mathbb{A}_2, \mathbb{A}_3$ is complete, but their ordinal sum $\mathbb{A}_1 \oplus \mathbb{A}_2 \oplus \mathbb{A}_3$ is not, because $S = R_1 \cup R_2 \cup R_3 \cup A_1 \times A_2 \times A_3 \cup A_2 \times A_3 \times A_1 \cup A_3 \times A_1 \times A_2$ and no permutation of the ordered triad (a, b, d) belongs to S .

18. Corollary. *The ordinal sum of n -ary ordered sets is an n -ary ordered set.*

19. Definition. Let $\mathbb{A} = (A, R)$, $\mathbb{B} = (B, S)$ be n -ary relational systems. The ordinal product $\mathbb{A} \circ \mathbb{B}$ of \mathbb{A} and \mathbb{B} is the n -ary relational system $\mathbb{C} = (C, T)$ where $C = A \times B$ and T is the n -ary relation defined as follows: For any $(a_1, \dots, a_n) \in A^n$ and $(b_1, \dots, b_n) \in B^n$, $((a_1, b_1), \dots, (a_n, b_n)) \in T \Leftrightarrow$ at least two of the elements a_1, \dots, a_n are different, $(a_1, \dots, a_n) \in R$, or $a_1 = \dots = a_n$, $(b_1, \dots, b_n) \in S$.

20. Theorem. *The ordinal product of discrete, reflexive, semisymmetric, antisymmetric, transitive and antisymmetric, or universal n -ary relational systems has the same property (properties).*

Proof. We show the fourth and fifth cases only. Assume the notation of 19.

(i) Antisymmetry: Let $((a_1, b_1), \dots, (a_n, b_n)) \in T$, let at least two of the elements $(a_1, b_1), \dots, (a_n, b_n)$ be different, and let $\omega \in O_n$. If at least two of the elements a_1, \dots, a_n are different, then $(a_1, \dots, a_n) \in R$, consequently $(a_{\omega(1)}, \dots, a_{\omega(n)}) \notin R$ and at least two of the elements $a_{\omega(1)}, \dots, a_{\omega(n)}$ are different, thus $((a_{\omega(1)}, b_{\omega(1)}), \dots, (a_{\omega(n)}, b_{\omega(n)})) \notin T$. Let $a_1 = \dots = a_n$. Then $(b_1, \dots, b_n) \in S$ and at least two of the elements b_1, \dots, b_n are different, hence again $((a_{\omega(1)}, b_{\omega(1)}), \dots, (a_{\omega(n)}, b_{\omega(n)})) \notin T$.

(ii) Transitivity and antisymmetry: Antisymmetry follows from (i). Let $((a_1, b_1), \dots, (a_n, b_n)) \in T$, $((a_1, b_1), \dots, (a_{n-2}, b_{n-2}), (a_n, b_n), (a_{n+1}, b_{n+1})) \in T$. First of all, let at least two of the elements a_1, \dots, a_n be different, and let at least two of the elements $a_1, \dots, a_{n-2}, a_n, a_{n+1}$ be different. Then $(a_1, \dots, a_n) \in R$, $(a_1, \dots, a_{n-2}, a_n, a_{n+1}) \in R$, consequently $(a_1, \dots, a_{n-1}, a_{n+1}) \in R$. If at least two of the elements $a_1, \dots, a_{n-1}, a_{n+1}$ are different, then $((a_1, b_1), \dots, (a_{n-1}, b_{n-1}), (a_{n+1}, b_{n+1})) \in T$. Admit that $a_1 = \dots = a_{n-1} = a_{n+1}$. Then $(a_1, \dots, a_1, a_n) \in R$, $(a_1, \dots, a_1, a_n, a_1) \in R$, $a_1 \neq a_n$, which is a contradiction to the antisymmetry of R . Let $a_1 = \dots = a_n = a_{n+1}$. Then $(b_1, \dots, b_n) \in S$, $(b_1, \dots, b_{n-2}, b_n, b_{n+1}) \in S$, so that $(b_1, \dots, b_{n-1}, b_{n+1}) \in S$ and $((a_1, b_1), \dots, (a_{n-1}, b_{n-1}), (a_{n+1}, b_{n+1})) \in T$. Let $a_1 = \dots = a_n \neq a_{n+1}$. Then $(a_1, \dots, a_1, a_{n+1}) \in R$, $a_1 \neq a_{n+1}$, so that $((a_1, b_1), \dots, (a_{n-1}, b_{n-1}), (a_{n+1}, b_{n+1})) \in T$. Finally, let $a_1 = \dots = a_{n-2} = a_n = a_{n+1} \neq a_{n-1}$. Then $(a_1, \dots, a_1, a_{n-1}, a_1) \in R$, $a_1 \neq a_{n-1}$, so that $((a_1, b_1), \dots, (a_{n-1}, b_{n-1}), (a_{n+1}, b_{n+1})) \in T$. \square

21. Remark. (i) For reflexivity of T , it is sufficient to assume reflexivity of S .

(ii) For transitivity of T , it is sufficient to assume transitivity of R and S and antisymmetry of R .

(iii) In contrast to the binary case, the ordinal product of complete n -ary relational systems need not be complete if $n \geq 3$. Let, for example, $A = \{a, b, c\}$, $B = \{d, e, f\}$, $R = \{(a, b, c)\}$, $S = \{(d, e, f)\}$, $\mathbb{A} = (A, R)$, $\mathbb{B} = (B, S)$. Then both \mathbb{A} and \mathbb{B} are

complete, but their ordinal product $\mathbb{A} \circ \mathbb{B}$ is not, because no permutation of the ordered triad $((a, d), (a, e), (b, f))$ belongs to T .

22. Corollary. *The ordinal product of n -ary ordered sets is an n -ary ordered set.*

23. Definition. Let $\mathbb{A} = (A, R)$ be a binary relational system, $\mathbb{B} = (B, S)$ an n -ary relational system. The ordinal power ${}^{\mathbb{A}}\mathbb{B}$ with the base \mathbb{B} and exponent \mathbb{A} is the n -ary relational system $\mathbb{C} = (C, T)$ where C is the set B^A of all mappings of A into B and T is the n -ary relation defined as follows: For any $f_1, \dots, f_n \in C$, $(f_1, \dots, f_n) \in T \Leftrightarrow$ for any $a \in A$, $(f_1(a), \dots, f_n(a)) \notin S$ there exists a $b \in A$ such that $(b, a) \in R$, $(f_1(b), \dots, f_n(b)) \in S$, at least two of the elements $f_1(b), \dots, f_n(b)$ are different, and $(f_1(x), \dots, f_n(x)) \in S$ for each $x \in A$, $(x, b) \in R$.

24. Theorem. *The ordinal power whose base is a discrete, reflexive, semisymmetric, antisymmetric, or universal n -ary relational system has the same property. The ordinal power whose base is transitive and antisymmetric and whose exponent is transitive is transitive as well.*

Proof. Let us show antisymmetry and transitivity only. Assume the notation of 23.

- (i) Antisymmetry: Let $(f_1, \dots, f_n) \in T$, $(f_{\omega(1)}, \dots, f_{\omega(n)}) \in T$ for some $\omega \in O_n$. Admit that there exists an $a \in A$ for which $(f_1(a), \dots, f_n(a)) \notin S$. Then there exists a $b \in A$ such that $(b, a) \in R$, $(f_1(b), \dots, f_n(b)) \in S$, at least two of the elements $f_1(b), \dots, f_n(b)$ are different. Consequently $(f_{\omega(1)}(b), \dots, f_{\omega(n)}(b)) \notin S$. There exists again a $c \in A$ such that $(c, b) \in R$, $(f_{\omega(1)}(c), \dots, f_{\omega(n)}(c)) \in S$, at least two of the elements $f_{\omega(1)}(c), \dots, f_{\omega(n)}(c)$ are different. But $\omega^{-1} \in O_n$ as well, so that $(f_1(c), \dots, f_n(c)) \notin S$. For each $x \in A$ such that $(x, b) \in R$ we have, however, $(f_1(x), \dots, f_n(x)) \in S$, which is a contradiction. Hence there is no $a \in A$ such that $(f_1(a), \dots, f_n(a)) \notin S$. Consequently $(f_1(a), \dots, f_n(a)) \in S$ for each $a \in A$. Similarly $(f_{\omega(1)}(a), \dots, f_{\omega(n)}(a)) \in S$ for each $a \in A$. As S is antisymmetric, we have $f_1(a) = \dots = f_n(a)$ for each $a \in A$, thus $f_1 = \dots = f_n$ and T is antisymmetric.
- (ii) Transitivity: Let $(f_1, \dots, f_n) \in T$, $(f_1, \dots, f_{n-2}, f_n, f_{n+1}) \in T$, and let $a \in A$ be such that $(f_1(a), \dots, f_{n-1}(a), f_{n+1}(a)) \notin S$. Then either $(f_1(a), \dots, f_n(a)) \notin S$ or $(f_1(a), \dots, f_{n-2}(a), f_n(a), f_{n+1}(a)) \notin S$ because of transitivity of S . Let the first case occur. As $(f_1, \dots, f_n) \in T$, there exists a $b \in A$ such that $(b, a) \in R$, $(f_1(b), \dots, f_n(b)) \in S$, at least two of the elements $f_1(b), \dots, f_n(b)$ are different, and for each $x \in A$, $(x, b) \in R$ we have $(f_1(x), \dots, f_n(x)) \in S$. Now, four possibilities (a), (b), (c), (d) can occur:
- (a) $(f_1(b), \dots, f_{n-2}(b), f_n(b), f_{n+1}(b)) \in S$, at least two of the elements $f_1(b), \dots, f_{n-2}(b), f_n(b), f_{n+1}(b)$ are different, and $(f_1(x), \dots, f_{n-2}(x), f_n(x), f_{n+1}(x)) \in S$ for

each $x \in A$, $(x, b) \in R$. Considering transitivity of S , we have $(f_1(b), \dots, f_{n-1}(b), f_{n+1}(b)) \in S$, and $(f_1(x), \dots, f_{n-1}(x), f_{n+1}(x)) \in S$ for each $x \in A$, $(x, b) \in R$. If at least two of the elements $f_1(b), \dots, f_{n-1}(b), f_{n+1}(b)$ are different, we have $(f_1, \dots, f_{n-1}, f_{n+1}) \in T$. Admit that $f_1(b) = \dots = f_{n-1}(b) = f_{n+1}(b)$. Then $f_n(b) \neq f_j(b)$ for each $j = 1, \dots, n-1, n+1$. Hence $(f_1(b), \dots, f_n(b)) = (f_1(b), \dots, f_1(b), f_n(b)) \in S$, $(f_1(b), \dots, f_{n-2}(b), f_n(b), f_{n+1}(b)) = (f_1(b), \dots, f_1(b), f_n(b), f_1(b)) \in S$, a contradiction to antisymmetry of S .

(b) $f_1(b) = \dots = f_{n-2}(b) = f_n(b) = f_{n+1}(b)$, $(f_1(x), \dots, f_{n-2}(x), f_n(x), f_{n+1}(x)) \in S$ for each $x \in A$, $(x, b) \in R$. Then $(f_1(b), \dots, f_{n-1}(b), f_{n+1}(b)) = (f_1(b), \dots, f_n(b)) \in S$, at least two of the elements $f_1(b), \dots, f_{n-1}(b), f_{n+1}(b)$ are different, $(f_1(x), \dots, f_{n-1}(x), f_{n+1}(x)) \in S$ for each $x \in A$, $(x, b) \in R$ due to transitivity of S , so that $(f_1, \dots, f_{n-1}, f_{n+1}) \in T$.

(c) $(f_1(b), \dots, f_{n-2}(b), f_n(b), f_{n+1}(b)) \notin S$. As $(f_1, \dots, f_{n-2}, f_n, f_{n+1}) \in T$, there exists a $c \in A$, $(c, b) \in R$ such that $(f_1(c), \dots, f_{n-2}(c), f_n(c), f_{n+1}(c)) \in S$, at least two of the elements $f_1(c), \dots, f_{n-2}(c), f_n(c), f_{n+1}(c)$ are different, and $(f_1(x), \dots, f_{n-2}(x), f_n(x), f_{n+1}(x)) \in S$ for each $x \in A$, $(x, c) \in R$. Considering transitivity of R we have $(c, a) \in R$. By the preceding, we have $(f_1(c), \dots, f_n(c)) \in S$ (for $(c, b) \in R$) and $(f_1(x), \dots, f_n(x)) \in S$ for each $x \in A$, $(x, c) \in R$ (by transitivity of R), consequently, regarding transitivity of S , $(f_1(c), \dots, f_{n-1}(c), f_{n+1}(c)) \in S$ and $(f_1(x), \dots, f_{n-1}(x), f_{n+1}(x)) \in S$ for each $x \in A$, $(x, c) \in R$. If at least two of the elements $f_1(c), \dots, f_{n-1}(c), f_{n+1}(c)$ are different, we are done. Admit that $f_1(c) = \dots = f_{n-1}(c) = f_{n+1}(c)$. Then $(f_1(c), \dots, f_n(c)) = (f_1(c), \dots, f_1(c), f_n(c)) \in S$, $(f_1(c), \dots, f_{n-2}(c), f_n(c), f_{n+1}(c)) = (f_1(c), \dots, f_1(c), f_n(c), f_1(c)) \in S$, $f_1(c) \neq f_n(c)$, which is a contradiction to antisymmetry of S .

(d) $(f_1(b), \dots, f_{n-2}(b), f_n(b), f_{n+1}(b)) \in S$, there exists an $x \in A$, $(x, b) \in R$ such that $(f_1(x), \dots, f_{n-2}(x), f_n(x), f_{n+1}(x)) \notin S$. As $(f_1, \dots, f_{n-2}, f_n, f_{n+1}) \in T$, there exists a $c \in A$, $(c, x) \in R$ such that $(f_1(c), \dots, f_{n-2}(c), f_n(c), f_{n+1}(c)) \in S$, at least two of the elements $f_1(c), \dots, f_{n-2}(c), f_n(c), f_{n+1}(c)$ are different, and $(f_1(y), \dots, f_{n-2}(y), f_n(y), f_{n+1}(y)) \in S$ for each $y \in A$, $(y, c) \in R$. Considering transitivity of R we have $(c, a) \in R$. By the preceding, we have $(f_1(c), \dots, f_n(c)) \in S$ (for $(c, b) \in R$) and $(f_1(y), \dots, f_n(y)) \in S$ for each $y \in A$, $(y, c) \in R$. Regarding transitivity of S , we obtain $(f_1(c), \dots, f_{n-1}(c), f_{n+1}(c)) \in S$ and $(f_1(y), \dots, f_{n-1}(y), f_{n+1}(y)) \in S$ for each $y \in A$, $(y, c) \in R$. If at least two of the elements $f_1(c), \dots, f_{n-1}(c), f_{n+1}(c)$ are different, we are done. The case of $f_1(c) = \dots = f_{n-1}(c) = f_{n+1}(c)$, however, leads to a contradiction to antisymmetry of S similarly as in (c). The case of $(f_1(a), \dots, f_{n-2}(a), f_n(a), f_{n+1}(a)) \notin S$ is analogous. \square

25. Corollary. *The ordinal power whose base is an n -ary ordered set and whose exponent is a transitive binary relational system is an n -ary ordered set.*

26. Definition. Let $\mathbb{A} = (A, R)$ be an n -ary relational system. The dual n -ary relational system $\tilde{\mathbb{A}}$ to \mathbb{A} is the n -ary relational system $\mathbb{B} = (A, S)$ where S is the n -ary relation defined as follows: For any $(a_1, \dots, a_n) \in A^n$,

$$(a_1, \dots, a_n) \in S \Leftrightarrow (a_n, \dots, a_1) \in R.$$

27. Theorem. *The dual n -ary relational system to a discrete, reflexive, semisymmetric, antisymmetric, semisymmetric and transitive, complete, universal, or diagonal n -ary relational system has the same property (properties).*

Proof. For brevity, we show the third and the fifth property only. Assume the notation of 26.

(i) Semisymmetry: Let $(a_1, \dots, a_n) \in S$, $\varepsilon \in E_n$. Then $(a_n, \dots, a_1) \in R$, thus $(a_{n+1-\varepsilon(1)}, \dots, a_{n+1-\varepsilon(n)}) \in R$, so that $(a_{\varepsilon(1)}, \dots, a_{\varepsilon(n)}) \in S$.

(ii) Semisymmetry and transitivity: Semisymmetry follows from (i). Let $(a_1, \dots, a_n) \in S$, $(a_1, \dots, a_{n-2}, a_n, a_{n+1}) \in S$. Then $(a_n, \dots, a_1) \in R$, $(a_{n+1}, a_n, a_{n-2}, \dots, a_1) \in R$. The case of $n = 2$ is evident. Let $n \geq 3$. Clearly $(3, 4, \dots, n, 1, 2) \in E_n$. Considering semisymmetry of R , we have $(a_{n-2}, \dots, a_1, a_n, a_{n-1}) \in R$, $(a_{n-2}, \dots, a_1, a_{n+1}, a_n) \in R$. Due to transitivity of R , it follows that $(a_{n-2}, \dots, a_1, a_{n+1}, a_{n-1}) \in R$, so that, by semisymmetry of R , $(a_{n+1}, a_{n-1}, \dots, a_1) \in R$. Hence $(a_1, \dots, a_{n-1}, a_{n+1}) \in S$. \square

28. Corollary. *The dual to a (complete) n -ary ordered set is a (complete) n -ary ordered set.*

29. Theorem. *For any n -ary relational systems \mathbb{A}, \mathbb{B} with disjoint carriers and for any n -ary relational system \mathbb{C} we have*

$$\begin{aligned} \mathbb{A} + \mathbb{B} &= \mathbb{B} + \mathbb{A}, \\ (\mathbb{A} + \mathbb{B})\mathbb{C} &= \mathbb{A}\mathbb{C} + \mathbb{B}\mathbb{C}, \\ \mathbb{C}(\mathbb{A} + \mathbb{B}) &= \mathbb{C}\mathbb{A} + \mathbb{C}\mathbb{B}. \end{aligned}$$

Proof. The first equality is trivial. The second and third are special cases of Theorem 4 in [8]. \square

30. Theorem. *For any n -ary relational systems $\mathbb{A}, \mathbb{B}, \mathbb{C}$ with pairwise disjoint carriers we have*

$$\mathbb{A} + (\mathbb{B} + \mathbb{C}) = (\mathbb{A} + \mathbb{B}) + \mathbb{C}.$$

Proof. It follows from [8], Theorem 1. \square

31. Theorem. For any n -ary relational systems $\mathbb{A}, \mathbb{B}, \mathbb{C}$ we have

$$\begin{aligned}\mathbb{A}\mathbb{B} &\sim \mathbb{B}\mathbb{A}, \\ \mathbb{A}(\mathbb{B}\mathbb{C}) &\sim (\mathbb{A}\mathbb{B})\mathbb{C}.\end{aligned}$$

Proof. In the first case, under the notation of 9, it suffices to take the natural bijection $\varphi: A \times B \rightarrow B \times A$ defined by $\varphi((a, b)) = (b, a)$ for any $a \in A, b \in B$; φ is clearly an isomorphism of $\mathbb{A}\mathbb{B}$ onto $\mathbb{B}\mathbb{A}$. The second case follows from [8], Theorem 2. \square

32. Theorem. For any n -ary relational systems \mathbb{B}, \mathbb{C} with disjoint carriers and for any n -ary relational system \mathbb{A} we have

$$\mathbb{A}^{\mathbb{B}+\mathbb{C}} \sim \mathbb{A}^{\mathbb{B}} \mathbb{A}^{\mathbb{C}}.$$

Proof. It follows from [8], Theorem 8. \square

33. Theorem. For any n -ary relational systems $\mathbb{A}, \mathbb{B}, \mathbb{C}$ we have

$$(\mathbb{A}\mathbb{B})^{\mathbb{C}} \sim \mathbb{A}^{\mathbb{C}} \mathbb{B}^{\mathbb{C}}.$$

Proof. It follows from [8], Theorem 7. \square

34. Theorem. Let $\mathbb{A}, \mathbb{B}, \mathbb{C}$ be n -ary relational systems such that \mathbb{A} is universal or \mathbb{A} is reflexive and both \mathbb{B} and \mathbb{C} are discrete. Then

$$\mathbb{A}^{\mathbb{B}\mathbb{C}} \sim (\mathbb{A}^{\mathbb{B}})^{\mathbb{C}}.$$

Proof. It follows from [8], Corollary of Theorem 5. \square

35. Theorem. Let $\mathbb{A}, \mathbb{B}, \mathbb{C}$ be n -ary relational systems such that \mathbb{B} and \mathbb{C} are reflexive. Then there exists an embedding of $\mathbb{A}^{\mathbb{B}\mathbb{C}}$ into $(\mathbb{A}^{\mathbb{B}})^{\mathbb{C}}$. If, moreover, \mathbb{A} is diagonal, then

$$\mathbb{A}^{\mathbb{B}\mathbb{C}} \sim (\mathbb{A}^{\mathbb{B}})^{\mathbb{C}}.$$

Proof. It follows from [8], Theorem 9. \square

36. Corollary. Let $\mathbb{A}, \mathbb{B}, \mathbb{C}$ be n -ary ordered sets such that \mathbb{A} is diagonal or both \mathbb{B} and \mathbb{C} are discrete. Then

$$\mathbb{A}^{\mathbb{B}\mathbb{C}} \sim (\mathbb{A}^{\mathbb{B}})^{\mathbb{C}}.$$

37. Theorem. For any n -ary relational systems \mathbb{A}, \mathbb{B} with disjoint carriers we have

$$\widetilde{\mathbb{A} + \mathbb{B}} = \widetilde{\mathbb{A}} + \widetilde{\mathbb{B}}.$$

Proof. It follows directly from the definitions. \square

38. Theorem. For any n -ary relational systems \mathbb{A}, \mathbb{B} we have

$$\begin{aligned}\widetilde{\mathbb{A}\mathbb{B}} &= \widetilde{\mathbb{A}}\widetilde{\mathbb{B}}, \\ \widetilde{\mathbb{A}^{\mathbb{B}}} &= \widetilde{\mathbb{A}}^{\widetilde{\mathbb{B}}}.\end{aligned}$$

Proof. Let us show the second equality only. Let $\mathbb{A} = (A, R)$, $\mathbb{B} = (B, S)$. It is easy to prove that $\text{Hom}(\mathbb{B}, \mathbb{A}) = \text{Hom}(\widetilde{\mathbb{B}}, \widetilde{\mathbb{A}})$. Hence both sides of the equality have the same carrier $\text{Hom}(\mathbb{B}, \mathbb{A})$. Denote $\widetilde{\mathbb{A}} = (A, T)$, $\mathbb{A}^{\mathbb{B}} = (\text{Hom}(\mathbb{B}, \mathbb{A}), U)$, $\widetilde{\mathbb{A}^{\mathbb{B}}} = (\text{Hom}(\mathbb{B}, \mathbb{A}), V)$, $\widetilde{\mathbb{A}}^{\widetilde{\mathbb{B}}} = (\text{Hom}(\mathbb{B}, \mathbb{A}), W)$. Let $(f_1, \dots, f_n) \in V$. Then $(f_n, \dots, f_1) \in U$, thus $(f_n(x), \dots, f_1(x)) \in R$ for each $x \in B$, consequently $(f_1(x), \dots, f_n(x)) \in T$ for each $x \in B$ and $(f_1, \dots, f_n) \in W$, so that $V \subseteq W$. Similarly $W \subseteq V$. \square

39. Theorem. For any n -ary relational systems $\mathbb{A}, \mathbb{B}, \mathbb{C}$ we have

$$\mathbb{A} \circ (\mathbb{B} \circ \mathbb{C}) \sim (\mathbb{A} \circ \mathbb{B}) \circ \mathbb{C}.$$

Proof. Denote $\mathbb{A} = (A, R)$, $\mathbb{B} = (B, S)$, $\mathbb{C} = (C, T)$, $\mathbb{A} \circ \mathbb{B} = (A \times B, U)$, $\mathbb{B} \circ \mathbb{C} = (B \times C, V)$, $\mathbb{A} \circ (\mathbb{B} \circ \mathbb{C}) = (A \times (B \times C), W)$, $(\mathbb{A} \circ \mathbb{B}) \circ \mathbb{C} = ((A \times B) \times C, X)$. Let φ be the natural bijection of $A \times (B \times C)$ onto $(A \times B) \times C$ defined by $\varphi((a, (b, c))) = ((a, b), c)$ for any $a \in A$, $b \in B$, $c \in C$. Let $((a_1, (b_1, c_1)), \dots, (a_n, (b_n, c_n))) \in W$. Then at least two of the elements a_1, \dots, a_n are different, $(a_1, \dots, a_n) \in R$ or $a_1 = \dots = a_n$, $((b_1, c_1), \dots, (b_n, c_n)) \in V$, so that at least two of the elements a_1, \dots, a_n are different, $(a_1, \dots, a_n) \in R$ or $a_1 = \dots = a_n$, at least two of the elements b_1, \dots, b_n are different, $(b_1, \dots, b_n) \in S$ or $a_1 = \dots = a_n$, $b_1 = \dots = b_n$, $(c_1, \dots, c_n) \in T$, consequently at least two of the elements $(a_1, b_1), \dots, (a_n, b_n)$ are different, $((a_1, b_1), \dots, (a_n, b_n)) \in U$ or $(a_1, b_1) = \dots = (a_n, b_n)$, $(c_1, \dots, c_n) \in T$, hence $(\varphi((a_1, (b_1, c_1))), \dots, \varphi((a_n, (b_n, c_n)))) = (((a_1, b_1), c_1), \dots, ((a_n, b_n), c_n)) \in X$ and φ is a homomorphism of $\mathbb{A} \circ (\mathbb{B} \circ \mathbb{C})$ onto $(\mathbb{A} \circ \mathbb{B}) \circ \mathbb{C}$. Similarly $(\varphi((a_1, (b_1, c_1))), \dots, \varphi((a_n, (b_n, c_n)))) \in X$ implies $((a_1, (b_1, c_1)), \dots, (a_n, (b_n, c_n))) \in W$ and φ is an isomorphism of $\mathbb{A} \circ (\mathbb{B} \circ \mathbb{C})$ onto $(\mathbb{A} \circ \mathbb{B}) \circ \mathbb{C}$. \square

40. Theorem. Let $\mathbb{A}_1, \dots, \mathbb{A}_n$ be n -ary relational systems with pairwise disjoint carriers. Then, for any n -ary relational system \mathbb{B} , we have

$$(\mathbb{A}_1 \oplus \dots \oplus \mathbb{A}_n) \circ \mathbb{B} = \mathbb{A}_1 \circ \mathbb{B} \oplus \dots \oplus \mathbb{A}_n \circ \mathbb{B}.$$

Proof. As the n -ary relational systems $\mathbb{A}_1, \dots, \mathbb{A}_n$ have pairwise disjoint carriers, the n -ary relational systems $\mathbb{A}_1 \circ \mathbb{B}, \dots, \mathbb{A}_n \circ \mathbb{B}$ have pairwise disjoint carriers as well. Let $\mathbb{A}_j = (A_j, R_j)$ for $j \in (n]$, $\mathbb{B} = (B, S)$, $\mathbb{A}_1 \oplus \dots \oplus \mathbb{A}_n = (\bigcup_{j=1}^n A_j, T)$, $(\mathbb{A}_1 \oplus \dots \oplus \mathbb{A}_n) \circ \mathbb{B} = ((\bigcup_{j=1}^n A_j) \times B, U)$, $\mathbb{A}_1 \circ \mathbb{B} \oplus \dots \oplus \mathbb{A}_n \circ \mathbb{B} = ((\bigcup_{j=1}^n A_j) \times B, V)$. Assume that $((a_1, b_1), \dots, (a_n, b_n)) \in U$. Then at least two of the elements a_1, \dots, a_n are different, $(a_1, \dots, a_n) \in T$ or $a_1 = \dots = a_n$, $(b_1, \dots, b_n) \in S$. Hence at least two of the elements a_1, \dots, a_n are different, $a_1, \dots, a_n \in A_j$, $(a_1, \dots, a_n) \in R_j$ for some $j \in (n]$ or $a_1 = \dots = a_n$, $(b_1, \dots, b_n) \in S$ or there exists an $\varepsilon \in E_n$ such that $a_i \in A_{\varepsilon(i)}$ for each $i \in (n]$ and $((a_1, b_1), \dots, (a_n, b_n)) \in V$. Similarly $V \subseteq U$. \square

41. Remark. The second distributive law, of course, does not hold in general, because it is even not valid for binary relational systems.

42. Theorem. For any binary relational systems \mathbb{A}, \mathbb{B} with disjoint carriers and for any n -ary relational system \mathbb{C} we have

$$\mathbb{A} + \mathbb{B} \mathbb{C} \sim (\mathbb{A} \mathbb{C})(\mathbb{B} \mathbb{C}).$$

Proof. Let $\mathbb{A} = (A, R)$, $\mathbb{B} = (B, S)$, $\mathbb{C} = (C, T)$, $\mathbb{A} \mathbb{C} = (C^A, U)$, $\mathbb{B} \mathbb{C} = (C^B, V)$, $\mathbb{A} + \mathbb{B} \mathbb{C} = (C^{A \cup B}, W)$, $(\mathbb{A} \mathbb{C})(\mathbb{B} \mathbb{C}) = (C^A \times C^B, X)$. Let φ be the natural bijection of $C^{A \cup B}$ onto $C^A \times C^B$ defined by $\varphi(f) = (f \upharpoonright A, f \upharpoonright B)$. Assume that $(f_1, \dots, f_n) \in W$. In order to show that $(\varphi(f_1), \dots, \varphi(f_n)) = ((f_1 \upharpoonright A, f_1 \upharpoonright B), \dots, (f_n \upharpoonright A, f_n \upharpoonright B)) \in X$, we must verify that $(f_1 \upharpoonright A, \dots, f_n \upharpoonright A) \in U$, $(f_1 \upharpoonright B, \dots, f_n \upharpoonright B) \in V$. Let $((f_1 \upharpoonright A)(a), \dots, (f_n \upharpoonright A)(a)) \notin T$ for some $a \in A$. Then $(f_1(a), \dots, f_n(a)) \notin T$ and there exists a $b \in A \cup B$ such that $(b, a) \in R \cup S$, $(f_1(b), \dots, f_n(b)) \in T$, at least two of the elements $f_1(b), \dots, f_n(b)$ are different, and $(f_1(x), \dots, f_n(x)) \in T$ for each $x \in A \cup B$, $(x, b) \in R \cup S$. But $(b, a) \in R \cup S$ implies $b \in A$ and $(x, b) \in R \cup S$ implies $x \in A$. Hence there exists a $b \in A$ such that $(b, a) \in R$, $((f_1 \upharpoonright A)(b), \dots, (f_n \upharpoonright A)(b)) \in T$, at least two of the elements $(f_1 \upharpoonright A)(b), \dots, (f_n \upharpoonright A)(b)$ are different, and $((f_1 \upharpoonright A)(x), \dots, (f_n \upharpoonright A)(x)) \in T$ for each $x \in A$, $(x, b) \in R$, so that $(f_1 \upharpoonright A, \dots, f_n \upharpoonright A) \in U$. Similarly $(f_1 \upharpoonright B, \dots, f_n \upharpoonright B) \in V$ and $(\varphi(f_1), \dots, \varphi(f_n)) \in X$. Analogously $(\varphi(f_1), \dots, \varphi(f_n)) \in X$ implies $(f_1, \dots, f_n) \in W$. \square

43. Theorem. For any binary relational systems \mathbb{A}, \mathbb{B} with disjoint carriers and for any reflexive n -ary relational system \mathbb{C} we have

$$\mathbb{A} \oplus \mathbb{B} \mathbb{C} \sim (\mathbb{A} \mathbb{C}) \circ (\mathbb{B} \mathbb{C}).$$

Proof. Let $\mathbb{A} = (A, R)$, $\mathbb{B} = (B, S)$, $\mathbb{C} = (C, T)$, $\mathbb{A} \oplus \mathbb{B} = (A \cup B, U)$, $\mathbb{A} \oplus \mathbb{B} \mathbb{C} = (C^{A \cup B}, V)$, $\mathbb{A} \mathbb{C} = (C^A, W)$, $\mathbb{B} \mathbb{C} = (C^B, X)$, $(\mathbb{A} \mathbb{C}) \circ (\mathbb{B} \mathbb{C}) = (C^A \times C^B, Y)$. Again, let φ be the natural bijection of $C^{A \cup B}$ onto $C^A \times C^B$ defined by $\varphi(f) = (f \upharpoonright A, f \upharpoonright B)$. Assume that $(\varphi(f_1), \dots, \varphi(f_n)) = ((f_1 \upharpoonright A, f_1 \upharpoonright B), \dots, (f_n \upharpoonright A, f_n \upharpoonright B)) \in Y$, i.e. either at least two of the elements $f_1 \upharpoonright A, \dots, f_n \upharpoonright A$ are different, $(f_1 \upharpoonright A, \dots, f_n \upharpoonright A) \in W$, or $f_1 \upharpoonright A = \dots = f_n \upharpoonright A$, $(f_1 \upharpoonright B, \dots, f_n \upharpoonright B) \in X$. Now, let there exist an $a \in A \cup B$ such that $(f_1(a), \dots, f_n(a)) \notin T$. If $a \in A$, then $((f_1 \upharpoonright A)(a), \dots, (f_n \upharpoonright A)(a)) \notin T$ and, because of reflexivity of T , at least two of the elements $f_1 \upharpoonright A, \dots, f_n \upharpoonright A$ are different, thus $(f_1 \upharpoonright A, \dots, f_n \upharpoonright A) \in W$. Hence there exists a $b \in A$ such that $(b, a) \in R$, $((f_1 \upharpoonright A)(b), \dots, (f_n \upharpoonright A)(b)) \in T$, at least two of the elements $(f_1 \upharpoonright A)(b), \dots, (f_n \upharpoonright A)(b)$ are different, and for each $x \in A$, $(x, b) \in R$ we have $((f_1 \upharpoonright A)(x), \dots, (f_n \upharpoonright A)(x)) \in T$. Consequently, there exists a $b \in A \cup B$ such that $(b, a) \in U$, $(f_1(b), \dots, f_n(b)) \in T$, at least two of the elements $f_1(b), \dots, f_n(b)$ are different, and for each $x \in A \cup B$, $(x, b) \in U$ we have $(f_1(x), \dots, f_n(x)) \in T$. If $a \in B$, then $((f_1 \upharpoonright B)(a), \dots, (f_n \upharpoonright B)(a)) \notin T$. If, moreover, at least two of the mappings $f_1 \upharpoonright A, \dots, f_n \upharpoonright A$ are different, then $(f_1 \upharpoonright A, \dots, f_n \upharpoonright A) \in W$. Let there exist a $b \in A$ such that $((f_1 \upharpoonright A)(b), \dots, (f_n \upharpoonright A)(b)) \notin T$. Then there exists a $c \in A$ such that $(c, b) \in R$, $((f_1 \upharpoonright A)(c), \dots, (f_n \upharpoonright A)(c)) \in T$, at least two of the elements $(f_1 \upharpoonright A)(c), \dots, (f_n \upharpoonright A)(c)$ are different, and for each $x \in A$, $(x, c) \in R$ we have $((f_1 \upharpoonright A)(x), \dots, (f_n \upharpoonright A)(x)) \in T$. Hence there exists a $c \in A \cup B$ such that $(c, a) \in U$, $(f_1(c), \dots, f_n(c)) \in T$, at least two of the elements $f_1(c), \dots, f_n(c)$ are different, and for each $x \in A \cup B$, $(x, c) \in U$ we have $(f_1(x), \dots, f_n(x)) \in T$. On the contrary, let there exist no $b \in A$ such that $((f_1 \upharpoonright A)(b), \dots, (f_n \upharpoonright A)(b)) \notin T$. Then for any $b \in A$ such that at least two of the elements $(f_1 \upharpoonright A)(b), \dots, (f_n \upharpoonright A)(b)$ are different, we have $b \in A \cup B$, $(b, a) \in U$, $(f_1(b), \dots, f_n(b)) \in T$, at least two of the elements $f_1(b), \dots, f_n(b)$ are different, and for each $x \in A \cup B$, $(x, b) \in U$ we have $(f_1(x), \dots, f_n(x)) \in T$. Finally, if $a \in B$, $((f_1 \upharpoonright B)(a), \dots, (f_n \upharpoonright B)(a)) \notin T$ and $f_1 \upharpoonright A = \dots = f_n \upharpoonright A$, then $(f_1 \upharpoonright B, \dots, f_n \upharpoonright B) \in X$, thus there exists a $b \in B$ such that $(b, a) \in S$, $((f_1 \upharpoonright B)(b), \dots, (f_n \upharpoonright B)(b)) \in T$, at least two of the elements $(f_1 \upharpoonright B)(b), \dots, (f_n \upharpoonright B)(b)$ are different, and for each $x \in B$, $(x, b) \in S$ we have $((f_1 \upharpoonright B)(x), \dots, (f_n \upharpoonright B)(x)) \in T$. Hence there exists a $b \in A \cup B$ such that $(b, a) \in U$, $(f_1(b), \dots, f_n(b)) \in T$, at least two of the elements $f_1(b), \dots, f_n(b)$ are different, and for each $x \in A \cup B$, $(x, b) \in U$ we have $(f_1(x), \dots, f_n(x)) \in T$ (for $x \in B$ this follows from the preceding, for $x \in A$ from the

reflexivity of T). In all cases we have $(f_1, \dots, f_n) \in V$. Similarly $(f_1, \dots, f_n) \in V$ implies $(\varphi(f_1), \dots, \varphi(f_n)) \in Y$. \square

44. Theorem. For any n -ary relational systems $\mathbb{A}_1, \dots, \mathbb{A}_n$ with pairwise disjoint carriers we have

$$\mathbb{A}_1 \oplus \dots \oplus \mathbb{A}_n = \widetilde{\mathbb{A}_1 \oplus \dots \oplus \mathbb{A}_n} = \widetilde{\mathbb{A}}_n \oplus \dots \oplus \widetilde{\mathbb{A}}_1.$$

Proof. Let $\mathbb{A}_j = (A_j, R_j)$, $\widetilde{\mathbb{A}}_j = (A_j, S_j)$ for $j \in (n)$, $\mathbb{A}_1 \oplus \dots \oplus \mathbb{A}_n = (\bigcup_{j=1}^n A_j, T)$, $\widetilde{\mathbb{A}_1 \oplus \dots \oplus \mathbb{A}_n} = (\bigcup_{j=1}^n A_j, U)$, $\widetilde{\mathbb{A}}_n \oplus \dots \oplus \widetilde{\mathbb{A}}_1 = (\bigcup_{j=1}^n A_j, V)$. Let $(a_1, \dots, a_n) \in U$. Then $(a_n, \dots, a_1) \in T$, thus either $(a_n, \dots, a_1) \in R_j$ for some $j \in (n)$ or there exists an $\varepsilon \in E_n$ such that $a_{n+1-j} \in A_{\varepsilon(j)}$ for each $j \in (n)$. In the first case we have $(a_1, \dots, a_n) \in S_j \subseteq V$. In the second case we have $a_j \in A_{\varepsilon(n+1-j)}$ for each $j \in (n)$. Hence $(a_1, \dots, a_n) \in A_{\varepsilon(n)} \times \dots \times A_{\varepsilon(1)} \subseteq V$. Consequently $U \subseteq V$. Similarly $V \subseteq U$. \square

45. Theorem. For any n -ary relational systems \mathbb{A}, \mathbb{B} we have

$$\widetilde{\mathbb{A} \circ \mathbb{B}} = \widetilde{\mathbb{A}} \circ \widetilde{\mathbb{B}}.$$

Proof. Let $\mathbb{A} = (A, R)$, $\mathbb{B} = (B, S)$, $\widetilde{\mathbb{A}} = (A, T)$, $\widetilde{\mathbb{B}} = (B, U)$, $\mathbb{A} \circ \mathbb{B} = (A \times B, V)$, $\widetilde{\mathbb{A} \circ \mathbb{B}} = (A \times B, W)$, $\widetilde{\mathbb{A}} \circ \widetilde{\mathbb{B}} = (A \times B, X)$. Let $((a_1, b_1), \dots, (a_n, b_n)) \in W$. Then $((a_n, b_n), \dots, (a_1, b_1)) \in V$, thus either at least two of the elements a_n, \dots, a_1 are different, $(a_n, \dots, a_1) \in R$ or $a_n = \dots = a_1$, $(b_n, \dots, b_1) \in S$. This implies that either at least two of the elements a_1, \dots, a_n are different, $(a_1, \dots, a_n) \in T$ or $a_1 = \dots = a_n$, $(b_1, \dots, b_n) \in U$, so that $((a_1, b_1), \dots, (a_n, b_n)) \in X$, and $W \subseteq X$. Similarly $X \subseteq W$. \square

46. Theorem. For any binary relational system \mathbb{A} and any n -ary relational system \mathbb{B} we have

$$\widetilde{\mathbb{A} \mathbb{B}} = \mathbb{A} \widetilde{\mathbb{B}}.$$

Proof. Let $\mathbb{A} = (A, R)$, $\mathbb{B} = (B, S)$, $\widetilde{\mathbb{B}} = (B, T)$, $\mathbb{A} \mathbb{B} = (B^A, U)$, $\widetilde{\mathbb{A} \mathbb{B}} = (B^A, V)$, $\mathbb{A} \widetilde{\mathbb{B}} = (B^A, W)$. Let $(f_1, \dots, f_n) \in V$. Then $(f_n, \dots, f_1) \in U$. Suppose that $(f_1(a), \dots, f_n(a)) \notin T$ for some $a \in A$. Then $(f_n(a), \dots, f_1(a)) \notin S$, thus there exists a $b \in A$ such that $(b, a) \in R$, $(f_n(b), \dots, f_1(b)) \in S$, at least two of the elements $f_n(b), \dots, f_1(b)$ are different, and for each $x \in A$, $(x, b) \in R$ we have $(f_n(x), \dots, f_1(x)) \in S$. Hence there exists a $b \in A$, $(b, a) \in R$ such that $(f_1(b), \dots, f_n(b)) \in T$, at least two of the elements $f_1(b), \dots, f_n(b)$ are different, and for each $x \in A$, $(x, b) \in R$ we have $(f_1(x), \dots, f_n(x)) \in T$. Consequently $(f_1, \dots, f_n) \in W$ and $V \subseteq W$. Similarly $W \subseteq V$. \square

References

- [1] *G. Birkhoff*: An extended arithmetic. *Duke Math. J.* *3* (1937), 311–316.
- [2] *G. Birkhoff*: Generalized arithmetic. *Duke Math. J.* *9* (1942), 283–302.
- [3] *G. Birkhoff*: *Lattice Theory*. Amer. Math. Soc., Providence, Rhode Island, Third Edition, 1973.
- [4] *M. M. Day*: Arithmetic of ordered systems. *Trans. Amer. Math. Soc.* *58* (1945), 1–43.
- [5] *V. Novák*: On a power of relational structures. *Czechoslovak Math. J.* *35* (1985), 167–172.
- [6] *V. Novák, M. Novotný*: Binary and ternary relations. *Math. Bohem.* *117* (1992), 283–292.
- [7] *V. Novák, M. Novotný*: Pseudodimension of relational structures. *Czechoslovak Math. J.* (submitted).
- [8] *J. Šlapal*: Direct arithmetics of relational systems. *Publ. Math. Debrecen* *38* (1991), 39–48.

Author's address: Jiří Karásek, Technical University, Technická 2, 616 69 Brno, Czech Republic.