

A REMARK ON BRANCH WEIGHTS IN COUNTABLE TREES

BOHDAN ZELINKA, Liberec

(Received February 27, 2003)

Abstract. Let T be a tree, let u be its vertex. The branch weight $b(u)$ of u is the maximum number of vertices of a branch of T at u . The set of vertices u of T in which $b(u)$ attains its minimum is the branch weight centroid $B(T)$ of T . For finite trees the present author proved that $B(T)$ coincides with the median of T , therefore it consists of one vertex or of two adjacent vertices. In this paper we show that for infinite countable trees the situation is quite different.

Keywords: branch weight, branch weight centroid, tree, path, degree of a vertex

MSC 2000: 05C05

In this paper we study the branch weight in countable infinite trees. We define it analogously to the case of finite trees.

Let T be a tree with the vertex set $V(T)$, let $u \in V(T)$. On the set $V(T) - \{u\}$ consider the binary relation β such that $(x, y) \in \beta$ if and only if the path connecting x and y in T does not contain u . The relation β is evidently an equivalence relation on $V(T) - \{u\}$. Each equivalence class of β induces a subtree of T which is called a branch of T at u . The set of all branches of T at u will be denoted by $\mathcal{B}(T, u)$.

The vertex set of every branch B will be denoted by $V(B)$, its cardinality by $|V(B)|$. If T is a countable tree, then $|V(B)|$ is either a positive integer, or the cardinal number \aleph_0 .

For each $u \in V(T)$ let $b(u) = \max(|V(B)| : B \in \mathcal{B}(T, u))$. The number $b(u)$ is called the branch weight of u in T .

The set of the vertices u of T in which $b(u)$ attains its minimum is called the branch weight centroid $B(T)$ of T (see e.g. [2], [3]). For finite trees the present author [4] proved, solving one problem from the book [1], that $B(T)$ coincides with the median of T ; therefore it consists either of one vertex, or of two adjacent vertices.

Here we will study countably infinite trees.

Theorem 1. *Let T be a countable infinite tree. If T contains an infinite path or at least two vertices of infinite degree, then $B(T) = V(T)$.*

Proof. Suppose that the condition is satisfied. Let $u \in V(T)$. If T contains at least two vertices of infinite degree, then at least one of them, say w , is in $V(T) - \{u\}$. As the vertex sets of branches of T at u form a partition of $V(T) - \{u\}$, the vertex w lies on some branch $B^* \in \mathcal{B}(T, u)$ and together with it an infinite number of neighbours of w lie in B^* . Therefore $|V(B^*)| = \aleph_0$. If T contains an infinite path P , then evidently there exists $B^{**} \in \mathcal{B}(T, u)$ which contains either the whole path P , or at least its one-way infinite subpath which does not contain u . Again $|V(B^{**})| = \aleph_0$. We have proved that $b(u) = \aleph_0$ for each vertex $u \in V(T)$. This is also the minimum of $b(u)$ over all $u \in V(T)$ and thus the branch weight centroid $B(T)$ is equal to the whole vertex set $V(T)$. \square

Theorem 2. *Let T be a countable infinite tree. Suppose that the following two conditions are satisfied:*

- (i) *The tree T contains no infinite path.*
- (ii) *The tree T contains exactly one vertex w of infinite degree.*

Then one of the following two assertions is true:

- (iii) *The branch weight centroid $B(T) = \{w\}$.*
- (iv) *The branch weight centroid $B(T)$ is undefined.*

Proof. Consider $\mathcal{B}(T, w)$ and let $B^0 \in \mathcal{B}(T, w)$. Suppose that B^0 is infinite. Then it is a locally finite subtree of T , because $V(B^0) \subseteq V(T) - \{w\}$ and w is the unique vertex of T of infinite degree. However, as is well-known, in that case B^0 contains an infinite path, which is a contradiction with (i). Hence all branches of T at w are finite. If there exists a finite upper bound of the values $|V(B)|$ for $B \in \mathcal{B}(T, w)$, then there exists a finite maximum of those values and this is $b(w)$. If that upper bound does not exist, that maximum does not exist, either and $b(w)$ is undefined. Analogously as in the proof of Theorem 1 we prove that $b(u) = \aleph_0$ for each $u \in V(T) - \{w\}$. Thus in the first case w is the unique vertex with the finite branch weight and $B(T) = \{w\}$. In the other the branch weight is not defined for all vertices of T and consequently $B(T)$ is undefined. \square

Corollary. *Let T be a countable infinite tree. Then either $B(T) = V(T)$, or T can be obtained by the following simple construction:*

Take a sequence $(T_n)_{n=1}^{\infty}$ of pairwise vertex-disjoint finite trees and a vertex w outside of all of them. In each tree T_n choose a vertex v_n . Join each vertex v_n by an edge with w (for $n = 1, 2, 3, \dots$).

A c k n o w l e d g e m e n t. This paper was supported by the Grant MSMT 245100302 of the Ministry of Education, Youth and Physical Culture of the Czech Republic.

References

- [1] *O. Ore*: Theory of Graphs. AMS Colloq. Publ., Providence, 1963.
- [2] *P. J. Slater*: Accretion centers: A generalization of branch weight centroids. *Discr. Appl. Math.* 3 (1981), 187–192.
- [3] *Z. Win, Y. Myint*: The cendian of a tree. *Southeast Asian Bull. Math.* 25 (2002), 757–767.
- [4] *B. Zelinka*: Medians and peripherians of trees. *Arch. Math. Brno* 4 (1968), 87–95.

Author's address: Bohdan Zelinka, Technická Univerzita Liberec, Pedagogická fakulta, katedra aplikované matematiky, Voroněžská 13, 460 01 Liberec, Czech Republic, e-mail: bohdan.zelinka@vslib.cz.