

## CERTAIN SUFFICIENT CONDITIONS FOR A SUBCLASS OF ANALYTIC FUNCTIONS ASSOCIATED WITH HOHLOV OPERATOR

S. Sivasubramanian, Thomas Rosy and K. Muthunagai

**Abstract.** Making use of the Hohlov operator, we obtain inclusion relations between the classes of certain normalized analytic functions. Relevant connections of our work with the earlier works are pointed out.

### 1. Introduction

Let  $\mathcal{A}$  be the class of functions  $f$  normalized by

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n, \quad (1.1)$$

which are analytic in the open unit disk

$$\mathbb{U} = \{z : z \in \mathbb{C} \text{ and } |z| < 1\}.$$

As usual, we denote by  $\mathcal{S}$  the subclass of  $\mathcal{A}$  consisting of functions which are also univalent in  $\mathbb{U}$ .

A function  $f \in \mathcal{A}$  is said to be starlike of order  $\alpha$  ( $0 \leq \alpha < 1$ ), if and only if

$$\Re \left( \frac{zf'(z)}{f(z)} \right) > \alpha \quad (z \in \mathbb{U}).$$

This function class is denoted by  $\mathcal{S}^*(\alpha)$ . We also write  $\mathcal{S}^*(0) =: \mathcal{S}^*$ , where  $\mathcal{S}^*$  denotes the class of functions  $f \in \mathcal{A}$  that are starlike in  $\mathbb{U}$  with respect to the origin.

A function  $f \in \mathcal{A}$  is said to be convex of order  $\alpha$  ( $0 \leq \alpha < 1$ ) if and only if

$$\Re \left( 1 + \frac{zf''(z)}{f'(z)} \right) > \alpha \quad (z \in \mathbb{U}).$$

---

*2010 Mathematics Subject Classification:* 33C45, 33A30, 30C45

*Keywords and phrases:* Starlike function; convolution, negative coefficients; coefficient inequalities; growth and distortion theorems.

This class is denoted by  $\mathcal{K}(\alpha)$ . Further,  $\mathcal{K} = \mathcal{K}(0)$ , the well-known standard class of convex functions. It is an established fact that

$$f \in \mathcal{K}(\alpha) \iff zf' \in \mathcal{S}^*(\alpha).$$

Let  $M(\lambda, \alpha)$  be a subclass of  $\mathcal{A}$  consisting of functions of the form that satisfy the condition

$$\Re \left( \frac{z(zf'(z))'}{(1-\lambda)zf'(z) + \lambda z(zf'(z))'} \right) > \alpha, \quad z \in \Delta$$

for some  $\alpha$  and  $\lambda$  where  $0 \leq \alpha < 1$  and  $0 \leq \lambda < 1$ . That is,  $M(\lambda, \alpha)$  be a subclass of  $\mathcal{S}$  consisting of functions of the form that satisfy the condition

$$\Re \left( \frac{f'(z) + zf''(z)}{f'(z) + \lambda zf''(z)} \right) > \alpha, \quad z \in \Delta$$

The class  $M(\lambda, \alpha)$  was introduced by Altıntaş and Owa [1] and also investigated very recently by Mostafa [12].

A function  $f \in \mathcal{A}$  is said to be in the class  $\mathcal{UCV}$  of uniformly convex functions in  $\mathbb{U}$  if and only if it has the property that, for every circular arc  $\delta$  contained in the unit disk  $\mathbb{U}$ , with center  $\zeta$  also in  $\mathbb{U}$ , the image curve  $f(\delta)$  is a convex arc. The function class  $\mathcal{UCV}$  was introduced by Goodman [7].

Furthermore, we denote by  $k\text{-UCV}$  and  $k\text{-ST}$ , ( $0 \leq k < \infty$ ), two interesting subclasses of  $\mathcal{S}$  consisting respectively of functions which are  $k$ -uniformly convex and  $k$ -starlike in  $\mathbb{U}$ . Namely, we have for  $0 \leq k < \infty$

$$k\text{-UCV} := \left\{ f \in \mathcal{S} : \Re \left( 1 + \frac{zf''(z)}{f'(z)} \right) > k \left| \frac{zf''(z)}{f'(z)} \right|, \quad (z \in \mathbb{U}) \right\}$$

and

$$k\text{-ST} := \left\{ f \in \mathcal{S} : \Re \left( \frac{zf'(z)}{f(z)} \right) > k \left| \frac{zf'(z)}{f(z)} - 1 \right|, \quad (z \in \mathbb{U}) \right\}.$$

The class  $k\text{-UCV}$  was introduced by Kanas and Wiśniowska [10], where its geometric definition and connections with the conic domains were considered. The class  $k\text{-ST}$  was investigated in [11]. In fact, it is related to the class  $k\text{-UCV}$  by means of the well-known Alexander equivalence between the usual classes of convex and starlike functions (see also the work of Kanas and Srivastava [9] for further developments involving each of the classes  $k\text{-UCV}$  and  $k\text{-ST}$ ). In particular, when  $k = 1$ , we obtain

$$k\text{-UCV} \equiv \mathcal{UCV} \quad \text{and} \quad k\text{-ST} = \mathcal{SP},$$

where  $\mathcal{UCV}$  and  $\mathcal{SP}$  are the familiar classes of uniformly convex functions and parabolic starlike functions in  $\mathbb{U}$  respectively (see for details, [7]). Indeed, by making use of a certain fractional calculus operator, Srivastava and Mishra [17] presented a systematic and unified study of the classes  $\mathcal{UCV}$  and  $\mathcal{SP}$ .

Let us denote (see [10,11])

$$P_1(k) = \begin{cases} \frac{8(\arccos k)^2}{\pi^2(1-k^2)} & \text{for } 0 \leq k < 1 \\ \frac{8}{\pi^2} & \text{for } k = 1 \\ \frac{\pi^2}{4\sqrt{t}(1+t)(k^2-1)\mathcal{K}(t)} & \text{for } k > 1 \end{cases} \quad (1.2)$$

where  $t \in (0, 1)$  is determined by  $k = \cosh(\pi\mathcal{K}'(t)/[4\mathcal{K}(t)])$ ,  $\mathcal{K}$  is the Legendre's complete elliptic integral of the first kind

$$\mathcal{K}(t) = \int_0^1 \frac{dx}{\sqrt{(1-x^2)(1-t^2x^2)}}$$

and  $\mathcal{K}'(t) = \mathcal{K}(\sqrt{1-t^2})$  is the complementary integral of  $\mathcal{K}(t)$ . Let  $\Omega_k$  be a domain such that  $1 \in \Omega_k$  and

$$\partial\Omega_k = \{w = u + iv : u^2 = k^2(u-1)^2 + k^2v^2\}, \quad 0 \leq k < \infty.$$

The domain  $\Omega_k$  is elliptic for  $k > 1$ , hyperbolic when  $0 < k < 1$ , parabolic when  $k = 1$ , and a right half-plane when  $k = 0$ . If  $p$  is an analytic function with  $p(0) = 1$  which maps the unit disc  $\mathbb{U}$  conformally onto the region  $\Omega_k$ , then  $P_1(k) = p'(0)$ .  $P_1(k)$  is strictly decreasing function of the variable  $k$  and its values are included in the interval  $(0, 2]$ .

Let  $f \in \mathcal{A}$  be of the form (1.1). If  $f \in k - \mathcal{UCV}$ , then the following coefficient inequalities hold true (cf. [10]):

$$|a_n| \leq \frac{(P_1(k))_{n-1}}{n!}, \quad n \in \mathbb{M} \setminus \{1\}. \quad (1.3)$$

Similarly, if  $f$  of the form (1.1) belongs to the class  $k - \mathcal{ST}$ , then (cf. [11])

$$|a_n| \leq \frac{(P_1(k))_{n-1}}{(n-1)!}, \quad n \in \mathbb{M} \setminus \{1\}.$$

A function  $f \in \mathcal{A}$  is said to be in the class  $\mathcal{R}^\tau(A, B)$ , ( $\tau \in \mathbb{C} \setminus \{0\}$ ,  $-1 \leq B < A \leq 1$ ), if it satisfies the inequality

$$\left| \frac{f'(z) - 1}{(A - B)\tau - B[f'(z) - 1]} \right| < 1 \quad (z \in \mathbb{U}).$$

The class  $\mathcal{R}^\tau(A, B)$  was introduced earlier by Dixit and Pal [5]. Two of the many interesting subclasses of the class  $\mathcal{R}^\tau(A, B)$  are worthy of mention here. First of all, by setting

$$\tau = e^{i\eta} \cos \eta \quad (-\pi/2 < \eta < \pi/2), \quad A = 1 - 2\beta \quad (0 \leq \beta < 1) \quad \text{and} \quad B = -1,$$

the class  $\mathcal{R}^\tau(A, B)$  reduces essentially to the class  $\mathcal{R}_\eta(\beta)$  introduced and studied by Ponnusamy and Rønning [14], where

$$\mathcal{R}_\eta(\beta) = \{f \in \mathcal{A} : \Re(e^{i\eta}(f'(z) - \beta)) > 0 \quad (z \in \mathbb{U}; -\pi/2 < \eta < \pi/2, 0 \leq \beta < 1)\}.$$

Secondly, if we put

$$\tau = 1, A = \beta \text{ and } B = -\beta \quad (0 < \beta \leq 1),$$

we obtain the class of functions  $f \in \mathcal{A}$  satisfying the inequality

$$\left| \frac{f'(z) - 1}{f'(z) + 1} \right| < \beta \quad (z \in \mathbb{U}; 0 < \beta \leq 1)$$

which was studied by (among others) Padmanabhan [13] and Caplinger and Causey [3], (see also the works [6, 15, 16, 18]).

The Gaussian hypergeometric function  $F(a, b; c; z)$  given by

$${}_2F_1(a, b; c; z) = F(a, b; c; z) = \sum_{n=0}^{\infty} \frac{(a)_n (b)_n}{(c)_n (1)_n} z^n \quad (z \in \mathbb{U})$$

is the solution of the homogenous hypergeometric differential equation

$$z(1-z)w''(z) + [c - (a+b+1)z]w'(z) - abw(z) = 0$$

and has rich applications in various fields such as conformal mappings, quasi conformal theory, continued fractions and so on.

By the Gauss Summation Theorem, we get

$$F(a, b; c; 1) = \sum_{n=0}^{\infty} \frac{(a)_n (b)_n}{(c)_n (1)_n} = \frac{\Gamma(c-a-b)\Gamma(c)}{\Gamma(c-a)\Gamma(c-b)} \quad \text{for } \Re(c-a-b) > 0.$$

Here,  $a, b, c$  are complex numbers such that  $c \neq 0, -1, -2, -3, \dots$ ,  $(a)_0 = 1$  for  $a \neq 0$ , and for each positive integer  $n$ ,  $(a)_n = a(a+1)(a+2)\dots(a+n-1)$  is the Pochhammer symbol. In the case of  $c = -k$ ,  $k = 0, 1, 2, \dots$ ,  $F(a, b; c; z)$  is defined if  $a = -j$  or  $b = -j$  where  $j \leq k$ . In this situation,  $F(a, b; c; z)$  becomes a polynomial of degree  $j$  with respect to  $z$ . Results regarding  $F(a, b; c; z)$  when  $\Re(c-a-b)$  is positive, zero or negative are abundant in the literature. In particular when  $\Re(c-a-b) > 0$ , the function is bounded. The hypergeometric function  $F(a, b; c; z)$  has been studied extensively by various authors and play an important role in Geometric Function Theory. It is useful in unifying various functions by giving appropriate values to the parameters  $a, b$  and  $c$ . We refer to [4, 6, 15, 16] and references therein for some important results.

For functions  $f \in \mathcal{A}$  given by (1.1) and  $g \in \mathcal{A}$  given by  $g(z) = z + \sum_{n=2}^{\infty} b_n z^n$ , we define the Hadamard product (or convolution) of  $f$  and  $g$  by

$$(f * g)(z) = z + \sum_{n=2}^{\infty} a_n b_n z^n, \quad z \in \mathbb{U}.$$

For  $f \in \mathcal{A}$ , we recall the operator  $I_{a,b,c}(f)$  of Hohlov [8] which maps  $\mathcal{A}$  into itself defined by means of Hadamard product as

$$I_{a,b,c}(f)(z) = zF(a, b; c; z) * f(z).$$

Therefore, for a function  $f$  defined by (1.1), we have

$$I_{a,b,c}(f)(z) = z + \sum_{n=2}^{\infty} \frac{(a)_{n-1}(b)_{n-1}}{(c)_{n-1}(1)_{n-1}} a_n z^n.$$

Using the integral representation,

$$F(a, b; c; z) = \frac{\Gamma(c)}{\Gamma(b)\Gamma(c-b)} \int_0^1 t^{b-1}(1-t)^{c-b-1} \frac{dt}{(1-tz)^a}, \quad \Re(c) > \Re(b) > 0,$$

we can write

$$[I_{a,b,c}(f)](z) = \frac{\Gamma(c)}{\Gamma(b)\Gamma(c-b)} \int_0^1 t^{b-1}(1-t)^{c-b-1} \frac{f(tz)}{t} dt * \frac{z}{(1-tz)^a}.$$

When  $f(z)$  equals the convex function  $\frac{z}{1-z}$ , then the operator  $I_{a,b,c}(f)$  in this case becomes  $zF(a, b; c; z)$ . If  $a = 1, b = 1 + \delta, c = 2 + \delta$  with  $\Re(\delta) > -1$  then the convolution operator  $I_{a,b,c}(f)$  turns into Bernardi operator

$$B_f(z) = [I_{a,b,c}(f)](z) = \frac{1 + \delta}{z^\delta} \int_0^1 t^{\delta-1} f(t) dt.$$

Indeed,  $I_{1,1,2}(f)$  and  $I_{1,2,3}(f)$  are known as Alexander and Libera operators, respectively.

To prove the main results, we need the following Lemmas.

LEMMA 1. [1] *A function  $f \in \mathcal{A}$  belongs to the class  $M(\lambda, \alpha)$  if*

$$\sum_{n=2}^{\infty} n(n - \lambda\alpha n - \alpha + \lambda\alpha) |a_n| \leq 1 - \alpha. \tag{1.4}$$

LEMMA 2. [5] *If  $f \in \mathcal{R}^\tau(A, B)$  is of form (1.1) then*

$$|a_n| \leq (A - B) \frac{|\tau|}{n}, \quad n \in \mathbb{N} \setminus \{1\}. \tag{1.5}$$

*The result is sharp.*

In this paper, we estimate certain inclusion relations involving the classes  $k - \mathcal{UCV}$ ,  $k - \mathcal{ST}$  and  $M(\lambda, \alpha)$ .

### 2. Main results

In this paper, we will study the action of the hypergeometric function on the classes  $k - \mathcal{UCV}$ ,  $k - \mathcal{ST}$ .

THEOREM 1. *Let  $a, b \in \mathbb{C} \setminus \{0\}$ . Also, let  $c$  be a real number such that  $c > |a| + |b| + 1$ . If  $f \in \mathcal{R}^\tau(A, B)$ , and if the inequality*

$$\begin{aligned} & \frac{\Gamma(c)\Gamma(c - |a| - |b| - 1)}{\Gamma(c - |a|)\Gamma(c - |b|)} [(1 - \lambda\alpha)|ab| + (1 - \alpha)(c - |a| - |b| - 1)] \\ & \leq (1 - \alpha) \left( \frac{1}{(A - B)|\tau|} + 1 \right) \end{aligned} \tag{2.1}$$

*is satisfied, then  $I_{a,b,c}(f) \in M(\lambda, \alpha)$ .*

*Proof.* Let  $f$  be of the form (1.1) belong to the class  $R^\tau(A, B)$ . By virtue of Lemma 1, it suffices to show that

$$\sum_{n=2}^{\infty} n(n - \lambda\alpha n - \alpha + \lambda\alpha) \left| \frac{(a)_{n-1}(b)_{n-1}}{(c)_{n-1}(1)_{n-1}} a_n \right| \leq 1 - \alpha.$$

Taking into account the inequality (1.5) and the relation  $|(a)_{n-1}| \leq (|a|)_{n-1}$ , we deduce that

$$\begin{aligned} & \sum_{n=2}^{\infty} (n - \lambda\alpha n - \alpha + \lambda\alpha) \left| \frac{(a)_{n-1}(b)_{n-1}}{(c)_{n-1}(1)_{n-1}} a_n \right| \\ & \leq (A - B)|\tau|(1 - \lambda\alpha) \sum_{n=2}^{\infty} n \left| \frac{(a)_{n-1}(b)_{n-1}}{(c)_{n-1}(1)_{n-1}} \right| \\ & \quad + (A - B)|\tau|\alpha(\lambda - 1) \sum_{n=2}^{\infty} \frac{(|a|)_{n-1}(|b|)_{n-1}}{(c)_{n-1}(1)_n} \\ & \leq (A - B)|\tau| \left\{ (1 - \lambda\alpha) \sum_{n=2}^{\infty} \frac{(|a|)_{n-1}(|b|)_{n-1}}{(c)_{n-1}(1)_{n-2}} + (1 - \alpha) \sum_{n=2}^{\infty} \frac{(|a|)_{n-1}(|b|)_{n-1}}{(c)_{n-1}(1)_{n-1}} \right\} \\ & = (A - B)|\tau| \left\{ (1 - \lambda\alpha) \frac{|ab|}{c} F(1 + |a|, 1 + |b|, 1 + c; 1) \right. \\ & \quad \left. + (1 - \alpha) (F(|a|, |b|, c; 1) - 1) \right\} \end{aligned}$$

where we use the relation

$$(a)_n = a(a + 1)_{n-1}. \tag{2.2}$$

The proof now follows by an application of Gauss summation theorem and (2.1). ■

For the choice of  $|b| = |a|$ , we have the following corollary.

**COROLLARY 1.** *Let  $a \in \mathbb{C} \setminus \{0\}$ . Also, let  $c$  be a real number such that  $c > 2|a| + 1$ . If  $f \in \mathcal{R}^\tau(A, B)$ , and if the inequality*

$$\begin{aligned} & \frac{\Gamma(c)\Gamma(c - 2|a| - 1)}{(\Gamma(c - |a|))^2} [(1 - \lambda\alpha)|a|^2 + (1 - \alpha)(c - 2|a| - 1)] \\ & \leq (1 - \alpha) \left( \frac{1}{(A - B)|\tau|} + 1 \right) \end{aligned}$$

*is satisfied, then  $I_{a,|a|,c}(f) \in M(\lambda, \alpha)$ .*

In the special case when  $b = 1$ , Theorem 1 immediately yields a result concerning the Carlson-Shaffer operator  $\mathcal{L}(a, c)(f) := I_{a, 1, c}(f)$  (see [4]).

**COROLLARY 2.** *Let  $a \in \mathbb{C} \setminus \{0\}$ . Also, let  $c$  be a real number such that  $c > |a| + 1$ . If  $f \in \mathcal{R}^\tau(A, B)$ , and if the inequality*

$$\frac{\Gamma(c)\Gamma(c - |a| - 2)}{\Gamma(c - |a|)\Gamma(c - 1)} [(1 - \lambda\alpha)|a| + (1 - \alpha)(c - |a| - 2)] \leq (1 - \alpha) \left( \frac{1}{(A - B)|\tau|} + 1 \right)$$

*is satisfied, then  $\mathcal{L}(a, c)(f) \in M(\lambda, \alpha)$ .*

**THEOREM 2.** Let  $a, b \in \mathbb{C} \setminus \{0\}$ . Also, let  $c$  be a real number and  $P_1 = P_1(k)$  be given by (1.2). If, for some  $k$  ( $0 \leq k < \infty$ ),  $f \in k - \mathcal{UCV}$ , and the inequality

$$(1 - \lambda\alpha) \frac{|ab|P_1}{c} {}_3F_2(1 + |a|, 1 + |b|, 1 + P_1; 1 + c, 2; 1) + \alpha(\lambda - 1) {}_3F_2(|a|, |b|, P_1; c, 1; 1) \leq 1 - \alpha\lambda \quad (2.3)$$

is satisfied, then  $I_{a,b,c}(f) \in M(\lambda, \alpha)$ .

*Proof.* Let  $f$  be given by (1.1). By (1.4), to show  $I_{a,b,c}(f) \in M(\lambda, \alpha)$ , it is sufficient to prove that

$$\sum_{n=2}^{\infty} n(n - \lambda\alpha n - \alpha + \lambda\alpha) \left| \frac{(a)_{n-1}(b)_{n-1}}{(c)_{n-1}(1)_{n-1}} a_n \right| \leq 1 - \alpha.$$

We will repeat the method of proving used in the proof of Theorem 1. Applying the estimates for the coefficients given by (1.3), and making use of the relations (2.2) and  $|(a)_n| \leq (|a|)_n$ , we get

$$\begin{aligned} & \sum_{n=2}^{\infty} n(n - \lambda\alpha n - \alpha + \lambda\alpha) \left| \frac{(a)_{n-1}(b)_{n-1}}{(c)_{n-1}(1)_{n-1}} a_n \right| \\ & \leq \sum_{n=2}^{\infty} [n(1 - \lambda\alpha) + \alpha(\lambda - 1)] \frac{(|a|)_{n-1}(|b|)_{n-1}(P_1)_{n-1}}{(c)_{n-1}(1)_{n-1}(1)_{n-1}} \\ & = (1 - \lambda\alpha) \sum_{n=2}^{\infty} \frac{|ab|P_1}{c} \frac{(1 + |a|)_{n-2}(1 + |b|)_{n-2}(1 + P_1)_{n-2}}{(1 + c)_{n-2}(1)_{n-2}(2)_{n-2}} \\ & \quad + \alpha(\lambda - 1) \sum_{n=2}^{\infty} \frac{(|a|)_{n-1}(|b|)_{n-1}(P_1)_{n-1}}{(c)_{n-1}(1)_{n-1}(1)_{n-1}} \\ & = (1 - \lambda\alpha) \frac{|ab|P_1}{c} {}_3F_2(1 + |a|, 1 + |b|, 1 + P_1; 1 + c, 2; 1) \\ & \quad + \alpha(\lambda - 1) [{}_3F_2(|a|, |b|, P_1; c, 1; 1) - 1] \\ & \leq 1 - \alpha, \end{aligned}$$

provided the condition (2.3) is satisfied. ■

For the choices of  $|b| = |a|$  and  $b = 1$ , we can deduce further corollaries of Theorem 2 as follows.

**COROLLARY 3.** Let  $a \in \mathbb{C} \setminus \{0\}$ . Suppose that  $|b| = |a|$ . Also, let  $c$  be a real number and  $P_1 = P_1(k)$  be given by (1.2). If  $f \in k - \mathcal{UCV}$  for some  $k$  ( $0 \leq k < \infty$ ) and the inequality

$$(1 - \lambda\alpha) \frac{|a|^2 P_1}{c} {}_3F_2(1 + |a|, 1 + |a|, 1 + P_1; 1 + c, 2; 1) + \alpha(\lambda - 1) {}_3F_2(|a|, |a|, P_1; c, 1; 1) \leq 1 - \alpha\lambda$$

is satisfied, then  $I_{a,|a|,c}(f) \in M(\lambda, \alpha)$ .

COROLLARY 4. Let  $a \in \mathbb{C} \setminus \{0\}$ . Also, let  $c$  be a real number and  $P_1 = P_1(k)$  be given by (1.2). If  $f \in k - \mathcal{UCV}$  for some  $k$  ( $0 \leq k < \infty$ ) and the inequality

$$(1 - \lambda\alpha) \frac{|a|P_1}{c} {}_3F_2(1 + |a|, 2, 1 + P_1; 1 + c, 2; 1) + \alpha(\lambda - 1) {}_3F_2(|a|, 1, P_1; c, 1; 1) \leq 1 - \alpha\lambda$$

is satisfied, then  $\mathcal{L}(a, c)(f) \in M(\lambda, \alpha)$ .

THEOREM 3. Let  $a, b \in \mathbb{C} \setminus \{0\}$ . Also, let  $c$  be a real number and  $P_1 = P_1(k)$  be given by (1.2). If  $f \in k - \mathcal{ST}$ , for some  $k$  ( $0 \leq k < \infty$ ), and the inequality

$$(1 - \lambda\alpha) \frac{|ab|P_1}{c} {}_3F_2(1 + |a|, 1 + |b|, 1 + P_1; 1 + c, 1; 1) + (2 - \alpha - \lambda\alpha) \frac{|ab|P_1}{c} {}_3F_2(1 + |a|, 1 + |b|, 1 + P_1; 1 + c, 2; 1) + (1 - \alpha) {}_3F_2(|a|, |b|, P_1; c, 1; 1) \leq 2(1 - \alpha) \tag{2.4}$$

is satisfied, then  $I_{a,b,c}(f) \in M(\lambda, \alpha)$ .

*Proof.* Let  $f$  be given by (1.1). We will repeat the method of proving used in the proof of Theorem 1. Applying the estimates for the coefficients given by (1.4), and making use of the relations (2.2) and  $|(a)_n| \leq (|a|)_n$ , we get

$$\begin{aligned} & \sum_{n=2}^{\infty} n(n - \lambda\alpha n - \alpha + \lambda\alpha) \left| \frac{(a)_{n-1}(b)_{n-1}}{(c)_{n-1}(1)_{n-1}} a_n \right| \\ & \leq \sum_{n=2}^{\infty} n[n(1 - \lambda\alpha) + \alpha(\lambda - 1)] \frac{(|a|)_{n-1}(|b|)_{n-1}(P_1)_{n-1}}{(c)_{n-1}(1)_{n-1}(1)_{n-1}} \\ & = \sum_{n=2}^{\infty} (n - 1)[(n - 1)(1 - \lambda\alpha) + (1 - \alpha)] \frac{(|a|)_{n-1}(|b|)_{n-1}(P_1)_{n-1}}{(c)_{n-1}(1)_{n-1}(1)_{n-1}} \\ & \quad + \sum_{n=2}^{\infty} [(n - 1)(1 - \lambda\alpha) + (1 - \alpha)] \frac{(|a|)_{n-1}(|b|)_{n-1}(P_1)_{n-1}}{(c)_{n-1}(1)_{n-1}(1)_{n-1}} \\ & = (1 - \lambda\alpha) \sum_{n=2}^{\infty} \frac{(|a|)_{n-1}(|b|)_{n-1}(P_1)_{n-1}}{(c)_{n-1}(1)_{n-2}(1)_{n-2}} + (1 - \alpha) \sum_{n=2}^{\infty} \frac{(|a|)_{n-1}(|b|)_{n-1}(P_1)_{n-1}}{(c)_{n-1}(1)_{n-1}(1)_{n-2}} \\ & \quad + (1 - \lambda\alpha) \sum_{n=2}^{\infty} \frac{(|a|)_{n-1}(|b|)_{n-1}(P_1)_{n-1}}{(c)_{n-1}(1)_{n-1}(1)_{n-2}} + (1 - \alpha) \sum_{n=2}^{\infty} \frac{(|a|)_{n-1}(|b|)_{n-1}(P_1)_{n-1}}{(c)_{n-1}(1)_{n-1}(1)_{n-1}} \\ & = (1 - \lambda\alpha) \sum_{n=2}^{\infty} \frac{(|a|)_{n-1}(|b|)_{n-1}(P_1)_{n-1}}{(c)_{n-1}(1)_{n-2}(1)_{n-2}} + (2 - \alpha - \lambda\alpha) \times \\ & \quad \times \sum_{n=2}^{\infty} \frac{(|a|)_{n-1}(|b|)_{n-1}(P_1)_{n-1}}{(c)_{n-1}(1)_{n-1}(1)_{n-2}} + (1 - \alpha) \sum_{n=2}^{\infty} \frac{(|a|)_{n-1}(|b|)_{n-1}(P_1)_{n-1}}{(c)_{n-1}(1)_{n-1}(1)_{n-1}} \\ & = (1 - \lambda\alpha) \frac{|ab|P_1}{c} {}_3F_2(1 + |a|, 1 + |b|, 1 + P_1; 1 + c, 1; 1) \\ & \quad + (2 - \alpha - \lambda\alpha) \frac{|ab|P_1}{c} {}_3F_2(1 + |a|, 1 + |b|, 1 + P_1; 1 + c, 2; 1) \\ & \quad + (1 - \alpha) [ {}_3F_2(|a|, |b|, P_1; c, 1; 1) - 1 ] \\ & \leq 1 - \alpha \end{aligned}$$

provided the condition (2.4) is satisfied. ■



If  $|b| = |a|$  we can rewrite the Theorem 3 as follows.

COROLLARY 5. *Let  $a, b \in \mathbb{C} \setminus \{0\}$ . Suppose that  $|b| = |a|$ . Also, let  $c$  be a real number and  $P_1 = P_1(k)$  be given by (1.2). If  $f \in k - \mathcal{ST}$  for some  $k$  ( $0 \leq k < \infty$ ) and the inequality*

$$(1 - \lambda\alpha) \frac{|a|^2 P_1}{c} {}_3F_2(1 + |a|, 1 + |a|, 1 + P_1; 1 + c, 1; 1) + (2 - \alpha - \lambda\alpha) \frac{|a|^2 P_1}{c} {}_3F_2(1 + |a|, 1 + |a|, 1 + P_1; 1 + c, 2; 1) + (1 - \alpha) {}_3F_2(|a|, |a|, P_1; c, 1; 1) \leq 2(1 - \alpha)$$

is satisfied, then  $I_{a, |a|, c}(f) \in M(\lambda, \alpha)$ .

In the special case when  $b = 1$ , Theorem 3 immediately yields a result concerning the Carlson-Shaffer operator  $\mathcal{L}(a, c)(f) = I_{a, 1, c}(f)$ .

COROLLARY 6. *Let  $a \in \mathbb{C} \setminus \{0\}$ . Also, let  $c$  be a real number and  $P_1 = P_1(k)$  be given by (1.2). If  $f \in k - \mathcal{UCV}$  for some  $k$  ( $0 \leq k < \infty$ ) and the inequality*

$$(1 - \lambda\alpha) \frac{|a| P_1}{c} {}_3F_2(1 + |a|, 2, 1 + P_1; 1 + c, 1; 1) + (2 - \alpha - \lambda\alpha) \frac{|a| P_1}{c} {}_3F_2(1 + |a|, 2, 1 + P_1; 1 + c, 2; 1) + (1 - \alpha) {}_3F_2(|a|, 1, P_1; c, 1; 1) \leq 2(1 - \alpha)$$

is satisfied, then  $\mathcal{L}(a, c)(f) \in M(\lambda, \alpha)$ .

REFERENCES

[1] O. Altıntaş, S. Owa, *On subclasses of univalent functions with negative coefficients*, Pusan Kyongnam Math. J. **4** (1988), 41-56.  
 [2] L. de Branges, *A proof of the Bieberbach conjecture*, Acta Math. **154** (1985), 137-152.  
 [3] T.R. Caplinger, W.M. Causey, *A class of univalent functions*, Proc. Amer. Math. Soc. **39** (1973), 357-361.  
 [4] B.C. Carlson, D.B. Shaffer, *Starlike and prestarlike hypergeometric functions*, SIAM J. Math. Anal. **15** (1984), 737-745.  
 [5] K.K. Dixit, S.K. Pal, *On a class of univalent functions related to complex order*, Indian J. Pure Appl. Math. **26** (1995), 889-896.  
 [6] A. Gangadharan, T.N. Shanmugam, H.M. Srivastava, *Generalized hypergeometric functions associated with  $k$ -uniformly convex functions*, Comput. Math. Appl. **44** (2002), 1515-1526.  
 [7] A.W. Goodman, *On uniformly convex functions*, Ann. Polon. Math. **56** (1991), 87-92.  
 [8] Y.E. Hohlov, *Operators and operations in the class of univalent functions*, Izv. Vysš. Učebn. Zaved. Matematika **10** (1978), 83-89 (in Russian).  
 [9] S. Kanas, H.M. Srivastava, *Linear operators associated with  $k$ -uniformly convex functions*, Integral Transform. Spec. Funct. **9** (2000), 121-132.  
 [10] S. Kanas, A. Wiśniowska, *Conic regions and  $k$ -uniform convexity*, J. Comput. Appl. Math. **105** (1999), 327-336.  
 [11] S. Kanas, A. Wiśniowska, *Conic regions and  $k$ -starlike functions*, Rev. Roumaine Math. Pures Appl. **45** (2000), 647-657.

- [12] A.O. Mostafa, *A study on starlike and convex properties for hypergeometric functions*, J.Inequal. Pure Appl. Math. **10** (3), Art. 87, (2009), 8 pp (electronic).
- [13] K.S. Padmanabhan, *On a certain class of functions whose derivatives have a positive real part in the unit disc*, Ann. Polon. Math. **23** (1970), 73–81.
- [14] S. Ponnusamy, F. Rønning, *Duality for Hadamard products applied to certain integral transforms*, Complex Variables Theory Appl. **32** (1997), 263-287.
- [15] T.N. Shanmugam, *Hypergeometric functions in the geometric function theory*, Appl. Math. Comput. **187** (2007), 433–444.
- [16] T.N. Shanmugam, S. Sivasubramanian, *On the Hohov convolution of the class  $S_p(\alpha, \beta)$* , Aust. J. Math. Anal. Appl. **2** Art. 7, (2005), 8 pp. (electronic).
- [17] H.M. Srivastava, A.K. Mishra, *Applications of fractional calculus to parabolic starlike and uniformly convex functions*, Comput. Math. Appl. **39** (2000), 57-69.
- [18] H.M. Srivastava, G. Murugusundaramoorthy, S. Sivasubramanian, *Hypergeometric functions in the parabolic starlike and uniformly convex domains*, Integral Transform. Spec. Funct. **18** (2007), 511–520.

(received 17.08.2011; in revised form 16.02.2012; available online 10.06.2012)

Department of Mathematics, University College of Engineering, Anna University of technology  
Chennai, Melpakkam-604 001, India

*E-mail:* sivasaisastha@rediffmail.com

Department of Mathematics, Madras Christian College Tambaram, Chennai-600 59, India

*E-mail:* thomas.rosy@gmail.com

Department of Mathematics, VIT university, Chennai Campus, India