

## A GENERALIZATION OF A THEOREM OF A. D. OTTO

Th. Exarchakos

**Abstract.** In this paper we prove that if  $G$  is a finite  $p$ -group of class  $c$  with  $G/G'$  of exponent  $p^r$  and  $L_i/L_{i+1}$  is cyclic of order  $p^r$  for  $i = 1, 2, \dots, c-1$ , where  $L_i, i = 0, 1, \dots, c$  is the lower central series of  $G$ , then the order of  $G$  divides the order of the group  $A(G)$  of automorphisms of  $G$ .

**Introduction and notation.** Let  $G$  be a finite  $p$ -group of class  $c$ . Let  $G = L_0 \supset L_1 \supset \dots \supset L_c = 1, 1 = Z_0 \subset Z_1 \subset \dots \subset Z_c = G$  be the lower and the upper central series of  $G$  respectively, where  $L_1 = G' = [G, G]$  and  $Z_1 = Z = Z(G)$ . If  $G$  has no non-trivial abelian direct factor, then  $G$  is called a  $PN$ -group. A. D. Otto in [1] proved that if  $G$  is a  $PN$ -group with  $|L_i/L_{i+1}| = p$  for all  $i = 1, 2, \dots, c-1$  and  $\exp(G/G') = p$ , then the order of  $G$  divides the order of the group of automorphisms of  $G$ . We generalize this result by showing that if  $G$  is any finite  $p$ -group with  $L_i/L_{i+1}$  cyclic of order  $p^r$  for all  $i = 1, 2, \dots, c-1$  and  $\exp(G/G') = p^r$ , then  $|G|$  divides  $A(G)$ . We also show that the same result holds if  $Z_i/Z_{i-1}$  is cyclic of order  $p^r, i = 1, 2, \dots, c-1$ , and  $L_j = Z_{c-j}$  for some  $j, 1 \leq j \leq c-1$ . Throughout this paper,  $G$  is a finite non-abelian  $p$ -group,  $|G|$  is the order of  $G, C(p^x)$  is the cyclic group of order  $p^x, A(G), I(G), A_c(G)$  are the groups of automorphisms, inner automorphisms, central automorphisms of  $G$ .

\* \* \*

We begin with

LEMMA 1. *Let  $G$  be a  $PN$ -group. If  $\exp(G/G') \leq |Z|$ , then  $|A_c(G)| \geq |C/G'|$ .*

*Proof.* Let  $|G/G'| = p^m$  and  $G/G' = C(p^{m_1}) \times C(p^{m_2}) \times \dots \times C(p^{m_t})$ , where  $m_1 \geq m_2 \geq \dots \geq m_t \geq 1$  and  $\sum_{j=1}^t m_j = m$ . Similarly let  $|Z| = p^k$  and  $Z = C(p^{k_1}) \times C(p^{k_2}) \times \dots \times C(p^{k_s})$  with  $k_1 \geq k_2 \geq \dots \geq k_s \geq 1$  and  $\sum_{i=1}^s k_i = k$ . If

$a_x$  is the number of times  $p^x$  appears in the invariants of  $G/G'$ , then  $\sum_{x \geq 1}^m x a_x = m$ . Since  $G$  is a  $PN$ -group  $|A_c(G)| = |\text{Hom}(G, Z)| = |\text{Hom}(G/G', Z)|$  [2]. So we have  $|A_c(G)| = |\text{Hom}(G/G', Z)| = |\text{Hom}(\prod_{j=1}^t (C(p^{m_j}), \prod_{i=1}^s C(p^{k_i}))|$ . Hence  $|A_c(G)| = \prod_{j,i}^{t,s} |\text{Hom}(C(p^{m_j}), C(p^{k_i}))| \prod_{j,i}^{t,s} p^{\min(m_j, k_i)} = p^A$  for some  $A$ . Summing powers over  $m_j = 1, 2, \dots, m_1$  for  $k_i = k_s, \dots, k_1$  we have

$$\begin{aligned} A &= \left( \sum_{x \geq 1}^{k_s} x a_x + k_s \sum_{x > k_s}^{m_1} a_x \right) + \dots + \left( \sum_{x \geq 1}^{k_1} x a_x + k_1 \sum_{x > k_1}^{m_1} a_x \right) = \\ &= \sum_{i=2}^s \left( \sum_{x \geq 1}^{k_i} x a_x + k_i \sum_{x > k_i}^{m_1} a_x \right) + \sum_{i=1}^s \left( k_i \sum_{x > k_1}^{m_1} a_x \right) + \sum_{x \geq 1}^{k_1} x a_x = \\ &= \sum_{i=2}^s \theta_i + k \sum_{x > k_1}^{m_1} a_x + \sum_{x \geq 1}^{k_1} x a_x, \text{ where } \theta_i = \sum_{x \geq 1}^{k_i} x a_x + k_i \sum_{x > k_i}^{m_1} a_x. \end{aligned}$$

Since  $k \geq m_1$  we have  $k \sum_{x > k_1}^{m_1} a_x \geq \sum_{x > k_1}^{m_1} x a_x$  and so  $A \geq k \sum_{i=2}^s \theta_i + \sum_{x > k_1}^{m_1} x a_x = \sum_{i=2}^s \theta_i + m$ .

LEMMA 2. [4]. *Let  $G$  be a finite non-abelian  $p$ -group. Let  $G = L_0 \supset L_1 \supset \dots \supset L_c = 1$ ,  $1 = Z_0 \subset Z_1 \subset \dots \subset Z_c = G$  be the lower and the upper central series of  $G$ . If  $L_i/L_{i+1}$  is cyclic of order  $p^r$  for all  $i = 1, 2, \dots, c-1$ . then  $L_i \cap Z_{c-i-1} = L_{i+1}$ ,  $i = 1, 2, \dots, c-1$ .*

LEMMA 3. [3]. *If  $G$  is a finite non-abelian group and  $Z_i/Z_{i-1}$  is cyclic of order  $p^r$  for all  $i = 1, 2, \dots, c-1$ , then  $[G, Z_{i+1}] = Z_i$  for  $i = 1, 2, \dots, c-1$ .*

THEOREM 1. *Let  $G$  be a finite group of order  $p^n$  and class  $c$ . If  $L_i/L_{i+1}$  is cyclic of order  $p^r$  for all  $i = 1, 2, \dots, c-1$  and  $\exp(G/G') = p^r$ , then  $|G|$  divides,  $|A(G)|$ .*

*Proof.* Consider the following:

**A:**  $G$  is a  $PN$ -group. Since  $L_i \subseteq Z_{c-i}$  and  $L_i \not\subseteq Z_{c-i-1}$  we have  $(Z_{c-i}/L_i) \cong (L_i Z_{c-i-1}/L_i) \cong Z_{c-i-1}/L_i \cap Z_{c-i-1} = Z_{c-i-1}/L_{i+1}$  (by Lemma 2). Hence

$$|Z_{c-i}/Z_{c-i-1}| \geq |L_i/L_{i+1}| = p^r$$

for all  $i = 1, 2, \dots, c-1$ . But  $|G/Z_{c-1}| = p^{2r}$  [4] and so  $|G/Z_2| \geq p^{(c-1)r}$  which gives  $|Z_2| \leq p^{n-(c-1)r}$ . If  $|Z| = p^k$  then  $|I(G)| = |G/Z| = p^{n-k}$  and  $|Z_2/Z| \leq p^{n-(c-1)r-k}$ . Since  $L_{c-1} \subseteq Z$  and  $|L_{c-1}| = p^r$  we have  $|Z| \geq p^r = \exp(G/G')$

and so by Lemma 1,  $|A_c(G)| \geq |G/G'|$ . But  $|L_i/L_{i+1}| = p^r$  for  $i = 1, 2, \dots, c-1$  which implies that  $|L_{c-i}| = p^{ir}$  and so  $|L_1| = p^{(c-1)r}$ . Therefore we have  $|G/G'| = |G/L_1| = p^{n-(c-1)r}$ , and so  $|A_c(G)| \geq p^{n-(c-1)r}$ . Since  $A_c(G)$  centralizes  $I(G)$  in  $A(G)$  we have  $|I(G) \cap A_c(G)| = |Z(I(G))| = |Z(G/Z)| = |Z_2/Z| \leq p^{n-(c-1)r-k}$ . Hence

$$\begin{aligned} |A(G)|_p &\geq |I(G)A_c(G)| = |I(G)| \cdot |A_c(G)|/|I(G) \cap A_c(G)| \geq \\ &\geq p^{n-k} p^{n-(c-1)r} / p^{n-(c-1)r-k} = p^n. \end{aligned}$$

**B:**  $G = H \times K$ , where  $H$  is abelian of order  $p^e$  and  $K$  is a  $PN$ -group. By [1],  $|A(G)|_p \geq p^e |A(K)|_p$ . Since  $G = H \times K$ ,  $|G'| = |K'|$ , and by induction  $|L_i(G)| = |L_i(K)|$  for all  $i = 1, 2, \dots, c$ . Hence  $L_i(K)/L_{i+1}(K)$  is cyclic of order  $p^r$  for  $i = 1, 2, \dots, c-1$ . Moreover  $G/G' = H \times K/K'$  and so  $\exp(K/K') \leq p^r$ . But  $\exp(L_i(K)/L_{i+1}(K)) = p^r$  and so  $\exp(K/K') = p^r$ . Therefore by A,  $|A(K)|_p \geq |K|$  and so  $|A(G)|_p \geq p^e$ .  $|K| = |G|$ .

**COROLLARY.** *Let  $G$  be a  $PN$ -group. If  $|L_i/L_{i+1}| = p$ ,  $i = 1, 2, \dots, c-1$ , and  $\exp(G/G') \leq |Z|$ , then  $|G|$  divides  $|A(G)|$ .*

**THEOREM 2.** *Let  $G$  be a finite  $p$ -group of order  $p^n$  and class  $c$ . If  $Z_{i+1}/Z_i$  is cyclic of order  $p^r$  for all  $i = 0, 1, \dots, c-2$ , and  $Z_{c-j} = L_j$  for some  $j$ ,  $1 \leq j \leq c-1$ , then  $|G|$  divides  $|A(G)|$ .*

*Proof.* By Lemma 3,  $L_{j+1} = [L_j, G] = [Z_{c-j}, G] = Z_{c-j-1}$  and so  $p^r = \exp(Z_{c-j}/Z_{c-j-1}) = \exp(L_j/L_{j+1}) \leq \exp(L_{j-1}/L_j) \leq |L_{j-1}/L_j| \leq |Z_{c-j-1}/Z_{c-j}| = p^r$ . Hence  $|L_{j-1}| = |Z_{c-j+1}|$  and since  $L_{j-1} \subseteq Z_{c-j+1}$  we have  $L_{j-1} = Z_{c-j+1}$ . Therefore  $L_j = Z_{c_j}$  for all  $j = 1, 2, \dots, c$ , and so  $L_j/L_{j+1}$  is cyclic of order  $p^r$  for all  $j$ . By [4],  $G/Z_{c-1} = p^{2r}$  and so  $|G/L_1| = |G/G'| = p^{2r}$ . Let  $p^{m_1} \geq p^{m_2} \geq \dots \geq p^{m_t}$  by the invariants of  $G/G'$ , if  $m_2 < r$ , then  $\exp(L_1/L_2) \geq p^{m_2} < p^r$ , which is a contradiction. Hence  $m_2 \geq r$  and so  $m_1 = m_2 = r$  and  $\exp(G/G') = p^r$ . The result follows from Theorem 1.

## REFERENCES

- [1] A. D. Otto, *Central automorphisms of finite  $p$ -group*, Trans, Amer. Math. Soc. **125** (1966), 280-287.
- [2] J. E. Adney, J. Yen, *Automorphisms of a  $p$ -group*, Illinois J. Math. **9** (1965), 137-143.
- [3] G. Baumslag, N. Blackburn, *Groups with cyclic upper central factors*, Proc. London Math. Soc. **10** (1960), 531-544.
- [4] J. A. Gallian, *Finite  $p$ -groups with homocyclic central factors*, Canad. J. Math. **26** (1974), 636-643.
- [5] N. Blackburn, *On special class of  $p$ -groups*, Acta Math **100** (1958), 45-92.

Marasleios Academy of Athens  
4. Marasli Street  
Athens, GREECE

(Received 07 04 1980)