

A PROPERTY OF CANONICAL GRAPHS

Aleksandar Torgašev

Abstract. A finite connected graph is called canonical if no two of its vertices have the same neighbours. In this paper we prove that in all but a sequence of exceptional cases, deleting of a suitable chosen vertex in a canonical graph also gives a connected canonical graph. This property can have applications in various hereditary problems in the spectral Theory of Graphs.

In this paper we consider only finite connected graphs without loops or multiple edges. The vertex set of a graph G is denoted by $V(G)$, and the number of its vertices by $|G|$. Relation $H \subseteq G$ will always mean that H is an induced subgraph of a graph G . For any two vertices u, v of G , $uv = 1$ will mean that u is adjacent to v , while $uv = 0$ will mean that u is nonadjacent to v .

The graph obtained by deleting a vertex $x \in V(G)$ from G is denoted by $G - x$. It can be connected or disconnected. But, as is well known, there is at least one vertex $x \in V(G)$ such that the corresponding graph $G - x$ is also connected.

Next, we say that two vertices $u, v \in V(G)$ are equivalent in G and we denote it by $u \sim v$ if we have

$$(1) \quad uv = 0 \quad \text{and} \quad ru = rv$$

for any vertex $r \in V(G) \setminus \{u, v\}$, thus if and only if u and v have the same neighbours in G . Relation \sim is obviously an equivalence relation on the vertex set $V(G)$. The corresponding quotient graph is denoted by g and called the *canonical graph* of G . This graph is also connected.


For instance, if $G = K_{m_1 \dots m_p}$ ($p \geq 2$) is a complete p -partite graph, then its canonical graph is the complete graph K_p . The canonical graph of the complete graph K_n is the same graph K_n .

We say that a graph G is *canonical* if $|G| = |g|$, i.e. if G has no two equivalent vertices.

If g is the canonical graph of G , $|g| = k$ and N_1, \dots, N_k are the corresponding sets of equivalent vertices in G , we denote


$$G = g(N_1, \dots, N_k).$$

We call N_1, \dots, N_k the characteristic sets of G . Obviously, each set $N_i \subseteq V(G)$ ($i = 1, \dots, k$) consists only of isolated vertices, and if at least one edge between the sets N_i and N_j ($i \neq j$) is present, then all possible edges between these sets are also present. Therefore, it is very convenient to display the sets N_i ($i = 1, \dots, k$) by white (that is, empty) circles, and all possible edges between the sets N_i and N_j ($i \neq j$) by only one edge between the corresponding circles. If, for example, G is the complete bipartite graph with characteristic sets N_1, N_2 , we can simply draw



We note that many hereditary problems in the Spectral Theory of Graphs can be reduced to finding firstly the corresponding sets of canonical graphs. Compare for instance the papers [3], [4], [5], [6], [7], [8], [9], [10], [11], or the monograph [2], where many results from these papers are presented. Therefore, the importance of the following hereditary property of canonical graphs is clear.

THEOREM. *In all but a sequence of exceptional cases, each canonical graph G with n vertices ($n \geq 2$) contains, as an induced subgraph, a connected and canonical subgraph H on $n - 1$ vertices. The exceptional graphs are*



where $a_j b_i = 1$ ($i \leq j$; $i, j = 1, \dots, m$). We obviously have that $T_0 \subseteq T_1 \subseteq T_2 \subseteq \dots$.

Proof. First, it is trivial to check that all graphs T_m ($m \geq 0$) are exceptional graphs. Graphs $G - a_0$ and $G - b_0$ are disconnected, while for any $m \geq 1$ and $i = 0, \dots, m - 1$ we have

$$\begin{aligned} a_0 \sim b_1 & \quad (\text{in } G - a), & b_0 \sim a_m & \quad (\text{in } G - b), \\ a \sim a_1 & \quad (\text{in } G - b_1), & b \sim b_m & \quad (\text{in } G - a_m), \\ a_i \sim a_{i+1} & \quad (\text{in } G - b_{i+1}), & b_i \sim b_{i+1} & \quad (\text{in } G - a_i). \end{aligned}$$

Next, we consider any connected canonical graph $G \neq T_m$ ($m = 0, 1, 2, \dots$), which is “bad” in the sense that deleting of any its vertex gives a disconnected or a

noncanonical graph. We shall prove that this assumption in all possible cases gives a contradiction, thus that such graphs really do not exist.

Firstly choose any vertex x of G such that the subgraph $G - x$ is connected. Then $G - x$ is a noncanonical graph. Let $G - x = f(M_1, \dots, M_k)$ ($k \geq 1$), where M_1, \dots, M_k are the characteristic sets of $G - x$, f is the corresponding canonical graph of $G - x$ (also connected), and at least one of the sets M_1, \dots, M_k is not a singleton. Obviously $k \geq 2$. Rearranging the vertices of f , we can assume that $|M_1| \geq 2$.

If $|M_i| \geq 2$ for some $i \in \{1, \dots, k\}$ and if $u, v \in M_i$ ($u \neq v$), the fact that G is a canonical graph implies that exactly one of u, v is adjacent to x . Therefore, we immediately get

$$(2) \quad |M_i| \leq 2 \quad (i = 1, \dots, k),$$

and consequently $|M_1| = 2$. We assume that $M_1 = \{y, z\}$, where $xy = 1$ and $xz = 0$.

Now assume that some $|M_j| = 2$ ($j > 1$), for instance that $M_2 = \{c, d\}$, where $xc = 1$ and $xd = 0$. Deleting then the vertex d from G we obviously get a connected graph. It is also easily seen that $G - d$ is a canonical graph, what is a contradiction. Hence we must have

$$(3) \quad |M_j| = 1 \quad (j = 2, \dots, k).$$

Thus, except y and z , there is no other equivalent pair of vertices in $G - x$.

Denoting $P = V(G) \setminus \{x, y, z\}$, we obviously have that $P \neq \emptyset$.

Deleting the vertex z from G we evidently get a connected subgraph. Since G is bad, we see that $G - z$ is a noncanonical graph. Since for any two vertices $r, s \in P$ we also have

$$r \not\sim s, \quad r \not\sim y \quad (\text{in } G - z),$$

we conclude that there is a vertex $t \in P$ such that

$$(4) \quad x \sim t \quad (\text{in } G - z).$$

Hence $xt = 0$, and we easily conclude that $yt = zt = 1$. Moreover, $rx = rt$ and $ry = rz$ for any other vertex $r \in P \setminus \{t\}$.

Since $G \neq xytz = T_0$, we have that $|P| \geq 2$. If $|P| = 2$, thus if $P = \{t, r\}$, we necessarily have that $rt = rx = ry = rz = 1$ since $f = try$ is a canonical graph, whence we obviously get that $G - r$ is a connected and canonical graph (a contradiction). Hence, we can assume that $|P| \geq 3$. Now delete the vertex y of G .

1° Assume first that $G - y$ is a disconnected graph.

Since f is a connected graph, it is easy to see that $rx = 0$ and consequently $rt = 0$ for any vertex $r \neq x, y, z, t$.

Denote next the vertex t by p_1 and the vertex z by q_1 .

Deleting now the vertex p_1 from G , we obviously get a connected graph $G - p_1$. Since G is bad, $G - p_1$ is a noncanonical graph. Discussing all the possible pairs

of vertices in $G - p_1$ as candidates for equivalent vertices, we conclude there is a vertex $q_2 \neq x, y, p_1, q_1$ such that

$$(5) \quad q_1 \sim q_2 \quad (\text{in } G - p_1).$$

Then $q_2x = q_2y = q_2p_1 = q_2q_1 = 0$, and $rx = rp_1 = 0$, $ry = rq_1 = rq_2$ for any vertex $r \neq x, y, p_1, q_1, q_2$. Making use of the last relation, we easily conclude that $G - q_2$ is a connected, thus a noncanonical graph. Discussing now all the possible pairs of vertices as candidates for two equivalent vertices in $G - q_2$, we conclude that there is a new vertex $p_2 \neq x, y, p_1, q_1, q_2$ such that

$$(6) \quad p_1 \sim p_2 \quad (\text{in } G - q_2).$$

Hence $p_2x = p_2p_1 = 0$, $p_2y = p_2q_1 = p_2q_2 = 1$, and $rp_2 = 0$ for any other vertex $r \neq x, y, p_1, q_1, p_2, q_2$.

Now, delete the vertex p_2 from G . If $G - p_2$ is disconnected, we conclude that $G = xyp_1q_1p_2q_2 = T_1$, which contradicts the assumption $G \neq T_m$ ($m = 0, 1, 2, \dots$). Thus, $G - p_2$ must be a connected (and hence a noncanonical) graph. Therefore, we conclude that there is a new vertex q_3 such that

$$(7) \quad q_2 \sim q_3 \quad (\text{in } G - p_2).$$

Then $q_3x = q_3y = q_3p_1 = q_3q_1 = q_3p_2 = q_3q_2 = 0$, and $rx = rp_1 = rp_2 = 0$, $ry = rq_1 = rq_2 = rq_3$ for any other vertex $r \neq x, y, p_1, q_1, p_2, q_2, q_3$. Deleting now the vertex q_3 from G , we conclude that $G - q_3$ is a connected (thus a noncanonical) graph. Discussing all the possible pairs of vertices in $G - q_3$ as candidates for a pair of equivalent vertices, we conclude that there is a new vertex p_3 such that

$$(8) \quad p_2 \sim p_3 \quad (\text{in } G - q_3).$$

Then $p_3x = p_3p_1 = p_3p_2 = 0$, $p_3y = p_3q_1 = p_3q_2 = p_3q_3 = 1$ and $rp_3 = 0$ for any other vertex $r \neq x, y, p_1, q_1, p_2, q_2, p_3, q_3$. Assuming that $G - p_3$ is a disconnected graph, we get the contradiction $G = xyp_1q_1p_2q_2p_3q_3 = T_2$. Hence, $G - p_3$ is a connected (thus a noncanonical) graph. Continuing this procedure, after finitely many steps, we conclude that there is a positive integer m such that

$$G = xyp_1q_1 \dots p_{m+1}q_{m+1} = T_m,$$

which is a contradiction again.

Hence, the case when $G - y$ is a disconnected graph is contradictory.

2° Now assume that $G - y$ is a connected (noncanonical) graph.

Since $G \neq T_0$, there is at least one vertex $r \neq x, y, z, t$ and we have $rx = rt$, $ry = rz$ for any such a vertex r . Since $G - y$ is a connected noncanonical graph, we may conclude that there is a new vertex $x_1 \neq x, y, z, t$ such that

$$(9) \quad x \sim x_1 \quad (\text{in } G - y).$$

Then $x_1x = x_1y = x_1z = x_1t = 0$ and $rx = rt = rx_1$, $ry = rz$ for any vertex $r \neq x, y, z, t, x_1$.

Since G is a connected graph, we see that there is at least one vertex $r \neq x, y, z, t, x_1$. By $rx = rt = rx_1$ for any such r , we may conclude that $G - x_1$ is a connected (thus a noncanonical) graph. Hence, one may conclude that there is a new vertex $y_1 \neq x, y, z, t, x_1$ such that

$$(10) \quad y_1 \sim y \quad (\text{in } G - x_1).$$

Therefore, we easily get $y_1x = y_1t = y_1x_1 = 1$, $y_1y = y_1z = 0$, and $rx = rt = rx_1$, $ry = rz = ry_1$ for any other vertex $r \neq x, y, z, t, x_1, y_1$.

Now we delete the vertex y_1 from G . If $G - y_1$ is a disconnected graph, we can put $x_1 = x'$, $y_1 = y'$, $y = z'$, $x = t'$, to get a contradiction, exactly as it has been done in the case 1°. Hence, we can assume that $G - y_1$ is a connected (thus a noncanonical) graph. Now, one can see that there must exist a new vertex $x_2 \neq x, y, z, t, x_1, y_1$ of G such that

$$(11) \quad x_2 \sim x_1 \quad (\text{in } G - y_1).$$

Then $x_2x = x_2y = x_2z = x_2t = x_2x_1 = x_2y_1 = 0$, and $rx = rt = rx_1 = rx_2$, $ry = rz = ry_1$ for any vertex $r \neq x, y, z, t, x_1, y_1, x_2$. Therefore, we easily conclude that $G - x_2$ is a connected (hence a noncanonical) graph. Moreover, one can conclude that, as the only possible case, there is a new vertex $y_2 \neq x, y, z, t, x_1, y_1, x_2$ such that

$$(12) \quad y_2 \sim y_1 \quad (\text{in } G - x_2).$$

Then $y_2x = y_2t = y_2x_1 = y_2x_2 = 1$, $y_2y = y_2z = y_2y_1 = 0$, and $rx = rt = rx_1 = rx_2$, $ry = rz = ry_1 = ry_2$ for any vertex $r \neq x, y, z, t, x_1, y_1, x_2, y_2$.

Continuing this procedure, we conclude that there is a positive integer m such that

$$G = xytzx_1y_1x_2y_2 \dots x_my_m = T_m,$$

which is again a contradiction.

This proves that case 2° is also impossible, hence our theorem is completely proved.

REFERENCES

- [1] D. Cvetković, M. Doob, H. Sachs, *Spectra of Graphs — Theory and Application*, Deutsch. Verl. Wiss., Berlin, 1980; Academic Press, New York, 1980.
- [2] D. Cvetković, M. Doob, I. Gutman, A. Torgašev, *Recent Results in the Theory of Graph Spectra*, North-Holland, Amsterdam, 1988.
- [3] M. Lepović, *On maximal canonical graphs with exactly 6 nonzero eigenvalues*, Glasnik Matem. (1989), (to appear).
- [4] M. Lepović, *On graphs whose energy does not exceed 4*, Publ. Inst. Math. (Beograd) **49** (63) (1991), 6–16.
- [5] A. Torgašev, *On infinite graphs with three and four nonzero eigenvalues*, Bull. Acad. Serbe Sci. Arts (Sci. Math.) (76) **11** (1981), 39–48.
- [6] A. Torgašev, *On infinite graphs with five nonzero eigenvalues*, Bull. Acad. Serbe Sci. Arts (Sci. Math.) (79) **12** (1982), 31–38.

- [7] A. Torgašev, *On graphs with exactly two negative eigenvalues*, Math. Nachr. **122** (1985), 135–140.
- [8] A. Torgašev, *On graphs with a fixed number of negative eigenvalues*, Discrete Math. **57** (1985), 311–317.
- [9] A. Torgašev, *On graphs with exactly three negative eigenvalues*, In: *Graph Theory, Proc. Sixth Yug. Sem. Graph Theory*, Dubrovnik, April 18–19, 1985, (Ed. R. Tošić, D. Aćeketa, V. Petrović), Inst. Math., Novi Sad, 1986, pp. 219–232.
- [10] A. Torgašev, *Graphs whose energy does not exceed 3*, Czech. Math. **36** (1986), 167–171.
- [11] A. Torgašev, *Maximal canonical graphs with three negative eigenvalues*, Publ. Inst. Math. (Beograd) **45 (59)** (1989), 7–10.

Matematički fakultet
Studentski trg 16, p.p. 550
11000 Beograd, Yugoslavia

(Received 07 02 1989)