

## THE STRONGLY ASYMMETRIC GRAPHS OF ORDER 6 AND 7

Mirko Lepović

**Abstract.** Let  $G$  be an arbitrary connected simple graph of order  $n$ .  $G$  is called strongly asymmetric graph if all induced overgraphs of  $G$  of order  $n + 1$  are nonisomorphic. We give the list of all strongly asymmetric graphs of order 6 and 7. Also we prove that there exist exactly 16 asymmetric graphs of order 7 which are not strongly asymmetric.

We consider only finite connected graphs having no loops or multiple edges. The vertex set of a graph  $G$  is denoted by  $V(G)$ , and its order (number of vertices) by  $|G|$ .

Let  $S$  be any (possibly empty) subset of the vertex set  $V(G)$ . Denote by  $G_S$  the graph obtained of the graph  $G$  by adding a new vertex  $x(x \in V(G))$ , which is adjacent exactly to vertices from  $S$ . Graph  $G$  is obviously an induced subgraph of the graph  $G_S$ , and  $G_S$  is an induced overgraph of  $G$ . Varying the set  $S \subseteq V(G)$  we get  $2^{|G|}$  such graphs  $G_S$ . The family of all such graphs is denoted by  $\mathcal{G}(G)$ . We shall call  $\mathcal{G}(G)$  the *overset* of  $G$ . We say that  $\mathcal{G}(G)$  is connected if every graph  $H \in \mathcal{G}(G)$  except  $G_\emptyset$  so does. It is easily seen that this happens if and only if the graph  $G$  is connected.

In [4] we introduced the following definition.

*Definition 1.* Graph  $G$  is called strongly asymmetric graph if for any two sets  $S_1, S_2 \subseteq V(G)$  ( $S_1 \neq S_2$ ), the graphs  $G_{S_1}, G_{S_2}$  are nonisomorphic.

Let  $\Gamma(G)$  be the automorphism group of a graph  $G$ . As is well known, if  $\Gamma(G)$  is trivial, then  $G$  is called asymmetric.

PROPOSITION 1. [4] *Every strongly asymmetric graph is also asymmetric.*

In [4] was proved that the class of all strongly asymmetric graphs is nonempty. In the same paper, many other properties of the strongly asymmetric graphs are also proved. We suppose that it can be also of some interest to describe all strongly

asymmetric graphs with small number of vertices. In [4] was proved that there are no strongly asymmetric graphs of order 3, 4, 5. In the following theorem we describe all strongly asymmetric graphs of order 6 and 7. We shall use here the list of all connected graphs with 6 vertices (published in [1]), and the corresponding list of all connected graphs with 7 vertices (published in [2]).

**THEOREM 1.** *There exist exactly 8 strongly asymmetric graphs of order 6. Their ordering numbers (according to [1]) are 46, 59, 60, 67, 77, 85, 87, 98.*

*There exist exactly 128 strongly asymmetric graphs of order 7. Their ordering numbers (according to [2]) are 3, 21, 39, 52, 64, 67, 68, 74, 75, 77, 98, 100, 126, 128, 130, 133, 136, 149, 150, 158, 159, 160, 161, 165, 167, 171, 177, 189, 194, 196, 201, 205, 226, 232, 234, 236, 243, 249, 255, 267, 268, 270, 272, 280, 281, 283, 284, 286, 288, 289, 291, 295, 296, 306, 307, 309, 314, 315, 322, 348, 367, 368, 370, 373, 376, 377, 380, 389, 394, 395, 399, 400, 404, 406, 407, 414, 416, 418, 419, 428, 442, 443, 448, 455, 458, 470, 473, 477, 502, 506, 517, 519, 520, 522, 527, 532, 533, 535, 544, 547, 552, 554, 559, 560, 578, 581, 584, 591, 606, 627, 628, 641, 645, 646, 650, 652, 655, 671, 674, 676, 692, 727, 745, 748, 757, 761, 785, 793.*

The proof is obtained by a straightforward use of computer.

We also note that there are asymmetric graphs which are not strongly asymmetric. Moreover, in [4] was proved that there are infinitely many such graphs.

It is not difficult to see that every asymmetric graph of order 6 is also strongly asymmetric. But, this is not so for graphs of order 7 (and for higher orders).





The following proposition describes all asymmetric graphs of order 7, which are not strongly asymmetric. It is also checked by the computer methods.

**PROPOSITION 2.** *There exist exactly 16 asymmetric graphs of order 7, which are not strongly asymmetric. They are represented in the figure.*

*Proof.* It is easy to check that all mentioned graphs are asymmetric. In order to prove that a graph  $G$  is not strongly asymmetric, it is sufficient to find two subset  $R, S \subseteq V(G)$  ( $R \neq S$ ) such that the corresponding overgraphs  $G_R, G_S \in \mathcal{G}(G)$  are isomorphic.

Indeed, for any graph  $G_i$  ( $i = 1, 2, \dots, 16$ ) from the figure we define the subsets  $R_i, S_i$  in the following way:

$$\begin{aligned}
 R_1 &= \{3, 6, 7\}, & S_1 &= \{1, 5, 6\}; & R_2 &= \{3, 5, 6\}, & S_2 &= \{2, 3, 7\}; \\
 R_3 &= \{5, 7\}, & S_3 &= \{4, 6\}; & R_4 &= \{2, 3, 6\}, & S_4 &= \{3, 4, 7\}; \\
 R_5 &= \{1, 6, 7\}, & S_5 &= \{2, 4, 7\}; & R_6 &= \{1, 3, 6, 7\}, & S_6 &= \{1, 2, 4, 7\}; \\
 R_7 &= \{2, 3, 5\}, & S_7 &= \{3, 6, 7\}; & R_8 &= \{2, 4, 7\}, & S_8 &= \{2, 3, 6\}; \\
 R_9 &= \{1, 2, 4\}, & S_9 &= \{2, 3, 7\}; & R_{10} &= \{4, 5\}, & S_{10} &= \{6, 7\}; \\
 R_{11} &= \{1, 3, 4\}, & S_{11} &= \{2, 3, 5\}; & R_{12} &= \{3, 4, 7\}, & S_{12} &= \{2, 4, 6\}; \\
 R_{13} &= \{2, 5, 6\}, & S_{13} &= \{1, 5, 7\}; & R_{14} &= \{1, 5, 7\}, & S_{14} &= \{3, 6, 7\}; \\
 R_{15} &= \{4, 5, 6, 7\}, & S_{15} &= \{1, 2, 6, 7\}; & R_{16} &= \{1, 5, 7\}, & S_{16} &= \{2, 5, 6\}.
 \end{aligned}$$



For any two graphs  $G_{R_i}, G_{S_i} \in \mathcal{G}(G_i)$  ( $i = 1, 2, \dots, 16$ ) the new vertex  $x$  added to the graph  $G_i$  is denoted by 8. The corresponding vertex sets of the graphs  $G_{R_i}, G_{S_i}$  are denoted by  $V(G_R^i)$  and  $V(G_S^i)$ , respectively. Now, we define the permutations  $f_i: V(G_R^i) \rightarrow V(G_S^i)$  in the following way:

$$\begin{aligned} f_1 &= (7, 6, 5, 8, 1, 2, 3, 4), & f_2 &= (8, 7, 4, 3, 2, 5, 6, 1), \\ f_3 &= (6, 8, 4, 5, 3, 7, 1, 2), & f_4 &= (7, 4, 2, 3, 8, 1, 6, 5), \\ f_5 &= (3, 1, 2, 7, 8, 4, 6, 5), & f_6 &= (3, 1, 2, 7, 8, 4, 6, 5), \\ f_7 &= (8, 7, 2, 3, 4, 5, 6, 1), & f_8 &= (2, 1, 7, 6, 8, 3, 4, 5), \\ f_9 &= (7, 5, 4, 1, 2, 8, 3, 6), & f_{10} &= (6, 8, 7, 1, 3, 4, 5, 2), \\ f_{11} &= (5, 4, 7, 1, 3, 8, 2, 6), & f_{12} &= (8, 4, 5, 7, 6, 3, 2, 1), \\ f_{13} &= (6, 7, 8, 1, 2, 4, 5, 3), & f_{14} &= (3, 8, 6, 7, 1, 5, 4, 2), \\ f_{15} &= (7, 6, 8, 2, 1, 4, 5, 3), & f_{16} &= (7, 6, 8, 5, 4, 1, 2, 3). \end{aligned}$$

By these permutations, we immediately get that graph  $G_{R_i}$  is isomorphic to the graph  $G_{S_i}$  ( $i = 1, 2, \dots, 16$ ), and consequently, the graphs from the figure are not strongly asymmetric.  $\square$

#### REFERENCES

- [1] D. Cvetković, M. Petrić, *A table of connected graphs on six vertices*, Discrete Math. **50** (1984), 37–49.
- [2] D. Cvetković, M. Doob, I. Gutman, A. Torgašev, *Recent Results in the Theory of Graph Spectra*, North-Holland, Amsterdam, 1988.
- [3] F. Harary, *Graph Theory*, Reading, Boston, Massachusetts, 1963.
- [4] M. Lepović, *On strongly asymmetric graphs*, Discrete Math., to appear.

Prirodno-matematički fakultet  
34001 Kragujevac, p.p.60  
Jugoslavija

(Received 12 01 1993)  
(Revised 03 02 1994)