

EQUITORSION GEODESIC MAPPINGS OF GENERALIZED RIEMANNIAN SPACES

Svetislav M. Minčić and Mića S. Stanković

Communicated by Mileva Prvanović

Abstract. We define an equitorsion geodesic mapping of two generalized Riemannian spaces and obtain some invariant geometric objects of this mapping, generalizing the Weil's tensor.

0. Introduction

A generalized Riemannian space GR_N in the sense of Eisenhart's definition [1] is a differentiable N -dimensional manifold, equipped with nonsymmetric basic tensor g_{ij} . Consider two N -dimensional generalized Riemannian spaces GR_N and $G\overline{R}_N$. Connexion coefficients of these spaces are generalized Cristoffel's symbols of second kind, respectively Γ_{jk}^i and $\overline{\Gamma}_{jk}^i$. Generally it is $\Gamma_{jk}^i \neq \Gamma_{kj}^i$.

One says that reciprocal one valued mapping $f : GR_N \rightarrow G\overline{R}_N$ is *geodesic* [5] (G-mapping), if geodesics of the space GR_N pass to geodesics of the space $G\overline{R}_N$. We can consider these spaces in the common by this mapping system of local coordinates. In the corresponding points $M(x)$ and $\overline{M}(x)$ we can put

$$(0.1) \quad \overline{\Gamma}_{jk}^i(x) = \Gamma_{jk}^i(x) + P_{jk}^i(x) \quad (i, j, k = 1, \dots, N),$$

where $P_{jk}^i(x)$ is the *deformation tensor* of the connection Γ of GR_N according to the mapping $f : GR_N \rightarrow G\overline{R}_N$.

A necessary and sufficient condition that the mapping $f : GR_N \rightarrow G\overline{R}_N$ be geodesic (see [5]) is that the deformation tensor P_{jk}^i from (0.1) at the mapping f has the form

$$(0.2) \quad P_{jk}^i(x) = \delta_j^i \psi_k(x) + \delta_k^i \psi_j(x) + \xi_{jk}^i(x),$$

AMS Subject Classification (1991): Primary 65M12

Key words and phrases: Geodesic mapping, generalized Riemannian space, equitorsion geodesic mapping, ET-projective parameter, ET-projective tensor

Supported by Grant 04M03D of RFNS through Math. Inst. SANU.

where

$$(0.3) \quad \psi_i(x) = \frac{1}{N+1}(\bar{\Gamma}_{i\alpha}^\alpha(x) - \Gamma_{i\alpha}^\alpha(x)), \quad \xi_{jk}^i(x) = P_{jk}^i = \frac{1}{2}(P_{jk}^i - P_{kj}^i).$$

Notice that in GR_N we have

$$(0.4) \quad \Gamma_{i\alpha}^\alpha = 0,$$

(eq. (2.10) in [5]).

In a generalized Riemannian space one can define four kinds of covariant derivatives [2, 3]. For example, for a tensor a_j^i in GR_N we have

$$\begin{aligned} a_{j1}^i|_m &= a_{j,m}^i + \Gamma_{\alpha m}^i a_j^\alpha - \Gamma_{jm}^\alpha a_\alpha^i, & a_{j2}^i|_m &= a_{j,m}^i + \Gamma_{m\alpha}^i a_j^\alpha - \Gamma_{mj}^\alpha a_\alpha^i, \\ a_{j3}^i|_m &= a_{j,m}^i + \Gamma_{\alpha m}^i a_j^\alpha - \Gamma_{mj}^\alpha a_\alpha^i, & a_{j4}^i|_m &= a_{j,m}^i + \Gamma_{m\alpha}^i a_j^\alpha - \Gamma_{jm}^\alpha a_\alpha^i. \end{aligned}$$

Denote by $\left|, \right|_{\frac{\theta}{\theta}}$ a covariant derivative of the kind θ in GR_N and $G\bar{R}_N$ respectively.

In the case of the space GR_N we have five independent curvature tensors [4] (in [4] R is denoted by \tilde{R}):

$$\begin{aligned} R_1^i{}_{jmn} &= \Gamma_{jm,n}^i - \Gamma_{jn,m}^i + \Gamma_{jm}^\alpha \Gamma_{\alpha n}^i - \Gamma_{jn}^\alpha \Gamma_{\alpha m}^i, \\ R_2^i{}_{jmn} &= \Gamma_{mj,n}^i - \Gamma_{nj,m}^i + \Gamma_{mj}^\alpha \Gamma_{n\alpha}^i - \Gamma_{nj}^\alpha \Gamma_{m\alpha}^i, \\ R_3^i{}_{jmn} &= \Gamma_{jm,n}^i - \Gamma_{nj,m}^i + \Gamma_{jm}^\alpha \Gamma_{n\alpha}^i - \Gamma_{nj}^\alpha \Gamma_{\alpha m}^i + \Gamma_{nm}^\alpha (\Gamma_{\alpha j}^i - \Gamma_{j\alpha}^i), \\ R_4^i{}_{jmn} &= \Gamma_{jm,n}^i - \Gamma_{nj,m}^i + \Gamma_{jm}^\alpha \Gamma_{n\alpha}^i - \Gamma_{nj}^\alpha \Gamma_{\alpha m}^i + \Gamma_{mn}^\alpha (\Gamma_{\alpha j}^i - \Gamma_{j\alpha}^i), \\ R_5^i{}_{jmn} &= \frac{1}{2}(\Gamma_{jm,n}^i + \Gamma_{mj,n}^i - \Gamma_{jn,m}^i - \Gamma_{nj,m}^i + \Gamma_{jm}^\alpha \Gamma_{\alpha n}^i + \Gamma_{mj}^\alpha \Gamma_{n\alpha}^i \\ &\quad - \Gamma_{jn}^\alpha \Gamma_{m\alpha}^i - \Gamma_{nj}^\alpha \Gamma_{\alpha m}^i). \end{aligned}$$

By virtue of the geodesic mapping $GR_N \rightarrow G\bar{R}_N$ we obtain tensors $\bar{R}_{\theta}^i{}_{jmn}$ ($\theta = 1, \dots, 5$), where for example

$$(0.5) \quad \bar{R}_1^i{}_{jmn} = \bar{\Gamma}_{jm,n}^i - \bar{\Gamma}_{jn,m}^i + \bar{\Gamma}_{jm}^\alpha \bar{\Gamma}_{\alpha n}^i - \bar{\Gamma}_{jn}^\alpha \bar{\Gamma}_{\alpha m}^i.$$

In the case of geodesic mapping $f : A_N \rightarrow \bar{A}_N$ of the symmetric affine connection spaces A_N and \bar{A}_N we have an invariant geometric object

$$(0.6) \quad W^i{}_{jmn} = R^i{}_{jmn} + \frac{2}{1+N} \delta_j^i R_{mn} + \frac{1}{N^2-1} [\delta_m^i (NR_{jn} + R_{nj}) - \delta_n^i (NR_{jm} + R_{mj})],$$

where R^i_{jmn} is Riemann-Cristoffel's curvature tensor of the space A_N , and R_{jm} Richi's tensor. For Riemannian space (0.6) reduces to [6, p. 80]

$$(0.6') \quad W^i_{jmn} = R^i_{jmn} + \frac{1}{N-1}(\delta^i_m R_{jn} - \delta^i_n R_{jm}).$$

The object W^i_{jmn} is called Weil's tensor, or a tensor of projective curvature [6]. Having a geodesic mapping of two generalized Riemannian spaces, we can not find a generalization of Weil's tensor as an invariant of geodesic mapping in general case. For that reason we define a special geodesic mapping.

A mapping $f : GR_N \rightarrow G\bar{R}_N$ is *equitorsion geodesic mapping* (ETG mapping) if the torsion tensor of the spaces GR_N and $G\bar{R}_N$ are equal. Then from (0.1) and (0.2)

$$(0.7) \quad \xi^h_{ij}(x) = 0.$$

1. ET-projective parameter of the first kind

Using (0.1) and (0.5), we get a relation between the first kind curvature tensors of the spaces GR_N and $G\bar{R}_N$

$$\bar{R}^i_{1jmn} = R^i_{1jmn} + P^i_{jm|n} - P^i_{jn|m} + P^{\alpha}_{jm} P^i_{\alpha n} - P^{\alpha}_{jn} P^i_{\alpha m} + 2\Gamma^{\alpha}_{m\nu} P^i_{j\alpha}.$$

Denoting $\psi_{mn} = \psi_{m|n} - \psi_m \psi_n$ ($\theta = 1, 2$) and substituting P with respect to (0.2), we obtain

$$(1.1) \quad \begin{aligned} \bar{R}^i_{1jmn} = R^i_{1jmn} &+ \delta^i_j (\psi_{mn} - \psi_{nm}) + \delta^i_m \psi_{jn} - \delta^i_n \psi_{jm} - \delta^i_m \xi^{\alpha}_{jn} \psi_{\alpha} \\ &+ \delta^i_n \xi^{\alpha}_{jm} \psi_{\alpha} + \xi^i_{jm|n} - \xi^i_{jn|m} + 2\psi_j \xi^i_{mn} + \xi^{\alpha}_{jm} \xi^i_{\alpha n} - \xi^{\alpha}_{jn} \xi^i_{\alpha m} \\ &+ 2\Gamma^i_{m\nu} \psi_j + 2\Gamma^{\alpha}_{m\nu} \psi_{\alpha} \delta^i_j + 2\Gamma^{\alpha}_{m\nu} \xi^i_{j\alpha}. \end{aligned}$$

From (0.7) and (1.1) we get

$$(1.2) \quad \bar{R}^i_{1jmn} = R^i_{1jmn} + \delta^i_j (\psi_{mn} - \psi_{nm}) + \delta^i_m \psi_{jn} - \delta^i_n \psi_{jm} + 2\Gamma^i_{m\nu} \psi_j + 2\Gamma^{\alpha}_{m\nu} \psi_{\alpha} \delta^i_j.$$

Contracting in (1.2) with respect to i and n and using (0.4), we obtain

$$(1.3) \quad \bar{R}_{jm} = R_{jm} - 2\psi_{jm} + (1-N)\psi_{jm} + 2\Gamma^{\alpha}_{mj} \psi_{\alpha}.$$

Here \bar{R}_{jm} i R_{jm} denote the first kind Richi tensors of the spaces $G\bar{R}_N$ and GR_N respectively.

From (1.3) we get

$$(1.4) \quad \bar{R}_{1 \vee}^{jm} = R_{1 \vee}^{jm} - (N+1)\psi_{1 \vee}^{jm} + 2\Gamma_{m \vee}^{\alpha} \psi_{\alpha}.$$

Substituting (0.3) in (1.4) we get

$$(1.5) \quad (1+N)\psi_{1 \vee}^{jm} = R_{1 \vee}^{jm} - \bar{R}_{1 \vee}^{jm} + \frac{2}{1+N}\Gamma_{m \vee}^{\alpha}(\bar{\Gamma}_{\alpha\beta}^{\beta} - \Gamma_{\alpha\beta}^{\beta}).$$

From (1.3), (0.3) and (1.5) we get

$$\begin{aligned} \bar{R}_{1 \vee}^{jm} &= R_{1 \vee}^{jm} - \frac{1}{1+N} \left[2R_{1 \vee}^{jm} - 2\bar{R}_{1 \vee}^{jm} + \frac{4}{1+N}\Gamma_{m \vee}^{\alpha}(\bar{\Gamma}_{\alpha\beta}^{\beta} - \Gamma_{\alpha\beta}^{\beta}) \right] \\ &\quad + (1-N)\psi_{1 \vee}^{jm} + \frac{2}{1+N}\Gamma_{m \vee}^{\alpha}(\bar{\Gamma}_{\alpha\beta}^{\beta} - \Gamma_{\alpha\beta}^{\beta}). \end{aligned}$$

Now it follows that

$$(1.6) \quad (1-N^2)\psi_{1 \vee}^{jm} = (N\bar{R}_{1 \vee}^{jm} + \bar{R}_{1 \vee}^{mj}) - (NR_{1 \vee}^{jm} + R_{1 \vee}^{mj}) + 2\Gamma_{m \vee}^{\alpha}(\bar{\Gamma}_{\alpha\beta}^{\beta} - \Gamma_{\alpha\beta}^{\beta})\frac{1-N}{1+N}.$$

Taking into account (0.3) and (1.6), we can write (1.2) in the form

$$\begin{aligned} &\bar{R}_{1 \vee}^{jmn} + \frac{2}{1+N}\delta_j^i \bar{R}_{1 \vee}^{mn} + \frac{1}{N^2-1} [\delta_m^i (N\bar{R}_{1 \vee}^{jn} + \bar{R}_{1 \vee}^{nj}) - \delta_n^i (N\bar{R}_{1 \vee}^{jm} + \bar{R}_{1 \vee}^{mj})] \\ &\quad - \frac{2}{(1+N)^2} \bar{\Gamma}_{\alpha\beta}^{\beta} (2\delta_j^i \bar{\Gamma}_{n \vee}^{\alpha} + \delta_m^i \bar{\Gamma}_{n \vee}^{\alpha} - \delta_n^i \bar{\Gamma}_{m \vee}^{\alpha}) - \frac{2}{1+N} \bar{\Gamma}_{\alpha\beta}^{\beta} (\bar{\Gamma}_{m \vee}^i \delta_j^{\alpha} + \bar{\Gamma}_{n \vee}^{\alpha} \delta_j^i) \\ &= R_{1 \vee}^{jmn} + \frac{2}{1+N}\delta_j^i R_{1 \vee}^{mn} + \frac{1}{N^2-1} [\delta_m^i (NR_{1 \vee}^{jn} + R_{1 \vee}^{nj}) - \delta_n^i (NR_{1 \vee}^{jm} + R_{1 \vee}^{mj})] \\ &\quad - \frac{2}{(1+N)^2} \Gamma_{\alpha\beta}^{\beta} (2\delta_j^i \Gamma_{n \vee}^{\alpha} + \delta_m^i \Gamma_{n \vee}^{\alpha} - \delta_n^i \Gamma_{m \vee}^{\alpha}) - \frac{2}{1+N} \Gamma_{\alpha\beta}^{\beta} (\Gamma_{m \vee}^i \delta_j^{\alpha} + \Gamma_{n \vee}^{\alpha} \delta_j^i). \end{aligned}$$

Therefore, the magnitude

$$(1.7) \quad \begin{aligned} E_{1 \vee}^{jmn} &= R_{1 \vee}^{jmn} + \frac{2}{1+N}\delta_j^i R_{1 \vee}^{mn} + \frac{1}{N^2-1} [\delta_m^i (NR_{1 \vee}^{jn} + R_{1 \vee}^{nj}) - \delta_n^i (NR_{1 \vee}^{jm} + R_{1 \vee}^{mj})] \\ &\quad - \frac{2}{(1+N)^2} \Gamma_{\alpha\beta}^{\beta} (2\delta_j^i \Gamma_{n \vee}^{\alpha} + \delta_m^i \Gamma_{n \vee}^{\alpha} - \delta_n^i \Gamma_{m \vee}^{\alpha}) \\ &\quad - \frac{2}{1+N} \Gamma_{\alpha\beta}^{\beta} (\Gamma_{m \vee}^i \delta_j^{\alpha} + \Gamma_{n \vee}^{\alpha} \delta_j^i), \end{aligned}$$

is invariant under an ETG mapping.

The magnitude (1.7) is not a tensor, and we call it *the ET-projective parameter of the first kind*.

2. ET-projective parameter of the second kind

For the second kind curvature tensors of the spaces GR_N and $G\bar{R}_N$ we get the relation

$$\bar{R}_{2jmn}^i = R_{2jmn}^i + P_{mj|n}^i - P_{nj|m}^i + P_{mj}^\alpha P_{n\alpha}^i - P_{nj}^\alpha P_{m\alpha}^i + 2\Gamma_{nm}^\alpha P_{\alpha j}^i,$$

i.e., using (0.2) one obtains

$$\begin{aligned} \bar{R}_{2jmn}^i &= R_{2jmn}^i + \delta_j^i (\psi_{mn} - \psi_{nm}) + \delta_m^i \psi_{jn} - \delta_n^i \psi_{jm} - \delta_m^i \xi_{nj}^\alpha \psi_\alpha + \delta_n^i \xi_{mj}^\alpha \psi_\alpha \\ &+ \xi_{mj|n}^i - \xi_{nj|m}^i + 2\psi_j \xi_{nm}^i + \xi_{mj}^\alpha \xi_{n\alpha}^i - \xi_{nj}^\alpha \xi_{m\alpha}^i + 2\Gamma_{nm}^i \psi_\alpha \delta_j^i + 2\Gamma_{nm}^\alpha \psi_j + 2\Gamma_{nm}^\alpha \xi_{\alpha j}^i. \end{aligned}$$

Now, analogously to previous case, we get the invariant magnitude of the ETG mapping $f : GR_N \rightarrow G\bar{R}_N$

$$\begin{aligned} E_{2jmn}^i &= R_{2jmn}^i + \frac{2}{1+N} \delta_j^i R_{2m\nu}^\alpha + \frac{1}{N^2-1} [\delta_m^i (NR_{2jn} + R_{2nj}) - \delta_n^i (NR_{2jm} + R_{2mj})] \\ &- \frac{2}{(1+N)^2} \Gamma_{\alpha\beta}^\beta (2\delta_j^i \Gamma_{m\nu}^\alpha + \delta_m^i \Gamma_{jn}^\alpha - \delta_n^i \Gamma_{mj}^\alpha) - \frac{2}{1+N} \Gamma_{\alpha\beta}^\beta (\Gamma_{nm}^i \delta_j^\alpha + \Gamma_{nm}^\alpha \delta_j^i). \end{aligned}$$

The magnitude E_{2jmn}^i is not a tensor and we call it *ET-projective parameter of the second kind*.

3. ET-projective parameter of the third kind

In the case of the third kind curvature tensors of the spaces GR_N and $G\bar{R}_N$ we get

$$\bar{R}_{3jmn}^i = R_{3jmn}^i + P_{jm|n}^i - P_{nj|1m}^i + P_{jm}^\alpha P_{n\alpha}^i - P_{nj}^\alpha P_{\alpha m}^i + 2P_{nm}^\alpha \Gamma_{\alpha j}^i + 2P_{nm}^\alpha P_{\alpha j}^i$$

i.e., in virtue of (0.2)

$$\begin{aligned} \bar{R}_{3jmn}^i &= R_{3jmn}^i + \delta_j^i (\psi_{mn} - \psi_{nm}) + \delta_m^i \psi_{jn} - \delta_n^i \psi_{jm} \\ (3.1) \quad &+ \psi_\alpha (\delta_n^i \xi_{jm}^\alpha - \delta_m^i \xi_{nj}^\alpha) + \xi_{jm|n}^i - \xi_{nj|1m}^i + \xi_{jm}^\alpha \xi_{n\alpha}^i - \xi_{jn}^\alpha \xi_{\alpha m}^i \\ &+ 2\psi_n (\Gamma_{mj}^i + \xi_{mj}^i) + 2\psi_m (\Gamma_{nj}^i + \xi_{nj}^i) + 2\xi_{nm}^\alpha (\Gamma_{\alpha j}^i + \xi_{\alpha j}^i). \end{aligned}$$

Also, it is satisfied

$$(3.2) \quad \psi_{2mn} = \psi_{1mn} + 2\Gamma_{m\nu}^\alpha \psi_\alpha.$$

From (3.1), (3.2) and (0.7) we get

$$(3.3) \quad \begin{aligned} \overline{R}_3^i{}_{jmn} &= R_3^i{}_{jmn} + \delta_j^i (\psi_{mn} - \psi_{nm}) + \delta_m^i \psi_{jn} - \delta_n^i \psi_{jm} \\ &\quad + 2\psi_n \Gamma_{m\check{\nu}}^i + 2\psi_m \Gamma_{n\check{\nu}}^i + 2\delta_j^i \Gamma_{m\check{\nu}}^\alpha \psi_\alpha + 2\delta_m^i \Gamma_{j\check{\nu}}^\alpha \psi_\alpha. \end{aligned}$$

Contracting (3.3) with respect to i and n , and alternating, we get

$$(3.4) \quad (1+N)\psi_{j\check{\nu}} = 2R_{j\check{\nu}} - 2\overline{R}_{j\check{\nu}} + \frac{4}{1+N}(\overline{\Gamma}_{\beta\alpha}^\alpha - \Gamma_{\beta\alpha}^\alpha)\Gamma_{\check{\nu}}^\beta,$$

From (0.3), (3.3) and (3.4) it follows

$$(3.5) \quad (1-N^2)\psi_{jm} = (N\overline{R}_{jm} + \overline{R}_{mj}) - (NR_{jm} + R_{mj}) + 2\Gamma_{m\check{\nu}}^\beta(\overline{\Gamma}_{\beta\alpha}^\alpha - \Gamma_{\beta\alpha}^\alpha)\frac{1-N}{1+N}.$$

Taking into account (0.3) and (3.5) we can write (3.3) in the form $\overline{E}_3^i{}_{jmn} = E_3^i{}_{jmn}$, where

$$\begin{aligned} E_3^i{}_{jmn} &= R_3^i{}_{jmn} + \frac{2}{1+N}\delta_j^i R_{m\check{\nu}} + \frac{1}{N^2-1}[\delta_m^i (NR_{jn} + R_{nj}) - \delta_n^i (NR_{jm} + R_{mj})] \\ &\quad - \frac{2}{(1+N)^2}\Gamma_{\alpha\beta}^\beta(2\delta_j^i \Gamma_{n\check{\nu}}^\alpha + \delta_m^i \Gamma_{n\check{\nu}}^\alpha - \delta_n^i \Gamma_{m\check{\nu}}^\alpha) \\ &\quad - \frac{2}{1+N}(\Gamma_{m\check{\nu}}^i \Gamma_{n\check{\nu}}^\alpha - \Gamma_{n\check{\nu}}^i \Gamma_{m\check{\nu}}^\alpha - \delta_j^i \Gamma_{m\check{\nu}}^\beta \Gamma_{\beta\alpha}^\alpha - \delta_m^i \Gamma_{j\check{\nu}}^\beta \Gamma_{\beta\alpha}^\alpha). \end{aligned}$$

The magnitude $E_3^i{}_{jmn}$ is an invariant of the ETG mapping. We call it *ET-projective parameter of the third kind*.

4. ET-projective parameter of the fourth kind

For curvature tensors of the fourth kind we get

$$\overline{R}_4^i{}_{jmn} = R_4^i{}_{jmn} + P_{jm|n}^i - P_{nj|m}^i + P_{jm}^\alpha P_{n\alpha}^i - P_{nj}^\alpha P_{\alpha m}^i + 2P_{mn}^\alpha \Gamma_{\alpha\check{\nu}}^i + 2P_{mn}^\alpha P_{\alpha\check{\nu}}^i$$

i.e.,

$$\begin{aligned} \overline{R}_4^i{}_{jmn} &= R_4^i{}_{jmn} + \delta_j^i (\psi_{mn} - \psi_{nm}) + \delta_m^i \psi_{jn} - \delta_n^i \psi_{jm} + \\ &\quad + \psi_\alpha (\delta_n^i \xi_{jm}^\alpha - \delta_m^i \xi_{nj}^\alpha) + \xi_{jm|n}^i - \xi_{nj|m}^i + \xi_{jm}^\alpha \xi_{n\alpha}^i - \xi_{jn}^\alpha \xi_{\alpha m}^i + \\ &\quad + 2\psi_n (\Gamma_{m\check{\nu}}^i + \xi_{mj}^i) + 2\psi_m (\Gamma_{n\check{\nu}}^i + \xi_{nj}^i) + 2\xi_{mn}^\alpha (\Gamma_{\alpha\check{\nu}}^i + \xi_{\alpha j}^i). \end{aligned}$$

In this case, analogously, we get an invariant magnitude of the ETG mapping in the form

$$\begin{aligned} E_4^i{}_{jmn} &= R_4^i{}_{jmn} + \frac{2}{1+N} \delta_j^i R_{4\sqrt{v}}{}^{mn} + \frac{1}{N^2-1} [\delta_m^i (NR_{4j}{}^n + R_{4nj}) - \delta_n^i (NR_{4j}{}^m + R_{4mj})] \\ &\quad - \frac{2}{(1+N)^2} \Gamma_{\alpha\beta}^\beta (2\delta_j^i \Gamma_{n\sqrt{v}}^\alpha + \delta_m^i \Gamma_{nj}^\alpha - \delta_n^i \Gamma_{mj}^\alpha) \\ &\quad - \frac{2}{1+N} (\Gamma_{m\sqrt{v}}^i \Gamma_{n\alpha}^\alpha - \Gamma_{nj}^i \Gamma_{m\alpha}^\alpha - \delta_j^i \Gamma_{m\sqrt{v}}^\beta \Gamma_{\beta\alpha}^\alpha - \delta_m^i \Gamma_{jn}^\beta \Gamma_{\beta\alpha}^\alpha). \end{aligned}$$

The magnitude $E_4^i{}_{jmn}$ is not a tensor and we call it *ET-projective parameter of the fourth kind* of the ETG mapping.

5. ET-projective curvature tensor

For the curvature tensors of the fifth kind of the spaces GR_N and $G\overline{R}_N$ we find the relation

$$\begin{aligned} \overline{R}_5^i{}_{jmn} &= \\ R_5^i{}_{jmn} &+ \frac{1}{2} (P_{jm|_3}^i - P_{jn|_4}^i + P_{mj|_4}^i - P_{nj|_3}^i + P_{jm}^\alpha P_{\alpha n}^i - P_{jn}^\alpha P_{m\alpha}^i + P_{mj}^\alpha P_{n\alpha}^i - P_{nj}^\alpha P_{\alpha m}^i) \end{aligned}$$

i.e.,

$$\begin{aligned} (5.1) \quad \overline{R}_5^i{}_{jmn} &= R_5^i{}_{jmn} \\ &+ \frac{1}{2} \delta_j^i (\psi_{mn} - \psi_{nm} + \psi_{mn} - \psi_{nm}) + \frac{1}{2} \delta_m^i (\psi_{jn} + \psi_{jn}) - \frac{1}{2} \delta_n^i (\psi_{jm} + \psi_{jm}) \\ &+ \frac{1}{2} (\xi_{jm|_3}^i - \xi_{jn|_4}^i + \xi_{mj|_4}^i - \xi_{nj|_3}^i + \xi_{jm}^\alpha \xi_{\alpha n}^i - \xi_{jn}^\alpha \xi_{m\alpha}^i + \xi_{mj}^\alpha \xi_{n\alpha}^i - \xi_{nj}^\alpha \xi_{\alpha m}^i) \end{aligned}$$

Substituting (0.6) in (5.1) we get

$$\begin{aligned} (5.2) \quad \overline{R}_5^i{}_{jmn} &= R_5^i{}_{jmn} + \frac{1}{2} (\psi_{mn} - \psi_{nm} + \psi_{mn} - \psi_{nm}) \\ &+ \frac{1}{2} \delta_m^i (\psi_{jn} + \psi_{jn}) - \frac{1}{2} \delta_n^i (\psi_{jm} + \psi_{jm}). \end{aligned}$$

Denoting $\psi_{jn} = \frac{1}{2} (\psi_{jn} + \psi_{jn})$, we can write (5.2) in the form

$$(5.3) \quad \overline{R}_5^i{}_{jmn} = R_5^i{}_{jmn} + \delta_j^i (\psi_{mn} - \psi_{nm}) + \delta_m^i \psi_{jn} - \delta_n^i \psi_{jm}.$$

Eliminating ψ_{mn} from (5.3), analogously to previous cases, we get

$$\overline{E}_5^i{}_{jmn} = E_5^i{}_{jmn},$$

where we denoted

$$E_{\frac{5}{5}}^i{}_{jmn} = R_{\frac{5}{5}}^i{}_{jmn} + \frac{2}{1+N} \delta_j^i R_{\frac{5}{5}mn} + \frac{1}{N^2-1} [\delta_m^i (NR_{\frac{5}{5}jn} + R_{\frac{5}{5}nj}) - \delta_n^i (NR_{\frac{5}{5}jm} + R_{\frac{5}{5}mj})].$$

The magnitude $E_{\frac{5}{5}}^i{}_{jmn}$ is an invariant of the ETG mapping. In contrast to the previous cases, when $E_{\theta}^i{}_{jmn}$ ($\theta = 1, \dots, 4$) are not tensors, the magnitude $E_{\frac{5}{5}}^i{}_{jmn}$ is a tensor. We call it *ET-projective curvature tensor*.

If $GR_N(G\bar{R}_N)$ reduces to $R_N(\bar{R}_N)$, then the magnitudes $E_{\theta}^i{}_{jmn}$ ($\theta = 1, \dots, 5$) reduce to Weil's tensor (0.6').

REFERENCES

- [1] L. P. Eisenhart, *Generalized Riemannian spaces I*, Proc. Nat. Acad. Sci. USA **37** (1951), 311-315.
- [2] S. M. Minčić, *Ricci identities in the space of non-symmetric affine connection*, Mat. Vesnik **10**(25) (1973), 161-172.
- [3] S. M. Minčić, *New commutation formulas in the non-symmetric affine connection space*, Publ. Inst. Math. (Beograd) (N. S) **22**(36) (1977), 189-199.
- [4] S. M. Minčić, *Independent curvature tensors and pseudotensors of spaces with non-symmetric affine connection*, Coll. Math. Soc. János Bolyai **31** (1979), 445-460.
- [5] S. M. Minčić, M. S. Stanković, *On geodesic mappings of general affine connection spaces and of generalized Riemannian spaces*, submitted.
- [6] Н. С. Синюков, *Геодезические отображения Римановых пространств*, Наука, Москва, 1987.

Matematički institut
Kneza Mihaila 2
11001 Beograd, p.p. 367
Yugoslavia

(Received 06 01 1997)

Filozofski fakultet
Ćirila i Metodija 2
18000 Niš
Yugoslavia