

ON GENERAL AND REPRODUCTIVE SOLUTIONS OF FINITE EQUATIONS

Sergiu Rudeanu

Communicated by Žarko Mijajlović

Abstract. Prešić and his followers have studied the general and reproductive solutions of a certain equation over a finite set. In this paper we provide simple characterizations of these solutions.

The concept of general or parametric solution of an equation is well known in various contexts. Schröder [21, vol. 1], introduced reproductive general solutions of Boolean equations, which were extensively studied by Löwenheim [13], [14] and his followers (the term “reproductive” was introduced by Löwenheim [14]; cf. Rudeanu [19]). Prešić [15] initiated the axiomatic study of general solutions and reproductive general solutions, i.e., for the most general concept of equation. This line of research was followed by Prešić [17], Božić [9], Banković [1], [2], [3], Rudeanu [20] and Chvalina [10]. In their monograph [12], unfortunately unpublished, Kečkić and Prešić showed that the concept of reproductive solution is very important in various fields of mathematics.

A further step was taken by Prešić [16], who considered the case of equations over a finite set, on which he introduced a certain algebraic structure yielding a reproductive solution in compact form; see also Ghilezan [11]. Then Prešić [18] introduced within this framework an equation which generalizes Boolean and Post equations in one unknown and for which he obtained all reproductive solutions. Other descriptions of all general solutions and all reproductive solutions were then obtained by Banković [4]–[8].

In this paper we provide simpler characterizations of the general and reproductive solutions of that equation.

We first recall all necessary prerequisites.

Let $T = \{t_0, t_1, \dots, t_m\}$ be a finite set and $0, 1$ two elements which may be either outside T or two distinguished elements of T . Define two binary partial operations $+$ and \cdot on $T \cup \{0, 1\}$ by the following conditions:

$$(1) \quad \begin{aligned} x + 0 &= 0 + x = x, \\ x \cdot 0 &= 0 \cdot x = 0, \\ x \cdot 1 &= 1 \cdot x = x, \end{aligned}$$

for every $x \in T \cup \{0, 1\}$; the operation \cdot is usually denoted by concatenation. Define also

$$(2) \quad x^y = \begin{cases} 1, & \text{if } x = y, \\ 0, & \text{if } x \neq y, \end{cases}$$

for every $x, y \in T \cup \{0, 1\}$, as well as a family $\sum_{i=0}^n$ of partial operations given by

$$(3) \quad \sum_{i=0}^0 x_i = x_0, \quad \sum_{i=0}^{n+1} x_i = \left(\sum_{i=0}^n x_i \right) + x_{n+1} \quad (n \in \mathbb{N}).$$

PROPOSITION 1. *Every function $f: T \rightarrow T$ can be written in the form*

$$(4) \quad f(x) = \sum_{i=0}^m f_i x^{t_i} \quad (\forall x \in T),$$

where the coefficients are uniquely determined by

$$(5) \quad f_k = f(t_k) \quad (k = 0, 1, \dots, m).$$

Proof. For each $k = 0, 1, \dots, m$, the equality

$$(6) \quad f(t_k) = \sum_{i=0}^m f_i t_k^{t_i}$$

holds because the right-hand side of (6) contains the term $f(t_k) t_k^{t_k} = f(t_k)$, while the other terms are 0. Conversely, (4) implies, for each $k = 0, 1, \dots, m$,

$$f(t_k) = \sum_{i=0}^m f_i t_k^{t_i} = f_k t_k^{t_k} = f_k.$$

The equation introduced by Prešić and studied by him and Banković is

$$(7) \quad a_0 x^{t_0} + a_1 x^{t_1} + \dots + a_m x^{t_m} = 0,$$

where $a_0, a_1, \dots, a_m \in \{0, 1\}$ and the unknown $x \in T$. Let us denote by S the (possibly empty!) set of solutions of equation (7).

Remark [18]. For $x = t_k$ the left-hand side of equation (7) reduces to a_k , hence

$$(8) \quad t_k \in S \Leftrightarrow a_k = 0 \quad (k = 0, 1, \dots, m),$$

therefore equation (7) is consistent if and only if $a_k = 0$ for some k . In other words, taking also into account the associativity of the operation \cdot on $\{0, 1\}$, we see that *the consistency condition for equation (7) is*

$$(9) \quad a_0 a_1 \dots a_m = 0.$$

The next definition is borrowed from the axiomatic framework introduced by Prešić [15].

Definition. Suppose equation (7) is consistent. A function $f: T \rightarrow T$ is called a *general solution* of equation (7) if $f(T) = S$. By a *reproductive general solution* or simply a *reproductive solution* of equation (7) is meant a general solution f such that $s = f(s)$ for all $s \in S$.

The next lemma is in fact valid for any equation over a finite set.

LEMMA 1 [4]. *A function $f: T \rightarrow T$ is a general solution of equation (7) if and only if it fulfils*

- (i) $f(T) \subseteq S$, and
- (ii) *there is a permutation β of $\{0, 1, \dots, m\}$ such that*

$$(10) \quad t_{\beta(k)} \in S \Rightarrow f(t_k) = t_{\beta(k)} \quad (k = 0, 1, \dots, m).$$

Proof. Sufficiency is obvious. Conversely, suppose f is a general solution. Define a map $\varphi_0: S \rightarrow T$ by choosing, for each element $t \in S$, an element $\varphi_0(t)$ such that $t = f(\varphi_0(t))$. Then φ_0 is obviously injective and it follows immediately that φ_0 can be extended to a bijection $\varphi: T \rightarrow T$. The defining property of φ_0 implies that $t_h = f(\varphi(t_h))$ for all $t_h \in S$. The latter property can be written in the form (10) for the permutation β of $\{0, 1, \dots, m\}$ defined by $\beta(k) = h \Leftrightarrow \varphi(t_h) = t_k$.

LEMMA 2. *The implication (10) in Lemma 1 can be written in the form*

$$(11) \quad f(t_k) = t_{\beta(k)} a_{\beta(k)}^0 + f(t_k) a_{\beta(k)}^1 \quad (k = 0, 1, \dots, m).$$

Proof. It follows from (11) that

$$(12) \quad a_{\beta(k)} = 0 \Rightarrow f(t_k) = t_{\beta(k)} \quad (k = 0, 1, \dots, m);$$

but (12) is equivalent to (10) by (8). Conversely, suppose (10) holds. Then (11) is checked immediately by considering the cases $a_{\beta(k)} = 1$ and $a_{(k)} = 0$ and by using again (8).

THEOREM 1. *A function $f: T \rightarrow T$ is a general solution of equation (7) if and only if it is of the form*

$$(13) \quad f(x) = \sum_{k=0}^m (t_{\beta(k)} a_{\beta(k)}^0 + c_k a_{\beta(k)}^1) x^{t_k} \quad (\forall x \in T)$$

where β is a permutation of $\{0, 1, \dots, m\}$ and

$$(14) \quad c_k \in S \quad (k = 0, 1, \dots, m)$$

Proof. Suppose (13) and (14) hold. Consider an arbitrary but fixed $k \in \{0, 1, \dots, m\}$. Taking into account Proposition 1, it follows that if $a_{\beta(k)} = 1$; then $f(t_k) = c_k \in S$, while $t_{\beta(k)} \in S \Leftrightarrow a_{\beta(k)} = 0 \Rightarrow f(t_k) = t_{\beta(k)}$. Therefore f is a general solution by Lemma 1. Conversely, necessity follows from Proposition 1 and Lemmas 1 and 2.

THEOREM 2. *A function $f: T \rightarrow T$ is a reproductive solution of equation (7) if and only if it is of the form*

$$(15) \quad f(x) = \sum_{k=0}^m (t_k a_k^0 + c_k a_k^1) x^{t_k} \quad (\forall x \in T)$$

where

$$(14) \quad c_k \in S \quad (k = 0, 1, \dots, m).$$

Proof. It follows from Theorem 1 and Proposition 1 that any function satisfying (15) and (14) is a general solution and

$$(16) \quad f(t_k) = t_k a_k^0 + c_k a_k^1 \quad (k = 0, 1, \dots, m),$$

therefore if $t_k \in S$ then $f(t_k) = t_k$ because $a_k = 0$. Conversely, suppose f is a reproductive solution of equation (7). Then f satisfies conditions (i) and (ii) in Lemma 1 with the identity in the role of β . Taking also into account Lemma 2 we see that relations (14) and (16) hold for $c_k = f(t_k)$. The proof is completed by Proposition 1.

Acknowledgement. I thank the referee for his/her useful remarks.

BIBLIOGRAPHY

- [1] D. Banković, *On general and reproductive solution of arbitrary equations*, Publ. Inst. Math. Beograd **26(40)** (1979), 31–33.
- [2] D. Banković, *Solving systems of arbitrary equations*, Discrete Math. **46** (1983), 305–309.
- [3] D. Banković, *Some remarks on reproductive solutions*, Publ. Inst. Math. Beograd **38(52)** (1985), 17–19.
- [4] D. Banković, *All general solutions of finite equations*, Publ. Inst. Math. Beograd **47(61)** (1990), 5–12.
- [5] D. Banković, *A new proof of Prešić's theorem on finite equations*, Publ. Inst. Math. Beograd **51(65)** (1992), 22–24.
- [6] D. Banković, *A note on finite equations*, Zb. Rad. Prir.-Mat. Fak. u Kragujevcu **14** (1993), 15–18.
- [7] D. Banković, *All solutions of finite equations*, Discrete Math. **137** (1995), 1–6.
- [8] D. Banković, *Finite equations in n unknowns*, Publ. Inst. Math. Beograd **61(75)** (1997), 1–5.
- [9] D. Božić, *A note on reproductive solutions*, Publ. Inst. Math. Beograd **19(33)** (1975), 33–35.
- [10] J. Chvalinna, *Characterizations of certain general and reproductive solutions of arbitrary equations*, Mat. Vesnik **39** (1987), 5–12.
- [11] C. Ghilezan, *Méthode à résoudre des relations dont les résolutions appartiennent à un ensemble fini*, Publ. Inst. Math. Beograd **10(24)** (1970), 21–23.
- [12] J.D. Kečkić, S.B. Prešić, *Reproductivity – a general approach to equations* (1984, unpublished).
- [13] L. Löwenheim, *Über das Auflösungsproblem im logischen Klassenkalkul*, Sitzungsber. Berl. Math. Gesellschaft **7** (1908), 89–94.
- [14] L. Löwenheim, *Über die Auflösung von Gleichungen im logischen Gebietenkalkul*, Math. Ann. **68** (1910), 169–207.
- [15] S.B. Prešić, *Une classe d'équations matricielles et l'équation fonctionnelle $f^2 = f$* , Publ. Inst. Math. Beograd **8 (22)** (1968), 143–148.
- [16] S.B. Prešić, *Une méthode de résolution des équations dont toutes les solutions appartiennent à un ensemble fini donné*, C.R. Acad. Sci. Paris **272** (1971), 654–657.
- [17] S.B. Prešić, *Ein Satz Über reproduktive Lösungen*, Publ. Inst. Math. Beograd **14(28)** (1972), 133–136.
- [18] S.B. Prešić, *All reproductive solutions of finite equations*, Publ. Inst. Math. Beograd **44(58)** (1988), 3–7.
- [19] S. Rudeanu, *Boolean functions and equations*, North-Holland Amsterdam 1974.
- [20] S. Rudeanu, *On reproductive solutions of arbitrary equations*, Publ. Inst. Math. Beograd **24(38)** (1978), 143–145.
- [21] E. Schröder, *Vorlesungen Über die Algebra der Logik*, Leipzig; vol. 1, 1890; vol. 2, 1891, 1905; vol. 3, 1895. Reprint 1966 (Chelsea, Bronx NY).

Facultatea de matematică
 Universitatea București
 Str. Academiei 14
 70109 București
 Romania

(Received 05 01 1998)