

ON SYLOW SUBGROUPS OF S_n

Radoš Bakić

Communicated by Žarko Mijajlović

ABSTRACT. We give a new approach to description of p -Sylow subgroup normalizers in the groups S_n (symmetric group on n letters).

1. Sylow subgroup normalizer in S_n

Let $P_n(p) = P_n$ and $N_n(p) = N_n$ denote p -Sylow subgroup in S_n and its normalizer respectively. For $\alpha, \beta \in S_n$, the product $\alpha\beta$ will mean $\beta(\alpha(x))$.

We shall briefly repeat the construction of P_n , which goes inductively. The set of letters will be the set of first n natural numbers. If $n = p$, where p is a prime, then $P_n = (12 \dots p)$. Let $n = p^k$. Set $T_i = \{(i-1)p^{k-1} + 1, (i-1)p^{k-1} + 2, \dots, ip^{k-1}\}$ for $1 \leq i \leq p$. Let $K = \langle \pi \rangle$, where $\pi \in S_n$ is such that $\pi(T_i) = T_{i+1}$ for $i < p$, and $\pi(T_p) = T_1$, and π is increasing on each T_i , i.e., $x < y \Rightarrow \pi(x) < \pi(y)$. Obviously, $K \cong C_p$. Further on, we define $H_1 = P_{p^{k-1}}$ and $H_i = \pi^{1-i} P_{p^{k-1}} \pi^{i-1}$. We take H_i as a permutation from S_n in natural way, fixing all the letters that are not in T_i . The subgroup K normalizes the group $H = H_1 H_2 \dots H_p = H_1 \times H_2 \times \dots \times H_p$, and finally we have $P_n = HK$. We are now going to describe N_n for $n = p^k$. Note that P_n is transitive on the set $\{1, 2, \dots, p^k\}$ and has T_i as a block system.

LEMMA 1. *Let $\{B_i \mid 1 \leq i \leq p\}$ be a partition of $\{1, 2, \dots, p^k\}$ such that $|B_i| = p^{k-1}$ for each i . Let $\alpha \in S_{p^k}$ have B_i as its block system. Then for $0 \leq i \leq p$ there exists unique α_i with the following properties:*

- 1) *if $x \notin B_i$, then $\alpha_i(x) = x$, for $i \neq 0$,*
- 2) *α_0 has B_i as a block system and is increasing on each B_i , $\alpha = \alpha_1 \alpha_2 \dots \alpha_p \alpha_0$.*

PROOF. Let $\alpha(B_i) = B_{\alpha(i)}$. Then there exists a unique increasing bijection $\beta_i : B_i \rightarrow B_{\alpha(i)}$, and let α_0 be the "union" of β_i . If we define $\alpha_i(x) = \alpha_0^{-1}(\alpha(x))$ for $x \in B_i$ and $\alpha_i(x) = x$ elsewhere, then we obtain the desired factors. The uniqueness follows easily since α_i and α_j commute for $i, j > 0$. \square

If $\alpha, \beta \in S_n$ both have the same block system B_i , then B_i is also a block system of $\alpha\beta$. So, if B_i is the block system as in the lemma above and if $\alpha = \alpha_1 \dots \alpha_p \alpha_0$ and $\beta = \beta_1 \dots \beta_p \beta_0$ are corresponding factorizations, then how can we get factorization of $\alpha\beta$? If $\alpha\beta = (\alpha\beta)_1 \dots (\alpha\beta)_p (\alpha\beta)_0$ one can easily check that $(\alpha\beta)_i = \alpha_i \alpha_0 \beta_{\alpha(i)} \alpha_0^{-1}$, where $\alpha(i)$ is defined by $\alpha(B_i) = B_{\alpha(i)}$, and $(\alpha\beta)_0 = \alpha_0 \beta_0$.

LEMMA 2. *Let B_i , $1 \leq i \leq p$, be a block system for P_n and $|B_i| = p^{k-1}$. Then for each i there exists j such that $B_i = T_j$.*

PROOF. Set $C_{i,j} = T_i \cap B_j$. Then, for $\alpha \in P_n$ the following holds: $\alpha(C_{i,j}) = \alpha(T_i) \cap \alpha(B_j) = T_{\alpha(i)} \cap B_{\alpha(j)} = C_{\alpha(i), \alpha(j)}$ so $C_{i,j}$ is also a block system for P_n . Suppose that $T_1 \neq B_j$ for all j . Then we can assume that $C_{1,1}$ and $C_{1,2}$ are not empty. Since $|C_{1,1}| < |B_1|$ then $C_{m,1}$ has to be also nonempty for some $m > 1$. Since $H_1 < P_n$ for $x \in C_{1,1}$ and $y \in C_{1,2}$, there exists an $\alpha \in H_1$ such that $\alpha(x) = y$ as H_1 is transitive. From $C_{1,1} \subseteq B_1$ and $C_{1,2} \subseteq B_2$ it follows that $\alpha(B_1) = B_2$, since $\alpha(B_1) = B_i$. Let $c \in C_{m,1}$. Since $C_{m,1} \cap T_1 = \emptyset$ and $\alpha \in H_1$ it follows that $\alpha(c) = c$. That means that $\alpha(c) \in B_1$, contradicting $\alpha(B_1) = B_2$. \square

LEMMA 3. *If $\alpha \in N_n$, then T_i is a block system for α .*

PROOF. Consider the set $\{\alpha(T_i) \mid 1 \leq i \leq p\}$, which is a partition of $\{1, 2, \dots, p^k\}$. For every $\beta \in P_n$ there exists $\gamma \in P_n$ such that $\beta\alpha = \alpha\gamma$, since $\alpha \in N_n$. Then we have $\beta(\alpha(T_i)) = \alpha(\gamma(T_i)) = \alpha(T_j)$, and this shows that $\alpha(T_i)$ is a block system for P_n . By Lemma 2, it follows that T_i is a block system for α , as required. \square

Now, let $\alpha \in S_n$ having T_i as its block system. So, by Lemma 1, we have $\alpha = \alpha_1 \dots \alpha_p \alpha_0$. We want to characterize $\alpha \in N_n$ in the terms of α_i . We say that α_1 is connected with α_i if $\alpha_i = \pi^{1-i} \alpha_1 \pi^{i-1}$. The condition $\alpha \in N_n$ is equivalent to $\alpha K \alpha^{-1} \in P_n$ and $\alpha H_i \alpha^{-1} \in P_n$ for all i . Hence, for arbitrary $\beta_i \in H_i$ the following should hold:

$$(1) \quad \alpha \pi^s \alpha^{-1} = \alpha_1 \dots \alpha_p \alpha_0 \pi^s \alpha_0^{-1} \alpha_p^{-1} \dots \alpha_1^{-1} \\ = \left(\prod_{i=1}^p \alpha_i \alpha_0 \pi^s \alpha_0^{-1} \alpha_{(\alpha \pi^s \alpha_0^{-1})(i)}^{-1} \alpha_0 \pi^{-s} \alpha_0^{-1} \right) \alpha_0 \pi^s \alpha_0^{-1} \in P_n$$

$$(2) \quad \alpha \beta_i \alpha^{-1} = \alpha_{\alpha_0^{-1}(i)} \alpha_0 \beta_i \alpha_0^{-1} \alpha_{\alpha_0^{-1}(i)}^{-1} \in P_n.$$

In the conditions above we applied factorization rule for a product of permutations.

If L is a subgroup of S_n consisting of permutations having T_i as a block system and being increasing on each T_i , then L is isomorphic to S_p and has K as its p -Sylow subgroup. By well-known fact then $|N_L(K)| = p(p-1)$.

If we analyze the conditions 1) and 2), then it is not difficult to see that $\alpha \in N_n$ iff $\alpha_0 \in N_L(K)$, $\alpha_1 \in N_{n/p}$ as a permutation of T_1 and each α_i , $i > 0$, is connected with some $\beta \in N_{n/p}$ such that $\beta \alpha_1^{-1} \in H_1$. From previous conditions we get a recurrent formula for $|N_{p^k}|$:

$$|N_{p^k}| = p(p-1)p^{p^{k-1}-1} |N_{p^{k-1}}|, \text{ and hence } |N_{p^k}| = (p-1)^k p^{\frac{p^k-1}{p-1}} = (p-1)^k |P_{p^k}|$$

Before considering the general case for P_n , we state a lemma.

LEMMA 4. Let $R = \{a_1, \dots, a_{p^k}\}$ be an increasing sequence of natural numbers, and $T = \{1, 2, \dots, p^k\}$. There are $|N_{p^k}|$ bijections $\Theta : T \rightarrow R$ such that $\Theta^{-1}P_{p^k}\Theta = P$, where P is a p -Sylow subgroup of S_{p^k} constructed on R , in the same way as P_{p^k} was constructed on T .

PROOF. The proof follows immediately since for every such Θ and some $\Omega \in N(P)$, we have $\Omega(a_i) = \Theta(i)$, for all i , $1 \leq i \leq p^k$. \square

We return now to the general case for S_n and P_n . In the general case the construction of P_n goes as follows. Let

$$n = a_0 + a_1p + a_2p^2 + \dots + a_kp^k$$

be p -adic decomposition of n . Now we make a partition $T_{i,j}$ of the set $\{1, 2, \dots, n\}$ such that $0 \leq i \leq k$, $1 \leq j \leq a_i$ and $|T_{i,j}| = p^i$. In that partition $T_{0,j} = \{j\}$, $T_{1,1}$ consists of the next p not taken numbers, while next p numbers are $T_{1,2}$ and so on. Let $P_{i,j} \in S_n$ be equal to the p -Sylow subgroup on the set $T_{i,j}$ constructed in the same way as P_{p^k} was on the set $\{1, 2, \dots, p^k\}$, with $P_{0,j}$ trivial group and $P_{i,j}(x) = x$ elsewhere. Then, as it is well known, P_n is the internal direct product of all $P_{i,j}$.

Let $h \in N_n$. If $f \in P_n$, then $f(h(T_{i,j})) = h(f_1(T_{i,j})) = h(T_{i,j})$ for some $f_1 \in P_n$. Suppose now that for some $T_{i,j}$ and $T_{m,l}$ we have $h(T_{i,j}) \cap T_{m,l} \neq \emptyset$. Then it must be $h(T_{i,j}) \supseteq T_{m,l}$. On the contrary, there would exist $x \in T_{m,l} \setminus h(T_{i,j})$ and $y \in T_{m,l} \cap h(T_{i,j})$. Because $P_{m,l}$ is transitive on $T_{m,l}$, there exists $f \in P_{m,l}$ with $f(y) = x$, contradicting $f(h(T_{i,j})) = h(T_{i,j})$. Therefore, $h(T_{i,j})$ is a union of some $T_{m,l}$. If $h(T_{i,j})$ contains at least two distinct $T_{m,l}$ (and of course $|T_{m,l}| < |h(T_{i,j})|$), then we have $p^i \leq a_0 + \dots + a_{i-1}p^{i-1}$ and this is false since, in general, $a_j < p$. So, it follows $h(T_{i,j}) = T_{i,l}$ and $hP_{i,l}h^{-1} = P_{i,j}$. Let $h_{i,j}$ be the restriction of h to $T_{i,j}$. Hence, $H_{i,j} : T_{i,j} \rightarrow h(T_{i,j})$. A permutation $h \in N_n$ is uniquely determined by its $H_{i,j}$. By Lemma 4, for each $H_{i,j}$ we have $|N_{p^i}|$ possibilities. We can conclude that

$$|N_n| = \prod_{i=0}^k |N_{p^i}|^{a_i} a_i!$$

Finally let us just notice that using the above description, N_n can be recursively generated.

References

- [1] B. Huppert, *Endliche Gruppen I*, Springer-Verlag, Berlin 1967.
- [2] Ya. G. Berkovic, *On p -subgroups of finite symmetric and alternating groups*, in: Isaacs, M. et al (eds.), *Representation Theory, Group Rings, and Coding Theory. Papers in Honor of S. D. Berman*, Contemp. Math. 93., American Mathematical Society Providence, RI, 1989, pp. 67–76.

Matematički institut
Kneza Mihaila 35
11000 Beograd, p.p. 367
Serbia

(Received 27 11 2001)
(Revised 12 12 2003)