

MONOTONE IMAGES OF W -SETS AND HEREDITARILY WEAKLY CONFLUENT IMAGES OF CONTINUA

Jonathan Hatch and Č. V. Stanojević*

Communicated by Rade Živaljević

ABSTRACT. A proper subcontinuum H of a continuum X is said to be a W -set provided for each continuous surjective function f from a continuum Y onto X , there exists a subcontinuum C of Y that maps entirely onto H . Hereditarily weakly confluent (HWC) mappings are those with the property that each restriction to a subcontinuum of the domain is weakly confluent. In this paper, we show that the monotone image of a W -set is a W -set and that there exists a continuum which is not in class W but which is the HWC image of a class W continuum.

1. Introduction

In what follows, a continuum is a compact, connected metric space, and the term map is used to denote a continuous function. It is known that monotone images of class W continua are in class W , as shown in [1]. In the summer of 2000, two questions arose related to this result. First, in personal communication, W. J. Charatonik asked whether HWC maps preserve membership in class W . We answer his question in the negative in Section 3. Second, while discussing approaching continuum theory from an analytical viewpoint and attempting to characterize continua which are not intrinsic W -sets, the idea of examining the preimages of such continua under certain types of maps arose. We give a related theorem in Section 4.

2. Definitions

A proper subcontinuum H of a continuum X is said to be a W -set provided for each continuous surjective function f from a continuum Y onto X , there exists

2000 *Mathematics Subject Classification*: Primary 54F15, 54C50.

Key words and phrases: W -sets, Class W , Monotone maps, HWC maps.

*The author would like to thank Dr Dragan Blagojević for subtle editorial touchings that improve appearance of the paper.




FIGURE 1. A two endpoint Knaster-type continuum and an appropriate quotient space. This provides the basis of Theorem 1.

a subcontinuum C of Y that maps entirely onto H . A continuum each proper subcontinuum of which is a W -set is said to be in class W . A weakly confluent map $f : X \rightarrow Y$ is one so that, for each subcontinuum K of the range, there is at least one component C of $f^{-1}(K)$ so that $f(C) = K$. A hereditarily weakly confluent (HWC) map $f : X \rightarrow Y$ is one so that for each subcontinuum K of X , $f|_K$ is weakly confluent. A monotone map $f : X \rightarrow Y$ is one so that $f^{-1}(y)$ is connected for each $y \in Y$. The Hausdorff distance between two compact sets A and B is defined to be

$$d_H(A, B) = \inf\{\epsilon > 0 \mid A \subseteq \mathcal{B}_\epsilon(B) \text{ and } B \subseteq \mathcal{B}_\epsilon(A)\}.$$

3. HWC maps and class W

W. J. Charatonik asked whether the HWC image of a class W continuum is necessarily in class W . The answer is no.

THEOREM 1. *There exist a a continuum M in class W , a continuum X , and a surjective HWC map $f : M \rightarrow X$ where X is a continuum not in class W .*

PROOF. First, consider the continuum formed when one takes a two endpoint Knaster-type continuum, which can be realized as an inverse limit on arcs with a three-pass bonding map, and joins the two endpoints. The result, which we will denote by X , is an indecomposable continuum homeomorphic to the one illustrated on the right in Figure 1. It is clear that X is not in class W , since for the quotient map itself, there is no continuum in the domain which is mapped onto the arc from $f(p')$ to $f(q')$.

Denote by C the composant of X containing $f(p)$, the joining point. In the strip $\mathbb{R} \times [0, 1]$, consider the collection of straight line segments of the following form: C_0 is the straight line segment from $(0, 1)$ to $(-1, 1/2)$. Then, for each positive integer n , let C_n be the straight line segment from $((-1)^n \cdot n, 1/(n+1))$ to $((-1)^{n+1} \cdot (n+1), 1/(n+2))$. Let $\hat{Y} = \bigcup_{n \geq 0} C_n$, and observe that \hat{Y} is a connected set containing all of $\mathbb{R} \times \{0\}$ in its closure.

To construct M , which will be a subset of $X \times [0, 1]$, first consider the straight line ℓ in the plane connecting the points a and b . There is a natural surjective and injective map \hat{g} from \mathbb{R} to C with the following properties: first, that $\hat{g}(0) = f(p)$ and second, that for each integer n in $\mathbb{R} \setminus \{0\}$, $\hat{g}(n) \in \ell$. Extend \hat{g} to $g : \mathbb{R} \times [0, 1] \rightarrow C \times [0, 1]$ by setting $g(x, t) = (\hat{g}(x), t)$. Let $Y = g(Y)$ and define $M = X \cup Y$.

Observe that \overline{Y} contains C , and since C is dense in X , $\overline{Y} = M$. It is easily verified that M is in class W (for example by using Theorem 67.1 of [3] and Proposition 4 of [4]). Let $\pi : X \times [0, 1] \rightarrow X$ be simple projection map, and we will now show that $\pi|_M$ is HWC. Let K be any subcontinuum of M . If $K \subseteq X$ or $X \subseteq K$, then since $\pi|_X$ is essentially the identity, $\pi|_K$ is clearly weakly confluent. If $K \not\subseteq X$ and $X \not\subseteq K$, then $K \subset Y$, since X is a C-set in M . For $K \subset Y$, K is an arc, and since arcs are in class W , $\pi|_K$ is weakly confluent. Hence $\pi|_M$ is HWC. Thus X is the HWC image of a class W continuum, and X is not in class W . \square

4. Monotone maps and W -sets

In investigating what properties of a subcontinuum imply that it is not an intrinsic W -set, the idea arose that perhaps considering the preimages of such a subcontinuum under various types of maps might be informative. One of the questions that was generated by that discussion was about the monotone preimage of a continuum which was not an intrinsic W -set. After answering this question, we realized that there was a different statement of the same result which might be more useful.

The proof of our Theorem 2 depends on the following theorem that will appear in [2]

THEOREM. [2, Theorem 7] *A subcontinuum H of a continuum X is a W -set in X if and only if for each $\epsilon > 0$ there is a pair H_1, H_2 of compact subsets of X so that any continuum C intersecting both H_1 and H_2 which is not separated by $H_1 \cup H_2$ has Hausdorff distance from H less than ϵ .*

THEOREM 2. *Let X be a continuum with W -set H , and let $f : X \rightarrow Y$ be a map of X to a continuum Y . If f is monotone and $f(H)$ a proper subcontinuum of $f(X)$, then $f(H)$ is a W -set in $f(X)$.*

PROOF. Let H be a W -set in continuum X , and let $f : X \rightarrow Y$ be a monotone map so that $f(H)$ is nondegenerate. Without loss of generality, assume that f is surjective. Given any $\epsilon > 0$ so that $\epsilon < \frac{1}{4} \text{diam}(f(H))$, there is a $\delta > 0$ so that if $x_1, x_2 \in X$ with $d(x_1, x_2) < \delta$, then $d(f(x_1), f(x_2)) < \epsilon$. Since H is a W -set, there exist two compact subsets, H_1 and H_2 , so that for any continuum C from H_1 to H_2 which is not separated by $H_1 \cup H_2$, $d_H(c, H) < \delta$.

Assume, for the sake of contradiction, that there is a point $p \in f(H_1) \cap f(H_2)$. Then $f^{-1}(p)$ is a continuum intersecting both H_1 and H_2 . $f^{-1}(p)$ must therefore contain a continuum C irreducible between H_1 and H_2 , which by thus must have $d_H(C, H) < \delta$. Therefore $d_H(\{p\}, f(H)) < \epsilon$, which implies that $f(H) \subset \mathcal{B}_\epsilon(p)$. This contradicts our choice of p , so $f(H_1)$ and $f(H_2)$ must be disjoint.

If C is a continuum from $f(H_1)$ to $f(H_2)$ not separated by their union, then $f^{-1}(C)$ is a continuum intersecting H_1 and H_2 . Define M and N as follows:

$$M = f^{-1} \left(\overline{C \setminus (f(H_1) \cup f(H_2))} \right), \quad N = \overline{f^{-1} (C \setminus (f(H_1) \cup f(H_2)))}.$$

Observe that N is a subcontinuum of M . Let P and Q be subcontinua of M so that P is irreducible between $H_1 \cap M$ and N and Q is irreducible between $H_2 \cap M$ and N . The continuum $N \cup P \cup Q$ intersects H_1 and H_2 but is not separated by their union, so $d_H(P \cup N \cup Q, H) < \delta$. Hence each point of $f(H)$ is within ϵ of $f(P \cup N \cup Q) \subseteq C$, and each point of C is either in $f(P \cup N \cup Q)$, and hence within ϵ of $f(H)$, or in $f(H_1) \cup f(H_2)$, which must be within ϵ of $F(H)$, since H_1 and H_2 must both be within δ of H . Thus $d_H(C, H) < \epsilon$.

This satisfies the conditions in Theorem [2, Theorem 7] for $f(H)$ to be a W -set in $f(X)$. \square

References

- [1] J. Grispolakis, S. B. Nadler, and E. D. Tymchatyn, *Some properties of hyperspaces with applications to continua theory*, *Canad. J. Math* **31**:1, (1979), 197–210.
- [2] Jonathan Hatch, *On a characterization of W -sets*, preprint.
- [3] A. Illanes, S. B. Nadler, *Hyperspaces; fundamentals and recent advances*, Marcel Dekker, New York, 1999.
- [4] W. J. Charatonik, *Some counterexamples concerning Whitney levels*, *Bull. Polish. Acad. Sci. Math.* **31** (1983), 385–391

Jonathan Hatch
 Department of Mathematical Sciences
 University of Delaware
 Newark, DE 19711
 jhatch@math.udel.edu

(Received 05 05 2003)

Č. V. Stanojević
 Department of Mathematics and Statistics
 University of Missouri – Rolla
 202 Rolla Building
 Rolla, MO 65409-0020
 iwaa@umr.edu