

ON THE CONNECTION BETWEEN AFFINE AND
PROJECTIVE FUNDAMENTAL GROUPS OF
LINE ARRANGEMENTS AND CURVES

by

David Garber

Abstract. — In this note we prove a decomposition related to the affine fundamental group and the projective fundamental group of a line arrangement and a reducible curve with a line component. We give some applications to this result.

Résumé (Sur le rapport entre les groupes fondamentaux d'arrangements affine et projectifs de droites et de courbes)

Dans cet article, nous montrons une décomposition reliée au groupe fondamental affine et au groupe fondamental projectif d'un arrangement de droites et d'une courbe réductible avec une composante linéaire. Nous donnons quelques applications de ce résultat.

1. Introduction

The fundamental group of complements of plane curves is a very important topological invariant with many different applications. There are two such invariants: the *affine fundamental group* of a plane curve, which is the fundamental group of its affine complement, and its *projective fundamental group*, which is the fundamental group of its projective complement.

Oka [7] has proved the following interesting result, which sheds new light on the connection between these two fundamental groups:

Theorem 1.1 (Oka). — *Let C be a curve in \mathbb{CP}^2 and let L be a general line to C , i.e. L intersects C in only simple points. Then, we have a central extension:*

$$1 \longrightarrow \mathbb{Z} \longrightarrow \pi_1(\mathbb{CP}^2 - (C \cup L)) \longrightarrow \pi_1(\mathbb{CP}^2 - C) \longrightarrow 1$$

2000 Mathematics Subject Classification. — Primary 14H30; Secondary 14N20,52C35.

Key words and phrases. — Fundamental group, line arrangement, plane curve, line component.

Partially supported by the Lady Davis postdoctoral fellowship.

Since L is a general line to C , then:

$$\pi_1(\mathbb{CP}^2 - (C \cup L)) = \pi_1(\mathbb{C}^2 - C).$$

Hence, we get the following interesting connection between the two fundamental groups:

$$1 \longrightarrow \mathbb{Z} \longrightarrow \pi_1(\mathbb{C}^2 - C) \longrightarrow \pi_1(\mathbb{CP}^2 - C) \longrightarrow 1$$

A natural question is:

Question 1.2. — *Under which conditions does this short exact sequence split? Notice that when it does, we have the following decomposition:*

$$\pi_1(\mathbb{C}^2 - C) \cong \pi_1(\mathbb{CP}^2 - C) \oplus \mathbb{Z}$$

A *real line arrangement* in \mathbb{C}^2 is a finite union of copies of \mathbb{C} in \mathbb{C}^2 , whose equations can be written by real coefficients. Some families of real line arrangements were already proved to satisfy this condition, see [2].

Here, we show that:

Theorem 1.3. — *If \mathcal{L} is a real line arrangement, then such a decomposition holds:*

$$\pi_1(\mathbb{C}^2 - \mathcal{L}) \cong \pi_1(\mathbb{CP}^2 - \mathcal{L}) \oplus \mathbb{Z}$$

Actually, one can see that the same argument holds for arbitrary line arrangements (see Theorem 2.3). Moreover, we give a different condition for this decomposition to hold: If C is a plane curve with a line component, *i.e.* $C = C' \cup L$ where L is a line, we have that $\pi_1(\mathbb{C}^2 - C) = \pi_1(\mathbb{CP}^2 - C) \oplus \mathbb{Z}$ too.

These results can be applied to the computation of the affine fundamental groups of line arrangements, since the projective fundamental group is an easier object to deal with than the affine fundamental group.

The paper is organized as follows. In Section 2 we prove Theorem 1.3. The other condition is discussed in Section 3, and in the last section we give some applications.

2. Proof of Theorem 1.3

In this section, we prove Theorem 1.3.

Proof of Theorem 1.3. — Let \mathcal{L} be a real line arrangement with n lines. Let L be an arbitrary line which intersects \mathcal{L} transversally. By the following remark, one can reduce the proof to the case where L is a line which intersects transversally all the lines in \mathcal{L} and all the intersection points of L with lines in \mathcal{L} are to the left of all the intersection points of \mathcal{L} (see Figure 1 for such an example, where the arrangement \mathcal{L} consists of L_1, L_2, L_3 and L_4).




FIGURE 1. An example

Remark 2.1. — We have proved in [3] that in a real line arrangement, if a line crosses a multiple intersection point from one side to its other side (see Figure 2), the fundamental groups remain unchanged (see [3, Theorem 4.13] for this property of the action \triangleq).




FIGURE 2. A line crosses a multiple intersection point.

By this argument, we can start with any transversal line L as “the line at infinity”. Then, by using this property repeatedly, we can push this line over all the intersection points of the arrangement \mathcal{L} , without changing the corresponding fundamental group. This process will be terminated when all the intersection points of L with \mathcal{L} are placed to the left of all the intersection points of \mathcal{L} , and this is the reduced case.

By this remark, we continue the proof of Theorem 1.3 for the reduced case, where all the intersection points of L with \mathcal{L} are placed to the left of all the intersection points of \mathcal{L} . We compute presentations for $\pi_1(\mathbb{C}\mathbb{P}^2 - \mathcal{L})$ and $\pi_1(\mathbb{C}\mathbb{P}^2 - (\mathcal{L} \cup L))$ by braid monodromy techniques (the Moishezon-Teicher algorithm) and the van Kampen Theorem. The original techniques are introduced in [6] and [5] respectively. Shorter presentations of these techniques can be found in [2] and [3].

We first have to compute the lists of Lefschetz pairs of the arrangements, which are the pairs of indices of the components intersected at the intersection point, where we numerate the components locally near the point (see [2]). Since \mathcal{L} is an arbitrary real line arrangement, its list of Lefschetz pairs is

$$([a_1, b_1], \dots, [a_k, b_k]),$$

where k is the number of intersection points in \mathcal{L} . If we assume that the additional line L crosses all the lines of \mathcal{L} transversally to the left of all the intersection points in \mathcal{L} , the list of Lefschetz pairs of $\mathcal{L} \cup L$ is

$$([a_1, b_1], \dots, [a_k, b_k], [n, n + 1], \dots, [1, 2])$$

By the braid monodromy techniques and the van Kampen Theorem, the group $\pi_1(\mathbb{CP}^2 - \mathcal{L})$ has n generators $\{x_1, \dots, x_n\}$ and $\pi_1(\mathbb{CP}^2 - (\mathcal{L} \cup L))$ has $n + 1$ generators $\{x_1, \dots, x_n, x_{n+1}\}$. Moreover, the first k relations of $\pi_1(\mathbb{CP}^2 - (\mathcal{L} \cup L))$ are equal to the k relations of $\pi_1(\mathbb{CP}^2 - \mathcal{L})$. Let us denote this set of relations by \mathcal{R} .

Now, we have to find out the relations induced by the n intersection points of the line L with the arrangement \mathcal{L} . Moreover, we have to add at last the appropriate projective relations.

First, we compute the relations induced by the n intersection points of the line L with the arrangement \mathcal{L} . The Lefschetz pair of the $(k + 1)$ th point (which is the first intersection point of L with \mathcal{L}) is $[n, n + 1]$. Hence, its corresponding initial skeleton is shown in Figure 3.




FIGURE 3. Initial skeleton of the $(k + 1)$ th point

Since the list of pairs $[a_1, b_1], \dots, [a_k, b_k]$ is induced by a line arrangement, it is easy to see that:

$$\Delta\langle a_k, b_k \rangle \cdots \Delta\langle a_1, b_1 \rangle = \Delta\langle 1, n \rangle,$$

where $\Delta\langle 1, n \rangle$ is the generalized half-twist on the segment $[1, n]$.

Therefore, applying this braid on the initial skeleton yields the resulting skeleton for this point which is presented in Figure 4.



FIGURE 4. Final skeleton of the $(k + 1)$ th point

By the van Kampen Theorem, the corresponding relation is:

$$[x_n x_{n-1} \cdots x_2 x_1 x_2^{-1} \cdots x_n^{-1}, x_{n+1}] = 1$$

Let P_i be the i th intersection point, where $k + 2 \leq i \leq k + n$. Its corresponding Lefschetz pair is $[n - (i - k) + 1, n - (i - k) + 2]$, and hence its initial skeleton is shown in Figure 5.

Now, we first have to apply on it the following braid:

$$\Delta\langle n - (i - k) + 2, n - (i - k) + 3 \rangle \cdots \Delta\langle n, n + 1 \rangle$$




FIGURE 5. Initial skeleton of the i th point

and afterwards we have to apply on the resulting skeleton $\Delta\langle 1, n \rangle$, which equals to

$$\Delta\langle a_k, b_k \rangle \cdots \Delta\langle a_1, b_1 \rangle$$

as before. Hence, Figure 6 presents the resulting skeleton for the i th point.




FIGURE 6. Computing the final skeleton of the i th point

Again, by the van Kampen Theorem, the corresponding relation is:

$$[x_n x_{n-1} \cdots x_{i-k+1} x_{i-k} x_{i-k+1}^{-1} \cdots x_n^{-1}, x_{n+1}] = 1$$

To summarize, we get that the set of relations induced by the intersection points of the additional line L is:

$$\{[x_n x_{n-1} \cdots x_{i-k+1} x_{i-k} x_{i-k+1}^{-1} \cdots x_n^{-1}, x_{n+1}] = 1 \mid 1 \leq i - k \leq n\}$$

One can easily see, by a sequence of substitutions, that actually this set of relations is equal to the following set:

$$\{[x_i, x_{n+1}] = 1 \mid 1 \leq i \leq n\}$$

Hence, we have the following presentations:

$$\begin{aligned} \pi_1(\mathbb{CP}^2 - \mathcal{L}) &= \langle x_1, \dots, x_n \mid \mathcal{R}; x_n x_{n-1} \cdots x_1 = 1 \rangle \\ \pi_1(\mathbb{CP}^2 - (\mathcal{L} \cup L)) &= \left\langle x_1, \dots, x_n, x_{n+1} \mid \mathcal{R}; [x_i, x_{n+1}] = 1, 1 \leq i \leq n; \right. \\ &\quad \left. x_{n+1} x_n \cdots x_1 = 1 \right\rangle \end{aligned}$$

It remains to show that these presentations imply that:

$$\pi_1(\mathbb{CP}^2 - (\mathcal{L} \cup L)) = \mathbb{Z} \oplus \pi_1(\mathbb{CP}^2 - \mathcal{L})$$

Denote by $\mathcal{R}(x_1 \leftarrow w)$ the set of relations \mathcal{R} after we substitute anywhere the generator x_1 by an element w in the corresponding group.

Hence, using the projective relations in both fundamental groups, we have the following new presentations:

$$\begin{aligned}\pi_1(\mathbb{CP}^2 - \mathcal{L}) &= \langle x_2, \dots, x_n \mid \mathcal{R}(x_1 \leftarrow x_2^{-1} \cdots x_n^{-1}) \rangle \\ \pi_1(\mathbb{CP}^2 - (\mathcal{L} \cup L)) &= \left\langle x_2, \dots, x_n, x_{n+1} \mid \begin{array}{l} \mathcal{R}(x_1 \leftarrow x_2^{-1} \cdots x_n^{-1} x_{n+1}^{-1}); \\ [x_i, x_{n+1}] = 1, \ 2 \leq i \leq n \end{array} \right\rangle\end{aligned}$$

Since all the relations in \mathcal{R} are either commutative relations or cyclic relations of the type

$$w_m w_{m-1} \cdots w_1 = w_{m-1} \cdots w_1 w_m = \cdots = w_1 w_m \cdots w_2,$$

where w_i are words in the generators x_1, \dots, x_n , then the sum of powers of the generator x_1 or its inverse are equal in any part of the relation. Hence, using the fact that x_{n+1} commutes with all the other generators, we actually can cancel x_{n+1} from the set of relations

$$\mathcal{R}(x_1 \leftarrow x_2^{-1} \cdots x_n^{-1} x_{n+1}^{-1}),$$

and therefore we have:

$$\mathcal{R}(x_1 \leftarrow x_2^{-1} \cdots x_n^{-1} x_{n+1}^{-1}) = \mathcal{R}(x_1 \leftarrow x_2^{-1} \cdots x_n^{-1}).$$

By this argument, we now have:

$$\begin{aligned}\pi_1(\mathbb{CP}^2 - (\mathcal{L} \cup L)) &\cong \left\langle x_2, \dots, x_n, x_{n+1} \mid \begin{array}{l} \mathcal{R}(x_1 \leftarrow x_2^{-1} \cdots x_n^{-1} x_{n+1}^{-1}); \\ [x_i, x_{n+1}] = 1, \ 2 \leq i \leq n \end{array} \right\rangle \\ &\cong \left\langle x_2, \dots, x_n, x_{n+1} \mid \begin{array}{l} \mathcal{R}(x_1 \leftarrow x_2^{-1} \cdots x_n^{-1}); \\ [x_i, x_{n+1}] = 1, \ 2 \leq i \leq n \end{array} \right\rangle \\ &\cong \langle x_2, \dots, x_n \mid \mathcal{R}(x_1 \leftarrow x_2^{-1} \cdots x_n^{-1}) \rangle \oplus \langle x_{n+1} \rangle \\ &\cong \pi_1(\mathbb{CP}^2 - \mathcal{L}) \oplus \mathbb{Z}\end{aligned}$$

Since L is transversal to \mathcal{L} , this is equivalent to:

$$\pi_1(\mathbb{C}^2 - \mathcal{L}) \cong \pi_1(\mathbb{CP}^2 - \mathcal{L}) \oplus \mathbb{Z}$$

and we are done. \square

Remark 2.2. — By the projective relation in $\pi_1(\mathbb{CP}^2 - (\mathcal{L} \cup L))$, we have that:

$$x_{n+1} = (x_n \cdots x_1)^{-1}.$$

Hence, we have that the element which corresponds to the loop around all the lines is a central element in $\pi_1(\mathbb{C}^2 - \mathcal{L}) = \pi_1(\mathbb{CP}^2 - (\mathcal{L} \cup L))$, as was shown by Oka [7] too.

Now, one can see that a similar argument holds for complex line arrangements too. Hence, one has the following generalization:

Theorem 2.3. — *Let \mathcal{L} be an arbitrary line arrangement. Then the following decomposition holds:*

$$\pi_1(\mathbb{C}^2 - \mathcal{L}) \cong \pi_1(\mathbb{CP}^2 - \mathcal{L}) \oplus \mathbb{Z}$$

3. A different condition for a decomposition

In this section we will show that the decomposition holds for any reducible curve with a line component.

Oka and Sakamoto [8] have proved the following decomposition theorem concerning the affine fundamental group:

Theorem 3.1 (Oka-Sakamoto). — *Let C_1 and C_2 be algebraic plane curves in \mathbb{C}^2 . Assume that the intersection $C_1 \cap C_2$ consists of distinct $d_1 \cdot d_2$ points, where d_i ($i = 1, 2$) are the respective degrees of C_1 and C_2 . Then:*

$$\pi_1(\mathbb{C}^2 - (C_1 \cup C_2)) \cong \pi_1(\mathbb{C}^2 - C_1) \oplus \pi_1(\mathbb{C}^2 - C_2)$$

Here is a parallel version for the projective fundamental group:

Lemma 3.2. — *Let C_1 and C_2 be plane curves in \mathbb{CP}^2 with respective degrees d_1 and d_2 . Let C'_i be the affine part of C_i with respect to a projective line L . If $C'_1 \cap C'_2$ consists of $d_1 d_2$ distinct simple points, then:*

$$\pi_1(\mathbb{CP}^2 - (C_1 \cup C_2 \cup L)) \cong \pi_1(\mathbb{CP}^2 - (C_1 \cup L)) \oplus \pi_1(\mathbb{CP}^2 - (C_2 \cup L)).$$

Proof. — When L becomes the line at infinity, the curve $C_1 \cup C_2$ becomes $C'_1 \cup C'_2$. Also, it is easy to see that if C' denotes the affine part of C with respect to L , then

$$\pi_1(\mathbb{CP}^2 - (C \cup L)) \cong \pi_1(\mathbb{C}^2 - C').$$

Now, we can compute:

$$\begin{aligned} \pi_1(\mathbb{CP}^2 - (C_1 \cup C_2 \cup L)) &\cong \pi_1(\mathbb{C}^2 - (C'_1 \cup C'_2)) \\ &\stackrel{[8]}{\cong} \pi_1(\mathbb{C}^2 - C'_1) \oplus \pi_1(\mathbb{C}^2 - C'_2) \\ &\cong \pi_1(\mathbb{CP}^2 - (C_1 \cup L)) \oplus \pi_1(\mathbb{CP}^2 - (C_2 \cup L)). \quad \square \end{aligned}$$

Hence, we get:

Corollary 3.3. — *Let C be a plane curve of degree d in \mathbb{CP}^2 and let \mathcal{L} be a line arrangement consisting of k lines meeting in a point outside C . Let L be an additional line which intersects both C and \mathcal{L} transversally. If $C \cap \mathcal{L}$ consists of dk distinct simple points, then:*

$$\pi_1(\mathbb{CP}^2 - (C \cup \mathcal{L} \cup L)) \cong \pi_1(\mathbb{CP}^2 - (C \cup L)) \oplus \mathbb{Z} \oplus \mathbb{F}_{k-1},$$

where \mathbb{F}_k is the free group with k generators. In particular, if $k = 1$, i.e. \mathcal{L} consists of one line L_1 , then:

$$\pi_1(\mathbb{CP}^2 - (C \cup L_1 \cup L)) \cong \pi_1(\mathbb{CP}^2 - (C \cup L)) \oplus \mathbb{Z}$$

Proof. — Substituting C instead of C_1 and \mathcal{L} instead of C_2 in the last lemma, yields the following equation:

$$\pi_1(\mathbb{CP}^2 - (C \cup \mathcal{L} \cup L)) \cong \pi_1(\mathbb{CP}^2 - (C \cup L)) \oplus \pi_1(\mathbb{CP}^2 - (\mathcal{L} \cup L)).$$

Since $\pi_1(\mathbb{CP}^2 - (\mathcal{L} \cup L)) = \mathbb{Z} \oplus \mathbb{F}_{k-1}$ (see for example [2]), the Corollary is proved. Fixing $k = 1$ yields the particular case. \square

Hence we got the following result:

Proposition 3.4. — *Let C be a plane curve with a line component, i.e. $C = C' \cup L$ where L is a line that intersects C' transversally. Then:*

$$\pi_1(\mathbb{C}^2 - C) = \pi_1(\mathbb{CP}^2 - C) \oplus \mathbb{Z}$$

Proof. — Using the case $k = 1$ in the last corollary, we get that:

$$\pi_1(\mathbb{CP}^2 - (C' \cup L_1 \cup L)) \cong \pi_1(\mathbb{CP}^2 - (C' \cup L)) \oplus \mathbb{Z}$$

But since L_1 intersects $C' \cup L$ transversally, we have that

$$\pi_1(\mathbb{CP}^2 - (C' \cup L_1 \cup L)) \cong \pi_1(\mathbb{C}^2 - (C' \cup L))$$

So we get:

$$\pi_1(\mathbb{C}^2 - (C' \cup L)) \cong \pi_1(\mathbb{CP}^2 - (C' \cup L)) \oplus \mathbb{Z}. \quad \square$$

Remark 3.5. — By a similar argument to the argument we have used to prove Theorem 1.3, and by the fact that the line at infinity is transversal to $C \cup L$, we can generalize Proposition 3.4 to the case that L does not have to be transversal to C .

4. Some applications

In this section, we present some immediate applications of the main result.

4.1. Structure of the affine fundamental groups for some real arrangements. — Fan [1] proved the following result:

Proposition 4.1 (Fan). — *Let Σ be an arrangement of n lines and $S = \{a_1, \dots, a_k\}$ be the set of all singularities of Σ with multiplicity ≥ 3 . Suppose that the subgraph of Σ , which contains only the higher singularities (i.e. with multiplicity ≥ 3) and their edges, is a union of trees. Then:*

$$\pi_1(\mathbb{CP}^2 - \Sigma) \cong \mathbb{Z}^r \oplus \mathbb{F}^{m(a_1)-1} \oplus \dots \oplus \mathbb{F}^{m(a_k)-1}$$

where $r = n + k - 1 - m(a_1) - \dots - m(a_k)$ and $m(a_i)$ is the multiplicity of the intersection point a_i .

Using Theorem 1.3, we have the following easy consequence:

Corollary 4.2. — *Under the same conditions of the last proposition, we have that:*

$$\pi_1(\mathbb{C}^2 - \Sigma) \cong \mathbb{Z}^{r+1} \oplus \mathbb{F}^{m(a_1)-1} \oplus \dots \oplus \mathbb{F}^{m(a_k)-1}$$

where $r = n + k - 1 - m(a_1) - \dots - m(a_k)$.

This generalizes Theorem 5.3 of [2].

4.2. The connection between the invariants of real arrangements. — In [3] and [4] we address the following question about the invariants of real arrangements:

Question 4.3. — *Does the incidence lattice determine the affine fundamental group and the projective fundamental group of real line arrangements?*

For complex line arrangements, the answer for this question is negative due to an example of Rybnikov (see [9]).

Our main result is (see [4]):

Theorem 4.4. — *For real line arrangements with up to 8 lines, the incidence lattice does determine the affine fundamental group and the projective fundamental group.*

By Theorem 1.3, we actually reduce this question to the projective case only, since we have:

Corollary 4.5. — *Let \mathcal{L}_1 and \mathcal{L}_2 be two real line arrangements such that:*

$$\pi_1(\mathbb{C}\mathbb{P}^2 - \mathcal{L}_1) \cong \pi_1(\mathbb{C}\mathbb{P}^2 - \mathcal{L}_2)$$

Then:

$$\pi_1(\mathbb{C}^2 - \mathcal{L}_1) \cong \pi_1(\mathbb{C}^2 - \mathcal{L}_2)$$

This simplifies much the computational aspect of this question, since during the computations we have performed in [4], it was turned out that the comparison of two projective fundamental groups was much easier than the comparison of the corresponding affine fundamental groups. The main reason for this difference is probably that the affine fundamental group has one more generator than the projective fundamental group (this generator is cancelled by the projective relation which implies that the multiplication of all the generators is the unit element).

Acknowledgments. — I wish to thank Mutsuo Oka and Pho Duc Tai for fruitful discussions. I also thank Ehud De-Shalit and Einstein Institute of Mathematics for hosting my stay.

References

- [1] K.M. FAN – Direct product of free groups as the fundamental group of the complement of a union of lines, *Michigan Math. J.* **44** (1997), no. 2, p. 283–291.
- [2] D. GARBER & M. TEICHER – The fundamental group’s structure of the complement of some configurations of real line arrangements, in *Complex Analysis and Algebraic Geometry* (T. Peternell & F.-O. Schreyer, eds.), de Gruyter, 2000, p. 173–223.
- [3] D. GARBER, M. TEICHER & U. VISHNE – Classes of wiring diagrams and their invariants, *J. Knot Theory Ramifications* **11** (2002), no. 8, p. 1165–1191.
- [4] ———, π_1 -classification of arrangements with up to eight lines, *Topology* **42** (2003), no. 1, p. 265–289.
- [5] E.R. VAN KAMPEN – On the fundamental group of an algebraic curve, *Amer. J. Math.* **55** (1933), p. 255–260.
- [6] B. MOISHEZON & M. TEICHER – Braid group techniques in complex geometry I, Line arrangements in $\mathbb{C}\mathbb{P}^2$, *Contemporary Math.*, vol. 78, American Mathematical Society, 1988, p. 425–555.
- [7] M. OKA – On the fundamental group of the complement of a reducible curve in \mathbb{P}^2 , *J. London Math. Soc. (2)* **12** (1976), p. 239–252.
- [8] M. OKA & K. SAKAMOTO – Product theorem of the fundamental group of a reducible curve, *J. Math. Soc. Japan* **30** (1978), no. 4, p. 599–602.
- [9] G. RYBNIKOV – On the fundamental group of the complement of a complex hyperplane arrangement, preprint, [math.AG/9805056](https://arxiv.org/abs/math/9805056), 1994.

D. GARBER, Einstein Institute of Mathematics, The Hebrew University, Givat Ram, 91904 Jerusalem, Israel • *E-mail* : garber@math.huji.ac.il • Department of Sciences, Holon Academic Institute of Technology, PO Box 305, Holon 58102, Israel • *E-mail* : garber@hait.ac.il