


Geometry & Topology
 Volume 3 (1999) 303{330
 Published: 13 September 1999



Piecewise Euclidean Structures and Eberlein's Rigidity Theorem in the Singular Case

Michael W Davis
 Boris Okun
 Fangyang Zheng

*Department of Mathematics, The Ohio State University
 Columbus, OH 43201, USA*

*Department of Mathematics, Vanderbilt University
 Nashville, TN 37400, USA*

*Department of Mathematics, The Ohio State University
 Columbus, OH 43201, USA*

Email: mdavis@math.ohio-state.edu, okun@math.vanderbilt.edu
 zheng@math.ohio-state.edu

Abstract

In this article, we generalize Eberlein's Rigidity Theorem to the singular case, namely, one of the spaces is only assumed to be a CAT(0) topological manifold. As a corollary, we get that any compact irreducible but locally reducible locally symmetric space of noncompact type does not admit a nonpositively curved (in the Aleksandrov sense) piecewise Euclidean structure. Any hyperbolic manifold, on the other hand, does admit such a structure.

AMS Classification numbers Primary: 57S30

Secondary: 53C20

Keywords: Piecewise Euclidean structure, CAT(0) space, Hadamard space, rigidity theorem

Proposed: Steve Ferry
 Seconded: Walter Neumann, David Gabai

Received: 19 December 1998
 Accepted: 27 August 1999

0 Introduction

Suppose that V and V' are compact locally symmetric manifolds with fundamental groups Γ and Γ' and with universal covers X and X' , respectively. Suppose further that the symmetric spaces X and X' have no compact or Euclidean factors and that no finite cover of V' splits off a hyperbolic 2-manifold as a direct factor. Let $\phi: \Gamma \rightarrow \Gamma'$ be an isomorphism. The Mostow Rigidity Theorem asserts that the isomorphism ϕ is induced by an isometry $V' \xrightarrow{\sim} V$, possibly after renormalizing the metric on V' . ("Renormalizing the metric" means that we are allowed to rescale the metric on each factor of X' by multiplying by a positive constant.)

In the early eighties, Eberlein, Gromov and Heintze independently proved generalizations of Mostow's Theorem in which the hypotheses on V' are weakened. They showed that the conclusion of Mostow's Theorem remains valid without the assumption that V' is locally symmetric. It is only necessary to assume that it is a closed Riemannian manifold of nonpositive sectional curvature, provided that the rank of X' is at least two. (If the rank of X' is one, there is no such generalization. For, if X' is of rank one, then the sectional curvature of V' is strictly negative and hence, the same will be true for any sufficiently small deformation of its metric; moreover, the new metric will not, in general, be isometric to the original one.)

Eberlein proved the above generalization of the Rigidity Theorem in the case where X' is reducible and the uniform lattice Γ' is irreducible. We shall sometimes refer to this as "Eberlein's case". Gromov proved the result in full generality; however, the argument has only appeared (in [2]) in the case where X' is irreducible.

In this paper we will prove a further generalization of Eberlein's case of the Rigidity Theorem by allowing a further weakening of the hypotheses on V' : the metric need not be smooth. To explain this, we first need to understand what "nonpositive curvature" means in this context.

About fifty years ago Aleksandrov showed that the notions of upper or lower curvature bounds make sense for a more general class of metric spaces than Riemannian manifolds, namely, they make sense for "geodesic spaces". A *geodesic segment* in a metric space Y is the image of an isometric embedding of an interval into Y ; the space Y is a *geodesic space* if any two points can be connected by a geodesic segment. A *triangle* in Y is a configuration of three points (the vertices) and three geodesic segments (the edges) connecting them. Following Aleksandrov, one defines Y to be *nonpositively curved* if the distance between

any two points in any small triangle in Y is no greater than the distance between the corresponding points of a comparison triangle in the Euclidean plane. (The precise definition will be given in Section 1.1, below.)

The weakened hypotheses on V will then be that it is a complete geodesic space, that it is nonpositively curved in the above sense and that it is a closed topological manifold.

Our interest in generalizing the Rigidity Theorem to the singular case arose from an attempt to answer the following basic question concerning nonpositively curved, polyhedral metrics on manifolds. Given a Riemannian manifold M of nonpositive sectional curvature, is it possible to find a piecewise Euclidean, polyhedral metric on M which is nonpositively curved in Aleksandrov's sense? An application of our rigidity result is that this is, in fact, impossible when $M = V$. Indeed, if V were a nonpositively curved polyhedron homeomorphic to V , then by the Rigidity Theorem it would be isometric to V , which is clearly impossible (since V contains open subsets which are flat and V does not). A further discussion of the problem of approximating nonpositively curved Riemannian metrics by nonpositively curved polyhedral ones will be given in Section 1.

Next we shall make a few remarks concerning the proofs of Mostow and Eberlein. Since V and V are Eilenberg-MacLane spaces, the isomorphism $\pi_1 V \cong \pi_1 V$ gives rise to a homotopy equivalence $\bar{f}: V \rightarrow V$. This lifts to a π_1 -equivariant map $f: X \rightarrow X$. By analyzing the maximal flats in X and X , Mostow first shows that f extends to a homeomorphism of the maximal (or Furstenberg) boundaries. (The maximal boundary of X is a homogeneous space of the form G/P , where G is the identity component of the isometry group $I(X)$ and P is a minimal parabolic subgroup.) Let $X = X_1 \times \dots \times X_k$ be the de Rham decomposition into irreducible factors so that $I(X) = I(X_1) \times \dots \times I(X_k)$. The proof of Mostow's Theorem is the easiest to finish in Eberlein's case (where X is reducible and X is irreducible). The key ingredient is the Borel Density Theorem (cf [22]). It implies that the image of f is dense in the identity component of each $I(X_i)$. It follows that the map of maximal boundaries is actually induced by an isomorphism of Lie groups $I_0(X) \cong I_0(X)$ (where I_0 denotes the identity component of the isometry group). From this, the result follows (cf Corollary 18.2, page 133, in [19]). (A historical remark: one of the earliest rigidity results was proved by Selberg in 1957. He considered the case where X is the product of $k > 1$ copies of the hyperbolic plane and where X is irreducible. He used the fact that $I_0(\mathbb{H}^2)$ was dense in each copy of $I_0(\mathbb{H}^2) = PSL(2; \mathbb{R})$ to show that any deformation of f in $I(X)$ had to be trivial. The details of this argument can be found in [23].)

There are two steps in Eberlein's generalization of Mostow's argument (to the case where X is not required to be symmetric). First one shows that X also splits isometrically as $X = X_1 \amalg \dots \amalg X_k$. (This is accomplished in [9].) In the second step (which was actually accomplished in the earlier papers [4] and [10]) one shows that each factor $I_0(X_i)$ is actually a simple Lie group and that each X_i is a symmetric space. Hence, one is reduced back to Mostow's Theorem. The arguments in the second step are mainly Lie group theoretic. The key ingredient is again a density property (which is closely connected to the Borel Density Theorem), namely, that $I_0(X_i)$ satisfies the "duality condition" of [4]. One shows that the image of $I_0(X_i)$ in $I(X_i)$ is not discrete and hence, that the identity component G_i of the closure of its image in $I(X_i)$ is a nontrivial Lie group. One then argues that each G_i must be semisimple since otherwise (and hence X) would contain a nontrivial normal abelian subgroup.

Our argument, in the case where X is no longer assumed to be Riemannian, follows the same general outline. We show, in Section 2.1, that Eberlein's arguments can be modified to prove that X splits isometrically, as before. As for the second step, we first must face the fact that we don't know, *a priori*, that the isometry groups $I(X_i)$ are Lie groups. (The Hilbert-Smith Conjecture is still open | if a topological group acts effectively on a topological manifold it is not known if this forces it to be a Lie group.) In fact, *a priori*, the spaces X_i , which are manifold factors, need not be manifolds. We get around this by showing, in Section 2.3, that $I(X_i)$ acts transitively on X_i . (This comes from the fact that $I_0(X_i)$ is dense in $I(X_i)$.) It then follows from a theorem of Montgomery and Zippin that $I(X_i)$ is a Lie group. Then, following the line of argument in [4] and [10] one can conclude that each $I_0(X_i)$ is a simple Lie group. At this stage we are still not done. Even though X is now a homogeneous space of the form G/K , the possibility remains that its G -invariant metric is not Riemannian. However, we show in Lemma 18 that the nonpositivity of the curvature excludes this possibility.

After writing a preliminary version of this paper we learned of B Leeb's recent preprint [17]. A corollary of his theorem is that the Rigidity Theorem also holds in the singular case whenever each factor of X has rank ≥ 2 . Leeb's approach generalizes Gromov's line of argument in [2]. Thus, the singular version of the Rigidity Theorem holds in complete generality. We should also mention that Bruce Kleiner can prove our result by a somewhat different argument.

Davis and Zheng were partially supported by NSF grants, Zheng was also partially supported by an Alfred P Sloan Fellowship. This project is also sponsored by the National Security Agency under grant # MDA 904-98-1-0036.

1 Piecewise Euclidean CAT(0) structures

1.1 Hadamard spaces

A metric space is a *length space* if the distance between any two points is the infimum of the lengths of all paths between them. The space is a *geodesic space* if the infimum is always realized by a path of minimum length.

Given any triangle Δ in a geodesic space X there is a triangle Δ' in the Euclidean plane with the same edge lengths and a well defined isometry $f: \Delta \rightarrow \Delta'$ which is denoted by $x \mapsto x'$. The CAT(0) inequality of [14] asserts that for any two points $x, y \in \Delta$, $d(x; y) \leq |x' - y'|$. (Thus, any triangle in X is "thinner" than the corresponding triangle in \mathbb{R}^2).

A complete geodesic space X is a *Hadamard space* (or a *CAT(0) space*) if each triangle satisfies the CAT(0) inequality.

A geodesic space is *nonpositively curved*, if the CAT(0) inequality holds locally. It can be proved (cf [1] or [3]) that if a geodesic space is complete and nonpositively curved, then its universal cover is a Hadamard space.

Example 1 (Aleksandrov and Toponogov) A complete, simply connected Riemannian manifold is a Hadamard space if and only if its sectional curvature is ≤ 0 .

A key fact is that, for any Hadamard space X , the metric $d: X \times X \rightarrow \mathbb{R}$ is a convex function, cf [14]. (This means that its restriction to any geodesic segment in $X \times X$ is a convex function.) It follows that any Hadamard space is contractible. (There is an unique geodesic segment from a given base point to any other point; one can then contract the space to the base point by shrinking the geodesic segments.)

A geodesic space has *extendible geodesics* if every geodesic segment can be extended to a larger interval. A geodesic space is *geodesically complete* if every geodesic segment can be extended to a geodesic line. An easy argument shows that if a complete geodesic space has extendible geodesics then it is geodesically complete.

Suppose that the geodesic space X is nonpositively curved. If X is a topological manifold (or even if its local homology $H_n(X; X - x)$ is nontrivial at each point $x \in X$), then it has extendible geodesics. Indeed, suppose $c: [0; d] \rightarrow X$ is a geodesic segment which does not extend past $c(0)$. Choose a small ball B

centered at $c(0)$ with $\delta < d$ and small enough so that B_δ is convex. Since every point in $B_\delta - c(0)$ can be joined by a geodesic segment to $c(0)$, $B_\delta - c(0)$ contracts to $c(0)$, contradicting the assumption that $H(B_\delta; B_\delta - c(0))$ is nonzero.

Given two subsets A and B of X , the *Hausdorff distance* between them, denoted by $Hd(A; B)$, is the smallest number r such that each subset is contained within an r -neighborhood of the other. The subsets A and B are *parallel* if their Hausdorff distance is finite. An r -flat in X is a subset which is isometric to r -dimensional Euclidean space.

A subset A of a geodesic space is *locally convex* if the geodesic segment between any two sufficiently close points in A actually lies in A . The subset A is *totally geodesic* if any geodesic line which contains two distinct points of A is contained in A .

Given a closed convex subset A in a Hadamard space X , there is a continuous *projection map* $p: X \rightarrow A$ which sends a point $x \in X$ to the nearest point in A .

If the Hadamard space X is locally compact, then it can be compactified by adding an "ideal boundary" $X(\infty)$. One way to define this is as the set of all geodesic rays emanating from a given basepoint. If X is Riemannian, then $X(\infty)$ is homeomorphic to a sphere, but in the general case it can be much more complicated.

Let T be an isometry of X . Its *translation distance* $d(T)$ is the infimum over all points $x \in X$ of $d(x; Tx)$. The isometry T is *semisimple* if there is at least one point x such that $d(T) = d(x; Tx)$; T is a *Cliﬀord translation* if $d(T) = d(x; Tx)$ for all $x \in X$. If T is semisimple, then its *minimum set* $MIN(T)$ is the set of x in X such that $d(T) = d(x; Tx)$. When X is Hadamard, it follows from the convexity of the metric that $MIN(T)$ is a convex subset.

A length space is *singular* if it is not isometric to a Riemannian manifold. An important class of singular metrics is the class of polyhedral metrics, defined below.

Suppose that K is a cell complex formed by gluing together convex polytopes in some Euclidean space via isometries between certain faces. Then K is said to have a *piecewise Euclidean structure*. For example, if the cell complex K is a piecewise linearly embedded subcomplex of Euclidean space (ie, if it is a cell complex in the classical sense), then it has such a structure. A piecewise Euclidean structure on K defines a length metric: the distance between two points is the infimum of the lengths of all piecewise linear paths between them. Such metrics are called *piecewise Euclidean* (or *polyhedral*).

For the further details concerning the definitions and basic properties of CAT(0) spaces, we refer the reader to the excellent books of Bridson{Haefliger [3] and Ballmann [1].

1.2 Convex hypersurfaces

It follows from the Gauss equation that a smooth, convex hypersurface in the Euclidean space has sectional curvature ≥ 0 . The singular analogue of this is the following. The boundary K of a convex polytope in Euclidean space has a piecewise Euclidean metric and this metric is nonnegatively curved in the sense that the reverse inequality to the CAT(0) inequality is satisfied. In fact, it was precisely for the purpose of studying polyhedral metrics on convex polyhedral surfaces that Aleksandrov initiated the study of length spaces with curvature bounded from below.

The Gauss equation remains valid for the hypersurfaces in manifolds equipped with an indefinite metric, the only change that is necessary is that one side of the equation must be multiplied by -1 when the inner product is negative definite in the normal direction. As we shall see, this sign change allows the possibility that a convex hypersurface in Minkowski space will be nonpositively curved.

Recall that *Minkowski space* $\mathbb{R}^{n,1}$ is an $(n + 1)$ -dimensional real vector space with coordinates $(x_0; x_1; \dots; x_n)$ equipped with a *Lorentzian* inner product of signature $(n; 1)$ (eg, $x \cdot x = -x_0^2 + x_1^2 + \dots + x_n^2$). A hyperplane in $\mathbb{R}^{n,1}$ is *spacelike* if its normal vector n is *timelike*, ie, if $n \cdot n < 0$. A smooth hypersurface in $\mathbb{R}^{n,1}$ is *spacelike* if its tangent space at each point is spacelike. Since the restriction of the inner product to any spacelike hyperplane is positive definite, any spacelike hypersurface inherits a Riemannian structure. By the Gauss equation any smooth, spacelike convex hypersurface in $\mathbb{R}^{n,1}$ is nonpositively curved.

By definition a *convex polyhedral hypersurface* in $\mathbb{R}^{n,1}$ is the boundary of an $(n + 1)$ -dimensional convex polyhedral set C in $\mathbb{R}^{n,1}$ (ie, C is defined by a discrete set of affine inequalities). The convex hypersurface is *spacelike* if each supporting hyperplane of C is spacelike. As in the Euclidean case, it has an induced piecewise Euclidean metric.

Example 2 [7] Any spacelike convex polyhedral hypersurface in $\mathbb{R}^{n,1}$ is a Hadamard space. (Recently, Moussong has extended this result to arbitrary spacelike convex hypersurfaces in $\mathbb{R}^{n,1}$.)

As a simple corollary we get the following example.

Example 3 [7] Every complete hyperbolic manifold M^n admits a nonpositively curved, piecewise Euclidean metric. The construction goes as follows. Identify M^n with \mathbb{H}^n/Γ where Γ is a discrete torsion-free subgroup of $O_0(n; 1)$, the isometry group of \mathbb{H}^n . Choose a net S in M^n (ie, S is a discrete subset of M^n such that each point of M lies within a bounded distance of S) and let \mathcal{S} denote the inverse image of S in \mathbb{H}^n . Let $\partial\mathcal{S}$ denote the boundary of the convex hull of \mathcal{S} in \mathbb{R}^{n+1} . Then $\partial\mathcal{S}$ is a $\{$ stable, spacelike, convex polyhedral hypersurface in \mathbb{R}^{n+1} . Hence, as in Example 2, it is CAT(0). Thus, $\partial\mathcal{S} = \mathcal{S}$ gives the desired piecewise Euclidean metric on M^n . (The details of this construction can be found in [7].) Similarly, the product of two hyperbolic manifolds can be given a piecewise Euclidean structure. However, if, for example, Γ is an irreducible uniform lattice in $I(\mathbb{H}^2 \times \mathbb{H}^2)$, then, as we shall see in the next section, the resulting 4-manifold is rigid.

Remark Gromov has suggested that the construction in Example 3 can be modified to show that any Riemannian manifold of sufficiently pinched negative sectional curvature can be given a nonpositively curved, piecewise Euclidean metric.

1.3 Questions and further examples

At this point we have two classes of examples of nonpositively curved geodesic metrics on manifolds: 1) Riemannian metrics and 2) piecewise Euclidean metrics on polyhedral manifolds. What is the intersection of these two classes? When can one type of metric be deformed (through nonpositively curved metrics) into the other type? Both classes contain the Riemannian manifolds of constant curvature 0 (ie, flat manifolds). Example 3 shows that a complete metric of constant curvature -1 can be deformed to a polyhedral metric of nonpositive curvature. Before continuing the discussion let us break our problem into two questions.

Question A Suppose M^n is a nonpositively curved Riemannian manifold. Does M^n admit a nonpositively curved piecewise Euclidean metric?

Question B Suppose M^n is a nonpositively curved piecewise Euclidean polyhedral manifold. Does M^n admit a Riemannian metric of nonpositive sectional curvature?

In dimension 2 both questions have affirmative answers. In dimension 3 the answers are not known (although it seems likely that the answer is "yes" in both cases). As we shall see in Example 6, below, in dimension 5, Question B has a negative answer, while in dimension 4 the answer is unknown. A corollary to the main result of this paper is that, in general, Question A also has a negative answer in dimensions ≥ 4 . As we have seen in Example 3, Question A has a positive answer when M^n is a real hyperbolic manifold. The authors speculate that the answer will be negative for the other locally symmetric manifolds, eg for complex and quaternionic hyperbolic manifolds.

Example 4 ([14], [5]) Suppose that V is a manifold with a (possibly singular) length metric, that W is a codimension-two submanifold and that V_k is a k -fold cyclic branched cover of V along W . (One way to insure that such branched cover exist, for any k , is to require V and W to be closed and oriented and the image $[W]$ of the orientation class to be zero in $H_{n-2}(V)$.)

Theorem A Let V , W and V_k be as the above. For the induced metric on V_k to be nonpositively curved it is necessary and sufficient that the following two conditions hold:

- (a) V is nonpositively curved, and
- (b) W is a locally convex subset of V .

Proof The necessity of the conditions (a) and (b) is obvious. As for the sufficiency, suppose \triangle is a small geodesic triangle in V_k and that \triangle denotes its image in V . If \triangle intersects the branch set W in a vertex or an edge or if it is contained in W , then condition (a) and (b) insure that the distance between two points of \triangle is the same as that of the corresponding points of \triangle . If the intersection of \triangle and W is one or more points in the interiors of the edges, then we can subdivide \triangle and apply a standard argument (cf Lemma 3.3, page 15 of [1]) to again conclude that the CAT(0)-inequality holds. \square

Next we consider some examples which show that in special cases both questions can have positive answers even when the curvature is not constant.

Example 5 [15] Using the arithmetic theory of quadratic forms, one can construct pairs $(V; W)$ as in Example 4, where V is a hyperbolic n -manifold. Consider the quadratic form, $-2x_0^2 + x_1^2 + \dots + x_n^2$. Let Γ_n denote the group of isometries of this form with coefficients in the ring of integers of $\mathbb{Q}(\sqrt{2})$.

Then Γ_n is a discrete cocompact subgroup of $O(n;1)$. If Γ is any torsion-free subgroup of finite index in Γ_n , then $V = \mathbb{H}^n$ is a closed hyperbolic n -manifold. Note that for each i , $1 \leq i \leq n$, the hyperbolic reflection r_i , defined by $(x_0, \dots, x_i, \dots, x_n) \mapsto (x_0, \dots, -x_i, \dots, x_n)$ lies in Γ_n . If Γ is required to be normal in Γ_n , then r_i descends to an isometric involution τ_i of V . Its fixed point set V^{τ_i} is then a locally convex, codimension-one submanifold of V . Consequently, $W = V^{\tau_1} \setminus V^{\tau_2}$ is a locally convex, codimension-two submanifold of V . If we further require Γ to lie in the identity component of $O(n;1)$, then both V and W will be orientable. Finally, by passing to a subgroup of index two if necessary, we may suppose, that each V^{τ_i} separates V into two components. This implies that $0 = [W] \geq H_{n-2}(V)$. Hence, we can obtain a pair $(V; W)$ satisfying all the conditions of Theorem A. It follows that the induced (singular) metric on V_k is nonpositively curved. This metric is naturally piecewise hyperbolic. In [15] Gromov and Thurston show that this metric on V_k can be smoothed to a Riemannian metric with pinched negative curvature. Moreover, if $n \geq 4$, V_k is not diffeomorphic to a hyperbolic n -manifold for infinitely many k .

The construction of Example 3 can be used to show that each Gromov-Thurston example also admits nonpositively curved, piecewise Euclidean metric. Simply choose the net S in V so that its inverse image \mathcal{S} in \mathbb{H}^n is Γ -stable. The convex hull construction then gives a piecewise Euclidean metric on V with the property that each τ_i acts isometrically. Hence, the fixed point sets V^{τ_1} and V^{τ_2} as well as their intersection W will be locally convex. The induced length metric on V_k is piecewise Euclidean and by Theorem A, it is nonpositively curved.

Example 6 [8, page 383] In each dimension $n \geq 5$ one can construct a nonpositively curved, piecewise Euclidean metric on a smoothable manifold M^n such that its universal cover X (a Hadamard space) is not simply connected at infinity and hence, not homeomorphic to \mathbb{R}^n . On the other hand, by the Cartan-Hadamard Theorem, if M^n admitted a nonpositively curved Riemannian metric, then its universal cover would be homeomorphic to \mathbb{R}^n . Thus, M^n admits no such metric.

2 Eberlein's Rigidity Theorem in the singular case

Throughout this section, we shall make the following assumptions.

Let $(X; d)$ be a reducible global Riemannian symmetric space with no compact or Euclidean factors and of rank $r \geq 2$, and let Γ be an irreducible torsion-free uniform lattice on X . Set $V = X/\Gamma$.

Let $(X; d)$ be a Hadamard space and let Γ be a discrete group of isometries on X such that Γ and Γ' are isomorphic as groups and such that $V = X/\Gamma$ is compact. We further assume that V is a topological manifold.

The goal of this section is to show that the rigidity holds even without the smoothness assumption, namely, one has the following.

Theorem B $(X; d)$ is isometric to $(X; d')$ for some d' which differs from d only by constant multiples on the factors of X .

By the singular analogue ([3], Chapter IV, Theorem 6.1) of the Schroeder's Splitting Theorem, we know that the theorem holds true if V is (the finite undercover of) the product of several such irreducible compact locally symmetric spaces.

The proof will be a modification of the one given by Eberlein ([9], [10], [4]).

The key to understanding the geometry of a symmetric space X (reducible or not) is the study of its maximal flats. Given a maximal flat F in X and a point $x \in F$, one considers the other maximal flats which contain x . It turns out that F intersects these other maximal flats in a union of hyperplanes. These hyperplanes cut F into simplicial cones, any one of which is called a *Weyl chamber*. Given a Weyl chamber and some other point $y \in X$, there is a unique maximal flat F' which contains y and a Weyl chamber parallel to the given one.

A geodesic line in X is *singular* if it is contained in more than one maximal flat. It is *maximally singular* if it is the intersection of the maximal flats which contain it, in other words, if it is an extremal line in a decomposition of a maximal flat into simplicial cones.

2.1 The image flats

In this subsection, we don't need the assumption that X is reducible.

First of all, choose a homotopy equivalence $\bar{f}: V \rightarrow V'$ and an inverse $\bar{g}: V' \rightarrow V$. Let f and g be their lifts to the universal covers. Then f and g are

{equivariant, uniformly continuous, and they are $(k; b)$ quasi-isometries, that is,

$$\frac{1}{k}d(x; y) \leq d(f(x); f(y)) \leq kd(x; y)$$

for any x, y in X with $d(x; y) \leq b$, and similarly for g . We may also choose a large constant A such that for any x and y in X ,

$$d(x; gf(x)) \leq A; \text{ and } d(f(x); f(y)) \leq kd(x; y) + A$$

and similarly for any x and y in X ,

$$d(x; fg(x)) \leq A; \text{ and } d(f(x); f(y)) \leq kd(x; y) + A:$$

The group Γ acts on X via the isomorphism $\Gamma \cong \mathbb{R}^r$. An r -flat F in X is called *compact* if $F \cap K$ is compact. This is equivalent to the condition that F contains an abelian subgroup L of rank r such that F is contained in the minimum set of L (which is defined as the intersection of the minimum sets of all g in L).

Since V is compact, all elements of Γ are semisimple on X . So L is an abelian group of semisimple isometries on X , therefore in the space X , the minimum set $MIN(L) = \mathbb{R}^r$ splits isometrically, with L acting as identity on the convex set C and as a lattice of translations on \mathbb{R}^r . Pick any $p \in C$ and let $F = \mathbb{R}^r + fp$, then it is clear that both $Hd(F; f(F))$ and $Hd(F; g(F))$ are finite, since all of the subsets involved are L -invariant and have compact quotients. Therefore, we have proved the following:

For any compact r -flat F in X , there exists r -flat F' in X such that $Hd(F; f(F')) < 1$.

Note that $Hd(gF; F) < 1$, and the r -flat F in X , which satisfies this is unique. So, by the Quasi-flat Theorem of Kleiner-Leeb [16] or Eskin-Farb (Corollary 7.4, [13]), we know that there exists a constant $R^0 > 0$, which is independent of the choice of flats, such that $Hd(gF; F) < R^0$. By letting $R = kR^0 + 2A$, we have the following lemma.

Lemma 1 *There exists a constant $R > 0$ such that for any r -flat F in X , there is a r -flat F' in X with Hausdorff distance $Hd(F; f(F')) < R$.*

Furthermore, if F_1 is any r -flat in X with $Hd(F_1; f(F)) < 1$, then $Hd(F_1; f(F)) < R$.

Proof From the above discussion, the statement holds when F is compact. Since the set of compact r -flats is dense in the set of all r -flats in X , and any sequence of r -flats in X which all meet a compact set will have a convergent subsequence, the general case follows. □

We will call such a flat F an *image flat* of F , and F the *preimage flat* of F . Note that the preimage flat of F is unique, while union of all the image flats of F forms a closed convex set which splits as a product $F \times I$, with I convex and having diameter $\leq 2R$.

2.2 The reducibility of X

In this subsection, we want to show that the reducibility result of [9] holds in the singular case as well. Most of the proof from [9] goes through, with the following modifications.

First, we need to establish the so-called "duality condition" of X . Recall that X satisfies the *duality condition* on X , if for any geodesic c in X , there exists a sequence F_n such that $c(n) \in F_n$ and $c(-n) \in F_n$. To establish this we need the following lemma.

Lemma 2 *Let c be a geodesic in X . Then there exists an image flat F of a flat $F \subset X$, such that $d(c(t); F)$ is constant.*

Proof Since there are r flats in X passing through any two given points, we may choose $F_n \subset X$ passing through $c(n)$ and $c(-n)$. Let F_n be an image flat of F , $n = 1, 2, \dots$. Since

$$d(c(-n); F_n) \leq A + d(c(n); F_n) + R = A + R$$

by the convexity of the metric, we have $d(c(0); F_n) \leq A + R$, so by passing to a subsequence, we may assume that $F_{n_k} \rightarrow F$ and F is an image flat of F , the limit of the F_{n_k} . We have $d(c(-n_k); F) \leq A + R$ is bounded for all k ; hence, $d(c(t); F)$ is constant, by the convexity of the metric. \square

Lemma 3 *Any finite index subgroup of Γ satisfies the duality condition on X .*

Proof Fix a geodesic c in X . By Lemma 2, there exists a flat F and an image flat F such that c has constant distance from F . Let F_n be a sequence of compact flats which converges to F . For each n , let F_n be an image flat of F , and denote by L_n the corresponding lattice on F_n or F_n . Passing to subsequence if necessary, we may assume that $F_n \rightarrow \mathbb{E}^2$, which is also an image flat of F , hence c has constant distance to \mathbb{E}^2 . Let ε be the projection of c onto \mathbb{E}^2 . We can choose a sequence of compact geodesics $c_n \subset F_n$ such

that $c_n \neq \epsilon$. That is, there exists a sequence $f_k^{(n)} g \in L_n$ such that for each n , $f_k^{(n)}(c_n(0)) \neq c_n(+1)$ and $(f_k^{(n)})^{-1}(c_n(0)) \neq c_n(-1)$ when $k \neq 1$. By taking a diagonal subsequence in $f_k^{(n)} g$, we know that there exists sequence $f_n g$ such that $f_n(x) \neq c(+1)$ and $f_n^{-1}(x) \neq c(-1)$ as $n \neq 1$. \square

Lemma 4 (cf Lemma 2.4a of [4]) *Let $x \in X(1)$ be arbitrary and let $y \in X(1)$ be a point that can be joined by a geodesic to x . If $z \in X(1)$ is any point that can be joined to x then $z \in \overline{(y)}$.*

Proof With the duality condition established as above, the proof of Lemma 2.4a in [4] can be easily modified to cover the singular case. All we have to do is to estimate $\angle_q(ny; np)$ the same way for a point p on γ and allow q to be an arbitrary but fixed point in X , then $\angle_q(w; z) = 0$ for any q would imply $w = z$. Here w is any accumulation point of ny . \square

With the duality condition established, let us walk through the proofs in sections 4 and 5 of [9]. Our metric space $(X; d)$ is no longer smooth and we don't have the uniqueness of continuation of geodesic segments. Our starting point is the following proposition.

Proposition 1 (cf Proposition 4.1 of [9]) *Let γ be a maximally singular geodesic of X and let F be an r -flat containing γ . Let F be an r -flat in X such that $Hd(f(F); F) < 1$. Then*

- (1) *the points $x = \lim_{t \rightarrow +1} (f^{-1} \circ \gamma)(t)$ and $y = \lim_{t \rightarrow -1} (f \circ \gamma)(t)$ exist and are distinct in $X(1)$*
- (2) *for any point p in F it follows that $\angle_p(x; y) = \pi$ and there exists the unique geodesic joining x and y passing through p , and γ is contained in F .*

Proof For Proposition 1, the proof given in the appendix of [9] works. The only place where care needs to be taken is with regard to the paragraph on page 73, where the angles are used. In a general Hadamard space, the angle function $\angle_x(y; z)$, while it is still continuous in y and z , is only upper semicontinuous with respect to the vertex x . Also, two different geodesics from x could form a zero angle (in fact they could even share a segment).

Note that given two distinct points u, v in $X(1)$, there always exists a point $q \in X$ such that $\angle_q(u; v) > 0$. Also, $\angle_q(u; v)$ is continuous in u or v on $\overline{X} = X \cup \{X(1)\}$, and satisfies the triangle inequality

$$\angle_q(u; v) \leq \angle_q(u; w) + \angle_q(w; v):$$

With these properties, the paragraph on page 73 of [9] can easily be modified and the proof is valid. \square

Now we consider the situation where $X = X_1 \cup X_2$. Define $S = X(1)$ to be the set of endpoints at infinity of the maximally singular geodesics of X and let $S_i = S \cap X_i$ for $i = 1, 2$. (S is the set of *maximally singular points* in $X(1)$.)

By Proposition 1 the map f defines an equivariant map $\bar{f}: S \rightarrow X(1)$. Let \bar{S}_i denote the closure of $\bar{f}(S_i)$ in $X(1)$.

Proposition 2 (cf Proposition 4.3 of [9]) *Let S_1, S_2 be as above. Then:*

- (1) *Each of the sets S_1, S_2 is invariant under \bar{f} .*
- (2) *A point $x \in S_i$ can only be joined by a geodesic to some other point in S_i .*

Proof (1) is obvious. (2) follows by taking closures from the fact that, for $x \in \bar{f}(S_i)$, the point y of Proposition 1 belongs to S_i , together with part (1) and Lemma 4. \square

Now let us continue with the proofs in Appendix 2 of [9]. We start with Lemma 5.1. Lemma 5.1 is needed in the proof of Proposition 5.1 and Sublemma 5.4b. We will state the following weaker version, which is sufficient for our purpose.

Lemma 5 (cf Lemma 5.1 of [9]) *Suppose (t) is a geodesic in the CAT(0) space X and write $x = (1)$, $y = (-1)$, $p = (0)$. Denote by $L(p; x)$ and $B(p; x)$ the horosphere and horoball, respectively, which are centered at x and pass through the point p . Let $C = C_{xy} = \{q \in X \mid \angle_q(x; y) = \frac{\pi}{2}\}$. Then $L(p; x) \cap L(p; y) = B(p; x) \cap B(p; y)$, C is the union of all the geodesics parallel to t , C is a closed convex subset of X , and C splits as $C = C^0 \cup C^\theta$ for a closed convex subset $C^\theta \subset L(p; x) \cap L(p; y)$.*

Proof In the singular case, it is still true that if two geodesic rays meet at q with angle $\frac{\pi}{2}$, then they form a geodesic. Also, two parallel geodesics bound a flat strip (the ‘Flat Strip Lemma’ or ‘Sandwich Lemma’, see for example [2] or [3]). By the definition of Busemann function (see [2]) and the Law of Cosine in a comparison triangle, it is not hard to see that if $q \in B(p; x)$, then $\angle_p(x; q) > \frac{\pi}{2}$. Also, if $q \in B(p; x) \cap L(p; x)$, then $\angle_p(x; q) < \frac{\pi}{2}$. From this it follows that $B(p; x) \cap B(p; y) \subset L(p; x) \cap L(p; y)$. \square

For each point $p \in X$, let B_p denote the smallest convex subset of X which contains all the maximally singular geodesics through p with endpoints in S_1 .

Proposition 3 *With notation as in Lemma 5, suppose $x \in S_2$. Then B_p is contained in an R^d -neighborhood of C_{xy} , for some constant R^d .*

Proof This is essentially the proof of Proposition 5.1 of [9], but since we do not assume that C_{xy} is a proper subset of X , we get a weaker result. \square

Proposition 4 (cf Proposition 5.2 of [9]) *If $r \in B_p$ then $B_r \subset B_p$.*

Proof The proof of Proposition 5.2 of [9] is valid in the singular case as well. \square

Before stating the next proposition we need the following four lemmas.

Lemma 6 (cf Lemma 5.5a of [9]) *Let p, r be any two points of X . Then $d(q; B_r) = d(p; B_r)$ for every point $q \in B_p$. Moreover suppose that $p \notin B_r$ and let $P: X \rightarrow B_r$ denote the nearest point projection. Then for every $x \in S_1$ and any geodesic γ , with $\gamma(0) = p$ and $\gamma(1) = x$, the projection $(P \circ \gamma)(t)$ is a unit speed geodesic and γ and $P \circ \gamma$ bound a flat strip in X .*

Lemma 7 ([1, page 25, Corollary 5.8 (i)]) *Let $p, q \in X$ and $x \in X(1)$. If $\angle_p(x; q) + \angle_q(x; p) = \pi$, then the three points span a flat strip.*

Lemma 8 (cf Sublemma 5.4b of [9]) *Let γ be a geodesic of X such that $\gamma(0) \in B_p$, where $p = \gamma(0)$. Let $z = \gamma(1)$. Then there exists $x \in S_1$ and $t \in \mathbb{R}$ such that $\angle_{\gamma(t)}(x; z) \leq \pi/2$.*

Proof The proof of Sublemma 5.4b of [9] holds true using Lemma 5 and Lemma 7, instead of Lemma 5.1 and Sublemma 5.4a of [9]. \square

Lemma 9 (cf Lemma 5.4 of [9]) *For every point $p \in X$ there exist a number $A > 0$ such that $d(p; B_r) < A$ for every $r \in B_p$.*

Proof Again, the proof of Lemma 5.4 of [9] works using Lemma 6 instead of Lemma 5.5a of [9] and the fact that the limit of a convergent sequence of (half) flats is still a (half) flat. \square

Proposition 5 (cf Proposition 5.4 of [9]) *There exists a point $p \in X$ such that $B_r = B_p$ for every point $r \in B_p$.*

Proposition 6 (cf Proposition 5.5 of [9]) *For every point $p \in X$ and every point $r \in B_p$, $B_r = B_p$.*

Proposition 7 (cf Proposition 5.6 of [9]) *For every point $p \in X$ the set B_p is complete, totally geodesic subspace of X .*

Proof The proof of Proposition 5.6 of [9] goes through without change, except for the last paragraph of the proof on page 63, where angles are again used. What is needed to complete the proof is the following lemma.

Lemma 10 *Consider a geodesic triangle with vertices $p, q,$ and r . Let $\gamma(t), t \in [0; d(p; q)]$, be the geodesic segment from p to q , with $\gamma(0) = p$. If there is a sequence t_i of small positive numbers that converges to 0, such that $d(r; p) < d(r; \gamma(t_i))$ for each i , then $\angle_p(q; r) < \frac{\pi}{2}$.*

Proof Let $\gamma(t)$ be the geodesic segment from p to r , with $\gamma(0) = p, \gamma(l) = r$, where $l = d(p; r)$. Denote by d_{st} the distance between $\gamma(s)$ and $\gamma(t)$. Then the function

$$f(t; s) = (t^2 + s^2 - d_{st}^2) = 2st \cos \theta$$

is monotonically decreasing in $s > 0$ and $t > 0$, and its limit when $s \rightarrow 0, t \rightarrow 0$ is the cosine of the angle $\angle_p(q; r)$.

Now assume that $\angle_p(q; r) < \frac{\pi}{2}$. Then there exist $t > 0$ and $s > 0$ such that $f(t; s) > 0$. By the monotonicity of f in t , we have

$$(t_i^2 + s^2 - d_{st_i}^2) = 2st_i \cos \theta$$

for any $0 < t_i < t$. So for t_i sufficiently small, we have $s > d_{st_i}$. Hence for t_i sufficiently small,

$$d(r; p) = d(r; \gamma(s)) + s > d(r; \gamma(s)) + d_{st_i} > d(r; \gamma(t_i))$$

which contradicts our assumption. □

To complete the proof of Proposition 7, we apply the lemma to the triangle with vertices q_1, q_2 and r . We have that two of the internal angles are at least $\frac{\pi}{2}$. So the sum of the internal angles is at least π , which implies that it must be a triangle in \mathbb{R}^2 , with two right internal angles. This is clearly impossible. □

Summarizing the above discussions, we can now state the Eberlein's result in [9] in the singular case as the following:

Theorem C Let $X = X_1 \cup \dots \cup X_k$ be the decomposition into irreducible factors, $k \geq 2$. For $1 \leq i \leq k$, denote by $S_i = X_i(1)$ the set at infinity of the maximally singular geodesics in X_i . Then the following holds:

- (1) For any maximally singular geodesic $c \subset X$, there exists a geodesic $c' \subset X$ such that the Hausdorff distance $Hd(c; c') \leq R_1$ for some uniform constant R_1 .
- (2) If c is the intersection of r flats F_1, \dots, F_m , and c_1 is a geodesic parallel to all the image flats F_1, \dots, F_m , then c_1 is parallel to c .
- (3) Let $S_i = X_i(1)$ be the image of S_i guaranteed by (1). For any $x \in X$, denote by $A_i(x)$ the union of all the geodesic rays from x to the points in S_i , and let $B_i(x)$ be the smallest closed convex subset of X that contains $A_i(x)$. Then each $B_i(x)$ is totally geodesic in X , and $B_i(x)$ is parallel to $B_i(y)$ for any $x, y \in X$.
- (4) For any $x \in X$ and $z \in S_j$, let c be a geodesic with $c(0) = x$ and $c(1) = z$. Then for $i \neq j$, $B_i(x) \cap N_{R^0}(P_c)$, the R^0 -neighborhood of P_c . Here P_c denotes the set of points $y \in X$ where there is a geodesic through y that is parallel to c .

2.3 Transitivity of the isometry group on X

Our next goal is to establish the fact that the isometry group $I(X)$ of X acts transitively on X , hence $I(X)$ is a Lie group by the theorem of Montgomery and Zippin [20].

Let $X = X_1 \cup \dots \cup X_k$ be the decomposition into irreducible factors. By assumption, $k \geq 2$. In order to distinguish the actions of $I(X)$ on X and X , we will denote the lattice on X by Γ . By passing to a sublattice if necessary, we may assume that the irreducible lattice $I_0(X) = I_0(X_1) \cup \dots \cup I_0(X_k)$, where I_0 stands for the identity component of the isometry group. So each Γ_i is in the form $\Gamma_i = (\Gamma_1, \dots, \Gamma_k)$. Denote by π_i the projection of Γ on X_i . Since Γ_i is assumed to be irreducible, by Borel Density Theorem, we know that for each $1 \leq i \leq k$, π_i is injective, and π_i is dense in $I_0(X_i)$.

From the discussion in the last subsection, we know that for each $1 \leq i \leq k$, X is foliated by parallel, totally geodesic leaves B_i . By the Sandwich Lemma,

for each i , X splits isometrically as $B_i \times Y_i$. Basically, we want to show that these foliations are mutually perpendicular, and hence, that X has a product structure corresponding to that of X . But we can only prove this in the orthogonal complement of the Euclidean factor of X .

Denote by $C(X)$ the set of all Clifford translations in $I(X)$, the isometry group of X . It is a normal abelian subgroup of $I(X)$. X splits isometrically $X = \mathbb{R}^s \times Y$, $s \geq 0$, such that each $\gamma \in C(X)$ has the form $(T; id)$ under this splitting with T a translation, and so that Y does not contain any Euclidean factor. Furthermore, any isometry of X respects this splitting: $I(X) = I(\mathbb{R}^s) \times I(Y)$. (cf (6.2) of [2]).

Our first goal is to show that, for each $1 \leq i \leq k$, the leaves of B_i all respect the product structure $X = \mathbb{R}^s \times Y$. To be more precise, for any $x = (x_0; y) \in X$, $B_i(x) = B_{i0}(x_0) \times B_{iY}(y)$, and B_{i0} (or B_{iY}) is a parallel totally geodesic foliation in \mathbb{R}^s (or Y).

As before, for each $1 \leq i \leq k$, let $S_i = X_i(1)$ denote the set at infinity of the maximal singular geodesics in X_i ($1 \leq i \leq k$), and $S_i = X(1)$ the image guaranteed by part (1) of Theorem C. Again denote by $A_i(x)$ the union of all geodesic rays from x to the points in S_i , and $B_i(x)$ the smallest closed convex set containing $A_i(x)$.

Now consider an image flat $F \subset X$. Let us denote by F_0 and F_Y the projection of F in \mathbb{R}^s and Y , respectively. If we fix a point $x = (x_0; y) \in F$ and regard $F_0 = \mathbb{R}^s \times \{y\}$, $F_Y = \{x_0\} \times Y$, then we have the following lemma.

Lemma 11 For any image flat F in X , $F = F_0 \times F_Y$.

Proof Since any image flat is parallel to an image flat that is the limit of a sequence of compact image flats, it suffices to prove the lemma for compact image flats. Let L be a rank r abelian subgroup which acts as a lattice on F . For any $\gamma \in L$, write $\gamma = (\gamma_0; \gamma_Y)$ under the splitting $X = \mathbb{R}^s \times Y$. Then γ_0 is a translation on F_0 . This is true because, for any geodesic c_0 in F_0 , c_0 is the projection of some geodesic c in F . Since γ maps c to a parallel geodesic in F , $\gamma_0(c_0)$ is parallel to c_0 . Let T be a translation in F_0 . As T commutes with γ_0 for all $\gamma \in L$, TF is also invariant under L . Since we have uniform bound in Hausdorff distance between $g(F)$ or $g(TF)$ and F , the preimage flat, we know that the Hausdorff distance between F and TF must be bounded by a uniform constant. Therefore, F_0 must be contained in F , since otherwise we can choose a translation in F_0 that sends F to a parallel TF with arbitrarily large $Hd(F; TF)$. □

As a corollary of this, we have the following.

Lemma 12 For each $1 \leq i \leq k$, the set $S_i = \mathbb{R}^S(1) \sqcup Y(1)$. So the foliation B_i is a product foliation $B_{i0} \times B_{iY}$. Furthermore, $Y = B_{1Y} \times \dots \times B_{kY}$.

Proof Since any maximally singular geodesic $c \subset X_i$ is an intersection of r flats, there exist r flats F_1, \dots, F_m with $c = F_1 \cap \dots \cap F_m$. Let F_j be an image flat of F_j , $1 \leq j \leq m$. Then part (1) of Theorem C says that there exists a geodesic $c \subset X$ which is parallel to all F_j , and any geodesic ε parallel to all F_j must be parallel to c . In the product space $X = \mathbb{R}^S \times Y$, c is parallel to F_j if and only if c_0 is parallel to F_0 and c_Y is parallel to F_Y . Here the subscripts denote the projections. So if c is not parallel to a leaf of \mathbb{R}^S or Y , then any line in the plane $c_0 \times c_Y$ would be parallel to all those F_j , which is impossible. So, c must be parallel to either \mathbb{R}^S or Y . That is, S_i is contained in the disjoint union of $\mathbb{R}^S(1)$ and $Y(1)$.

From this and the definition of B_i , it is clear that for each i , $B_i = B_{i0} \times B_{iY}$ is a product foliation.

Next we claim that for $i \neq j$, B_{iY} and B_{jY} are perpendicular. Take any geodesic c with $c(1) \in S_j \setminus Y(1)$. Then any leaf $B_i(x)$ is contained in the neighborhood $N_{R^0}(P_c)$ by part (4) of Theorem C. Here P_c stands for the union of all geodesics parallel to c . So if the projection τ of c onto B_{iY} is not a point, any point in B_{iY} will be within distance R^0 to another point in B_{iY} where there is a geodesic parallel to τ passing through. That is, $B_{iY} = N_{R^0}(P)$, where P is the union of all geodesics in B_{iY} that is parallel to τ . Since B_{iY} is geodesically complete, and $P \subset B_{iY}$ is convex, we know that $B_{iY} = P$. That is, B_{iY} is foliated by geodesics parallel to τ , so B_{iY} , hence Y will contain a Euclidean factor, a contradiction. So geodesics from $x \in B_{iY}$ to any point in $S_j \setminus Y(1)$ must be perpendicular to B_{iY} . So $A_j(x)$ must be contained in $Z(x)$, the leaf through x of the orthogonal complement of B_{iY} in Y , where $Y = B_{iY} \times Z$ by the Sandwich lemma. Since $Z(x)$ is convex, we have $B_{jY} \subset Z$, hence B_{iY} is perpendicular to B_{jY} . On the other hand, since any geodesic in X is parallel to an image flat, which is contained the span of B_1 through B_k , so Y is spanned by those B_{iY} , $1 \leq i \leq k$. \square

Note that if X_i is of rank 1, then S_i , hence S_i is connected, so either $S_i = \mathbb{R}^S(1)$ or $S_i = Y(1)$. In this case B_i is entirely in \mathbb{R}^S or Y .

Write $X = \mathbb{R}^S \times B_{1Y} \times \dots \times B_{kY}$. Each \mathcal{Z} can be written in the form $\mathcal{Z} = (0; 1Y; \dots; kY) = (0; Y)$, since it preserves all the foliations B_i and

respects the product structure $X = \mathbb{R}^s \times Y$. Denote by Γ_Y the group of all such γ . Set $\Gamma = \prod_{i=1}^k \Gamma_i$, where Γ_i is the projection of Γ onto the irreducible factor X_i . The group Γ is dense in $I_0(X)$.

We want to construct a surjective, continuous map $f: X \rightarrow Y$ which is $(\Gamma; \Gamma)$ -equivariant. Then Γ_Y will have a dense orbit in Y . Because Y is locally compact, any sequence of isometries $f_n g$ on Y such that $f_n(y)$ converges for some $y \in Y$ will have a subsequence that converges to an element in $I(Y)$. Therefore, $I(Y)$ acts transitively on Y . Hence, it is a Lie group by the theorem of Montgomery and Zippin [20]. Thus, $I(X)$ is also a Lie group.

If Q is a bounded subset in a Hadamard space Y , we will denote by r_Q the inimum of the radii of the closed balls containing Q . There exists an unique point $y \in Y$ such that $Q \subset B_{r_Q}(y)$. This point is called the *circumcenter* of Q and is denoted by $\tau(Q)$. We refer the reader to page 26 of [1] for more details.

Lemma 13 *Suppose P and Q are two bounded subsets in a Hadamard space Y with Hausdorff distance $Hd(P; Q) = h$. Then their circumcenters satisfy*

$$d(\tau(P); \tau(Q)) \leq \frac{h}{r_P + r_Q + h}:$$

Proof Write $x = \tau(P)$ and $y = \tau(Q)$. Then we have $P \subset B_{r_P}(x) \setminus B_{r_Q+h}(y)$. Let z be the mid point of the geodesic segment from x to y . Then since Y is Hadamard, any $p \in P$ will satisfy

$$d^2(p; z) \leq \frac{1}{2}d^2(p; x) + \frac{1}{2}d^2(p; y) - \frac{1}{4}d^2(x; y) \leq \frac{1}{2}r_P^2 + \frac{1}{2}(r_Q + h)^2 - \frac{1}{4}d^2(x; y):$$

Since this is true for any $p \in P$, the far right hand side of the above inequality must be bigger than $\frac{r_P^2}{2}$, hence

$$d^2(x; y) < 2r_Q^2 - 2r_P^2 + 4r_Qh + 2h^2:$$

Add this with the similar inequality obtained by reversing the role of P and Q , we get

$$d^2(x; y) < 2h(r_P + r_Q + h):$$

If one replace z by the points on the geodesic segment from x to y that are very closed to x or y , then the coefficient 2 in the right hand side can be removed. □

We are now ready to construct f . Let π be the projection map from X onto B_{1Y} . For any $x \in X_1$, write $Q(x) = \pi x \cap X_2 \cap \dots \cap X_k$, and $\tau(x) = \tau(Q(x))$. Define $f_1(x) = \tau(Q(x))$ to be the circumcenter of $Q(x)$. For

this we need to show that $Q(x)$ is always a bounded subset of B_{1Y} . Fix a point $p \in Q$. For any $q \in Q$, there exists a r -flat $F \subset X$ containing both p and q . So, if r_1 denotes the rank of X_1 , then there are $m = r - r_1$ maximally singular geodesics in F , denoted by l_1, \dots, l_m , each is perpendicular to X_1 , $p \in l_1$, $q \in l_m$, and $l_i \perp l_{i+1}$ for $1 \leq i \leq m - 1$. For each l_i , by Theorem C, there exists geodesic l_i in X perpendicular to B_{1Y} , such that the Hausdorff distance between l_i and $f(l_i)$ is bounded by an uniform constant R_1 . Therefore $d(f(q), f(p)) \leq 2mR_1 \leq 2rR_1$, and $Q \subset B_{2rR_1}$. Since f is uniformly continuous, and $Hd(Q(x), Q(y)) \leq d(x, y)$, so Lemma 6 implies that $f_1: X_1 \rightarrow B_{1Y}$ is continuous. It is also clear that for any $\epsilon > 0$, we have $f_1(\epsilon_1(x)) = \epsilon_Y(f_1(x))$. Now if we define $f_i: X_i \rightarrow B_{iY}$ similarly for each $1 \leq i \leq k$ and then take the product, we get a continuous map $f: X \rightarrow Y$ which is $(f; \gamma)$ -equivariant, where $f = \prod_{i=1}^k f_i$. This implies the following.

Lemma 14 *The map $f: X \rightarrow Y$ is surjective. Hence $\overline{\rho_Y}$ acts transitively on Y , and $I(X)$ is a Lie group.*

Note that for any $x \in X$, we have $d(f(x), \gamma f(x)) \leq \rho_Y k 2rR_1$. So for any $y \in Y$,

$$d(y, f \circ g(0; y)) \leq A + 2r \rho_Y k R_1.$$

Therefore f is surjective by the following lemma.

Lemma 15 *If a Hadamard space Y is a topological manifold and if $h: Y \rightarrow Y$ is a continuous map within bounded distance from the identity map, then h is surjective.*

Proof The geodesic segment from $h(y)$ to y gives a proper homotopy between h and the identity map. Hence, h induces the identity map on the top dimensional cohomology group with compact support, $H_c^n(Y)$, $n = \dim(Y)$. The result follows. □

2.4 Proof of the Rigidity Theorem

Now we can apply the techniques in [10], [11], [12] and [4] to finish the proof of Theorem B. From the previous subsection, $I(X)$ is Lie group. This implies that X has no Euclidean factor, that is, we have the following lemma.

Lemma 16 $X = Y = B_1 \times \dots \times B_k$.

Proof First, notice that X does not contain any Clifford translations. This is because the set of Clifford translations in X forms a normal abelian subgroup. By the Main Theorem of [11], X does not contain any nontrivial normal abelian subgroup, since X has no Euclidean de Rham factor.

Now let us assume that $X = \mathbb{R}^s \times Y$ for $s > 0$. We want to derive a contradiction. Note that Y is not a point, for otherwise X would contain a translation by the Bieberbach Theorem.

Denote by $\pi_{\mathbb{R}}$ and π_Y the projection of X on \mathbb{R}^s and Y , respectively. Since $I(X)$ is a Lie group, the proof of Lemma A in [11] says that π_Y is discrete. Let N be the kernel of π_Y . The proof of Theorem 4.1 of [12] says that either $\pi_{\mathbb{R}}$ is discrete or N contains a Clifford translation. Since X is assumed to be irreducible, $\pi_{\mathbb{R}}$ can not be discrete. Since X contain no Clifford translation, so does the subgroup N . This completes the proof. (Note that in this argument, Proposition 2.3 of [4] is used. But it is obviously valid in the singular case.) \square

In our next argument we also need an extension of Theorem 2.4 of [4] to singular spaces. The proof there is based on four lemmas. We already established singular version of Lemma 2.4a | see Lemma 4. The same sort of modification works for Lemma 2.4c. The proof of Lemma 2.4b holds without change. For the proof of Theorem 2.4 the following replacement of Lemma 2.4d suffices.

Lemma 17 *Let $A \leq I(X)$ be a nontrivial abelian subgroup such that normalizer D of A in $I(X)$ satisfies the duality condition. Then $I(X)$ contains a Clifford translation.*

Proof (cf proof of Lemma 2.4d of [4]) Let $L(A) \subset X(1)$ denote the limit set of A . By Lemma 2.4b of [4], $L(A)$ is not empty. Let $x \in L(A)$, then by Lemma 2.4c there exist a unique point $y \in L(A)$ such that x can be joined to y . If $z \in H(1)$ is a point that can be joined to y then by Lemma 4 $z \in \overline{D(x)} \subset L(A)$ and therefore $z = x$. So X admits Clifford translations along the geodesics joining x to y by Sandwich Lemma. \square

Denote by $\pi_i = \pi_{iY}$ the projection of X on B_i , and write $G_i = (\overline{\pi_i})_0 = I_0(B_i)$ the identity component of its closure. By the proof of Lemma 3.1 of [10], which essentially uses Theorem 2.4 of [4] (cf Lemma 17), we know that each G_i is a centerless semisimple Lie group of noncompact type. Fix a maximal compact subgroup $K \subset G_i$. By taking the circumcenter of an orbit, we know that K will fix some point in B_i . Let $\rho \in B_i$ be a fixed point of K . Since B_i is locally compact, the isotropy subgroup of G_i at ρ is compact, and equals to K since

K is maximal compact. So the transitivity of the action of G_i on B_i gives a homeomorphism between B_i and G_i/K . The metric d on B_i is a G_i -invariant Hadamard metric. By the following lemma, d must be smooth, so $(B_i; d)$ is a global Riemannian symmetric space of noncompact type. Therefore $(X; d)$ is symmetric, and isometric to $(X; d)$ by the theorem of Mostow [19]. This completes the proof of Theorem B.

Lemma 18 *Suppose G is a semisimple Lie group of noncompact type and that K a maximal compact subgroup. If d is a Hadamard metric on G/K that is G -invariant, then $(G/K; d)$ is Riemannian symmetric space of noncompact type.*

Proof If G is not simple, we may take a lattice $\Gamma \subset G$ and consider $X = G/\Gamma K$ equipped with d symmetric and $X = G/K$ equipped with d . By the previous arguments, we know that $(X; d)$ will be a product space of $(G_i/K_i; d_i)$, where those G_i are the simple factors of G . Therefore we may assume that G is simple.

First consider the case when the rank of G is 1. In this case, the geodesic sphere in d is a single K orbit, so it coincides with a geodesic sphere with the same center under d , the symmetric metric. For any two points $p, q \in X$, let m be the midpoint under the metric d . Then there are two geodesic spheres under d , centered at p and q , respectively, so that m is their unique intersection point. This implies that m must be on the geodesic segment from p to q with respect to the metric d . So the geodesics in d and d have the same images.

Fix p and q with $d(p; q) = 1$. Let d be the symmetric metric on $X = G/K$ such that $d(p; q) = 1$. Let γ_t be a one parameter subgroup of G such that $\gamma(1) = q$ and $\gamma(0) = p$ is the unit speed geodesic under d with $\gamma(0) = p$.

By considering $\gamma_{\frac{1}{2^n}}$, we know that $d(\gamma(0); \gamma(t)) = d(\gamma(0); \gamma(t))$ for any rational number t , whose denominator is a power of two. Hence, it also holds for any $t \in \mathbb{R}$ by continuity. For any $p^\theta, q^\theta \in X$, and for any $\gamma \in G$ such that $\gamma(p^\theta) = p$ and such that $\gamma(q^\theta)$ lies in the image of γ , we have $d(p^\theta; q^\theta) = d(p; q)$. So d is symmetric in this case.

Now assume G is simple with rank $r \geq 2$. We need the following.

Sublemma *Suppose G is a connected, centerless, semisimple Lie group of noncompact type, and K a maximal compact subgroup. Let $\mathfrak{g} = \mathfrak{k} + \mathfrak{p}$ be the Cartan decomposition of the Lie algebras. Suppose \mathfrak{a} is a maximal abelian subalgebra in \mathfrak{p} , and $A = \exp(\mathfrak{a})$. Denote by $M = Z_K(A)$ the centralizer of A*

in K , and denote by Y the fixed point set in $G=K$ of M . Then Y is a totally geodesic submanifold in $G=K$ containing the maximal flat $F = Ax_0$, where $x_0 = K$. Denote by r the rank of $G=K$.

- (1) $M = \mathbb{H}^2$ if and only if $G=K$ is the product of r copies of the hyperbolic plane \mathbb{H}^2 .
- (2) If G is simple, then either $Y = (\mathbb{H}^2)^r$, or $Y = F = \mathbb{R}^r$, depending on whether $G=K$ is Hermitian symmetric or not.

Now let us use Sublemma to finish the proof of Lemma 18. Let $F \subset B = G=K$ be a maximal flat under d , with K the isotropy subgroup at $x_0 \in F$. First we claim that F is d -convex.

Let $\mathfrak{g} = \mathfrak{k} + \mathfrak{p}$ be the Cartan decomposition of the Lie algebras of G and K , and \mathfrak{h} a maximal abelian subalgebra contained in \mathfrak{p} . Write $A = \exp \mathfrak{h}$ so that $F = Ax_0$. As in Sublemma, denote by M the centralizer of A in K , and denote by Y the fixed point set of M in B . Then $Y \subset B$ is totally geodesic in B under d .

Note that $Y = G_1=K_1$ is a symmetric space of nonpositive curvature, with $G_1 \subset G$ and $K_1 \subset K$. Since Y is the fixed point set of d -isometries, so it is d -convex, and G_1 acting transitively on Y as both d and d -isometries.

By Sublemma, either $Y = F$, in which case F is a d -flat, or $Y = (\mathbb{H}^2)^r$. In this case, the arguments in the previous subsections implies that $(Y; d)$ must split as the product of r surfaces, each is symmetric since $(\mathbb{H}^2; d)$ has rank 1. So d is a symmetric metric on Y , and F is a d -flat.

Now consider any maximal flat $F \subset B$ passing through x_0 . The restriction of d to each flat F is a Euclidean metric which is invariant under the Weyl group, which acts irreducibly on F since G is assumed to be simple. So $d|_F = (F)d|_F$. Since any two flats through x_0 can be joined by finitely many flats through x_0 so that each one intersects the next at more than one point, $(F)d|_F$ must be constant, and $d = d|_F$ is symmetric. This completes the proof of Lemma 18 assuming Sublemma. □

Proof of Sublemma Let us start with part (1). Without loss of generality, we may assume that G is simple. Assume $M = 1$, we want to conclude that $G=K$ is the hyperbolic plane. Let r be the rank of $G=K$.

Since the Lie algebra \mathfrak{m} of M is trivial, $\mathfrak{h}_{\mathbb{C}}$ is a Cartan subalgebra for $\mathfrak{g}_{\mathbb{C}}$, the complexification of \mathfrak{g} . Denote by G^{θ} the real points of $G_{\mathbb{C}}$ under the conjugation

of \mathfrak{g} in $\mathfrak{g}_{\mathbb{C}}$. Then $G = \text{Ad}(G)$ is the identity component of G^{θ} . Denote by K^{θ} the maximal compact subgroup of G^{θ} with identity component K .

Let $P = (\mathbb{Z}_2)^r$ be the group of 2-torsion elements in $H_{\mathbb{C}} = \exp(\mathfrak{h}_{\mathbb{C}})$. Then P is contained in $Z_{K^{\theta}}(A)$, the centralizer of A in K^{θ} .

Note that the number of connected components in G^{θ} is bounded by the order of the outer automorphism group $Q = \text{Aut}(G)/\text{Int}(G)$. So the assumption $M = Z_K(A) = 1$ implies that $2^r \leq |Q|$.

By the Table 10 on page 156 of [18], we see that this is impossible unless $\mathfrak{g}_{\mathbb{C}}$ is $\mathfrak{a}_1 = \mathfrak{sl}(2; \mathbb{C})$ or $\mathfrak{d}_4 = \mathfrak{so}(8; \mathbb{C})$. In the \mathfrak{d}_4 case, in fact in all four classical Lie algebra cases, it can be easily checked that $M \neq 1$ except for \mathfrak{a}_1 . This completes the proof of part (1).

For part (2), let us apply part (1) to the symmetric space (which may contain a Euclidean factor now) Y , since $M_1 = Z_{K_1}(A)$ is trivial, by part (1), we know that $Y = (\mathbb{H}^2)^s \times \mathbb{R}^{r-s}$. We want to show that s is either 0 or r if G is simple.

Let $M^{\theta} = N_K(A)$ be the normalizer of A in K . Then M^{θ} (or M) consists of elements in K that stabilize F (fix every point in F). M is a normal subgroup of finite index in M^{θ} , and $W = M^{\theta}/M$ is Weyl group which acts irreducibly on F . For each $g \in M^{\theta}$, since $gMg^{-1} = M$, we have $g(Y) = Y$. Now since any isometry of Y preserves the Euclidean factor, we know that the \mathbb{R}^{r-s} part of F is invariant under W . So either $s = 0$ or $s = r$. This completes the proof of Sublemma. \square

Remark Note that the CAT(0) assumption on d in Lemma 18 is necessary. When rank $r \geq 2$, there are G -invariant length metric on G/K which are geodesic. The simplest example would be taking product of two copies of symmetric spaces, and let $d = d_1 + d_2$ where d_i are the standard metrics on the factors. Then clearly d is G -invariant but not Riemannian. More generally, Planché showed [21] that the set of G -invariant Finsler metrics on G/K is in one-to-one correspondence with the set of W -invariant norms on a maximal flat F . (Here W denotes the Weyl group.) Some of these metrics are even *uniquely* geodesic, that is, any two points in G/K can be joined by a unique geodesic segment.

References

- [1] **W Ballmann**, *Lectures on Spaces of Nonpositive Curvature*, Birkhäuser (1995)
- [2] **W Ballmann, M Gromov, V Schroeder**, *Manifolds of Nonpositive Curvature*, Birkhäuser (1985)
- [3] **M Bridson, A Haefliger**, *Spaces of nonpositive curvature*, Springer-Verlag, Berlin, Heidelberg, New York (1999)
- [4] **S-S Chen, P Eberlein**, *Isometry groups of simply connected manifolds of non-positive curvature*, Illinois Journal of Math. 24 (1980) 73{103
- [5] **R Charney, M W Davis**, *Singular metrics of nonpositive curvature on branched covers of Riemannian manifolds*, Amer. J. Math. 115 (1993) 929{1009
- [6] **R Charney, M W Davis**, *The polar dual of a convex polyhedral set in hyperbolic space*, Michigan Math. J. 42 (1995) 479{510
- [7] **R Charney, M W Davis, G Moussong**, *Nonpositively curved piecewise Euclidean structures on hyperbolic manifolds*, Michigan Math. J. 44 (1997) 201{208
- [8] **M W Davis, T Januszkiewicz**, *Hyperbolization of polyhedra*, J. Di . Geom. 34 (1991) 347{388
- [9] **P Eberlein**, *Rigidity of lattices of nonpositive curvature*, Ergodic Theory & Dynamical Systems, 3 (1983) 47{85
- [10] **P Eberlein**, *Isometry groups of simply connected manifolds of non-positive curvature, II*, Acta Math. 149 (1982) 41{69
- [11] **P Eberlein**, *Euclidean de Rham factor of a lattice of nonpositive curvature*, J. Di . Geometry, 18 (1983) 209{220
- [12] **P Eberlein**, *Lattices in spaces of nonpositive curvature*, Ann. Math. 111 (1980) 435{476
- [13] **A Eskin, B Farb**, *Quasi-flats and rigidity in higher rank symmetric spaces*, J. Amer. Math. Soc. 10 (1997) 653{692
- [14] **M Gromov**, *Hyperbolic groups*, Essays in group theory (SM Gersten, ed.), MSRI Publ. 8, Springer-Verlag, Berlin, Heidelberg, New York (1987) 75{264
- [15] **M Gromov, W Thurston**, *Pinching constants for hyperbolic manifolds*, Invent. Math. 89 (1987) 1{12
- [16] **B Kleiner, B Leeb**, *Rigidity of quasi-isometries for symmetric spaces and Euclidean buildings*, Inst. Hautes Etudes Sci. Publ. Math. No. 86 (1997) 115{197
- [17] **B Leeb**, *A characterization of irreducible symmetric spaces and Euclidean buildings of higher rank by their asymptotic geometry*, preprint (1997)
- [18] **O Loos**, *Symmetric spaces: Volume II*, W.A. Benjamin, Inc. (1969)
- [19] **G Mostow**, *Strong rigidity of locally symmetric spaces*, Annals of Math. Studies 78, Princeton University Press (1973)

- [20] **D Montgomery, L Zippin**, *Topological Transformation Groups*, Interscience, New York (1955)
- [21] **P Planche**, *Structures de Finsler invariantes sur les espaces symetriques*, CR. Acad. Sci. Paris, 321 (1995) 1455{1458
- [22] **M S Raghunathan**, *Discrete subgroups of Lie groups*, Springer{Verlag, Berlin, Heidelberg, New York (1972)
- [23] **A Selberg**, *Recent developments in the theory of discontinuous groups of motion of symmetric spaces*, Springer Lecture Notes in Math. 118, Springer{Verlag, Berlin, Heidelberg, New York (1969)