


Geometry & Topology
 Volume 4 (2000) 451{456
 Published: 26 November 2000



Di eomorphisms, symplectic forms and Kodaira brations

Claude LeBrun

*Department of Mathematics, SUNY at Stony Brook
 Stony Brook, NY 11794-3651, USA*

Email: cl aude@math. sunysb. edu

Abstract

As was recently pointed out by McMullen and Taubes [7], there are 4 {manifolds for which the di eomorphism group does not act transitively on the deformation classes of orientation-compatible symplectic structures. This note points out some other 4 {manifolds with this property which arise as the orientation-reversed versions of certain complex surfaces constructed by Kodaira [3]. While this construction is arguably simpler than that of McMullen and Taubes, its simplicity comes at a price: the examples exhibited herein all have large fundamental groups.

AMS Classi cation numbers Primary: 53D35

Secondary: 14J29, 57R57

Keywords: Symplectic manifold, complex surface, Seiberg{Witten invariants

Proposed: Ronald Stern
 Seconded: Yasha Eliashberg, Ronald Fintushel

Received: 11 June 2000
 Accepted: 21 November 2000

Let M be a smooth, compact oriented 4-manifold. If M admits an orientation-compatible symplectic form, meaning a closed 2-form ω such that $\omega \wedge \omega$ is an orientation-compatible volume form, one might well ask whether the space of such forms is connected. In fact, it is not difficult to construct examples where the answer is negative. A more subtle question, however, is whether the group of orientation-preserving diffeomorphisms $M \rightarrow M$ acts transitively on the set of connected components of the orientation-compatible symplectic structures of M . As was recently pointed out by McMullen and Taubes [7], there are 4-manifolds M for which this subtler question also has a negative answer. The purpose of the present note is to point out that many examples of this interesting phenomenon arise from certain complex surfaces with Kodaira fibrations.

A *Kodaira fibration* is by definition a holomorphic submersion $f: M \rightarrow B$ from a compact complex surface to a compact complex curve, with base B and fiber $F_z = f^{-1}(z)$ both of genus $g \geq 2$. (In C^1 terms, f is thus a locally trivial fiber bundle, but nearby fibers of f may well be non-isomorphic as complex curves.) One says that M is a *Kodaira-bered surface* if it admits such a fibration f . Now any Kodaira-bered surface M is algebraic, since $K_M = f^* K_B + (2g-2)F$ is obviously positive for sufficiently large g . On the other hand, recall that a holomorphic map from a curve of lower genus to a curve of higher genus must be constant.¹ If $f: M \rightarrow B$ is a Kodaira fibration, it follows that M cannot contain any rational or elliptic curves, since composing f with the inclusion would result in a constant map, and the curve would therefore be contained in a fiber of f ; contradiction. The Kodaira-Enriques classification [2] therefore tells us that M is a minimal surface of general type. In particular, the only non-trivial Seiberg-Witten invariants of the underlying oriented 4-manifold M are [8] those associated with the canonical and anti-canonical classes of M . Any orientation-preserving self-diffeomorphism of M must therefore preserve $f \cdot c_1(M)g$.

We have just seen that M is of Kähler type, so let ω denote some Kähler form on M , and observe that ω is then of course a symplectic form compatible with the usual 'complex' orientation of M . Let ν be any area form on B , compatible with its complex orientation, and, for sufficiently small $\epsilon > 0$, consider the closed 2-form

$$\omega_\epsilon = \omega - \epsilon f^* \nu;$$

¹Indeed, by Poincaré duality, a continuous map $h: X \rightarrow Y$ of non-zero degree between compact oriented manifolds of the same dimension must induce inclusions $h_*: H^j(Y; \mathbb{R}) \rightarrow H^j(X; \mathbb{R})$ for all j . Such a map h therefore cannot exist whenever $b_j(X) < b_j(Y)$ for some j .

Then

$$\frac{! \wedge !}{\mu} = -2(f^{-1})^* \wedge + \mu^* \wedge = (\mu - hf^{-1}; i)^* \wedge ;$$

where the inner product is taken with respect to the Kähler metric corresponding to μ . Now $hf^{-1}; i$ is a positive function, and, because M is compact, therefore has a positive minimum. Thus, for a sufficiently small $\mu > 0$, $! \wedge !$ is a volume form compatible with the *non-standard* orientation of M ; or, in other words, $!$ is a symplectic form for the reverse-oriented 4-manifold \overline{M} . For related constructions of symplectic structures on fiber-bundles, cf [6].

It follows that \overline{M} carries a unique deformation class of almost-complex structures compatible with $!$. One such almost-complex structure can be constructed by considering the (non-holomorphic) orthogonal decomposition

$$TM = \ker(f^*) \oplus f^*(TB)$$

induced by the given Kähler metric, and then reversing the sign of the complex structure on the 'horizontal' bundle $f^*(TB)$. The first Chern class of the resulting almost-complex structure is thus given by

$$c_1(\overline{M}; !) = c_1(M) - 4(1 - g)F;$$

where g is the genus of B , and where F now denotes the Poincare dual of a fiber of f . For further discussion, cf [4, 5, 9].

Of course, the product $B \times F$ of two complex curves of genus $g \geq 2$ is certainly Kodaira fibered, but such a product also admits orientation-reversing diffeomorphisms, and so, in particular, has signature $\sigma = 0$. However, as was first observed by Kodaira [3], one can construct examples with $\sigma > 0$ by taking *branched covers* of products; cf [1, 2].

Example Let C be a compact complex curve of genus $k \geq 2$, and let B_1 be a curve of genus $g_1 = 2k - 1$, obtained as an unbranched double cover of C . Let $\tau: B_1 \rightarrow B_1$ be the associated non-trivial deck transformation, which is a free holomorphic involution of B_1 . Let $\rho: B_2 \rightarrow B_1$ be the unique unbranched cover of order 2^{4k-2} with $\rho^{-1}(B_2) = \ker[\tau_1(B_1) \rightarrow H_1(B_1; \mathbb{Z}_2)]$; thus B_2 is a complex curve of genus $g_2 = 2^{4k-1}(k - 1) + 1$. Let $\pi: B_2 \rightarrow B_1$ be the union of the graphs of ρ and $\tau \circ \rho$. Then the homology class of π is divisible by 2. We may therefore construct a ramified double cover $M \rightarrow B_2 \rightarrow B_1$ branched over π . The projection $f_1: M \rightarrow B_1$ is then a Kodaira fibration, with fiber F_1 of genus $2^{4k-2}(4k - 3) + 1$. The projection $f_2: M \rightarrow B_2$ is also a Kodaira fibration, with fiber F_2 of genus $4k - 2$. The signature of this doubly Kodaira-fibered complex surface is $\sigma(M) = 2^{4k}(k - 1)$.

We now axiomatize those properties of these examples which we will need.

Definition Let M be a complex surface equipped with two Kodaira fibrations $f_j: M \rightarrow B_j$, $j = 1, 2$. Let g_j denote the genus of B_j , and suppose that the induced map

$$f_1 - f_2: M \rightarrow B_1 \times B_2$$

has degree $r > 0$. We will then say that $(f_1; f_2)$ is a *Kodaira double-fibration* of M if $c_1(M) \neq 0$ and

$$(g_2 - 1) \neq r(g_1 - 1):$$

In this case, $(M; f_1; f_2)$ will be called a *Kodaira doubly-bered surface*.

Of course, the last hypothesis depends on the ordering of $(f_1; f_2)$, and is automatically satisfied, for fixed r , if $g_2 > g_1$. The latter may always be arranged by simply replacing M and B_2 with suitable covering spaces.

Note that $r = 2$ in the explicit examples given above.

Given a Kodaira doubly-bered surface $(M; f_1; f_2)$, let \overline{M} denote M equipped with the non-standard orientation, and observe that we now have two different symplectic structures on \overline{M} given by

$$\begin{aligned} \omega_1 &= \omega - f_1^* \omega_1 \\ \omega_2 &= \omega - f_2^* \omega_2 \end{aligned}$$

for any given area forms ω_j on B_j and any sufficiently small $\epsilon > 0$.

Theorem 1 *Let $(M; f_1; f_2)$ be any Kodaira doubly-bered complex surface. Then for any self-diffeomorphism $\phi: M \rightarrow M$, the symplectic structures ω_1 and ω_2 are deformation inequivalent.*

That is, ω_1 , $-\omega_1$, ω_2 , and $-\omega_2$ are always in different path components of the closed, non-degenerate 2-forms on \overline{M} . (The fact that ω_1 and $-\omega_1$ are deformation inequivalent is due to a general result of Taubes [10], and holds for any symplectic 4-manifold with $b^+ > 1$ and $c_1 \neq 0$.)

Theorem 1 is actually a corollary of the following result:

Theorem 2 *Let $(M; f_1; f_2)$ be any Kodaira doubly-bered complex surface. Then for any self-diffeomorphism $\phi: M \rightarrow M$,*

$$[c_1(\overline{M}; \omega_2)] \neq c_1(\overline{M}; \omega_1):$$

Proof Because $c_1(M) \neq 0$, any self-diffeomorphism of M preserves orientation. Now M is a minimal complex surface of general type, and hence, for the standard ‘complex’ orientation of M , the only Seiberg–Witten basic classes [8] are $\pm c_1(M)$. Thus any self-diffeomorphism of M satisfies

$$[c_1(M)] = \pm c_1(M);$$

Letting F_j be the Poincaré dual of the fiber of f_j , and letting g_j denote the genus of B_j , we have

$$c_1(\overline{M}; !_j) = c_1(M) + 4(g_j - 1)F_j$$

for $j = 1, 2$. The adjunction formula therefore tells us that

$$[c_1(\overline{M}; !_j)] \cdot [c_1(M)] = (2g_j + 3)(M) - 2c_1(M) = 3c_1(M) \neq 0;$$

where the intersection form is computed with respect to the ‘complex’ orientation of M .

If we had a diffeomorphism $\phi : M \rightarrow M$ with $[c_1(\overline{M}; !_2)] = [c_1(\overline{M}; !_1)]$, this computation would tell us that that

$$[c_1(M)] = \pm c_1(M) \implies [c_1(\overline{M}; !_2)] = c_1(\overline{M}; !_1)$$

and that

$$[c_1(M)] = -c_1(M) \implies [c_1(\overline{M}; !_2)] = -c_1(\overline{M}; !_1);$$

In either case, we would then have

$$4(g_1 - 1)F_1 = c_1(\overline{M}; !_1) - c_1(M) = \pm [c_1(\overline{M}; !_2) - c_1(M)] = \pm 4(g_2 - 1)F_2;$$

On the other hand, $F_1 \cdot F_2 = r$, so intersecting the previous formula with F_2 yields

$$4(g_1 - 1)r = \pm 4(g_2 - 1)F_1 \cdot F_2 = \pm 4(g_2 - 1)r;$$

and hence

$$(g_2 - 1) = \pm (g_1 - 1);$$

in contradiction to our hypotheses. The assumption that $[c_1(\overline{M}; !_1)] = [c_1(\overline{M}; !_2)]$ is therefore false, and the claim follows. \square

Theorem 1 is now an immediate consequence, since the first Chern class of a symplectic structure is deformation-invariant.

Acknowledgment This work was supported in part by NSF grant DMS-0072591.

References

- [1] **M F Atiyah**, *The signature of fibre-bundles*, from \Global Analysis (Papers in honor of K Kodaira)", Univ. Tokyo Press, Tokyo (1969) 73{84
- [2] **W Barth C Peters, A V de Ven**, *Compact complex surfaces*, Springer{Verlag (1984)
- [3] **K Kodaira**, *A certain type of irregular algebraic surfaces*, J. Analyse Math. 19 (1967) 207{215
- [4] **D Kotschick**, *Signatures, monopoles and mapping class groups*, Math. Res. Lett. 5 (1998) 227{234
- [5] **N C Leung**, *Seiberg{Witten invariants and uniformizations*, Math. Ann. 306 (1996) 31{46
- [6] **D McDuff, D Salamon**, *Introduction to symplectic topology*, Oxford Science Publications, The Clarendon Press and Oxford University Press, New York (1995)
- [7] **C T McMullen, C H Taubes**, *4-manifolds with inequivalent symplectic forms and 3-manifolds with inequivalent fibrations*, Math. Res. Lett. 6 (1999) 681{696
- [8] **J Morgan**, *The Seiberg{Witten equations and applications to the topology of smooth four-manifolds*, vol. 44 of Mathematical Notes, Princeton University Press (1996)
- [9] **J Petean**, *Indefinite Kähler{Einstein metrics on compact complex surfaces*, Comm. Math. Phys. 189 (1997) 227{235
- [10] **C H Taubes**, *More constraints on symplectic forms from Seiberg{Witten invariants*, Math. Res. Lett. 2 (1995) 9{14