


Geometry & Topology
Volume 7 (2003) 645{711
Published: 28 October 2003



Calculus III: Taylor Series

Thomas G Goodwillie

*Department of Mathematics, Brown University
Box 1917, Providence RI 02912{0001, USA*

Email: tomg@math.brown.edu

Abstract

We study functors from spaces to spaces or spectra that preserve weak homotopy equivalences. For each such functor we construct a universal n -excisive approximation, which may be thought of as its n -excisive part. Homogeneous functors, meaning n -excisive functors with trivial $(n - 1)$ -excisive part, can be classified: they correspond to symmetric functors of n variables that are reduced and 1-excisive in each variable. We discuss some important examples, including the identity functor and Waldhausen's algebraic K -theory.

AMS Classification numbers Primary: 55P99

Secondary: 55U99

Keywords: Homotopy functor, excision, Taylor tower

Proposed: Haynes Miller
Seconded: Ralph Cohen, Gunnar Carlsson

Received: 8 November 2002
Accepted: 20 October 2003

Introduction

This paper should have been finished many years ago. It is a continuation of [9] and [10] (which were themselves a little late in coming). The author has no excuse for this procrastination and wishes to apologize to anyone who has been inconvenienced, especially his students and former students.

As in [9] and [10] we are concerned with functors from \mathcal{C} , which may be the category of spaces or more generally spaces over a fixed space, to \mathcal{D} , which may be either the category of spaces or the category of spectra. It is clear that one could extend the ideas to many other settings. The functors will always be *homotopy functors* in the sense that they preserve weak homotopy equivalences. We are also concerned with (natural) maps between such functors. We call such a map of functors $F \dashv G$ a weak equivalence (or for emphasis an objectwise weak equivalence, or for brevity an equivalence) if for every object X the map $F(X) \dashv G(X)$ is a weak equivalence. The goal is to shed light on homotopy functors in general (and on particular ones) by systematically comparing them with homotopy functors of some very special kinds, namely those which satisfy " n^{th} order excision".

In taking this point of view we are led naturally to speak of the category $\mathcal{F} = \mathcal{F}(\mathcal{C}; \mathcal{D})$ of homotopy functors. Some of our conclusions refer to the homotopy category $h\mathcal{F}$ of homotopy functors, meaning the universal example of a category equipped with a functor from \mathcal{F} that takes every weak equivalence to an isomorphism. For example, some of our results are most neatly expressed as statements to the effect that such and such a functor from one category of functors to another induces an equivalence of homotopy categories. Homotopy categories in this sense can be made by the "generators and relations" construction: a morphism in the homotopy category is an equivalence class of words, where each letter in a word is either a forward arrow (a morphism in \mathcal{F}) or a backward arrow (a formal inverse to a weak equivalence in \mathcal{F}) and two words are declared to represent the same morphism if they are related by a sequence of basic moves (composing two forward or two backward arrows, deleting an identity arrow, cancelling a forward arrow and its formal inverse).

The author has recently been reminded that there are set-theoretic objections to bandying this kind of language about too freely, but he does not want to stop doing so. Some ways of resolving this difficulty will be discussed at the end of this introduction.

Of course, if one wants to prove theorems about a homotopy category then one should be prepared to do most of the work in the original category. For example, commutative diagrams in the category of functors will be a ubiquitous

tool here, but commutative diagrams in the homotopy category are relatively useless.

We now recall the definition of n^{th} order excision, from section 3 of [10]. Whereas ordinary, or first order, excision has to do with the behavior of a functor on certain square diagrams, n^{th} order excision has to do with the behavior of a functor on certain cubical diagrams of dimension $n + 1$. We briefly recall the definitions. The reader can consult the opening sections of [10] for details, including the specific models we are using for homotopy limits and colimits and the basic definitions and facts about cubical diagrams.

The homotopy functor F is *excisive* if it takes homotopy pushout squares (also called homotopy cocartesian squares) to homotopy pullback squares (also called homotopy cartesian squares). It is *reduced* if it takes the final object to an object weakly equivalent to the final object. It is *linear* if it is both excisive and reduced.

A cubical diagram is called strongly homotopy cocartesian if all of its two-dimensional faces are homotopy pushouts. It is called homotopy cartesian if it is a homotopy pullback in the sense that the final object is equivalent (by the obvious map) to the homotopy limit of all the others. In this paper we have generally omitted the word "homotopy" in the expressions "(strongly) homotopy (co)cartesian". F is said to be *n-excisive*, or to satisfy n^{th} order excision, if it takes strongly homotopy cocartesian $(n + 1)$ -cubical diagrams to homotopy cartesian diagrams. An n -excisive functor is always $(n + 1)$ -excisive.

In [9] we studied the approximation of homotopy functors by excisive functors and codified this in the notion of (first) derivative of a functor. We calculated the derivative, first in some basic examples and then in the example which had given rise to the whole project: stable pseudoisotopy theory (and with it Waldhausen's algebraic K -theory functor A).

In [10] we introduced the class of *analytic* functors. These are homotopy functors whose deviation from being n -excisive is bounded in a certain way for all n . Some functors are more analytic than others; an analytic functor is ϵ -analytic for some integer ϵ , where a smaller ϵ means stronger bounds. We showed that ϵ -analytic functors behave rather rigidly on the category of ϵ -connected spaces (and more generally on the category of spaces equipped with $(\epsilon + 1)$ -connected maps to a fixed space), in the sense, roughly, that if a map between two such functors induces an equivalence of first derivatives everywhere then its homotopy fiber is constant up to homotopy within the category of ϵ -connected spaces.

In [5] these results were used to relate A to some functors built out of the free loop space.

Here we move from excisive approximation to n -excisive approximation, obtaining functors $P_n F$ which can be thought of as Taylor polynomials of F . We show that if F is \mathbb{A}^1 -analytic and X is \mathbb{A}^1 -connected then as n tends to infinity the n^{th} approximation converges to $F(X)$ in the strong sense that the connectivity of a certain map $F(X) \rightarrow (P_n F)(X)$ tends to infinity. In other words, the number ρ_n gives a sort of radius of convergence.

What we were doing in [10] was like showing that a function $f(x)$ is determined, in some region, by $f(0)$ and $f'(x)$. Continuing with the same analogy, what we are doing here is like showing that $f(x)$ is determined by $f(0)$, $f'(0)$, $f''(0)$ and so on.

Here is a sketch of the main results of this paper, presented in a rather different order from the one in which they will be proved.

First consider homotopy functors $T \rightarrow \text{Sp}$ from based spaces to spectra. For any such functor, and for any $n \geq 0$, we can make an n -excisive functor $T \rightarrow \text{Sp}$ with a map $F \rightarrow P_n F$ that is universal (in a homotopy category) among maps from F to n -excisive functors. The functors $\{P_n F\}_{n \geq 0}$ together form a tower, and F maps into the limit of the tower.

If the role of n^{th} Taylor polynomial is being played by $P_n F$, then the role of n^{th} term in the series is played by the (homotopy) fiber of the map $P_n F \rightarrow P_{n-1} F$, which will be denoted by $D_n F$. We refer to $P_n F$ as the n^{th} stage of the tower and to $D_n F$ as the n^{th} layer. These *homogeneous polynomial functors* are the focus of much of the paper.

The constant term $(P_0 F)(X)$ is the same, up to natural weak equivalence, as the constant functor $F(\ast)$.

Up to the same kind of equivalence, the linear (or homogeneous of degree one) functor $(D_1 F)(X)$ necessarily has the form $C_1 \wedge X$ for some fixed spectrum C_1 , at least when restricted to finite complexes X . The coefficient spectrum C_1 , which of course is $(D_1 F)(S^0)$, is called the derivative of F at the one-point space.

Likewise the homogeneous quadratic functor $(D_2 F)(X)$ is necessarily given by

$$(C_2 \wedge (X \wedge X))_{h_2}$$

where h_2 is the symmetric group, C_2 is some spectrum with h_2 -action, $X \wedge X$ has the obvious action, and the subscript h_2 denotes homotopy orbit spectrum. The coefficient spectrum C_2 , with its h_2 -action, is called the second derivative of F at the one-point space.

The explanation of why homogeneous quadratic functors always have this form involves *symmetric bilinear* functors. If H is any 2-excisive functor (still from

based spaces to spectra), then its second order cross-effect, defined as the total fiber (= homotopy fiber of homotopy fibers) of the square diagram

$$\begin{array}{ccc}
 H(X_1; X_2) & \longrightarrow & H(X_1) \\
 \downarrow \cong & & \downarrow \cong \\
 H(X_2) & \longrightarrow & H(\ast);
 \end{array}$$

will be a functor $L(X_1; X_2)$ of two variables, linear in each variable, and symmetric with respect to interchanging the variables. It turns out that the homogeneous quadratic part of H can be recovered as $L(X; X)_{h_2}$. A bilinear functor necessarily has the form

$$L(X_1; X_2) = C \wedge (X_1 \wedge X_2);$$

and a symmetry on L means a \mathbb{Z}_2 -action on C .

This pattern persists. For any $n \geq 1$ an n^{th} degree homogeneous functor H must have the form

$$H(X) = L(X; \dots; X)_{h_n}$$

where L , the n^{th} order cross-effect of H , is a symmetric multilinear functor of n variables. Thus $(D_n F)(X)$ will have the form

$$(C_n \wedge (X \wedge \dots \wedge X))_{h_n}$$

at least for finite X . The object C_n , a spectrum with an action of \mathbb{Z}_n , will be called the n^{th} derivative of F at the one-point space. We also refer to it as the coefficient of the multilinear functor, or of the homogeneous functor.

It is worth emphasizing what brand of stable equivariant homotopy theory is appearing here. Let G be a finite group. To do serious homotopy theory in the category of G -spaces one usually takes the weak equivalences to be those equivariant maps which induce weak equivalences on spaces of H -fixed points for all subgroups H . (For CW objects this is equivalent to saying that the map has an inverse up to G -homotopy.) There is also a very weak notion of equivalence: equivariant maps which are nonequivariantly weak equivalences. Using these as the weak equivalences leads to a theory that is easier and less interesting, but which has its uses. For example, a very weak equivalence of G -spaces always induces a weak equivalence of homotopy orbit spaces and also of homotopy fixed point spaces. Every G -space is equivalent in the very weak sense to a free G -space (product with EG), and for free G -spaces the two kinds of equivalence coincide.

In the course of investigating homogeneous functors we encounter spectra with finite groups acting on them, and this leads us to an obvious stable analogue

of this easier brand of equivariant homotopy theory: Make a category whose objects are spectra equipped with a G -action and whose maps are maps of spectra respecting the action, and call such an equivariant map a (very) weak equivalence if it is nonequivariantly a weak equivalence of spectra. It is these objects (with $G = \mathbb{Z}/n$) up to this kind of equivalence which correspond to homogeneous functors of degree n up to weak natural equivalence.

In order to extend the main ideas of *serious* equivariant homotopy theory to the stable setting, May and his collaborators created a beautiful and powerful theory of G -spectra [12]. We mention this only to say that we do not need it here. From that sophisticated point of view our spectra with G -action are what are sometimes called *naive* G -(pre)spectra, namely G -spectra involving only trivial actions of G on suspension coordinates. The G -spectrum expert will know what else to say about the place of the naive theory in the sophisticated one, but neither the expert nor the novice should have to think about the sophisticated theory to read this paper (unless the expert just cannot help thinking about it).

Returning to the towers, a simple and familiar example is the "Snaith splitting" of the functor $F(X) = \mathbb{Z}/n[X]$. Its n^{th} homogeneous part is $(D_n F)(X) = \mathbb{Z}/n[X^{\wedge n}]$, and its tower splits: $P_n F = \mathbb{Z}/n \oplus \mathbb{Z}/n \oplus \dots \oplus \mathbb{Z}/n = \bigoplus_k D_k F$. The limit of the tower is the product of all the layers. If X is connected then the tower converges to $F(X)$ in the sense that the map from $F(X)$ to the homotopy limit of the tower is a weak equivalence, and in fact in the stronger sense that the map from $F(X)$ to $(P_n F)(X)$ has a connectivity tending to infinity with n .

A related example is $F(X) = \mathbb{Z}/n[X]$. Here we have

$$(D_n F)(X) = \mathbb{Z}/n[X^{\wedge n}];$$

(We will give a quick proof of this using the previous example and the general fact $P_n(F \oplus G) = (P_n F) \oplus (P_n G)$. See Example 1.20 below.) The n^{th} derivative is the wedge of n factorial copies of the spectrum S^{-n} permuted transitively by \mathbb{Z}/n . The tower does not split. It converges (in the strong sense again) when the space X is 1-connected.

A more general example is $F(X) = \mathbb{Z}/n \text{Map}(K; X)$ where K is a based finite complex and Map means the space of based maps. Here the n^{th} derivative is the S -dual of a based complex with \mathbb{Z}/n -action, namely the quotient of the smash product $K^{\wedge n}$ by the fat diagonal. F is $\dim(K)$ -analytic, and the tower converges to $F(X)$ if X is $\dim(K)$ -connected. Arone [1] thoroughly explored this class of examples, giving a concrete description of the stages $P_n F$ and not just the layers $D_n F$.

The n^{th} coefficient of Waldhausen's $A(X)$ is again the S -dual of a based finite complex with \mathbb{Z}/n -action, namely $(\mathbb{Z}/n)_+ \wedge_{C_n} S^{n-1}$, where $C_n = \mathbb{Z}/n$ is the

transitive cyclic subgroup of order n and the sphere S^{n-1} is the one-point compactification of the reduced regular representation of C_n .

So far we have been concerned with functors from based spaces to spectra. We now discuss three different variants of this setup: functors from unbased spaces, functors from spaces over a fixed space, and functors to spaces. In these new settings the Taylor tower construction goes through with no change, but some additional work is needed to understand what a homogeneous functor looks like.

The switch from functors $T \dashv Sp$ of based spaces to functors $U \dashv Sp$ of unbased spaces is fairly innocuous. If $T \dashv U$ is the forgetful functor then $(P_n F) \cong P_n(F)$ and $(D_n F) \cong D_n(F)$.

A snag appears when one tries to relate homogeneous functors to symmetric multilinear functors, since the definition of the cross-effect requires basepoints. The good news is that in the end this does not matter: homogeneous functors $T \dashv Sp$ of any degree extend uniquely (in an appropriate up-to-natural-weak-equivalence sense) to U , and the same is true for symmetric multilinear functors in any number of variables. The proof involves the left adjoint of Σ , in other words the functor that adds a disjoint basepoint to an unbased space. For example, although the usual definition of the suspension spectrum of a space X requires X to be based, there is a well-known extension to unbased spaces. It associates to X the homotopy fiber of

$$\Sigma^{-1}(X) \dashv \Sigma^{-1}(\ast):$$

If X is based then this is naturally (with respect to based maps) equivalent to $\Sigma^{-1}X$, and on the other hand if $U \dashv Sp$ is any linear functor such that L is (naturally equivalent to) Σ^{-1} then L must be as defined above. This was explained in [9] and is generalized to higher degrees here.

Note that for inhomogeneous functors this goes very wrong. For example, the 1-excisive functor

$$\Sigma^{-1}(\ast) \text{ -- fiber of } (\Sigma^{-1}(X) \dashv \Sigma^{-1}(\ast)):$$

from U to Sp is genuinely different from $\Sigma^{-1}(X)$, but the difference disappears after restriction to T . (The empty set is the only space X at which the functors disagree, but even for nonempty spaces the equivalence cannot be chosen to be natural with respect to unbased maps.) An example of a 2-excisive functor $T \dashv Sp$ which does not extend to U at all is the suspension spectrum of

$$F(X) = ((X \wedge X) =_X)_2;$$

the orbit space for the \mathbb{Z}_2 -action on the quotient of $X \wedge X$ by the diagonal. If X is the disjoint union of a point and a circle then the rational homology of $F(X)$ depends on where the basepoint is placed in X .

The second switch, from functors of spaces to functors of spaces over a fixed space, is something like the switch from MacLaurin series to general Taylor series. Instead of building, for each space X , a tower that has $F(\ast)$ at the bottom and attempts to converge to $F(X)$, one can build, for each map of spaces $X \rightarrow Y$, a tower that has $F(Y)$ at the bottom and attempts to converge to $F(X)$. If we fix Y and think of everything in sight as a functor of spaces over Y then the n^{th} stage of the tower is n -excisive. As in the case $Y = \ast$, there are two options, each with its own technical advantages: the category U_Y of plain spaces over Y and the category T_Y of spaces over Y equipped with a section. We sometimes refer to the latter as fiberwise based spaces over Y .

The correspondence between homogeneous and symmetric multilinear functors works the same for spaces over Y as it does for spaces, but the business of describing a multilinear functor by coefficient spectra becomes more complicated when Y has more than one point. As a trivial example, to describe a linear functor of spaces over the two-point space $\{y_1, y_2\}$ one needs two spectra. In general a linear functor L of spaces over Y determines a spectrum for each point $y \in Y$, namely the coefficient $L(Y \rightarrow_y S^0)$ of the linear functor $Z \rightarrow L(Y \rightarrow_y Z)$ from T to Sp , where $Y \rightarrow_y Z$ means the wedge sum of $(Y; y)$ with the based space Z , viewed as a space over Y . This spectrum depends "continuously" on the point y in some sense. If Y is path-connected then up to weak homotopy equivalence the spectrum is independent of the point; but there is a twisting that must not be overlooked.

In [9] we defined the differential $D_Y F$ of a functor $U \rightarrow Sp$ at a space Y to be a linear functor from spaces over Y to spectra, the linear approximation to the functor

$$(X \rightarrow Y) \rightarrow \text{ho } \text{ber}(F(X) \rightarrow F(Y))$$

We also defined the derivative to be a spectrum $@_y F(Y)$ depending on a space Y and a point $y \in Y$, namely $(D_Y F)(Y \rightarrow_y S^0)$; it might be called the y coefficient of $D_Y F$ or the partial derivative of F at Y in the y direction. The relationship between the differential $D_Y F$ (a linear functor of spaces over Y) and the derivative $@_y F(Y)$ (a spectrum) is something like the relationship between the differential of a function (a linear function on the tangent space) and a partial or directional derivative (a number, which specifies the behavior of that linear function on a certain one-dimensional tangent subspace). The spectrum $@_y F(Y)$ records the derivative of F at Y in the "direction" given by y . Here we make the multilinear generalization, defining the n^{th} differential

to be the symmetric multilinear functor corresponding to the n^{th} layer of the Taylor tower and defining the n^{th} derivative

$$D_n^{(n)} F(Y)$$

to be its value at $(Y_{-y_1} S^0; \dots; Y_{-y_n} S^0)$.

The third switch, from spectrum-valued to space-valued functors, brings a real surprise. The construction of the Taylor series goes through with no change. If the functor F happens to be $\Sigma^{-1} G$ for some spectrum-valued functor G , then we have $P_n F = \Sigma^{-1} P_n G$ and $D_n F = \Sigma^{-1} D_n G$. The surprise is that although most functors are not of the form $\Sigma^{-1} G$ this is not reflected at the homogeneous level: every homogeneous functor $T \in \mathcal{T}^n$ of degree $n - 1$ is infinitely deloopable, in fact in a unique and functorial way. Thus $D_n F$ always has the form $\Sigma^{-1} ((C_n \wedge X^{\wedge n})_{h_n})$ and even in the space-valued case we can introduce a spectrum called the n^{th} derivative of the functor, or the coefficient spectrum of the n^{th} homogeneous layer.

A central example of a functor $T \in \mathcal{T}^n$ that is interesting from this point of view is the identity. Let us call it I . Its first derivative is the sphere spectrum. It is easy to see, in any of a number of ways, that the n^{th} derivative is equivalent to the wedge sum of $(n - 1)!$ copies of the $(1 - n)$ -sphere spectrum, with Σ_n acting in such a way that the subgroup Σ_{n-1} freely permutes the summands. To identify $D_n I$ one has to know the action of the full group Σ_n . Johnson [11] gave an explicit finite complex with Σ_n -action whose S-dual is the answer. Arone and Mahowald [4] gave a different answer of that kind, showed that it was equivalent to Johnson's, and used it to make some very interesting calculations.

A basic example of an inhomogeneous functor that does not deloop is $P_2 I$, the 2-excisive approximation of the identity functor. The functor $P_1 I$ is of course $Q = \Sigma^{-1} \Sigma^{-1}$. $(P_2 I)(X)$ is another standard object, the homotopy fiber of the James-Hopf map

$$QX \rightarrow Q((X \wedge X)_{h_2}):$$

(which is not induced by a spectrum map from $\Sigma^{-1} X$ to $\Sigma^{-1} (X \wedge X)_{h_2}$).

One notable feature of this last example turns out to be quite general: the fibration sequence

$$D_n F \rightarrow P_n F \rightarrow P_{n-1} F$$

can always be extended one step to the right, as long as the constant functor $P_0 F$ is contractible. This falls out of the proof of (and is essentially equivalent to) the deloopability of homogeneous functors.

The construction of the Taylor tower also goes through for functors taking values in unbased instead of based spaces, but in that case it is nonsense to

we speak of the layers $D_n F$ as functors, since that would involve using the fiber of a map of unbased spaces. This small fussy point is really not so small. The identity functor I above was from based spaces to based spaces. Let J be the identity functor from unbased spaces to unbased spaces. Then $(P_1 J)(X)$ is the homotopy fiber of

$$Q(X_+) \rightarrow QS^0;$$

but not with respect to the usual basepoint in QS^0 . In particular $(P_1 J)(\cdot)$ is empty, so that there is no natural basepoint in $(P_1 J)(X)$. This seriously interferes with defining $D_2 J$. $P_1 J$ is excisive and reduced, but perhaps too badly twisted to be rightly called linear. For such reasons we hesitate to even speak of homogeneous functors to unbased spaces.

The paper is organized as follows:

Section 1 defines the tower $fP_n Fg$ in the general case, proves that $P_n F$ is n -excisive, establishes the universal mapping property of $P_n F$, and notes the convergence of the tower in the case of an analytic functor F .

Section 2 shows that homogeneous space-valued functors can be functorially delooped and concludes that they correspond precisely to homogeneous spectrum-valued functors.

Section 3 establishes the correspondence between homogeneous functors and symmetric multilinear functors in the case of functors from fiberwise based spaces over Y to spectra. By Section 2 this result extends to space-valued functors.

Section 4 shows that for homogeneous functors, and also for symmetric multilinear functors, it does not matter whether the domain category is fiberwise based or unbased spaces, so that some of the results of Section 3 extend to functors of plain spaces over Y .

Section 5 establishes notation for the symmetric multilinear functors that (according to sections 2{4) encode the homogeneous layers of a Taylor tower, and develops the idea of coefficient spectra for multilinear functors.

Section 6 establishes a useful tool for working out examples. The key point is that the n^{th} cross effect of the homogeneous functor $D_n F$ can be made by "multilinearizing" the n^{th} cross effect of F itself.

Section 7 works out the n^{th} derivatives of functors like $\text{Map}(K; -)$.

Section 8 recalls and discusses known results on the Taylor tower of the identity.

Section 9 indicates how to get the n^{th} derivative of Waldhausen's A , taking the first derivative as starting point.

A few words about set theory are in order. We all know that we must not speak of the set of all sets or the set of all topological spaces; set theory, as formulated to avoid Russell-type paradoxes, does not allow it. And since the category \mathcal{C} of spaces is not small, there may be objections to speaking of the category of all functors, or homotopy functors, from \mathcal{C} to \mathcal{D} in that case. Even if \mathcal{C} and \mathcal{D} are both small, the functor category will not be small in general, so that it is illegal to make a new category by using generators and relations to invert some morphisms.

We choose to dodge this as follows. As usual when basing mathematics on set theory, we begin by fixing a universe U of sets. Now by topological spaces we mean those whose point sets are sets in this strict sense. The category of such spaces is not a small category, any more than the set of all (U -)sets is a (U -)set. Nevertheless, if we are willing to work in two universes [14; I.6], we are not in such bad shape. Introduce a larger universe U^0 in which the set of all U -sets is a set. The category of all (U -)spaces is then U^0 -small. In U^0 there can be no objection to speaking of the category of all functors from spaces to spaces, or of the full subcategory of homotopy functors. To invert the weak equivalences by generators and relations, one can always pass to a third universe in which the category of homotopy functors is small.

This solution will not suit all tastes. It may be a bit wasteful and it may be a bit crude. For some more refined purposes it will be inadequate. For example, if one wants to introduce some sort of space of maps between two homotopy functors such that the set of components will be the set of morphisms in the homotopy category, then it will certainly be a drawback to find that this "function space" is not a space in the original sense. In general, if one wants to work very seriously with the homotopy category of functors then one will probably want to introduce a closed model structure on the functor category, with all the benefits that that brings. (In fact, although we have not attempted to do so, there are certainly many reasons for reworking this whole theory in the context of closed model categories. The objects of study should be functors from one (small?) model category to another, subject to some mild axioms, and the category of homotopy functors should turn out to be a model category, too.)

On the other hand, in this paper the role of the homotopy categories is a modest one: they are used here mainly as shorthand, to make some sentences briefer and more readily comprehensible than they otherwise would be. The "multiple universes" device allows us to use this shorthand without telling any lies, but without making the category of spaces artificially small.

1 The Taylor tower

Let $C \xrightarrow{F} D$ be a homotopy functor, where $C = C_Y$ is either U_Y or T_Y and D is either U , T , or Sp . Let $n \geq 0$.

The n -excisive approximation $P_n F$ will be constructed by the infinite iteration of another construction T_n which is designed to bring the functor F a little closer to being n -excisive. The special case when $n = 1$ was treated in [9; Def. 1.10], where T_1 and P_1 were called T and P . Thus if $Y = \ast$ and $F(\ast) = F$ then $T_1 F = F$ and $P_n F = \text{hocolim}_{i \geq 0} T^i F$.

The definition of T_n uses the *berwise join* over Y , as introduced in the proof of [10; 5.1]. Let X be a space over Y and let U be a space. (In most applications U will be a discrete finite set.) The *berwise join* is the space

$$X \underset{Y}{\vee} U = \text{hocolim}(X \times X \times U \rightrightarrows Y \times U)$$

considered as a space over Y . The name signifies that the functor "*join with U* " is being applied to all the fibers of $X \rightarrow Y$.

If U has one element, then $X \underset{Y}{\vee} U \rightarrow Y$ is the *berwise cone* of X over Y (the mapping cylinder of $X \rightarrow Y$ considered as a space over Y). If U has two elements then $X \underset{Y}{\vee} U \rightarrow Y$ is the *berwise suspension* and will sometimes be denoted by $\underset{Y}{\vee} X$.

It should be noted that this construction has the best of both worlds, in that on the one hand each fiber of $X \underset{Y}{\vee} U \rightarrow Y$ is homeomorphic to the join with U of the corresponding fiber of $X \rightarrow Y$, and on the other hand each homotopy fiber is weakly homotopy equivalent to the join with U of the corresponding homotopy fiber of $X \rightarrow Y$.

Recall that even if C is T_Y rather than U_Y then $X \underset{Y}{\vee} U$ is still in C ; it inherits a section from X . In other words, if X is a *berwise based space* over Y then $X \underset{Y}{\vee} U$ is also canonically *berwise based*, without for example choosing a point in U .

The object $X \underset{Y}{\vee} U$ depends (bi-)functorially on X and U . Note also that there is a natural isomorphism of spaces over Y :

$$(X \underset{Y}{\vee} U) \underset{Y}{\vee} V = X \underset{Y}{\vee} (U \vee V)$$

where $U \vee V$ is the ordinary join of two spaces.

Let $P(\underline{n+1})$ be the poset of subsets of $\underline{n+1} = \{1, \dots, n+1\}$. Any object $X \in C$ yields an $(n+1)$ -dimensional cubical diagram in C .

$$\begin{array}{c} P(\underline{n+1}) \rightarrow C \\ U \rightarrow X \underset{Y}{\vee} U \end{array}$$

Consider the composed functor $U \dashv F(X \dashv U)$, a cubical diagram in D . The homotopy limit of its restriction to $P_0(\underline{n+1})$, the poset of nonempty subsets of $\underline{n+1}$, will be called $(T_n F)(X)$. Clearly this yields a homotopy functor $C \xrightarrow{T_n F} D$. There is a natural map $F \xrightarrow{t_n F} T_n F$, since any cubical diagram determines a map from the "initial" object to the homotopy limit of the others (see [10; Def. 1.2]):

$$F(X) = F(X \dashv ;) \xrightarrow{t_n F} \operatorname{holim}_{U \subseteq P_0(\underline{n+1})} F(X \dashv U) = (T_n F)(X)$$

Let $(P_n F)(X)$ be the sequential homotopy colimit of the diagram

$$F(X) \xrightarrow{(t_n F)(X)} (T_n F)(X) \xrightarrow{(t_n T_n F)(X)} (T_n^2 F)(X) \xrightarrow{(t_n T_n^2 F)(X)} \dots$$

Clearly $P_n F$ is a homotopy functor and we have a natural map $F \xrightarrow{p_n F} P_n F$.

The cubical diagram $U \dashv X \dashv U$ is strongly cocartesian for all X . Therefore if the functor F is n -excisive then the maps $t_n F$ and $p_n F$ will be weak equivalences for all X . In this sense n -excisive functors are unchanged by T_n and P_n .

We will see below that $P_n F$ is always n -excisive, and that (in a homotopy category) it is the best n -excisive approximation to F in a categorical sense.

1.1 Remark If the map $X \dashv Y$ is m -connected, then for nonempty U the map $X \dashv U \dashv Y$ is $(m+1)$ -connected, and therefore $(T_n^i F)(X)$ depends only on the behavior of F on objects whose structural maps to Y are $(m+i)$ -connected. In this sense the Taylor approximations of a functor $C_Y \dashv D$ depend only on the restriction of the functor to objects "arbitrarily close to Y " (just as the Taylor expansion of $f(x)$ in powers of $x - y$ depends only on the behavior of $f(x)$ in an arbitrarily small neighborhood of y). A related comment is that P_n commutes (up to natural isomorphism) with fiberwise suspension:

$$P_n(F \dashv Y) = (P_n F) \dashv Y$$

This follows from the natural isomorphism

$$Y \dashv (X \dashv U) = (Y \dashv X) \dashv U$$

Before giving the general proof that $P_n F$ is n -excisive, we give a different proof under the added assumption that F is *stably n -excisive* [10; 4.1], and in this case we show that $P_n F$ approximates F not only in a categorical sense but also from the point of view of connectivity.

The first task is to show that if F is stably n -excisive then $T_n F$ is more nearly n -excisive than F is, and that $T_n F$ agrees with F to n^{th} order in the following sense:

1.2 Definition A map $u: F \rightarrow G$ between two functors from \mathcal{C}_Y to D is said to satisfy $O_n(c; \ell)$ if, for every $k \geq \ell$, for every object X of \mathcal{C}_Y such that the map $X \rightarrow Y$ to the final object is k -connected, the map $u_X: F(X) \rightarrow G(X)$ is $(-c + (n+1)k)$ -connected. We say that F and G agree to order n (via the map u) if this holds for some constants c and ℓ .

1.3 Remark The letter O stands for "osculation". This condition on (a map between) two functors is analogous to a condition on two functions (say real functions of one or several variables). The functions f and g can be said to agree to n^{th} order at y if there are constants C and K such that for every x such that $|x - y| \leq K$ we have $|f(x) - g(x)| \leq C|x - y|^{n+1}$.

The condition $E_n(c; \ell)$, stable n^{th} order excision, was defined in [10: Def. 4.1].

1.4 Proposition If F satisfies $E_n(c; \ell)$, then

- (1) $T_n F$ satisfies $E_n(c-1; \ell-1)$ and
- (2) $t_n F: F \rightarrow T_n F$ satisfies $O_n(c; \ell)$.

Proof The second conclusion is immediate from the definitions. For the first, note that the functor $X \mapsto X \times_Y U$ from \mathcal{C} to \mathcal{C} preserves strongly cocartesian cubes and (unless U is empty) increases the connectivity of maps. It follows immediately that for each nonempty U the functor $X \mapsto F(X \times_Y U)$ from \mathcal{C} to D satisfies $E_n(c - (n+1); \ell - 1)$. By [10; 1.20], $T_n F$ satisfies $E_n(c-1; \ell-1)$.

1.5 Proposition If F is stably n -excisive, then

- (1) $P_n F$ is n -excisive and
- (2) F agrees with $P_n F$ to order n (via $p_n F$).

Proof Suppose that F satisfies $E_n(c; \ell)$. By 1.4(1) and induction, $T_n^i F$ satisfies $E_n(c-i; \ell-i)$. This easily implies that $P_n F$ is n -excisive. We also find, by 1.4(2), that $t_n T_n^i F$ satisfies $O_n(c-i; \ell-i)$, and in particular $O_n(c; \ell)$, for all i . It follows easily that $O_n(c; \ell)$ is also satisfied by the composed maps $(t_n T_n^i F) \rightarrow t_n F$ and in the limit by $p_n F$. \square

1.6 Proposition Let $F \xrightarrow{u} G$ be a map between homotopy functors. If F and G agree to n^{th} order via u , then the induced map $P_n F \xrightarrow{P_n u} P_n G$ is an equivalence. The converse holds if F and G are stably n -excisive.

Proof Suppose u satisfies $O_n(c; \cdot)$. For each nonempty finite set U the resulting map of functors

$$F(- \times U) \rightarrow G(- \times U)$$

satisfies $O_n(c - (n + 1); \cdot - 1)$; again we have used the fact that the functor $X \mapsto X \times U$ from \mathcal{C} to \mathcal{C} preserves strongly cocartesian cubical diagrams and increases the connectivity of maps. Therefore, using [10; 1.20] as in the proof of 1.4, $T_n F \xrightarrow{T_n u} T_n G$ satisfies $O_n(c - 1; \cdot - 1)$. By induction, $T_n^i F \xrightarrow{T_n^i u} T_n^i G$ satisfies $O_n(c - i; \cdot - i)$. It follows, letting i tend to infinity, that $P_n F \xrightarrow{P_n u} P_n G$ is an equivalence. For the converse, use 1.5(2) and the commutative diagram

$$\begin{array}{ccc} F & \xrightarrow{u} & G \\ \wr & & \wr \\ \rho_n F & & \rho_n G \\ P_n F & \xrightarrow{P_n u} & P_n G \end{array} \quad \square$$

Note that when F is stably n -excisive $P_n F$ can be characterized, up to natural equivalence, as the only n -excisive functor that agrees to n^{th} order with F .

The remaining results of Section 1 (except for the last sentence of 1.13) have nothing to do with connectivity. In particular, functors are not assumed to satisfy any kind of stable excision hypothesis.

1.7 Proposition *Up to equivalence,*

- (1) T_n commutes with holim
- (2) P_n commutes with finite holim
- (3) T_n and P_n commute with homotopy fiber.
- (4) T_n and P_n commute with iterated (in particular sequential) hocolim.
- (5) for spectrum-valued functors, both T_n and P_n commute with any hocolim.

Proof and explanation The main point is that holim commutes with holim, hocolim commutes with hocolim, and, up to equivalence, finite holim commutes with iterated hocolim.

(1) This one is true up to isomorphism: If $f: F \rightarrow G$ is any sort of diagram of homotopy functors $\mathcal{C} \rightarrow \mathcal{D}$ and F is given by $F(X) = \text{holim}_{U \in \mathcal{C}(X)} F(X \times U)$ then we have

$$\begin{aligned} (T_n F)(X) &= \text{holim}_U \text{holim}_{U'} F(X \times U \times U') \\ &= \text{holim}_U \text{holim}_{U'} F(X \times U) \\ &= \text{holim}(T_n F)(X) \end{aligned}$$

(2) In the same situation we have a natural map

$$\begin{aligned} (P_n F)(X) &= \operatorname{hocolim}_i (T_n^i F)(X) \\ &= \operatorname{hocolim}_i \operatorname{holim} (T_n^i F)(X) \\ &\simeq \operatorname{holim}_i \operatorname{hocolim} (T_n^i F)(X) \\ &= \operatorname{holim} (P_n F)(X) \end{aligned}$$

It is an equivalence in the case when $F \rightarrow G$ is a finite diagram.

(3) This follows from (1) and (2). We have

$$\begin{aligned} P_n \operatorname{hocolim} (F \rightarrow G) &= P_n \operatorname{holim} (F \rightarrow G) \\ &= \operatorname{holim} (P_n F \rightarrow P_n G) \\ &= \operatorname{holim} (P_n F \rightarrow P_n G) \\ &= \operatorname{hocolim} (P_n F \rightarrow P_n G) \end{aligned}$$

and likewise for T_n .

(4) T_n is a finite holim and P_n is a hocolim of finite holim's.

(5) Since hocolims commute, hocolim preserves cocartesian cubes. Thus hocolim of spectra preserves cartesian cubes. This implies (using [10; 1.19]) that T_n commutes with hocolim up to equivalence. Therefore the same holds for $P_n = \operatorname{hocolim}_i T_n^i$. \square

Let $F(C; D)$ be the category whose objects are the homotopy functors from C to D , and whose morphisms are the natural maps. Let $hF(C; D)$ be its *homotopy category*; it has the same objects and is obtained by formally inverting the equivalences. Because the functors T_n and P_n from $F(C; D)$ to itself take equivalences to equivalences, they give rise to functors from the homotopy category to itself. Morphisms in $hF(C; D)$ are called *weak maps*. Any functor weakly isomorphic to an n -excisive functor is n -excisive.

1.8 Theorem *For any homotopy functor $C \xrightarrow{f} D$, the functor $P_n F$ is n -excisive. In the homotopy category $hF(C; D)$, $p_n F$ is the universal map from F to an n -excisive functor.*

The key to 1.8 is the following:

1.9 Lemma *Let X be any strongly cocartesian $(n+1)$ -cube in C , and let F be any homotopy functor. Then the map of cubes*

$$F(X) \xrightarrow{(p_n F)(X)} (P_n F)(X)$$

factors through some cartesian cube.

Proof of 1.8, assuming 1.9 Let X be any strongly cocartesian $(n + 1)$ -cube in \mathcal{C} , and consider the diagram of cubes

$$F(X) \dashrightarrow (T_n F)(X) \dashrightarrow (T_n^2 F)(X) \dashrightarrow \dots$$

which leads by hocolim to the cube $(P_n F)(X)$. By 1.9 each of the maps of cubes displayed above factors through some cartesian cube. Therefore the cube $(P_n F)(X)$ is equivalent to a sequential hocolim of cartesian cubes, and so it is itself cartesian. This shows that $P_n F$ is n -excisive.

For the existence half of the universal mapping property, let $F \dashrightarrow P$ be any weak map to an n -excisive functor. The diagram

$$\begin{array}{ccc} F & \xrightarrow{u} & P \\ \wr & & \wr \\ \rho_n F & \dashrightarrow & \rho_n P \end{array}$$

$$P_n F \xrightarrow{P_n u} P_n P$$

shows that u factors through $\rho_n F$, since $\rho_n P$ as a weak map is invertible.

For the uniqueness we must show that if P is n -excisive then a weak map $P_n F \dashrightarrow P$ is determined by the composition $v \circ \rho_n F$. It suffices if in the diagram of weak maps

$$\begin{array}{ccccc} F & \xrightarrow{\rho_n F} & P_n F & \xrightarrow{v} & P \\ \wr & & \wr & & \wr \\ \rho_n F & \dashrightarrow & \rho_n P_n F & \dashrightarrow & \rho_n P \end{array}$$

$$P_n F \xrightarrow{P_n(\rho_n F)} P_n P_n F \xrightarrow{P_n v} P_n P$$

the marked (\wr) arrows are invertible, for then v is determined by $P_n v$, which is determined by $P_n v \circ P_n(\rho_n F) = P_n(v \circ \rho_n F)$, which is of course determined by $v \circ \rho_n F$.

The marked vertical arrows are equivalences because $P_n F$ and P are n -excisive. In order for $P_n(\rho_n F)$ to be an equivalence, it will be enough (by 1.7(4)) if $P_n(t_n F)$ is an equivalence. Define the functor $J_U F$ by $(J_U F)(X) = F(X \vee U)$. In the diagram

$$P_n F \xrightarrow{P_n(t_n F)} P_n T_n F = P_n(\operatorname{holim}_{U \in \mathcal{P}_0(n+1)} J_U F)$$

$$\dashrightarrow \operatorname{holim}_{U \in \mathcal{P}_0(n+1)} P_n J_U F = \operatorname{holim}_{U \in \mathcal{P}_0(n+1)} J_U P_n F$$

the second arrow is an equivalence by 1.7(2). The composition is also an equivalence; this simply means that for every X the functor $P_n F$ takes the cube $fX \vee Ug$ to a cartesian cube, and it is true because $P_n F$ is n -excisive. It follows that $P_n(t_n F)$ is an equivalence. \square

The fact that $P_n(p_n F) = p_n P_n F$ in the homotopy category follows from 1.8; if we had had it before 1.8, then we could have skipped the last part of the proof of 1.8.

1.10 Remark Theorem 1.8 is one of several statements in this paper for which we have two different proofs: an older one which requires some kind of stable excision hypothesis, and a newer one which has nothing to do with connectivity. Another such statement is Theorem 2.1, which also depends (near the end of the proof of 2.2) on 1.9. Another is 3.1, and another is 6.1. The older proofs have the advantage of a certain common-sense quality, and if what we really care about here is convergent Taylor towers then the older proofs are good enough; but the fact that the theorems are still true without connectivity hypotheses is striking and the newer proofs are perhaps worth looking at, too.

Proof of 1.9 (Any reader who, like the author, finds this proof a little opaque, may wish to look at it again after reading the proof of 3.2, which is related but simpler.) Let us write n instead of $n+1$. For subsets $T; U_1; \dots; U_n$ of \underline{n} , define $\hat{X}(T; U_1; \dots; U_n)$ to be

$$\text{hocolim}_{1 \leq s \leq n} (X(T) \cup_{U_s}) \neq \text{hocolim}_{1 \leq s \leq n} (X(T \setminus \{s\}) \cup_{U_s});$$

This can also be described as the union, along $X(T)$, of the spaces $X(T) \cup_{X(T \setminus \{s\})} U_s$.

Clearly \hat{X} is a functor from $P(\underline{n}) \times P(\underline{n})^n$ to \mathcal{C} . We have $X(T) = \hat{X}(T; \dots; \dots)$, and there is a natural map $\hat{X}(T; U; \dots; U) \rightarrow X(T) \cup U$ that corresponds to the identity map when $U = \emptyset$.

Let E be the set of all $(U_1; \dots; U_n) \in P_0(\underline{n})^n$ such that, for at least one $s \in \underline{n}$, $s \in U_s$. Since E contains the image of the diagonal map $P_0(\underline{n}) \rightarrow P_0(\underline{n})^n$, we can factor the map $t_{n-1} F(X(T))$ as follows:

$$\begin{aligned} F(X(T)) &\rightarrow \text{holim}_{(U_1; \dots; U_n) \in E} F(\hat{X}(T; U_1; \dots; U_n)) \\ &\rightarrow \text{holim}_{U \in P_0(\underline{n})} F(\hat{X}(T; U; \dots; U)) \\ &\rightarrow \text{holim}_{U \in P_0(\underline{n})} F((X(T) \cup U)) \\ &= (T_{n-1} F)(X(T)); \end{aligned}$$

We claim that the cube

$$T \rightarrow \text{holim}_{(U_1; \dots; U_n) \in E} F(\hat{X}(T; U_1; \dots; U_n))$$

is cartesian (for any homotopy functor F).

Let E be the set of all $(U_1; \dots; U_n) \in P_0(\mathbb{R})^n$ such that, for some s , $U_s = fs_g$. This poset is left cofinal in E (exercise; see [10; page 298] for the definition), and it follows that the restriction map from the holim over E to the holim over E is an equivalence. Therefore it is enough if, whenever $(U_1; \dots; U_n) \in E$, the cube

$$T \not\rightarrow F(\hat{X}(T; U_1; \dots; U_n))$$

is cartesian. In fact, it is cartesian for a very basic reason: if $U_{s_0} = fs_{0g}$ then the map

$$\hat{X}(T; U_1; \dots; U_n) \rightarrow \hat{X}(T [fs_{0g}; U_1; \dots; U_n])$$

is an equivalence. To see this, examine the diagram

$$\begin{array}{ccccc} X(T) & \longrightarrow & {}_s(X(T) \quad U_s) & \longrightarrow & {}_s(X(T [fs_g] \quad U_s)) \\ \downarrow \cong & & \downarrow \cong & & \downarrow \cong \\ X(T [fs_{0g}]) & \longrightarrow & {}_s(X(T [fs_{0g}] \quad U_s)) & \longrightarrow & {}_s(X(T [fs; s_{0g}] \quad U_s)) \end{array}$$

The fact that the induced map from hocolim of top row to hocolim of bottom row is an equivalence follows from the trivial fact that for $s \neq s_0$ the square

$$\begin{array}{ccc} X(T) \quad U_s & \longrightarrow & X(T [fs_g] \quad U_s \\ \downarrow \cong & & \downarrow \cong \\ X(T [fs_{0g}] \quad U_s) & \longrightarrow & X(T [fs; s_{0g}] \quad U_s \end{array}$$

is cocartesian, plus the even more trivial fact that the square

$$\begin{array}{ccc} X(T) & \longrightarrow & X(T) \quad U_{s_0} \\ \downarrow \cong & & \downarrow \cong \\ X(T [fs_{0g}]) & \longrightarrow & X(T [fs_{0g}] \quad U_{s_0} \end{array}$$

is cocartesian. □

1.11 Corollary *If $0 \leq m \leq n$ then the map*

$$P_m F \xrightarrow{P_m(P_n F)} P_m P_n F$$

is an equivalence.

Proof This is formal; using the universal mapping properties of P_m and P_n and the fact that m -excisive functors are n -excisive, one sees that $P_m P_n F$ has the universal mapping property that characterizes $P_m F$. □

We now collect the "Taylor polynomials" $P_n F$ into a "Taylor series" by showing that there is a natural map $P_n F \xrightarrow{q_n} P_{n-1} F$ satisfying $q_n F \circ \rho_n F = \rho_{n-1} F$.

An effortless way to produce $q_n F$ as a weak map would be to reason as in the proof of 1.11: $\rho_n F$ is the universal map from F to an n -excisive functor and $(n - 1)$ -excisive functors are n -excisive, so there is a unique map $q_n F$ in $hF(C; D)$ such that $q_n F \circ \rho_n F = \rho_{n-1} F$. On the other hand, it is desirable to define $q_n F$ in such a way that $q_n F \circ \rho_n F = \rho_{n-1} F$ on the nose and not just weakly. Moreover, the explicit construction for $q_n F$ will be useful in its own right in proving Lemma 2.2.

We will make a (commutative) diagram

$$(1.12) \quad \begin{array}{ccccccc} F & \xrightarrow{t_n F} & T_n F & \xrightarrow{t_n T_n F} & T_n^2 F & \xrightarrow{t_n T_n^2 F} & \dots \\ \downarrow \text{?} & & \downarrow \text{?} & & \downarrow \text{?} & & \\ \mathcal{Y} & & q_{n,1} \mathcal{Y} & & q_{n,2} \mathcal{Y} & & \\ & & & & & & \\ F & \xrightarrow{t_{n-1} F} & T_{n-1} F & \xrightarrow{t_{n-1} T_{n-1} F} & T_{n-1}^2 F & \xrightarrow{t_{n-1} T_{n-1}^2 F} & \dots \end{array}$$

and then define $q_n F$ as the induced map of horizontal homotopy colimits. We must define $q_{n,i}$ and then verify that the squares commute.

Notice that $T_n^i F$ is naturally isomorphic to a homotopy limit indexed by a product of i copies of the partially ordered set $P_0(n + 1)$:

$$(T_n^i F)(X) = \operatorname{holim}_{(U_1, \dots, U_i) \in P_0(n+1)^i} F(X \times (U_1 \times \dots \times U_i))$$

From this point of view there is an obvious map $T_n^i F \xrightarrow{q_n} T_{n-1}^i F$, induced by the inclusion of $P_0(n)^i$ in $P_0(n + 1)^i$. The first square in (1.12) now obviously commutes. The $(i + 1)^{\text{st}}$ square will commute if both squares commute in

$$\begin{array}{ccc} T_n^i F & \xrightarrow{t_n T_n^i F} & T_n^{i+1} F \\ \downarrow \text{?} & & \downarrow \text{?} \\ q_{n,i} F \mathcal{Y} & & T_n q_{n,i} F \mathcal{Y} \end{array} \quad \begin{array}{ccc} & \xrightarrow{\quad} & T_n^{i+1} F \\ & & \downarrow \text{?} \\ & & q_{n,i+1} F \mathcal{Y} \end{array}$$

$$\begin{array}{ccc} T_{n-1}^i F & \xrightarrow{t_n T_{n-1}^i F} & T_n T_{n-1}^i F \\ \downarrow \text{?} & & \downarrow \text{?} \\ q_{n-1,i} F \mathcal{Y} & & q_{n,1} T_{n-1}^i F \end{array} \quad \begin{array}{ccc} & \xrightarrow{\quad} & T_{n-1}^{i+1} F \\ & & \downarrow \text{?} \\ & & q_{n-1,i+1} F \mathcal{Y} \end{array}$$

The left square commutes because $t_n F$ is natural in F . The other commutes because it is induced by a commutative diagram of posets

$$\begin{array}{ccc} P_0(n+1)^{i+1} & \xrightarrow{\quad} & P_0(n+1)^{i+1} \\ \downarrow \text{?} & & \downarrow \text{?} \\ P_0(n+1) & \xrightarrow{\quad} & P_0(n)^{i+1} \end{array}$$

Summing up, we have:

1.13 Theorem A homotopy functor F from spaces over Y (with or without section) to either spaces, based spaces, or spectra, determines a tower of such functors

$$\begin{array}{c}
 \vdots \\
 \wr \\
 Y \\
 \\
 P_n F \\
 \wr \\
 Y_{q_n} F \\
 \\
 P_{n-1} F \\
 \wr \\
 Y_{q_{n-1}} F \\
 \\
 \vdots \\
 \wr \\
 Y_{q_2} F \\
 \\
 P_1 F \\
 \wr \\
 Y_{q_1} F \\
 \\
 P_0 F
 \end{array}$$

and a map $F \xrightarrow{f_{p_n F}} \lim_n P_n F$. Each P_n is a homotopy functor (from homotopy functors to homotopy functors), and p_n and q_n are natural. The functor $P_n F$ is always n -excisive, and (in the homotopy category of homotopy functors) $p_n F$ is universal among maps from F to n -excisive functors. If F is q -analytic and the structural map $X \rightarrow Y$ is $(q + 1)$ -connected, then the connectivity of the map $F(X) \xrightarrow{p_n F} (P_n F)(X)$ tends to $+1$ with n , so that $F(X)$ is equivalent to the homotopy limit $(P_1 F)(X \rightarrow Y)$ of the tower.

Proof The last statement is the only new one. Recall from [10; 4.2] that q -analyticity means that there is a number q such that, for all $n \geq 0$, F satisfies $E_n(n - q; +1)$. If $X \rightarrow Y$ is such that the map $X \rightarrow Y$ is k -connected with $k > q$ then by the proof of 1.5 the connectivity of the map $F(X) \xrightarrow{p_n F} (P_n F)(X)$ is at least $(q + k + n(k - q))$, which tends to $+1$ with n . □

If $P_n F$ is analogous to an n^{th} Taylor polynomial, then the homotopy fiber

$$D_n F = \text{ho fiber}(P_n F \xrightarrow{q_n F} P_{n-1} F)$$

is analogous to the n^{th} term in a Taylor series. Notice that the definition of $D_n F$ is meaningful if the category D is either based spaces or spectra, but not if it is unbased spaces. From $P_n(F \dashv \gamma) = (P_n F) \dashv \gamma$ (Remark 1.1) we have

$$(1.14) \quad D_n(F \dashv \gamma) = (D_n F) \dashv \gamma:$$

1.15 Definition A homotopy functor $F : C \rightarrow D$ is called n -reduced if $P_{n-1} F = *$. It is called n -homogeneous, or homogeneous of degree n , if it is both n -excisive and n -reduced.

Thus 1-reduced means reduced, and 1-homogeneous means linear.

1.16 Remark If $n > 1$ then it is not easy in general to test whether a functor F is n -reduced. Perhaps the main difficulty is that $P_{n-1} F = *$ does not imply $T_{n-1} F = *$. On the other hand, by 1.6 a sufficient condition for F to be n -reduced is that F agrees to order $n - 1$ with the constant functor $*$, in other words that the connectivity of $F(X)$ tends to infinity at least n times faster than the connectivity of the map $X \rightarrow Y$. (If F is analytic, or even just stably n -excisive, then this condition is also necessary for F to be n -reduced.) Thus, for example, for any spectrum C that is bounded below (k -connected for some k) the n -excisive functors $X \mapsto C \wedge X^{\wedge n}$ from based spaces to spectra and $X \mapsto \Omega^{-1}(C \wedge X^{\wedge n})$ from based spaces to based spaces are homogeneous. In fact this holds for all spectra C , either by expressing C as a homotopy colimit of bounded-below spectra or by 3.1 below.

1.17 Proposition $D_n F$ is always n -homogeneous.

Proof The functor $D_n F = \text{holim}(P_n F \dashv P_{n-1} F \dashv \dots)$ is n -excisive because it is a homotopy limit of n -excisive functors. To see that it is also n -reduced, use 1.7(4) to identify $P_{n-1} D_n F$ with the homotopy fiber of the map

$$P_{n-1}(P_n F) \xrightarrow{P_{n-1}(q_n F)} P_{n-1}(P_{n-1} F);$$

by (1.11) this map is an equivalence. \square

1.18 Proposition Up to equivalence,

- (1) D_n commutes with finite holim.
- (2) D_n commutes with homotopy fiber.
- (3) D_n commutes with iterated hocolim.
- (4) for spectrum-valued functors, D_n commutes with arbitrary hocolim.

Proof This follows easily from 1.7. □

1.19 Example Let the functors F_a and F_b be a -homogeneous and b -homogeneous respectively, with $a < b$, and let $F_a \xrightarrow{f} F_b$ be any natural map. A simple example is the diagonal inclusion $QX \hookrightarrow Q(X \wedge X)$. Then the functor

$$F(X) = \text{ho } \text{ber}(F_a \xrightarrow{f} F_b)$$

is b -excisive. Its tower has only two nontrivial layers $D_a F \simeq F_a$ and $D_b F \simeq F_b$; we have

$$\begin{aligned} P_n F &\simeq F_a & n < a \\ P_n F &\simeq F & a \leq n < b \\ P_n F &\simeq F & n \geq b \end{aligned}$$

All of this follows from $P_n F \simeq \text{ho } \text{ber}(P_n F_a \xrightarrow{f} P_n F_b)$. Note that when $a > b$ there can be no interesting natural map $F_a \rightarrow F_b$, since by (1.8) any such map factors (in $hF(\mathcal{C}; D)$) through $P_{a-1} F_a$.

1.20 Example There is a weak equivalence ("Snaith splitting")

$$(1.21) \quad \mathbb{1} \rightarrow X \rightarrow \prod_{n=1}^{\infty} X^{\wedge n}$$

for based connected spaces X . The functor $\mathbb{1} \rightarrow X^{\wedge n}$ of X is n -homogeneous. The m^{th} Taylor polynomial of the right-hand side of 1.21 is $\mathbb{1} \rightarrow \prod_{n=1}^m X^{\wedge n}$, by 1.6, since this finite product agrees with the infinite product to order m . The same therefore holds for the left-hand side, by 1.1. In particular the n^{th} homogeneous layer of $\mathbb{1} \rightarrow X$ is $\mathbb{1} \rightarrow X^{\wedge n}$. We will find later that this is enough to determine the n^{th} homogeneous layer of $F(X) = \mathbb{1} \rightarrow X$. In fact, using 1.14 we have

$$(D_n F)(X) \simeq (D_n(\mathbb{1} \rightarrow \prod_{n=1}^{\infty} X^{\wedge n}))(X) \simeq \mathbb{1} \rightarrow X^{\wedge n} \simeq (X^{\wedge n})^{\wedge n};$$

and by 3.8 this will imply

$$(D_n F)(X) \simeq \mathbb{1} \rightarrow X^{\wedge n};$$

Incidentally, the naturality of Taylor towers gives a quick way to get from the James model of $\mathbb{1} \rightarrow X$ to that splitting of $\mathbb{1} \rightarrow X$. The space JX , free monoid on the based space X , is naturally filtered by word length as an increasing union $J_0 X \hookrightarrow J_1 X \hookrightarrow \dots$ in such a way that the subquotient $J_n = J_{n-1} \rightarrow J_n$ is $X^{\wedge n}$. It follows that for each n there is a natural filtration sequence

$$\mathbb{1} \rightarrow J_{n-1} X \hookrightarrow \mathbb{1} \rightarrow J_n X \hookrightarrow \mathbb{1} \rightarrow X^{\wedge n}$$

One sees by induction that $\tau_{\leq n} J_n$ is n -excisive, since it fibers over a (homogeneous) n -excisive functor and the fiber is $((n - 1)$ -excisive, hence) n -excisive. The sequence above must split. Indeed, any natural filtration sequence

$$F(X) \rightarrow G(X) \rightarrow H(X)$$

of spectra in which H is n -homogeneous and F is $(n - 1)$ -excisive must split: a retraction from G to F (in the homotopy category of homotopy functors) is given by the diagram

$$\begin{array}{ccc} F & \xrightarrow{\simeq} & G \\ \wr \downarrow & & \wr \downarrow \\ P_{n-1}F & \xrightarrow{\simeq} & P_{n-1}G \end{array}$$

The left arrow is an equivalence because F is $(n - 1)$ -excisive; the lower arrow because $P_{n-1}H$ is contractible.

1.22 Remark The Taylor tower construction extends easily to functors

$$C_{Y_1} \rightarrow C_{Y_n} \xrightarrow{\simeq} D$$

of several variables. Let us say that F is $(d_1; \dots; d_n)$ -excisive [resp. $(d_1; \dots; d_n)$ -reduced] if for $1 \leq j \leq n$ it is d_j -excisive [resp. d_j -reduced] as a functor of the j^{th} variable. There is an n -variable Taylor polynomial construction which will be denoted $P_{d_1, \dots, d_n} F$, and which gives the universal $(d_1; \dots; d_n)$ -excisive functor under F (in the homotopy category of homotopy functors). It may be defined either as $P_{d_1}^{(1)} \dots P_{d_n}^{(n)} F$ where $P_d^{(j)}$ is P_d with respect to the j^{th} variable, or directly as the homotopy colimit of $(T_{d_1, \dots, d_n})^k F$ where

$$(T_{d_1, \dots, d_n} F)(X_1; \dots; X_n) = \text{hocolim}_{(U_1; \dots; U_n) \in \mathcal{P}_0(d_j+1)} F(X_1 \times_{Y_1} U_1; \dots; X_n \times_{Y_n} U_n);$$

these are naturally equivalent. A functor of n variables will be called multilinear if it is linear in each variable, in other words both $(1; \dots; 1)$ -excisive and $(1; \dots; 1)$ -reduced. If F is $(1; \dots; 1)$ -reduced then the functor $P_{1, \dots, 1} F$ is multilinear and may be called the multilinearization of F . Loosely, this is the homotopy colimit of

$$k_1 + \dots + k_n F \left(\prod_{Y_1}^{k_1} X_1; \dots; \prod_{Y_n}^{k_n} X_n \right)$$

as $(k_1; \dots; k_n) \neq (1; \dots; 1)$.

2 Delooping homogeneous functors

We will show that all homogeneous space-valued functors arise from homogeneous spectrum-valued functors. The main step is to show that they have natural deloopings.

Let $H_n(\mathcal{C}; D)$ be the category of homogeneous functors of degree n from \mathcal{C} to D , a full subcategory of $F(\mathcal{C}; D)$. The homotopy category $hH_n(\mathcal{C}; D)$ is obtained from $H_n(\mathcal{C}; D)$ by formally inverting the (objectwise) equivalences.

The functor $S\rho \xrightarrow{1} T$ preserves both weak equivalences and cartesian cubes, and therefore composing with it yields a functor

$$H_n(\mathcal{C}; S\rho) \xrightarrow{1} H_n(\mathcal{C}; T)$$

which itself takes weak equivalences to weak equivalences.

2.1 Theorem *The functor $H_n(\mathcal{C}; S\rho) \xrightarrow{1} H_n(\mathcal{C}; T)$ has an inverse up to weak equivalence.*

Proof The key is to get a functor $F \dashv BF$ from $H_n(\mathcal{C}; T)$ to itself such that BF is naturally equivalent to F . This will be given by the next lemma.

Assume for now that we have this. Then any object F of $H_n(\mathcal{C}; T)$ yields a sequence $fB^p Fg$ of such objects related by equivalences $B^p F \simeq B^{p+1} F$. Call the resulting spectrum-valued functor $B^{-1} F$. The fact that $B^{-1} F$ is an object of $H_n(\mathcal{C}; S\rho)$ follows easily from the fact that each $B^p F$ is an object of $H_n(\mathcal{C}; T)$. The fact that the functor B^{-1} takes equivalences to equivalences follows from the fact that each functor B^p does so. Clearly $B^{-1} B^{-1}$ is naturally equivalent to the identity. To check that this is also true for $B^{-1} B^{-1}$, let F be an object of $H_n(\mathcal{C}; S\rho)$ and write $F(X) = fF_q(X)g$. The bispectrum $fB^p F_q(X)g$ shows that the two spectra

$$fB^p F_0(X)g = fB^{p-1} F(X)g = B^{-1} B^{-1} F(X)$$

and

$$fB^0 F_q(X)g = fF_q(X)g = F(X)$$

are naturally equivalent. □

The next result provides the desired delooping of F , namely $R_n F$. As usual \mathcal{C}_Y is either U_Y or T_Y .

2.2 Lemma *Let $n > 0$. If $F : \mathcal{C}_Y \rightarrow \mathcal{T}$ is any reduced ($F(Y) = 0$) homotopy functor, then up to natural equivalence there is a fibration sequence*

$$P_n F \xrightarrow{q_n F} P_{n-1} F \longrightarrow R_n F$$

in which the functor $R_n F$ is n -homogeneous.

Proof More precisely, we will obtain a natural diagram of homotopy functors

$$(2.3) \quad \begin{array}{ccc} P_n F & \xrightarrow{q_n F} & P_{n-1} F \\ \downarrow \simeq & & \downarrow \simeq \\ \downarrow ? & & \downarrow ? \\ P_n F & \longrightarrow & P_{n-1} F \\ \downarrow \simeq & & \downarrow \simeq \\ \downarrow \gamma & & \downarrow \gamma \\ K_n F & \longrightarrow & R_n F \end{array}$$

in which the marked arrows are equivalences, the lower square is cartesian, $R_n F$ is n -homogeneous, and $K_n F$ is contractible. (If F were not reduced, then in fact $K_n F$ would be equivalent to the constant functor $F(Y)$.)

The proof is based on a close examination of the maps $q_{n,i} F$ which were used in defining $q_n F$. The first step is to define, for each $i \geq 0$, a diagram

$$(2.4(i)) \quad \begin{array}{ccc} T_n^i F & \longrightarrow & S_{n-1}^i F \\ \downarrow \simeq & & \downarrow \simeq \\ \downarrow ? & & \downarrow ? \\ \downarrow \gamma & & \downarrow \gamma \\ K_{n,i} F & \longrightarrow & R_{n,i} F \end{array}$$

Roughly, this will become the lower square of (2.3) when i goes to infinity.

Define posets

$$B_n = P_0(\underline{n+1}) - \text{ffn} + 1g \\ A_{n,i} = P_0(\underline{n+1})^i - P_0(\underline{n})^i:$$

Define the functor $S_{n-1} F$ by

$$(S_{n-1} F)(X) = \text{holim}_{U \in 2B_n} F(X \times_Y U)$$

Note that the inclusions $P_0(\underline{n+1}) \rightarrow B_n \rightarrow P_0(\underline{n})$ induce maps

$$(2.5) \quad T_n F \longrightarrow S_{n-1} F \longrightarrow T_{n-1} F$$

whose composition is $q_{n,1}F$. The second map is an equivalence because $P_0(\underline{n})$ is left co-nal in B_n . Now let (2.4(i)) be obtained from the posets and inclusions:

$$(2.6(i)) \quad \begin{array}{ccc} P_0(\underline{n+1})^i & \longrightarrow & B_n^i \\ \downarrow \wr & & \downarrow \wr \\ A_{n,i} & \longrightarrow & A_{n,i} \setminus B_n^i \end{array}$$

by forming the homotopy limit of $(U_1; \dots; U_i) \not\rightarrow F(X \vee (U_1 \dots U_i))$ over each poset.

Diagram (2.4(i)) is cartesian by [[10]; 1.9], since $A_{n,i}$ and B_n^i are concave and their union is $P_0(\underline{n+1})^i$.

2.7 Claim $K_{n,i}F$.

Proof Compare $(K_{n,i}F)(X)$, a holim over $A_{n,i}$, with the corresponding holim over the smaller poset

$$A_{n,i} = P_0(\underline{n+1})^i - B_n^i$$

On the one hand, the comparison map is an equivalence, because $A_{n,i}$ is left co-nal in $A_{n,i}$. On the other hand, the holim of $F(X \vee (U_1 \dots U_i))$ over $A_{n,i}$ is contractible: for each $(U_1; \dots; U_i) \in A_{n,i}$, we have $\sum |U_j| = 1$ for some j , so that $U_1 \dots U_i$ is contractible and

$$F(X \vee (U_1 \dots U_i)) \simeq F(Y) \quad \square$$

2.8 Claim $R_{n,i}F$ is n -reduced.

Proof There is a natural equivalence

$$P_{n-1}R_{n,i}F \simeq R_{n,i}P_{n-1}F$$

by 1.7, so it will be enough if $R_{n,i}F$ is contractible whenever F is $(n-1)$ -excisive (and reduced).

There is an isomorphism of posets

$$A_{n,i} \setminus B_n^i = P_0(\underline{n})^i \simeq P_0(\underline{i})$$

given by

$$\begin{aligned} (U_1; \dots; U_i) &\not\rightarrow (V_1; \dots; V_i; W) \\ V_i &= U_i - f_{n+1}g \\ W &= f_i : n+1 \geq U_i g \end{aligned}$$

Therefore $(R_{n,i}F)(X)$ can be written as

$$\text{holim}_{(V_1, \dots, V_i; W) \in 2P_0(\underline{n})^i \times P_0(\underline{j})} F(X \times_{\gamma} (U_1 \times \dots \times U_i))$$

Analyze this as an iterated homotopy limit: First $\times (V_2; \dots; V_i; W)$ and take homotopy limit with respect to V_1 . Since F is $(n - 1)$ -excisive, this yields, up to equivalence, $F(X \times_{\gamma} (e_1 \times U_2 \times \dots \times U_i))$ where e_1 is $n + 1$ g or γ ; according as 1 is or is not in W . Next take the homotopy limit with respect to V_2 , then V_3 , and so on through V_i , obtaining

$$(R_{n,i}F)(X) \cong \text{holim}_{W \in 2P_0(\underline{j})} F(X \times_{\gamma} (e_1 \times \dots \times e_i))$$

This is contractible because for each $W \in 2P_0(\underline{j})$ the space $e_1 \times \dots \times e_i$ is contractible. (It is a simplex of dimension $jWj - 1 \geq 0$.) □

The next step ought to be to take the homotopy colimit of (2.4(i)) as i tends to ∞ . This is not possible, since we do not have natural maps $K_{n,i}F \rightarrow K_{n,i+1}F$ or $R_{n,i}F \rightarrow R_{n,i+1}F$. We do, however, have maps defined up to homotopy, and with a little care these will suffice.

Here are two variations on (2.6(i+1)):

$$(2.9(i+1)) \quad \begin{array}{ccc} P_0(\underline{n+1})^{i+1} & \longrightarrow & P_0(\underline{n+1}) \times B_n^i \\ \downarrow \times & & \downarrow \times \\ \downarrow ? & & \downarrow ? \end{array}$$

$$(2.10(i+1)) \quad \begin{array}{ccc} P_0(\underline{n+1}) \times A_{n,i} & \longrightarrow & P_0(\underline{n+1}) \times (A_{n,i} \setminus B_n^i) \\ \downarrow \times & & \downarrow \times \\ \downarrow ? & & \downarrow ? \end{array}$$

$$A_{n,i+1} \longrightarrow A_{n,i+1} \setminus (P_0(\underline{n+1}) \times B_n^i)$$

We have maps of square diagrams of posets

$$(2.9(i+1)) \rightarrow (2.10(i+1)) \rightarrow (2.6(i+1)):$$

From each of the three diagrams we get a square diagram of functors by taking the holim of $F(X \times_{\gamma} (U_1 \times \dots \times U_i))$. From (2.6(i+1)) we get (2.4(i+1)). From (2.9(i+1)) we get what might be called $T_n(2.4(i))$. From (2.10(i+1)) we get something new; call it (2.11(i+1)). These are related by maps (of square diagrams)

$$\dots \rightarrow 2.4(i) \rightarrow T_n(2.4(i)) \rightarrow 2.11(i+1) \rightarrow 2.4(i+1) \rightarrow \dots$$

Now consider the (pointwise) homotopy colimit of this, another square diagram. This will be the lower half of (2.3). In view of the following, it is essentially a limit of the cartesian squares $2.4(i)$ and therefore it is itself cartesian:

2.12 Claim *The backwards arrow $T_n(2:4(i)) \rightarrow 2:11(i+1)$ is an equivalence (in all four corners of the square).*

Proof In the upper corners it is an isomorphism. In the lower corners it is induced by the inclusions

$$P_0(\underline{n+1}) \hookrightarrow A_{n;i} \xrightarrow{\quad} A_{n;i+1}$$

$$P_0(\underline{n+1}) \hookrightarrow (A_{n;i} \setminus B_n^i) \xrightarrow{\quad} A_{n;i+1} \setminus (P_0(\underline{n+1}) \setminus B_n^i)$$

In each of the two cases the larger poset is the union of the smaller one, which is concave (in the sense of [10; page 298]), with the concave set

$$Q = A_{n;1} \setminus B_n^i$$

and in each case the intersection is

$$Q^0 = A_{n;1} \setminus (A_{n;i} \setminus B_n^i)$$

Thus by [10; 0.2] it will be enough if the inclusion $Q^0 \hookrightarrow Q$ induces an equivalence of holims. But each of these holims is contractible. (The argument is as in the proof of 2.7; replace Q by $A_{n;1} \setminus B_n^i$ and Q^0 by $A_{n;1} \setminus (A_{n;i} \setminus B_n^i)$.) \square

It follows from 2.7 and 2.12 that $K_n F$ is contractible; it is the hocolim of a diagram

$$\cdots \hookrightarrow K_{n;i} F \xrightarrow{t_n} T_n K_{n;i} F \xrightarrow{\quad} ? \xrightarrow{\quad} K_{n;i+1} F \xrightarrow{t_n} \cdots$$

of contractible objects.

It follows from 2.8 and 2.12 that $R_n F$ is n -reduced, being the hocolim of the functors

$$\cdots \hookrightarrow R_{n;i} F \xrightarrow{t_n} T_n R_{n;i} F \xrightarrow{\quad} ? \xrightarrow{\quad} R_{n;i+1} F \xrightarrow{t_n} \cdots$$

Moreover, $R_n F$ is n -excisive, by Lemma 1.9; if X is any strongly cocartesian $(n+1)$ -cube in \mathcal{C} then $(R_n F)(X)$ is cartesian because for each i the map

$$(R_{n;i} F)(X) \xrightarrow{t_n} (T_n R_{n;i} F)(X)$$

factors through a cartesian cube.

Finally we construct the upper half of (2.3). Note that $\hat{P}_n F$ is the hocolim of

$$\cdots \hookrightarrow T_n^i F \xrightarrow{t_n} T_n^{i+1} F \xrightarrow{=} T_n^{i+1} F \xrightarrow{=} T_n^{i+1} F \xrightarrow{t_n} \cdots$$

Eliminating the identity maps we obtain an equivalence from $\hat{P}_n F$ to the hocolim of

$$\cdots \hookrightarrow T_n^i F \xrightarrow{t_n} T_n^{i+1} F \xrightarrow{t_n} \cdots$$

which is $P_n F$. This is the upper left arrow in (2.3). Likewise $P_{n-1} F$ is the hocolim of

$$\cdots \rightarrow S_{n-1}^i F \xrightarrow{t_n} T_n S_{n-1}^i F \xrightarrow{=} T_n S_{n-1}^i F \rightarrow S_{n-1}^{i+1} F \xrightarrow{t_n} \cdots$$

and so has an equivalence to the hocolim of the upper row in

$$\begin{array}{ccccccc} \cdots & \longrightarrow & S_{n-1}^i F & \xrightarrow{t_n} & T_n S_{n-1}^i F & \longrightarrow & S_{n-1}^{i+1} F \longrightarrow \cdots \\ & & \downarrow \text{?} & & \downarrow \text{?} & & \downarrow \text{?} \\ \cdots & \longrightarrow & T_{n-1}^i F & \xrightarrow{t_n} & T_n T_{n-1}^i F & \xlongequal{\quad} & T_{n-1}^{i+1} F \longrightarrow \cdots \end{array}$$

This in turn has an equivalence (2.7) to the hocolim of the lower row, and hence to $P_{n-1} F$. The composed map is the upper right arrow in (2.3). The square commutes. This concludes the proof of (2.2). \square

3 Symmetric multilinear functors

Let $C \rightarrow C^n$ be the diagonal functor. It was shown in [10; 3.4] that the composed functor $F \circ C$ is $(d_1 + \dots + d_n)$ -excisive if the functor $C^n \rightarrow D$ is d_j -excisive in the j^{th} variable, or in the language of 1.22 $(d_1; \dots; d_n)$ -excisive. (The latter is easier to write but harder to pronounce.) In particular $F \circ C$ is n -excisive if F is $(1; \dots; 1)$ -excisive.

3.1 Lemma *If $C^n \rightarrow D$ is a $(1; \dots; 1)$ -reduced homotopy functor, then $F \circ C$ is n -reduced. Thus L is n -homogeneous if $C^n \rightarrow D$ is multilinear.*

The proof resembles that of 1.8. The key is:

3.2 Lemma *If $C^n \rightarrow D$ is a $(1; \dots; 1)$ -reduced homotopy functor, then for any $X \in C$ the map*

$$(F \circ C)(X) \xrightarrow{t_{n-1}(F \circ C)} T_{n-1}(F \circ C)(X)$$

factors through a (weakly) contractible object.

Proof of 3.1, assuming 3.2 Let $X \in C$. The object $P_{n-1}(F \circ C)(X)$ is defined as the sequential hocolim of a diagram whose $(i + 1)^{st}$ map is

$$T_{n-1}^i(F \circ C)(X) \xrightarrow{t_{n-1} T_{n-1}^i(F \circ C)} T_{n-1}^{i+1}(F \circ C)(X)$$

It is enough if each of these maps factors through a weakly contractible object. Lemma 3.2 takes care of this, not only for the first map but also for the others, since the functor $T_{n-1}(F \circ C)$ is equal to G for a functor G that satisfies the same hypotheses as F . \square

Proof of 3.2 Let E and E' be as in the proof of 1.9. Since E contains the image of the diagonal map $P_0(\underline{n}) \rightarrow P_0(\underline{n})^n$, the map $t_{n-1}(F \rightarrow G)(X)$ can be factored

$$\begin{aligned} F(X_1, \dots, X_n) &= F(X_1 \times \dots \times X_n) \\ &\rightarrow \operatorname{holim}_{(U_1, \dots, U_n) \in E} F(X_1 \times U_1, \dots, X_n \times U_n) \\ &\rightarrow \operatorname{holim}_{U \in P_0(\underline{n})} F(X_1 \times U, \dots, X_n \times U) \end{aligned}$$

But

$$\operatorname{holim}_{(U_1, \dots, U_n) \in E} F(X_1 \times U_1, \dots, X_n \times U_n) \rightarrow \operatorname{holim}_{(U_1, \dots, U_n) \in E} F(X_1 \times U_1, \dots, X_n \times U_n)$$

and this is a holim of weakly contractible objects: if $(U_1, \dots, U_n) \in E$ then some U_s is a one-element set, giving $X_s \times U_s \rightarrow Y$ and

$$F(X_1 \times U_1, \dots, X_n \times U_n) \rightarrow F(Y) \quad \square$$

The functor $L : \mathcal{C}^n \rightarrow \mathcal{D}$ is *symmetric* if it has additional structure consisting of isomorphisms $L(\sigma) : L(X_1, \dots, X_n) \rightarrow L(X_{\sigma(1)}, \dots, X_{\sigma(n)})$ for all $\sigma \in S_n$, with $L(\sigma^{-1}) = L(\sigma)^{-1}$. (In other words L is extended from \mathcal{C}^n to a wreath product category.) If L is symmetric and multilinear then the homogeneous functor L has a S_n -action. In the case of spectrum-valued functors (that is, when $\mathcal{D} = Sp$), the object

$$(S_n L)(X) = L(X_1, \dots, X_n)_{S_n}$$

is then again an n -homogeneous functor of X , by 1.7(5), since homotopy orbit space (or spectrum) is a special case of hocolim. We are headed toward proving that *all* n -homogeneous functors arise in just this way.

The inverse of $(S_n)_*$ will be provided by a construction called the n^{th} cross-effect, which takes a homotopy functor F and produces a symmetric homotopy functor $cr_n F$ of n variables. To see how this inverse construction should go, we recall an algebraic analogue. $(S_n)_*$ is analogous to the construction in algebra which uses a symmetric multilinear function $l(x_1, \dots, x_n)$ to make a homogeneous function $f(x) = l(x, \dots, x) = n!$. For example, if f is a degree two polynomial then from the bilinear form

$$l(x_1, x_2) = f(x_1 + x_2) - f(x_1) - f(x_2) + f(0)$$

we recover the purely quadratic part of f as $l(x, x) = 2f(x) - f(0)$. The second cross-effect $cr_2 F$ of a functor F of based spaces will take the based spaces X_1 and X_2 to the total homotopy fiber of the diagram

$$\begin{array}{ccc} F(X_1 \times X_2) & \longrightarrow & F(X_1) \\ \downarrow \wr & & \downarrow \wr \\ F(X_2) & \longrightarrow & F(*) \end{array}$$

Strictly speaking, in order to make cr_2F preserve weak equivalences we must first replace each X_i by an equivalent object having nondegenerate basepoint, perhaps by using "whiskers". In the general case (functors of n -wise based spaces over Y) the wedge $X_1 \vee X_2$ is replaced by a categorical sum in T_Y . Again, before forming the sum we should use n -wise whiskers, replacing each object X by the mapping cylinder of the structural coretraction $Y \rightarrow X$.

Thus the n^{th} cross-effect is defined as follows: Let $T_Y \xrightarrow{F} D$ be a homotopy functor, $D = T$ or Sp . For objects (X_1, \dots, X_n) of T_Y , let $S(X_1, \dots, X_n)$ be the evident n -cube taking $\underline{n} \rightarrow T$ to the (whiskered) sum, in T_Y , of the objects X_s for $s \geq T$. Define the cross-effect by first applying F to the cube and then taking the total homotopy fiber:

$$(cr_n F)(X_1, \dots, X_n) = \text{t.fiber } F(S(X_1, \dots, X_n)).$$

It is easy to see that $cr_n F$ is a homotopy functor (in each variable), and symmetric and $(1, \dots, 1)$ -reduced.

The first cross-effect is the "reduced functor":

$$(cr_1 F)(X) = \text{fiber}(F(X) \rightarrow F(Y)).$$

The 0^{th} cross-effect, should we ever need it, is a functor of no variables: the object $F(Y)$.

3.3 Proposition *If F is n -excisive then for $0 \leq m \leq n$ the functor $cr_{m+1}F$ is $(n-m)$ -excisive in each variable. In particular, the n^{th} cross-effect of an n -excisive functor is symmetric multilinear and the n^{th} cross-effect of an $(n-1)$ -excisive functor is trivial (equivalent to a point).*

Proof Induction on m . The case $m = 0$ is clear. To pass from $m-1$ to m , write

$$(cr_{m+1}F)(X_1, \dots, X_m; A) = (cr_m F_{+A})(X_1, \dots, X_m);$$

where $F_{+A}(X) = \text{ho fiber}(F(X+A) \rightarrow F(X))$ and $+$ denotes whiskered sum in T_Y . Use the rather obvious fact that F_{+A} is $(n-1)$ -excisive if F is n -excisive. \square

In general, then,

$$\begin{aligned} cr_n D_n F &= cr_n \text{ho fiber}(P_n F \rightarrow P_{n-1} F) \\ &= \text{ho fiber}(cr_n P_n F \rightarrow cr_n P_{n-1} F) \\ &= cr_n P_n F. \end{aligned}$$

Thus if F is n -excisive then $cr_n F$ is naturally equivalent to $cr_n D_n F$; the n^{th} cross-effect of an n -excisive functor "sees" only the n -homogeneous part. The following simple result suggests that it sees it quite clearly, at least in the spectrum-valued case.

3.4 Proposition *If F and G are n -homogeneous functors $T_Y \nrightarrow Sp$, then any (natural) map $F \nrightarrow G$ that induces an equivalence $cr_n F \nrightarrow cr_n G$ must be an equivalence itself.*

Proof Let H be the homotopy fiber of $F \nrightarrow G$. Thus H is n -homogeneous and $cr_n H$ is the homotopy fiber of $cr_n F \nrightarrow cr_n G$. Since a map of spectra must be an equivalence if its homotopy fiber is contractible, we have only to show that, for an n -homogeneous functor $T_Y \nrightarrow Sp$, $cr_n H \simeq *$ implies $H \simeq *$. In fact we will do a little better: assuming only that H is n -excisive and $cr_n H \simeq *$, we will show that H is $(n - 1)$ -excisive.

If X is any strongly cocartesian n -cube in T_Y , we must show that $H(X)$ is cartesian. Because we are dealing with spectra, it will be enough if we show that $\text{tiber } H(X) \simeq *$.

By assumption, this holds in the case where X is the cube $S(X_1; \dots; X_n)$ for objects $X_1; \dots; X_n$ of T_Y .

It also holds for the related cube, call it $S(X_1; \dots; X_n)$, which has the same objects but with reversed arrows, sending T rather than $\underline{n} - T$ to the sum of the objects X_s for $s \geq T$. In fact we have

$$\begin{aligned} \text{tiber } H(S(X_1; \dots; X_n)) &\simeq {}^n \text{tiber } H(S(X_1; \dots; X_n)) \\ &= {}^n (cr_n H)(X_1; \dots; X_n): \end{aligned}$$

This takes care of all cubes X in which $X(\cdot) \simeq Y$, because such a cube is naturally equivalent to $S(X(f1g); \dots; X(fng))$ (see [10; 2.2]).

Given an arbitrary strongly cocartesian cube X , put

$$X^0(T) = \text{hocolim}(Y \leftarrow X(\cdot) \rightarrow X(T)):$$

The obvious map of n -cubes $X \rightarrow X^0$ is a strongly cocartesian $(n + 1)$ -cube, so the resulting cube $H(X) \rightarrow H(X^0)$ is cartesian. The cube $H(X^0)$ is cartesian by the case already treated. Therefore by [9; 1.6] $H(X)$ is cartesian. \square

Let $L_n(C; D)$ be the category of symmetric multilinear functors $L : C^n \rightarrow D$. The maps are the natural maps that respect the symmetry. As usual the homotopy category $hL_n(C; D)$ means the category obtained by inverting the (objectwise) equivalences.

If D is either Sp or T then there is the cross-effect functor

$$H_n(T_Y; D) \xrightarrow{cr_n} L_n(T_Y; D)$$

If C is either U_Y or T_Y then there is the functor

$$L_n(C; Sp) \xrightarrow{!} H_n(C; Sp)$$

going the other way. Both cr_n and $!$ preserve weak equivalences and so induce functors on homotopy categories.

3.5 Theorem *The functors*

$$H_n(T_Y; Sp) \xrightarrow{cr} L_n(T_Y; Sp)$$

$$L_n(T_Y; Sp) \xrightarrow{-f} H_n(T_Y; Sp)$$

are mutual inverses up to natural (weak) equivalence.

Proof of 3.5 To prove that the composition

$$L_n(T_Y; Sp) \xrightarrow{-f} H_n(T_Y; Sp) \xrightarrow{cr} L_n(T_Y; Sp)$$

is equivalent to the identity, we look to the algebraic analogue. If l is a symmetric multilinear function of n variables and f is the homogeneous polynomial

$$f(x) = l(x, \dots, x)_{=n!};$$

then l can be recovered from f . It is given by

$$l(z_1, \dots, z_n) = f(z_1 + \dots + z_n) - f(z_1 + \dots + z_{n-1}) - \dots + (-1)^n f(0)$$

(an alternating sum of 2^n terms). One sees this, of course, by expanding the expression

$$l(x_1 + \dots + x_n, \dots, x_1 + \dots + x_n)$$

as a sum of n^n terms, cancelling all except the permutations of $l(x_1, \dots, x_n)$ and dividing by n factorial.

Here is a corresponding categorical argument: We have

$$\begin{aligned} (cr_n \circ nL)(X_1, \dots, X_n) &= \text{t ber}((nL) \circ S(X_1, \dots, X_n)) \\ &= \text{t ber}(\underline{n} - T \int L(\sum_{s \in T} X_s, \dots, \sum_{s \in T} X_s)_{h_n}) \\ &= (\text{t ber}(X))_{h_n}; \end{aligned}$$

where $X(\underline{n} - T) = L(\sum_{s \in T} X_s, \dots, \sum_{s \in T} X_s)$ and \int denotes whiskered sum in T_Y . The obvious equivalence

$$X(\underline{n} - T) \xrightarrow{\int} L(X_{(1)}, \dots, X_{(n)})$$

is natural with respect to T . It can be interpreted as an equivalence of cubes $X \xrightarrow{\int} Y$ where

$$Y(\underline{n} - T) = L(X_{(1)}, \dots, X_{(n)}); \text{ if } \underline{n} = T$$

$$Y(\underline{n} - T) = \dots; \text{ otherwise.}$$

For any \underline{n} that is not a permutation and therefore not surjective, the cube Y is cartesian. (Viewed in one way it is an isomorphism of $(n - 1)$ -cubes: if

$s \geq n$ then all maps $Y(T) \rightarrow Y(T \circ fsg)$ are isomorphisms.) For any permutation σ we have

$$\text{t ber}(Y) = L(X_{(1)}; \dots; X_{(n)})$$

Therefore

$$\text{t ber}(X) \xrightarrow{\cong} L(X_{(1)}; \dots; X_{(n)})$$

This map respects the symmetry if the group is made to permute the factors of the right-hand side, and so it leads to an equivalence of homotopy orbit spectra

$$(cr_n \circ nL)(X_1; \dots; X_n) \xrightarrow{\cong} L(X_{(1)}; \dots; X_{(n)})_{h_n} \cong L(X_1; \dots; X_n)$$

Note for future reference that an explicit inverse equivalence

$$L(X_1; \dots; X_n) \xrightarrow{\cong} (cr_n \circ nL)(X_1; \dots; X_n)$$

is the map of total homotopy fibers induced by an obvious map of cubes

$$Y_1 \rightarrow X \rightarrow X_{h_n};$$

where $\underline{n} \xrightarrow{\cong} \underline{n}$ is the identity map and the map $Y_1(\cdot) \rightarrow X(\cdot)$ is the map

$$L(X_1; \dots; X_n) \xrightarrow{\cong} L(Z; \dots; Z)$$

induced by the inclusions $X_j \rightarrow Z = \prod_{j=1}^n X_j$. The following diagram commutes:

$$\begin{array}{ccc} L(X_1; \dots; X_n) & \xrightarrow{\cong} & L(Z; \dots; Z) \\ \downarrow \cong & & \downarrow \cong \\ (cr_n \circ nL)(X_1; \dots; X_n) & \xrightarrow{\cong} & L(Z; \dots; Z)_{h_n} \end{array}$$

Here, and throughout this proof, $\downarrow \cong$ denotes the projection from the total homotopy fiber of a cubical diagram to the "initial" object in the diagram.

The remaining task is to exhibit an equivalence

$$cr_n \circ nF \xrightarrow{\cong} F$$

for any n -homogeneous $T_Y \xrightarrow{\cong} Sp$. In fact we will define γ for any n -excisive F and then show that it induces an equivalence

$$cr_n \circ ncr_n F \xrightarrow{\cong} cr_n F:$$

By 3.4 this suffices.

To define $\hat{}$ we use a map $\hat{}$, defined for $Z \in T_Y$ as the composition

$$(cr_n F)(Z; \dots; Z) \xrightarrow{\hat{}} F(\prod_{j=1}^n Z) \xrightarrow{F(f)} F(Z)$$

where f is the "fold" map which takes each copy of Z identically to Z . The map $\hat{}$ is equivariant with respect to the obvious S_n -actions. (The action on $F(Z)$ is trivial.) Define $\hat{}$ as the composition

$$(\hat{} cr_n F)(Z) = ((cr_n F)(Z; \dots; Z))_{h_n} \xrightarrow{\hat{}} F(Z)_{h_n} = F(Z) \wedge (B_{n+}) \xrightarrow{\hat{}} F(Z)$$

where the last arrow is induced by the nontrivial based map $(B_{n+}) \rightarrow S^0$.

To see that $cr_n(\hat{})$ is an equivalence we examine the composition

$$cr_n F \xrightarrow{\hat{}} cr_n (\hat{} cr_n F) \xrightarrow{cr_n(\hat{})} cr_n F$$

It will be enough if it coincides with the identity, at least on the level of homotopy groups. In fact it will be enough if the composition

$$(cr_n F)(X_1; \dots; X_n) \xrightarrow{cr_n(\hat{})} (cr_n F)(X_1; \dots; X_n) \xrightarrow{\hat{}} F(Z)$$

is equal to $\hat{}$, since $\hat{}$ is a split injection. (Here again $Z = \prod_{j=1}^n X_j$.)

That it is is a direct consequence of the following facts: The diagram

$$\begin{array}{ccc} (cr_n F)(X_1; \dots; X_n) & \xrightarrow{\hat{}} & (cr_n F)(Z; \dots; Z) \\ \downarrow \hat{} & & \downarrow \hat{} \\ (cr_n (\hat{} cr_n F))(X_1; \dots; X_n) & \xrightarrow{\hat{}} & (cr_n F)(Z; \dots; Z)_{h_n} \\ \downarrow \hat{} & & \downarrow \hat{} \\ (cr_n F)(X_1; \dots; X_n) & \xrightarrow{\hat{}} & F(Z); \end{array}$$

commutes; the composition of the right-hand vertical maps above is $\hat{}$, which is also the composition of the lower horizontal maps below; the diagram

$$\begin{array}{ccc} cr_n F(X_1; \dots; X_n) & \xrightarrow{\hat{}} & F(Z) \\ \downarrow \hat{} & & \downarrow \hat{} \\ cr_n F(Z; \dots; Z) & \xrightarrow{\hat{}} & F(\prod_{j=1}^n Z) \xrightarrow{F(f)} F(Z); \end{array}$$

commutes, where $Z \xrightarrow{D} \prod_{j=1}^n Z$ sends the copy of X_i in Z to the copy of X_i in the i^{th} copy of Z ; and the composition $f \circ D$ is the identity map. \square

3.6 Corollary The functor $H_n(T_Y; T) \xrightarrow{cr_n} L_n(T_Y; T)$ has an inverse up to weak equivalence.

Proof of 3.6 Using the commutative diagram

$$\begin{array}{ccc} H_n(T_Y; T) & \xrightarrow{1} & H_n(T_Y; Sp) \\ \wr \uparrow \text{cr}_n & & \wr \uparrow \text{cr}_n \\ L_n(T_Y; T) & \xrightarrow{1} & L_n(T_Y; Sp) \end{array}$$

this will follow from 3.5, 2.1, and the next result. □

3.7 Proposition The functor $L_n(C; Sp) \xrightarrow{1} L_n(C; T)$ has an inverse up to weak equivalence.

Proof As in proving 2.1, we need a delooping functor from $L_n(C; T)$ to itself. This is much easier than 2.2.

We refer to Remark 1.22 for notation. If L is symmetric multilinear then there are natural equivalences

$$L(X_1; \dots; X_n) \cong (T_{1, \dots, 1} L)(X_1; \dots; X_n) \cong {}^n L(\vee X_1; \dots; \vee X_n):$$

We will need these equivalences to respect the n -symmetry. In order for this to be true, permutations of the loop coordinates must be built into the symmetry of the last expression; a better expression is $V_n L(\vee X_1; \dots; \vee X_n)$, where V_n is the standard n -dimensional representation of Σ_n . Since this contains a trivial one-dimensional representation, we have what we need: if $V_n = \mathbb{R} \oplus V_n'$ then $L \cong BL$ where BL is defined by

$$(BL)(X_1; \dots; X_n) = V_n L(\vee X_1; \dots; \vee X_n): \quad \square$$

We can now justify an assertion made in 1.20. Let D be either T or Sp .

3.8 Corollary If $T_Y \xrightarrow{F} D$ is n -homogeneous then F is determined by $F \vee \gamma$.

Proof According to 3.5 and 3.6 F is determined by the cross-effect $cr_n F$, and this satisfies

$$(cr_n F)(\vee X_1; \dots; \vee X_n) = (cr_n(F \vee \gamma))(X_1; \dots; X_n):$$

On the other hand, for any symmetric multilinear functor L we have

$$L(X_1; \dots; X_n) \cong V_n L(\vee X_1; \dots; \vee X_n);$$

so that

$$\begin{aligned} (cr_n F)(X_1; \dots; X_n) &\cong V_n (cr_n F)(\vee X_1; \dots; \vee X_n) \\ &= V_n cr_n(F \vee \gamma)(X_1; \dots; X_n) \end{aligned} \quad \square$$

4 The role of the base point

The cross-effect construction applies to functors $T_Y \dashv D$ but not to functors $U_Y \dashv D$. In spite of this we now show that the classification of homogeneous functors extends without change to the U_Y case.

Let $T_Y \dashv U_Y$ be the forgetful functor. Because \dashv preserves equivalences and cocartesian square diagrams, composition with \dashv yields functors

$$\begin{aligned}
 H_n(U_Y; D) &\dashv H_n(T_Y; D) \\
 L_n(U_Y; D) &\dashv L_n(T_Y; D):
 \end{aligned}$$

4.1 Theorem *If D is T or Sp then both of the two functors \dashv above have inverses up to weak equivalence.*

4.2 Corollary *The functor $L_n(U_Y; Sp) \dashv H_n(U_Y; Sp)$ has an inverse up to weak equivalence.*

Proof of 4.2 Use 3.5, 4.1, and the diagram

$$\begin{array}{ccc}
 L_n(U_Y; Sp) & \dashv & H_n(U_Y; Sp) \\
 \downarrow \cong & & \downarrow \cong \\
 L_n(T_Y; Sp) & \dashv & H_n(T_Y; Sp): \quad \square
 \end{array}$$

Proof of 4.1 We can assume $D = Sp$, since by 2.1 and 3.7 the spectrum-valued case of 4.1 implies the space-valued case.

Let $U_Y \dashv T_Y$ be the left adjoint of \dashv , so that if X is a space over Y then $\dashv(X)$ is the disjoint union of X and Y viewed as a fiberwise based space over Y . Like \dashv , this functor preserves equivalences and cocartesian square diagrams. Therefore if $T_Y \dashv Sp$ is n -excisive then $U_Y \dashv Sp$ will be n -excisive. Unlike \dashv , \dashv does not preserve the final object Y , so $\dashv F$ need not be homogeneous if F is. We will show that

$$H_n(T_Y; Sp) \xrightarrow{D_n} H_n(U_Y; Sp)$$

is an inverse, up to weak equivalence, for

$$H_n(U_Y; Sp) \dashv H_n(T_Y; Sp):$$

The unit map $X \dashv \dashv X$ of the adjoint pair induces a map $F \dashv \dashv F$. For any morphism $X \dashv \dashv X^0$ in U_Y the square diagram

$$\begin{array}{ccc} X & \xrightarrow{\quad} & X \\ \wr & & \wr \\ Y & & Y \\ X^0 & \xrightarrow{\quad} & X^0 \end{array}$$

is cocartesian. Therefore if X is any strongly cocartesian n -cube in U_Y the unit yields a strongly cocartesian $(n + 1)$ -cube

$$X \dashv \dashv X:$$

It follows that if $U_Y \dashv \dashv Sp$ is n -excisive then the fiber of $F \dashv \dashv F$ is $(n - 1)$ -excisive. Thus if F is n -homogeneous there are natural equivalences

$$F \dashv \dashv P_n F \dashv \dashv D_n F \dashv \dashv D_n \dashv \dashv F$$

(The last map is an equivalence because its fiber is contractible. This implication relies on the fact that these are spectrum-valued functors.)

The counit $X \dashv \dashv X$, like the unit, yields a cocartesian square for every map and a strongly cocartesian $(n + 1)$ -cube for every strongly cocartesian n -cube.

Therefore for any n -homogeneous $U_Y \dashv \dashv Sp$ it yields equivalences

$$F \dashv \dashv P_n F \dashv \dashv D_n F \dashv \dashv D_n \dashv \dashv F = \dashv \dashv D_n \dashv \dashv F:$$

This completes the proof in the homogeneous case.

We sketch the proof in the multilinear case, which is much the same. The inverse of

$$L_n(U_Y; Sp) \dashv \dashv L_n(T_Y; Sp)$$

is

$$L_n(T_Y; Sp) \xrightarrow{\quad r \quad} L_n(U_Y; Sp);$$

where r is the operation of "reducing" a symmetric homotopy functor of n variables in all variables simultaneously. For example, when $n = 2$ then $(rL)(X_1; X_2)$ is the total homotopy fiber of

$$\begin{array}{ccc} L(X_1; X_2) & \xrightarrow{\quad} & L(X_1; Y) \\ \wr & & \wr \\ Y & & Y \\ L(Y; X_2) & \xrightarrow{\quad} & L(Y; Y): \end{array}$$

The key point is that the maps $L \dashv \dashv L$ and $L \dashv \dashv L$ induce equivalences $rL \dashv \dashv r \dashv \dashv L = \dashv \dashv r \dashv \dashv L$ and $r \dashv \dashv L \dashv \dashv rL$. For example, in the composition

$$L(X_1; X_2) \dashv \dashv L(X_1; X_2) \dashv \dashv L(X_1; X_2)$$

the fiber of the first map becomes contractible upon reducing with respect to X_2 while the fiber of the second map becomes contractible upon reducing with respect to X_1 , and so they both become contractible upon applying r . \square

The proof of 4.1 suggests a variant of the notion of multilinear functor, related to it as unreduced homology is related to reduced homology.

4.3 Definition A functor $U_Y^n \rightarrow Sp$ is unreduced-multilinear if it is 1-exciseive in each variable and if it satisfies $L(X_1, \dots, X_n) = 0$ whenever X_s is the initial object \ast for some s . The category of symmetric unreduced-multilinear functors is $\hat{L}_n(U_Y; Sp)$.

Since $(\cdot)_Y = Y$, the functor r maps $L_n(T_Y; Sp)$ into $\hat{L}_n(U_Y; Sp)$.

4.4 Proposition The functor $L_n(T_Y; Sp) \xrightarrow{r} \hat{L}_n(U_Y; Sp)$ has an inverse up to weak equivalence.

Proof The proof of 4.1 shows that r is an inverse. \square

4.5 Warning Although Definition 4.3 could be extended verbatim to functors into based spaces, the resulting category $\hat{L}_n(U_Y; T)$ would not have the expected property: the corresponding variant of Proposition 4.4 would be false. An instructive example is obtained by adding a disjoint basepoint to the excisive functor $U \xrightarrow{P_1} U$ mentioned near the end of the introduction.

4.6 Remark 3.8 is valid for functors of unbased spaces, in view of 4.1 and the fact that r commutes with γ up to isomorphism.

5 The n^{th} differential and the n^{th} derivative

We can summarize the main results of Sections 2 through 4 by saying that the following eight categories of functors are equivalent at the homotopy category level:

$$\begin{array}{ccc} H_n(U_Y; Sp) & \xrightarrow{r} & H_n(U_Y; T) \\ \underbrace{\quad}_{\gamma} & & \underbrace{\quad}_{\gamma} \\ H_n(T_Y; Sp) & \xrightarrow{r} & H_n(T_Y; T) \end{array}$$

$$\begin{array}{ccc}
 L_n(U_Y; Sp) & \xrightarrow{1} & L_n(U_Y; T) \\
 \downarrow \text{?} & & \downarrow \text{?} \\
 L_n(T_Y; Sp) & \xrightarrow{1} & L_n(T_Y; T)
 \end{array}$$

In addition to the arrows displayed, there is also cr_n from the upper square to the lower in each of the two $(T_Y; -)$ cases and \downarrow_n from the lower square to the upper in each of the $(-; Sp)$ cases. We have explicitly inverted enough of these arrows to show that all of them are invertible: We inverted the four called $\xrightarrow{1}$ in 2.1 and 3.7; we inverted the left hand \downarrow of each square in 4.1; and in 3.1 we connected the two squares by showing that cr_n and \downarrow_n are inverses in the $(T_Y; Sp)$ case.

Let $C_Y \xrightarrow{F} D$ be a homotopy functor from spaces over Y to either based spaces or spectra, and suppose that we wish to describe the homogeneous layer $D_n F$ in its Taylor tower for some $n \geq 1$. By the results above, knowing $D_n F$ is the same as knowing a certain symmetric multilinear functor. This will be called the n -fold differential of F .

5.1 Definition The object in $L_n(C_Y; D)$ corresponding to the homogeneous functor $D_n F \in H_n(C_Y; D)$ is called the n -fold differential of F and is denoted by $D^{(n)} F$.

Specifically, $D^{(n)} F$ determines $D_n F$ by the rule:

$$(D_n F)(X) = (D^{(n)} F)(X; \dots; X)_{h_n}$$

in the case when $D = Sp$, or

$$(D_n F)(X) = B^{-1}((B^1 D^{(n)} F)(X; \dots; X)_{h_n})$$

in the case when $D = T$. Here B^{-1} is the inverse (up to natural weak equivalence) to

$$L_n(C_Y; Sp) \xrightarrow{1} L_n(C_Y; T)$$

provided by 3.6.

Conversely, to obtain $D^{(n)} F$ from $D_n F$ one simply takes the n^{th} cross-section in the case when $C_Y = T_Y$, while in the case when $C_Y = U_Y$ one knows that $D^{(n)} F$ determines $D_n F$ by 4.2 and is given by

$$D^{(n)} F = D^{(n)} F = cr_n D_n F:$$

In practice we often work with a functor $U_Y \xrightarrow{F} D$ that is the restriction of a functor $U \xrightarrow{F} D$, meaning the composition of F with the forgetful functor $U_Y \leftarrow U$.

5.2 Definition In this case $D^{(n)}F_Y$ is called the n -fold differential of F at Y and denoted by $D_Y^{(n)}F$.

The next theme to be developed is the description of multilinear functors by their "coefficients". We discuss this briefly in the important special case when Y is the one-point space, and then a little more elaborately in the general case.

If C is a spectrum then the functor

$$L(X_1; \dots; X_n) = C \wedge (X_1 \wedge \dots \wedge X_n)$$

from $(n$ -tuples of) based spaces to spectra is multilinear, and if C has an action of the symmetric group Σ_n then L is symmetric.

Conversely, if L is a multilinear functor from based spaces to spectra then, taking C to be $L(S^0; \dots; S^0)$, we have (essentially, see [9; page 5]) a natural assembly map

$$C \wedge (X_1 \wedge \dots \wedge X_n) \rightarrow L(X_1; \dots; X_n):$$

This is an equivalence when $X_j = S^0$ for all j , and it follows (see 5.8 below), using the multilinearity of both functors, that it is an equivalence when the X_j are finite complexes. If L satisfies a suitable limit axiom (5.10 below), then this even holds for all X_j . If L is symmetric then C gets a Σ_n -action and the assembly map respects the symmetry.

In short, symmetric multilinear functors from finite based spaces to spectra correspond precisely, in the sense of an equivalence of homotopy categories, to spectra with Σ_n -action. The spectrum (together with its Σ_n -action) is called the *coefficient* of the symmetric multilinear functor, or of the corresponding homogeneous functor.

Thus if F is a homotopy functor from based spaces to based spaces or spectra then for every $n > 0$ the n -homogeneous layer $D_n F$ of its Taylor tower is governed by a certain spectrum with Σ_n -action. This will be called the n^{th} derivative of F at the one-point space and denoted by $@^{(n)}F(\)$. In the spectrum-valued case we have

$$@^{(n)}F(\) \simeq (D^{(n)}F)(S^0; \dots; S^0)$$

and

$$(D_n F)(X) \simeq (@^{(n)}F(\) \wedge X^{\wedge n})_{h \Sigma_n}$$

In the space-valued case we have

$$^1 @^{(n)}F(\) \simeq (D^{(n)}F)(S^0; \dots; S^0)$$

and

$$(D_n F)(X) \simeq ^1 ((@^{(n)}F(\) \wedge X^{\wedge n})_{h \Sigma_n})$$

To fully describe $@^{(n)}F(\cdot)$ in the space-valued case, we examine the proof of 3.7 and see that the i^{th} space in the spectrum is

$$iV_n(D^{(n)}F)(S^i; \dots; S^i)$$

5.3 Remark By the proof of 3.8 we have

$$@^{(n)}F(\cdot) = V_n @^{(n)}(F(\cdot))(\cdot)$$

5.4 Remark If a subgroup $G \subset_n$ acts on a spectrum C then the functor $(C \wedge X^{\wedge n})_{hG}$ is homogeneous. When written in standard form it is

$$((\mathbb{Z}/n)_+ \wedge_G C \wedge X^{\wedge n})_{h_n};$$

the coefficient is the "induced spectrum" $(\mathbb{Z}/n)_+ \wedge_G C$.

We now pass from functors of spaces to functors of spaces over Y while doing our best to retain the principle that a multilinear functor is determined by its behavior on a small class of objects.

Here is some notation for naming special objects of T_Y : If Z is a based space and y is a point in Y then let $Y_{-y}Z$ be the union of Y and Z with y identified to the basepoint of Z . This is to be viewed as a space over Y with all of Z being mapped to y . More generally if several points $y_1; \dots; y_n$ and several based spaces $Z_1; \dots; Z_n$ are given then $Y_{-y_1}Z_1 \dots_{-y_n}Z_n$ is the space obtained from Y by attaching Z_j at y_j for all j . Again this is to be viewed as an object of T_Y .

Let F be a homotopy functor from unbased spaces to either based spaces or spectra.

5.6 Definition Let Y be a space and let $y_1; \dots; y_n$ be points in Y . The n^{th} derivative of F at $(Y; y_1; \dots; y_n)$, denoted

$$@_{y_1; \dots; y_n}^{(n)}F(Y);$$

is the coefficient spectrum of the multilinear functor

$$(Z_1; \dots; Z_n) \mapsto (D_Y^{(n)}F)(Y_{-y_1}Z_1; \dots; Y_{-y_n}Z_n);$$

Thus in the spectrum-valued case we have

$$\partial_{y_1, \dots, y_n}^{(n)} F(Y) \simeq (D_Y^{(n)} F)(Y_{-y_1} S^0; \dots; Y_{-y_n} S^0);$$

$\partial_{y_1, \dots, y_n}^{(n)} F$ has a symmetry with respect to permutations of $(y_1; \dots; y_n)$ and is also functorial in Y . To be more precise, we have here a functor whose domain is the category in which an object is a "space with n basepoints" $(Y; y_1; \dots; y_n)$ and a morphism $(Y; y_1; \dots; y_n) \rightarrow (W; w_1; \dots; w_n)$ is a pair $(f; \sigma)$ with $Y \xrightarrow{f} W$ being a continuous map and σ a permutation such that $f(y_{\sigma(j)}) = w_j$ for all j . The points y_j are not assumed distinct.

It is true in various senses, beginning with 5.9 below, that the n^{th} derivative of a functor determines the behavior of the n^{th} differential, at least on "finite" objects.

5.7 Definition An object $Y \xrightarrow{f} X \xrightarrow{g} Y$ of T_Y is "finite" if $(X; i(Y))$ is a "finite CW pair". An object $X \xrightarrow{f} Y$ of U_Y is "finite" if the space X is "finite CW".

5.8 Proposition Let $L_1 \xrightarrow{L} L_2$ be a map between symmetric multilinear functors from T_Y to Sp . In order that

$$L_1(X_1; \dots; X_n) \xrightarrow{L} L_2(X_1; \dots; X_n)$$

should be an equivalence whenever the objects X_j are "finite", it is enough if g is an equivalence in the special case when each X_j is $Y_{-y_j} S^0$ for an arbitrary point $y_j \in Y$.

Proof The symmetry is irrelevant here, and it is clear that the case $n = 1$ implies the general case. We therefore give a proof for the case of a map $L_1 \xrightarrow{L} L_2$ between linear functors from T_Y to spectra.

Define $T_Y \xrightarrow{L} Sp$ by letting $L(X)$ be the homotopy fiber of $L_1(X) \xrightarrow{L} L_2(X)$. The functor L "vanishes" at $Y_{-y} S^0$, for every point y in Y , in the sense that $L(Y_{-y} S^0) \simeq *$. We also know that L is linear; it preserves weak equivalences, it takes cocartesian squares to cartesian squares, and it vanishes at Y . We must show that L vanishes at every "finite" object X .

We name some more objects of T_Y : If Z is a space and $Z \xrightarrow{f} Y$ is a map then let $Y +_f Z$ be the disjoint union of Y and Z , considered as an object of T_Y in the evident way.

If X is an object of T_Y obtained by attaching an m -cell to another object X^0 , then there is a cocartesian diagram

$$\begin{array}{ccc} Y +_f j_m S^{m-1} & \longrightarrow & X^0 \\ \uparrow \scriptstyle{?} & & \uparrow \scriptstyle{?} \\ Y & & Y \\ Y +_f D^m & \longrightarrow & X \end{array}$$

for some $D^m \xrightarrow{f} Y$, and thus a cartesian diagram

$$\begin{array}{ccc} L(Y +_f j_m S^{m-1}) & \longrightarrow & L(X^0) \\ \uparrow \scriptstyle{?} & & \uparrow \scriptstyle{?} \\ Y & & Y \\ L(Y +_f D^m) & \longrightarrow & L(X) \end{array}$$

$L(X)$ will be contractible if the other three spectra are. Thus an induction on the number of cells in $X - i(Y)$ will be possible as soon as we have dealt with the cases $X = Y +_f D^m$ and $X = Y +_f S^m$. The sphere case follows from the disk case by induction on m (beginning with the case $Y +_f S^{-1} = Y$), and the disk case is taken care of by a weak equivalence

$$Y -_y S^0 \xrightarrow{\simeq} Y +_f D^n$$

obtained by choosing a point $y \in f(D^n)$. □

Note that if Y is path-connected then it is only necessary to verify the hypothesis of 5.8 for one choice of (y_1, \dots, y_n) , since a path $I \xrightarrow{f} Y$ from y to y^0 in Y yields a diagram of equivalences in T_Y :

$$Y -_y S^0 \xrightarrow{\simeq} Y +_f I \xrightarrow{\simeq} Y -_{y^0} S^0$$

There is a variant of 5.8, with the same proof, for unreduced-multilinear functors from U_Y to spectra, with a point over y_j replacing $Y -_{y_j} S^0$. The proof of 5.8 also generalizes rather easily to prove that a map between d -excisive homotopy functors from T_Y [resp. U_Y] to $S\rho$ must be an equivalence for all objects if it is an equivalence for all objects of the form $Y +_f S$ [resp. S] where S is a discrete set of at most d points with a map to X .

For us the main consequence of 5.8 is:

Corollary 5.9 *If a map $F \xrightarrow{\simeq} G$ of homotopy functors $U \xrightarrow{\simeq} T$ induces an equivalence of n^{th} derivatives*

$$\text{@}_{y_1, \dots, y_n} F(Y) \xrightarrow{\simeq} \text{@}_{y_1, \dots, y_n} G(Y)$$

for every point $(y_1, \dots, y_n) \in Y^n$ then it induces an equivalence of n^{th} derivatives

$$D_n F_Y(X) \xrightarrow{\simeq} D_n G_Y(X)$$

for every finite object X of U_Y or T_Y .

Proof Apply 5.8 to $D^{(n)}F_Y \neq D^{(n)}G_Y$, noting that the behavior of $D_n F_Y$ on finite objects is determined by the behavior of $D^{(n)}F_Y$ on finite objects. \square

These statements definitely require spectra rather than spaces as the output of the functors. For example, if C is a spectrum and L is the linear functor from based spaces to based spaces given by

$$L(X) = \pi^{-1}(C \wedge X);$$

then the statement that $L(S^0)$ is contractible means only that the homotopy groups $\pi_j(C)$ are trivial for $j \neq 0$.

The restriction to finite objects in the results above can be removed if F satisfies a suitable limit axiom.

5.10 Definition A homotopy functor $C \xrightarrow{F} D$ is *continuity* if it preserves filtered homotopy colimits up to weak equivalence, that is, if the natural map

$$\text{hocolim } F(X_i) \rightarrow F(\text{hocolim } X_i)$$

is a weak equivalence whenever $f: X \rightarrow g$ is a diagram in C indexed by a filtering category.

In the case of linear functors from spaces to spaces, this condition means that the corresponding homology theory satisfies Milnor's wedge axiom. It is clear that the n^{th} Taylor approximation of a continuity functor is itself continuity.

5.11 Remark This was called the *limit axiom* in [10], but it seems useful to have an adjective available. The term "continuous functor" has been used, but we would rather reserve that for something else (a functor that behaves continuously on morphisms).

Since every space is equivalent to a filtered hocolim of finite CW complexes, a continuity functor of spaces is determined (up to natural weak equivalence) by its behavior on finite complexes. More generally a continuity functor of objects in C_Y is determined by its behavior on objects that are finite in the sense of 5.7.

According to 5.9 the n^{th} differential (which knows all about the n^{th} layer of the Taylor tower) is in some sense controlled by the n^{th} derivative. This is only a weak sense, however, since 5.9 cannot produce an equivalence between the differentials of F and G unless a map between F and G is already given. It would be better to have a way of building the n^{th} differential from the n^{th} derivative.

The n^{th} derivative of F at Y gives a spectrum for each ordered n -tuple of points in Y . To identify the n^{th} differential, one needs just a little more information. We will sketch one of several possible answers to the following vague question:

5.13 Vague question How can $D^{(n)}F$ be assembled from the spectra $@_{y_1, \dots, y_n}^{(n)} F(Y)$?

Answers in the case $n = 1$ tend to generalize easily to the general case, so we will concentrate on that case.

A useful point of view is that a (unitary) linear functor of spaces over Y corresponds to a "coefficient system" on Y which assigns a spectrum E_y to each point y . The spectrum E_y must depend on the point y continuously in some sense, so the linear functor cannot really be specified by merely giving E_y for each y . (That would be like trying to specify a vector bundle by giving all of its fibers, a mere collection of vector spaces.)

One can give rigorous sense to this idea by making the following definition: A *system of spectra* on Y consists of objects $fE_n g$ in T_Y such that the structural maps $E_n \rightarrow Y$ are fibrations, related by maps

$$Y \rightarrow E_n \rightarrow E_{n+1}$$

We may write $E_{y,n}$ for the fiber of $E_n \rightarrow Y$ over $y \in Y$. The spaces $fE_{y,n} g$ for fixed y constitute a spectrum E_y , the fiber of E over y .

Such a system E determines a prespectrum whose n^{th} space is the homotopy cofiber of the structural map $Y \rightarrow E_n$. The associated spectrum, which may be thought of as the homology of Y with coefficients in E , will be denoted by

$$\int_{y \in Y} E_y dy$$

Of course this notation is deceptive, since it appears not to matter how the various spectra E_y are related.

The integral signs are not meant to suggest antiderivation.

We draw attention to the familiar special case when E is a "trivial bundle of spectra" over Y : Let C be a spectrum and take E_n to be $Y \times C_n$. Then

$$\int_{y \in Y} E_y dy = \int_{y \in Y} C dy = C \wedge Y$$

In the general case there is a spectral sequence of Atiyah-Hirzebruch type, with

$$E_{p,q}^2 H_p(Y; \pi_q(E_y)) \Rightarrow_{p+q} \int_{y \in Y} E_y dy$$

One constructs it by taking the direct limit over n of a spectral sequence with

$$E_{p,q}^2 = H_p(Y; \pi_{q+n}(E_{n,y})) \Rightarrow_{p+q+n} \text{hoco fiber}(Y \rightarrow E_n)$$

Systems of spectra pull back: a map $X \xrightarrow{f} Y$ and a system E on Y determine a system f^*E on X given by $(f^*E)_n = E_n \times_Y X$, whose fiber at $x \in X$ is isomorphic to $E_{f(x)}$. Thus a system E on Y gives a functor from U_Y to Sp :

$$X \mapsto \prod_{x \in X} E_{f(x)}$$

This is a homotopy functor, and it is multiplicative and 1-excisive; all of these assertions can be proved by spectral sequence comparison arguments. This functor vanishes at the empty set, so it is not linear but rather unreduced-linear (4.5). Reducing it, one gets a linear functor, which sends X to the homotopy fiber of

$$\prod_{x \in X} E_{f(x)} \rightarrow \prod_{y \in Y} E_y$$

It is fairly clear that one could classify the multiplicative linear functors on T_Y along these lines, but we will not pursue that here. The generalization to $n > 1$ involves systems of spectra over Y^n with a Σ_n -symmetry, or alternatively systems of spectra over Y^n / Σ_n .

If Y is equivalent to the classifying space of a group G then systems of spectra on Y are the same (at the homotopy category level) as spectra with G -action. This idea can be extended to simplicial groups G , so that it applies to all based connected Y .

The idea of constructing linear functors by systems of spectra over a space was implicitly present in sections 2 and 3 of [9]. The systems that arose there were mainly "fiberwise suspension spectra" in the following sense: If W is an object of T_Y whose structural map $W \rightarrow Y$ is a fibration, then the repeated fiberwise suspensions $\Sigma^q W$ form a system E whose fiber E_y is the suspension spectrum of the fiber of W over y .

There is also a dual construction, in which E is used as coefficients for twisted cohomology rather than twisted homology. If E is a system of spectra on a CW complex K then the spectrum

$$\prod_{k \in K} E_k$$

is defined by letting the n^{th} space be the space of sections of the fibration $E_n \rightarrow Y$. If K is locally compact then there is also the compactly supported version

$$\int_c \prod_{k \in K} E_k$$

made out of spaces of compactly supported sections.

If $X \not\rightarrow E(X)$ is a homotopy functor from spaces to systems of spectra on the finite complex K then "differentiation under the integral" is valid; there is an equivalence

$$\int_{@_y} \int_{k \in K} E_k(Y) dk \cong \int_{@_y} \int_{k \in K} E_k(Y) dk:$$

The proof is by induction on the number of cells in K . This extends to the compactly supported case if for example the one-point compactification of K is a finite complex.

We will also need the following simple principle: If $P \rightarrow B$ is a principal G -bundle (G being a topological group) and E is a system of spectra on the locally compact space B , then when G is made to act on the spectrum

$$\int_{p \in P} E_{(p)} dp$$

in the obvious way the homotopy fixed point spectrum is equivalent to

$$\int_{b \in B} E_b db:$$

Confession We should really distinguish between (co)unreduced suspension and (co)reduced suspension. The former, (co)join with a two-point set, is what we ordinarily denote by Σ here, and it has the pleasant feature that it takes fibrations to fibrations. The latter, on the other hand, is much better for making spectra. When they are inequivalent, we have to choose the former, but this must be paid for by growing whiskers. We have not systematically imposed a solution of this small technical difficulty on the reader, because we do not have a neat solution.

6 Multilinearized cross-effects

Before working on some examples, we need one more tool.

6.1 Theorem Let D be T or Sp . The n^{th} differential $D^{(n)}F = cr_n D_n F$ of a homotopy functor $T_Y \dashv D$ is (naturally, weakly) equivalent to the multilinearization of the n^{th} cross-effect $cr_n F$ of F itself.

This means that $(D^{(n)} F)(X_1; \dots; X_n)$ is essentially the homotopy colimit of

$${}^{i_1+\dots+i_n}(cr_n F)(\ {}^i_1 X_1; \dots; \ {}^i_n X_n)$$

over $(i_1; \dots; i_n)$. The latter may also be described as

$$\text{hocolim}_i \ {}^{iV_n}(cr_n F)(\ {}^i_1 X_1; \dots; \ {}^i_n X_n):$$

In particular the n^{th} derivative $@_{y_1, \dots, y_n}^{(n)} F(Y)$ is equivalent to the associated spectrum of a prespectrum whose i^{th} space is

$${}^{iV_n}(cr_n F)(Y_{-y_1} S^i; \dots; Y_{-y_n} S^i):$$

6.1 can be a key tool for identifying $D_n F$ in examples. While the definitions of $P_n F$ and $D_n F$ are difficult to use for explicit calculation when n is greater than one, we do know how to recover $D_n F$ from $cr_n D_n F$, and 6.1 says that this in turn can be obtained rather directly from F itself.

6.2 Remark Like 1.7 and 2.1, 6.1 has an easier proof in the case when the functor F is analytic. In fact, in that case using 1.5(2) one sees easily that if the maps $X_j \not\rightarrow Y$ are all k -connected then the map

$$(cr_n F)(X_1; \dots; X_n) \not\rightarrow (cr_n P_n F)(X_1; \dots; X_n)$$

induced by $p_n F$ is $((n + 1)k - c^0)$ -connected for some constant c^0 , from which it follows that the canonical map from

$${}^{iV_n}(cr_n F)(\ {}^i_1 X_1; \dots; \ {}^i_n X_n)$$

to

$${}^{iV_n}(cr_n P_n F)(\ {}^i_1 X_1; \dots; \ {}^i_n X_n) \rightarrow (cr_n P_n F)(X_1; \dots; X_n)$$

has a connectivity tending to ∞ with i . For that matter, in applying 6.1 to a particular F one often uses this same kind of reasoning again: One identifies the multilinearization of $cr_n F$ with a given functor L by exhibiting a (natural and symmetry-preserving) map

$$(cr_n F)(X_1; \dots; X_n) \not\rightarrow L(X_1; \dots; X_n)$$

and checking that it is $((n + 1)k - c^0)$ -connected for some constant c^0 when the $X_j \not\rightarrow Y$ are k -connected. Therefore some readers may prefer to skip to Section 7 after the proof of 6.3.

A consequence of 6.1 is that what we are calling a second derivative (that is, the coefficient of the bilinear functor corresponding to a 2-homogeneous layer) can actually be seen as the derivative of a derivative. More generally, we have the following useful interpretation of 6.1:

6.3 Corollary

$$\partial_{y_1, \dots, y_{p+q}}^{(p+q)} F(Y) = \partial_{y_1, \dots, y_p}^{(p)} \partial_{y_{p+1}, \dots, y_{p+q}}^{(q)} F(Y):$$

Explanation and sketch of proof If F is a homotopy functor from spaces to either based spaces or spectra, then, as we have already observed, the spectrum $\partial_{y_{p+1}, \dots, y_{p+q}}^{(q)} F(Y)$ depends functorially on $Y = (Y; y_{p+1} \dots y_{p+q})$ with appropriate definitions. In stating 6.3 we are extending the "partial derivative" notation to cover functors of "spaces with several base points". Thus, for example, if F is a functor of based spaces then $\partial_y F(Y; y_0)$ depends functorially on $(Y; y; y_0)$ and is defined as the coefficient spectrum of the linear part of the functor

$$Z \vee F(Y \underset{y}{-} Z):$$

Here $y_0 \in Y$ is serving as basepoint in $Y \underset{y}{-} Z$ for purposes of applying the functor F while y is serving as basepoint in Y for wedging with Z .

In view of 6.1, to prove 6.3 one has only to see that for a functor of $p+q$ based spaces, such as

$$(Z_1; \dots; Z_{p+q}) \vee F(Y \underset{y_1}{-} Z_1; \dots; Y \underset{y_n}{-} Z_{p+q}):$$

the following two processes are equivalent: (1) reducing in all variables followed by multilinearizing in all variables, (2) reducing and multilinearizing in the last q variables, followed by reducing and multilinearizing in the first p variables. This is easy. □

Another consequence of 6.1 is:

6.4 Corollary F is m -reduced if and only if F is reduced and for every $0 < n < m$ the multilinearization of the n^{th} cross-effect of F is contractible.

Proof This is clear from 6.1 and the fact that $cr_n D_n F$ determines $D_n F$. □

The proof of 6.1 is connected with the idea of "multivariable Taylor series" (Remark 1.22). A key observation is that the product category $C_{Y_1} \times \dots \times C_{Y_n}$ is itself the category of spaces over a space: it is equivalent to $C_{Y_1 \sqcup \dots \sqcup Y_n}$ where the \sqcup denotes disjoint union. In this equivalence of categories, n -tuples of weak equivalences correspond to weak equivalences and n -tuples of cocartesian cubes correspond to cocartesian cubes. When it is necessary to distinguish between a functor

$$C_{Y_1} \times \dots \times C_{Y_n} \xrightarrow{F} D$$

of n variables and the associated functor

$$C_{Y_1, \dots, Y_n} \dashv D$$

of one variable, we will denote the latter by F .

6.1 is proved using the following statement, which will be proved at the end of Section 6:

6.5 Lemma *If $C_{Y_1, \dots, Y_n} \dashv D$ is $(1; \dots; 1)$ -reduced then $P_{1; \dots; 1} G \cong P_n G$.*

Proof of 6.1 Applying 6.5 in the special case $Y = Y_1 = \dots = Y_n$, with $G = cr_n F$, we find that what we need is a natural equivalence

$$P_n cr_n F \cong cr_n P_n F$$

This will follow from a natural equivalence $T_n cr_n F \cong cr_n T_n F$, which in turn will follow from a natural equivalence $J_U cr_n F \cong cr_n J_U F$, where (as in the proof of 1.8) J_U is composition with the n -ary join with a finite set U . cr_n is a three-step process: Compose with the (whiskered) sum

$$T_Y^n \dashv T_Y$$

then reduce in all variables (this was called r in the proof of 4.2), then compose with the equivalence of categories

$$T_Y \dashv T_Y^n$$

Each of these steps commutes with J_U up to natural equivalence. \square

To get to 6.5 we must revisit and strengthen some results that were discussed in Section 3. The following is a strengthening of [10; 3.4].

6.6 Lemma *If a homotopy functor $C_{Y_1, \dots, Y_n} \dashv D$ is $(d_1; \dots; d_n)$ -excisive then F is $(d_1 + \dots + d_n)$ -excisive.*

Proof In fact the proof of [10; 3.4] becomes a proof of 6.6 if one uses $(X_1; \dots; X_n)$ in place of $(X; \dots; X)$ throughout. \square

In particular F is n -excisive if F is $(1; \dots; 1)$ -excisive. We need to know also that F is n -homogeneous if F is multilinear. The following is a strengthening of 3.1.

6.7 Lemma If $C_{Y_1} \xrightarrow{F} C_{Y_n}$ and D is $(1; \dots; 1)$ -reduced, then F is n -reduced.

Proof This follows from the next statement as 3.1 followed from 3.2. \square

6.8 Lemma If $C_{Y_1} \xrightarrow{F} C_{Y_n}$ and D is $(1; \dots; 1)$ -reduced, then the map

$$F \xrightarrow{T_{n-1}} T_{n-1} \circ F$$

factors through a weakly contractible functor.

Proof Again, the proof of 3.2 applies with no change except $(X_1; \dots; X_n)$ for $(X; \dots; X)$. \square

We need this partial converse to 6.6:

6.9 Lemma If $C_{Y_1} \xrightarrow{L} C_{Y_n}$ and D is $(1; \dots; 1)$ -reduced and L is n -excisive, then L is multilinear.

Proof We have to show that L is 1-excisive in each variable. It is sufficient to consider the last variable. Fix an object X_j of C_{Y_j} for each $1 \leq j \leq n-1$ and let

$$\begin{array}{ccc} A & \longrightarrow & B \\ \downarrow & & \downarrow \\ Y & & Y \\ C & \longrightarrow & D \end{array}$$

be any cocartesian square in C_{Y_n} . We have to show that the square

$$\begin{array}{ccc} L(X_1; \dots; X_{n-1}; A) & \longrightarrow & L(X_1; \dots; X_{n-1}; B) \\ \downarrow & & \downarrow \\ L(X_1; \dots; X_{n-1}; C) & \longrightarrow & L(X_1; \dots; X_{n-1}; D) \end{array}$$

is cartesian.

Define an $(n+1)$ -cube in $C_{Y_1} \times \dots \times C_{Y_n}$ as follows: For each subset S of $\{1; \dots; n-1\}$, let $X_j(S)$ be Y_j if $j \in S$ and X_j if $j \notin S$.

Then $f(X_1(S); \dots; X_{n-1}(S))g$ constitutes an $(n-1)$ -cube in $C_{Y_1} \times \dots \times C_{Y_{n-1}}$ and our $(n+1)$ -cube will consist of the squares

$$\begin{array}{ccc} (X_1(S); \dots; X_{n-1}(S); A) & \longrightarrow & (X_1(S); \dots; X_{n-1}(S); B) \\ \downarrow & & \downarrow \\ (X_1(S); \dots; X_{n-1}(S); C) & \longrightarrow & (X_1(S); \dots; X_{n-1}(S); D) \end{array}$$

The $(n + 1)$ -cube is strongly cocartesian, so L yields a cartesian cube. On the other hand, for each nonempty S the square

$$\begin{array}{ccc}
 L(X_1(S); \dots; X_{n-1}(S); A) & \longrightarrow & L(X_1(S); \dots; X_{n-1}(S); B) \\
 \downarrow \text{?} & & \downarrow \text{?} \\
 L(X_1(S); \dots; X_{n-1}(S); C) & \longrightarrow & L(X_1(S); \dots; X_{n-1}(S); D)
 \end{array}$$

is cartesian; in fact it is made up of contractible objects because L is reduced in each variable. It follows by 1.6 of [10] that the square corresponding to $S = \emptyset$ is also cartesian. □

Proof of 6.5 Now it is convenient to drop the distinction between F and \bar{F} . $P_{1, \dots, 1} F$ is the universal example of a $(1; \dots; 1)$ -excisive functor under F , and is also n -excisive (by 6.6). $P_n F$ is the universal example of an n -excisive functor under F , and is also $(1; \dots; 1)$ -excisive (by 6.9). It follows that they are the same. □

The reader, looking at 6.6 and 6.7, might have wondered about:

6.10 Lemma *If a homotopy functor $C_{Y_1} \xrightarrow{\bar{F}} C_{Y_n} \xrightarrow{D}$ is $(d_1; \dots; d_n)$ -reduced then \bar{F} is $(d_1 + \dots + d_n)$ -reduced.*

In fact this is true, and it can be deduced from 6.4.

7 Example: Suspension spectra of mapping spaces

For an unbased space X let ${}^1_+ X$ be the suspension spectrum of the based space X_+ obtained by adding an extra point to X . We will call this the unreduced suspension spectrum (and hope that this does not lead anyone to confuse the unreduced suspension of X with $S^1 \wedge X_+$). Similarly, if X is a space fibered over Y we can speak of its unreduced fiberwise suspension spectrum, meaning the fiberwise suspension spectrum of the fiberwise based space X , where as in Section 4 this means the disjoint union of X and Y considered as an object of T_Y .

For a finite CW complex K , the functor $F(X) = {}^1_+ X^K$ is analytic by [10; 4.4], and its first derivative was found in [9; 2.4]. We now find its n^{th} derivative. We begin by recalling what the formula for the first derivative is and where that formula came from.

In the notation of Section 5 the formula for the first derivative is

$$(7.1) \quad \partial_y \int_+ Y^K \stackrel{Z}{\int}_{k \geq K} \int_+ (Y; y)^{(K;k)} dk:$$

Recall that this means that the linearization of the functor

$$(7.2) \quad \int_+ \text{hoco ber}(\int_+ (Y \dashv_y Z)^K \dashv \int_+ Y^K) \dashv \int_+ \text{hoco ber}(Y^K \dashv (Y \dashv_y Z)^K)$$

is the functor

$$\int_+ \int_+ Z \wedge \stackrel{Z}{\int}_{k \geq K} \int_+ (Y; y)^{(K;k)} dk:$$

Because K is finite, this last can also be written

$$(7.3) \quad \int_+ \int_+ \int_{k \geq K} \int_+ (Z \wedge (Y; y)^{(K;k)} \dashv_+) dk:$$

Implicitly in 7.1 we are using a certain system of spectra on K , namely the unreduced fiberwise suspension spectrum of a certain space over K , let us call it W , whose fiber over $k \geq K$ is $W_k = (Y; y)^{(K;k)}$; W is the subspace of $Y^K \dashv K$ consisting of pairs $(f; k)$ such that $f(k) = y$. Likewise in 7.3 we are using the fiberwise suspension spectrum of a space over K , call it $Z \wedge_K W$, whose fiber is $Z \wedge W_{k+}$, namely

$$\text{colim}(Z \dashv W \dashv W \dashv K):$$

7.4 Remark A formula for the differential and not just the derivative was given in [9]. In the present notation it says that the unreduced-linear functor corresponding to $D_Y F$ takes the object $X \dashv Y$ to

$$\int_+ \int_+ \int_{k \geq K} \int_+ (Y; f(x))^{(K;k)} dk dx:$$

In other words, the linear functor is given by a coefficient system on Y which may be obtained by "integration over the fiber" from a system on $K \dashv Y$, the fiberwise unreduced suspension spectrum of the fibration $K \dashv Y^K \dashv K \dashv Y$ whose fiber over $(k; y)$ may be identified with $(Y; y)^{(K;k)}$.

The method of proof for 7.1 was this: First give a natural map from 7.2 to 7.3, then show that its connectivity is roughly twice that of the space Z .

To produce the map, it was enough to give a natural map of based spaces

$$\text{hoco ber}(Y^K \dashv (Y \dashv_y Z)^K) \dashv \int_+ \int_+ \int_{k \geq K} \int_+ (Z \wedge (Y; y)^{(K;k)} \dashv_+) dk:$$

This was done by means of a tautological map from $(Y \rightarrow Z)^K$ to the space of sections of $Z \wedge_K W \rightarrow K$. To specify this map we say where it sends the map $K \xrightarrow{f} Y \rightarrow Z$. Let f be the composed map

$$K \xrightarrow{f} Y \rightarrow Z \rightarrow Y$$

Then f is sent to the section whose value at k is $f(k) \wedge f \in Z \wedge (Y; y)^{(K; k)}_+$ if $f(k) \in Z$ and otherwise is the (otherwise) basepoint.

It is a good precaution to add a "whisker" to Z , replacing $Y \rightarrow Z$ by $Y \rightarrow Y \rightarrow Z$, before making the construction just described, to insure that the map $Z \wedge_Y W \rightarrow K$ is a fibration.

The proof that the resulting map from 7.2 to 7.3 is highly connected will not be repeated here.

There is a variant of 7.1 for spaces of based maps: If K has a basepoint k_0 , then we obtain

$$\int_c \int_{k_2 \in K - f k_0 g} \int_{y_0}^1 (Y; y_0)^{(K; k_0)} \int_{y_0}^1 (Y; y; y_0)^{(K; k; k_0)} dk$$

(This time the whisker is even more important, because it insures that even if $y = y_0$ the section being constructed will have compact support.)

Now we are in a position to compute a second derivative, using 6.2. We have:

$$\begin{aligned} & \int_{y_1, y_2} \int_{y_0}^1 Y^K \int_{y_1} \int_{k_2 \in K} \int_{y_0}^1 (Y; y_2)^{(K; k_2)} dk_2 \\ & \int_{y_1} \int_{k_2 \in K} \int_{y_0}^1 (Y; y_2)^{(K; k_2)} dk_2 \\ & \int_{k_2 \in K} \int_{k_1 \in K - f k_2 g} \int_{y_0}^1 (Y; y_1; y_2)^{(K; k_1; k_2)} dk_1 dk_2 \\ & \int_c \int_{(k_1; k_2) \in K^{(2)}} \int_{y_0}^1 (Y; y_1; y_2)^{(K; k_1; k_2)} d(k_1; k_2): \end{aligned} \tag{7.5}$$

The differentiation under the integral sign depends on the hypothesis that K is finite. $K^{(2)}$ is the complement of the diagonal in $K \times K$. The last step, replacing a double integral by a single integral, is a tautology.

The expression in the last line implicitly refers to a system of spectra on $K^{(2)}$ whose fiber at $(k_1; k_2)$ is $\int_{y_0}^1 (Y; y_1; y_2)^{(K; k_1; k_2)}$, namely the unreduced fiberwise suspension spectrum of a certain fibration

$$W^{[2]} \rightarrow K^{(2)}:$$

The space $W^{[2]}$ is the subspace of $Y^K \times K^{(2)}$ consisting of all $(f; k_1; k_2)$ such that $f(k_1) = y_1$ and $f(k_2) = y_2$, so that the fiber $W_{(k_1; k_2)}^{[2]}$ over $(k_1; k_2)$ is $(Y; y_1; y_2)^{(K; k_1; k_2)}$.

This is not enough to determine the quadratic functor D_2F_Y , because we do not yet know the \mathbb{Z}_2 -symmetry in the second derivative. We can make a good guess about that, because the last expression in 7.5 has an obvious symmetry. To verify the guess, we can proceed as follows:

We have to study the second-order cross-section of F_Y as applied to objects $Y \rightarrow Y$, in other words the total homotopy fiber of

$$(7.6) \quad \begin{array}{ccc} \underset{\substack{? \\ \dot{Y}}}{(Y \rightarrow_{y_1} Z_1 \rightarrow_{y_2} Z_2)^K} & \longrightarrow & \underset{\substack{? \\ \dot{Y}}}{(Y \rightarrow_{y_1} Z_1)^K} \\ \underset{\substack{? \\ \dot{Y}}}{(Y \rightarrow_{y_2} Z_2)^K} & \longrightarrow & \underset{\substack{? \\ \dot{Y}}}{(Y)^K} \end{array}$$

We know that the bilinearization of this is equivalent to

$$Z_1 \wedge_c Z_2 \wedge_{\mathbb{Z}_{(k_1; k_2)2K^{(2)}}} \underset{\substack{? \\ \dot{Y}}}{(Y; y_1; y_2)^{(K; k_1; k_2)}} \rightarrow d(k_1; k_2);$$

or equivalently

$$(7.7) \quad \underset{\substack{? \\ \dot{Y}}}{Z_{(k_1; k_2)2K^2} \wedge_c} \rightarrow \underset{\substack{? \\ \dot{Y}}}{(Z_1 \wedge Z_2 \wedge (Y; y_1; y_2)^{(K; k_1; k_2)})_+} \rightarrow d(k_1; k_2);$$

This last expression refers to the fiberwise suspension spectrum of a certain space over $K^{(2)}$ whose fibers are $Z_1 \wedge Z_2 \wedge W_{(k_1; k_2)_+}^{[2]}$: Call it $(Z_1 \wedge Z_2) \wedge_{K^{(2)}} W^{[2]}$.

We are seeking to show that that equivalence can be chosen to preserve the \mathbb{Z}_2 -symmetry, so we should look for a symmetry-preserving map from the total homotopy fiber of 7.6 to 7.7.

The total homotopy fiber of 7.6 can be rewritten as the suspension spectrum of the total cofiber of

$$(7.8) \quad \begin{array}{ccc} \underset{\substack{? \\ \dot{Y}}}{(Y)^K} & \longrightarrow & \underset{\substack{? \\ \dot{Y}}}{(Y \rightarrow_{y_2} Z_1)^K} \\ \underset{\substack{? \\ \dot{Y}}}{(Y \rightarrow_{y_2} Z_2)^K} & \longrightarrow & \underset{\substack{? \\ \dot{Y}}}{(Y \rightarrow_{y_1} Z_1 \rightarrow_{y_2} Z_2)^K} \end{array}$$

There is a tautological map from this total cofiber to the zeroth space of 7.7. It arises from a tautological map from $(Y \rightarrow_{y_1} Z_1 \rightarrow_{y_2} Z_2)^K$ to the space of

compactly supported sections of $(Z_1 \wedge Z_2) \wedge_Y W^{[2]} \neq K^{(2)}$. To specify this we say where it sends the map $K \xrightarrow{f} Y_{-y_1} Z_1 \rightarrow_{-y_2} Z_2$. Let f be the composed map

$$K \xrightarrow{f} Y_{-y_1} Z_1 \rightarrow_{-y_2} Z_2 \neq Y$$

Then f is sent to the section whose value at $(k_1; k_2)$ is $f(k_1) \wedge f(k_2) \wedge (f; k_1; k_2)$ if $f(k_1) \in Z_1$ and $f(k_2) \in Z_2$ and otherwise the (berwise) basepoint.

We claim, leaving the remaining details to the reader, that this results in a map

$$\text{@}_{y_1; y_2}^{(2)} \quad \Gamma(Y^K) \neq \int_c \int_{(k_1; k_2) \in K^{(2)}} \Gamma(Y; y_1; y_2)^{(K; k_1; k_2)} d(k_1; k_2)$$

that corresponds to 7.5 under 6.2 and is therefore an equivalence.

The same method gives the n^{th} derivative. The conclusion is:

7.10 Theorem *For a finite complex K we have a symmetry-preserving equivalence*

$$\text{@}_{y_1; \dots; y_n}^{(n)} \quad \Gamma(Y^K) \neq \int_c \int_{k \in K^{(n)}} \Gamma(Y; y_1; \dots; y_n)^{(K; k_1; \dots; k_n)} dk$$

Here $K^{(n)}$ is the space of all ordered n -tuples $k = (k_1; \dots; k_n)$ of distinct points in K .

It is interesting to work out what this says in the case when K is a finite set of cardinality m , and to compare it with the formula

$$m(m-1) \dots (m-n+1) y^{m-n};$$

for the n^{th} derivative of y^m in ordinary calculus.

In the case when Y is a one-point space, the right-hand side of 7.10 becomes

$$\int_c \int_{k \in K^{(n)}} \Gamma(S^0) dk \quad \text{Map}(K^{(n)c}; \Gamma(S^0)) = (K^{(n)c})$$

In other words, the n^{th} derivative of $F(X) = \Gamma(X^K)$ at a point is the S -dual $(K^{(n)c})$ of a certain based n -space $K^{(n)c}$, the one-point compactification of $K^{(n)}$ (or the quotient of K^n by the fat diagonal $K^n - K^{(n)}$). The n^{th} homogeneous functor is

$$(D_n F)(X) = ((K^{(n)c}) \wedge X^{\wedge n})_{h_n};$$

and this can be identified with

$$\text{Map}(K^{(n)c}; \Gamma(X^{\wedge n}))_{h_n}$$

because $K^{(n)c}$ is finite. It can also be identified with

$$\text{Map}(K^{(n)c}; {}^1 X^{\wedge n})^n$$

because the group action on $K^{(n)c}$ is free.

The analogous conclusion for based K says

$$\int_c \int_{k \in (K - fk_0g)^{(n)}} \int_{y_1, \dots, y_n} {}^1 (Y; y_0)^{(K; k_0)} \int_{y_1, \dots, y_n} {}^1 (Y; y_0; y_1; \dots; y_n)^{(K; k_0; k_1; \dots; k_n)} dk:$$

When Y is a point we find that the n^{th} coefficient of $F(X) = \int_+ (X; x_0)^{(K; k_0)}$ is the S -dual of the one-point compactification of $(K - fk_0g)^{(n)}$.

The case when K is a based circle is particularly simple. Since $(S^1 - fk_0g)^{(n)}$ is the disjoint union of open n -cells freely and transitively permuted by the group, the formula for the n^{th} homogeneous layer of $\int_+ X$ becomes $\int_+ (X^{\wedge n})^n$, as already pointed out in 1.20.

8 Example: The identity functor

The identity functor $\mathcal{T} \not\downarrow \mathcal{T}$ from based spaces to based spaces is a central example, and its n^{th} derivative $\int^{(n)} I(\)$ is a basic object in homotopy theory. This spectrum with \int_n -action turns out to be S -dual to a certain finite complex with \int_n -action. We summarize and discuss some known results.

Note that the problem of determining the n^{th} derivative of the identity is equivalent to that of determining the n^{th} derivative of the functor \int_+ . In fact, by 5.3 we have

$$\int^{(n)} I(\) \simeq S^{V_n} \wedge \int^{(n)} I(\)$$

and therefore

$$\int^{(n)} I(\) \simeq V_n \int^{(n)} I(\)$$

To begin with the obvious, the first derivative of I is the sphere spectrum.

The second derivative can be identified by using the second cross-effect. The total homotopy fiber of

$$\begin{array}{ccc} X_1 & \xrightarrow{\int_+} & X_2 & \xrightarrow{\int_+} & X_1 \\ \int_+ \downarrow & & \int_+ \downarrow & & \int_+ \downarrow \\ \int_+ X_1 & & \int_+ X_2 & & \int_+ X_1 \\ \int_+ \downarrow & & \int_+ \downarrow & & \int_+ \downarrow \\ X_2 & \xrightarrow{\int_+} & X_2 & \xrightarrow{\int_+} & X_2 \end{array}$$

is the homotopy fiber of $X_1 \times X_2 \rightarrow X_1 \wedge X_2$, and the bilinearization of this is $Q(X_1 \wedge X_2)$, with the obvious Σ_2 -symmetry (trivial action on the loop coordinate), simply because there is a map (natural and symmetry-preserving)

$$\text{hofiber}(X_1 \times X_2 \rightarrow X_1 \wedge X_2) \rightarrow Q(X_1 \wedge X_2)$$

that is approximately $3k$ -connected when X_1 and X_2 are k -connected. It follows that $@^{(2)}I(\)$ is the -1 -sphere with trivial action.

For the n^{th} derivative, partial information can be obtained from the Hilton-Milnor theorem [15]. Recall that this describes the space $(X \times Y)$ as a weak product (direct limit of finite products) of factors each of which has the form $(X^{\wedge a} \wedge Y^{\wedge b})$. The factors for a given pair $(a; b)$ correspond to certain nested commutator expressions, an integral basis for the bidegree $(a; b)$ summand of a free Lie ring on two generators whose bidegrees are $(1; 0)$ and $(0; 1)$. Iteration yields a description of $(X_1 \times \dots \times X_n)$ as a weak product of factors of the form $(X_1^{\wedge a_1} \wedge \dots \wedge X_n^{\wedge a_n})$. The cross-effect $(cr_n I)(X_1; \dots; X_n)$ is the product of those factors for which $a_j = 1$ for all j . The multilinearized cross-effect sees only those factors for which $a_j = 1$ for all j . Thus the n^{th} differential of the functor $@^{(n)}I(\)$ is a product of copies of $Q(X_1 \wedge \dots \wedge X_n)$, and the n^{th} derivative is the product of a corresponding number of copies of the sphere spectrum. The number of copies is $(n-1)!$, and they correspond to a basis for the group $Lie(n)$ generated by all Lie monomials in n variables such that each variable occurs just once (and of course considered modulo the Jacobi identity and antisymmetry). A standard choice of basis for $Lie(n)$ consists of the monomials

$$[X_{(1)}; [X_{(2)}; [\dots; [X_{(n-1)}; X_n] \dots]]]$$

one for each permutation σ belonging to the subgroup Σ_{n-1} . It follows that the n^{th} derivative of $@^{(n)}I(\)$, regarded as a spectrum with Σ_{n-1} -action, is the suspension spectrum of the finite set $(\Sigma_{n-1})_+$.

Of course for determining $D_n I$ one needs the full Σ_n -action. The method outlined above even identifies the action of Σ_n on the homology of the spectrum $@^{(n)}I(\)$ (a free abelian group of rank $(n-1)!$ concentrated in degree $1-n$); it is the obvious action of Σ_n on $Lie(n)$, twisted by signs of permutations. But this is still insufficient for determining the homogeneous functor.

Johnson [11] defined a based finite complex K_n with Σ_n -action whose S -dual is $@^{(n)}I(\)$. Her K_n was designed to admit an interesting map

$$(cr_n I)(X_1; \dots; X_n) \rightarrow \text{Map}(K_n; X_1 \wedge \dots \wedge X_n);$$

both natural and symmetry-preserving, and she showed that after multilinearization this map leads to an equivalence

$$(D^{(n)} I)(X_1; \dots; X_n) \rightarrow \text{Map}(K_n; Q(X_1 \wedge \dots \wedge X_n));$$

Arone and Mahowald [4] later came up with another answer to the same question. It is defined in terms of the poset of all partitions (equivalence relations) on the set $\underline{n} = \{1, \dots, n\}$. The poset has a maximal element (the trivial partition) and a minimal element (the improper partition), so that the nerve of the poset is (for two reasons) contractible. The Arone-Mahowald version of K_n can be described as the double suspension of the nerve of the poset of proper nontrivial partitions, or alternatively as the quotient of the nerve of all partitions by the union of the nerve of the nontrivial partitions and the nerve of the proper partitions.

In [4] this model is justified by proving directly that it is (equivariantly) homotopy equivalent to Johnson's space, but it was actually discovered from a very different point of view, which is worked out in detail in [3]. There is a cosimplicial functor from spaces to spaces which has in degree d the functor Q^{d+1} , iterated composition of Q with itself. When applied to 1-connected spaces it serves as a resolution of the identity functor. (In general it gives the Bousfield-Kan integral completion functor.) For each d the functor Q^{d+1} has a split Taylor tower, which can be read off from the Snaithe splitting formula. It is rather easy to see that the n^{th} coefficient of the d^{th} functor is (functorially in d , up to homotopy) the S -dual of the (discrete) space of d -simplices in K_n . (The details worked out in [3] dispose of that unfortunate "\up to homotopy".)

Arone and Mahowald use this description of K_n to investigate the mod p homology of the spectrum $(\mathbb{Z}/p)_* \langle X \wedge \mathbb{Z}/p \rangle_n$ whose zeroth space is $(D_n I)(X)$. Their main results concern the case when X is a sphere. For simplicity take it to be an odd sphere. When n is not a power of p they find that the layer $(D_n I)(S^{2m-1})$ is p -locally trivial. (This is equivalent to the statement that the homology of \mathbb{Z}/p with coefficients in $Lie(n)$ localized at p is trivial.) When $n = p^k$ they use Dyer-Lashof operations to calculate the homology as a module over the Steenrod algebra, finding in particular that the layer $(D_{p^k} I)(S^{2m-1})$ has trivial v_j -periodic homotopy when $j < k$.

For further insight into these matters, see [2].

We close with some remarks about the derivatives of the identity at a general space Y . Recall from the discussion following 6.3 that the n^{th} derivative of $T \not\leftarrow T$ will be a functor of spaces with $n + 1$ basepoints, symmetric with respect to permutations of the last n points. It is clear that

$$\mathbb{Z}/p_*(Y; y_0) \cong \mathbb{Z}/p_* P_{y_0}^Y$$

where $P_{y_0}^Y$ is the path space $(Y; y_0; y)^{(l; 0; 1)}$. In general, the n^{th} derivative of the identity at an arbitrary space may be described in terms of the n^{th}

derivative of the identity at a point. For example, given a space Y and points y_0, y_1, y_2 , the total homotopy fiber (with respect to y_0) of

$$\begin{array}{ccc} Y_{-y_1} Z_1 \xrightarrow{?} Y_{-y_1} Z_1 & \xrightarrow{!} & Y_{-y_1} Z_1 \\ \downarrow \scriptstyle{?} & & \downarrow \scriptstyle{?} \\ Y_{-y_2} Z_2 & \xrightarrow{!} & Y \end{array}$$

is equivalent to the total homotopy fiber of

$$\begin{array}{ccc} (P_{y_0}^{y_1} Y_+ \wedge Z_1) \xrightarrow{?} (P_{y_0}^{y_2} Y_+ \wedge Z_2) & \xrightarrow{!} & P_{y_0}^{y_1} Y_+ \wedge Z_1 \\ \downarrow \scriptstyle{?} & & \downarrow \scriptstyle{?} \\ P_{y_0}^{y_2} Y_+ \wedge Z_2 & \xrightarrow{!} & \end{array}$$

(The second square consists essentially of the homotopy fibers over $y_0 \in Y$ of the spaces in the first square.) Bilinearizing with respect to Z_1 and Z_2 , we find that

$$(D_{(Y; y_0)}^{(2)} I)(Y_{-y_1} Z_1; Y_{-y_2} Z_2) = (D^{(2)} I)(P_{y_0}^{y_1} Y_+ \wedge Z_1; P_{y_0}^{y_2} Y_+ \wedge Z_2):$$

The same argument applies for any n and yields a natural and symmetrical equivalence

$$@_{y_1, \dots, y_n}^{(n)}(Y; y_0) = P_{y_0}^{y_1} Y_+ \wedge \dots \wedge P_{y_0}^{y_n} Y_+ \wedge K_n$$

These observations can be used to give an alternative to the Hilton-Milnor argument above; if one is willing to settle for $(n-1)$ -symmetry rather than n -symmetry, then by 6.3 one can write

$$@_{y_1, \dots, y_n}^{(n)}(Y; y_0) = @_{y_1, \dots, y_{n-1}}^{(n-1)} + P_{y_0}^{y_n} Y_+$$

which by another very slight generalization of 7.10 is equivalent to

$$\int_c \sum_{k \geq 2(l-f_0; 1g)^{(n-1)}} + (Y; y_0; y_1, \dots, y_{n-1}; y_n)^{(l; 0; k_1, \dots, k_{n-1}; 1)} dk;$$

In the case $Y = \mathbb{S}^n$ this becomes the S -dual of the one-point compactification of $(l - f_0; 1g)^{(n-1)}$, in other words the S -dual of a wedge of $(n - 1)$ -spheres freely and transitively permuted by Σ_{n-1} .

In fact, by taking a different point of view, one can see that the n^{th} differential or derivative of the identity has a $(n+1)$ -symmetry and not just a n -symmetry, just as if the identity were itself a derivative. We hope to return to this point in a future paper.

9 Example: Waldhausen K-theory

Let A be Waldhausen's algebraic K -theory functor from spaces to spectra [15]. In [9] it was shown that

$$(9.1) \quad @_y A(Y) \quad \overset{1}{+} \quad Y;$$

We will give similar formulas for the higher derivatives of A .

Really what was shown in [9] was

$$(9.2) \quad @_y P^{Diff}(Y) \quad \overset{2}{+} \quad \overset{1}{+} \quad Y;$$

where P^{Diff} is stable smooth pseudoisotopy theory. Then 9.1 was a corollary in view of Waldhausen's relation

$$(9.3) \quad A(X) \quad Wh^{Diff} X \quad \overset{1}{+} \quad X;$$

where the Whitehead functor Wh^{Diff} satisfies $\overset{2}{+} \quad \overset{1}{+} \quad Wh^{Diff} X \quad P^{Diff} X$.

A natural map

$$P^{Diff} X \quad \overset{2}{+} \quad Q(X^{S^1} = X)$$

played a key role in obtaining 9.2, where $X^{S^1} = X$ is the quotient of the free loop space X^{S^1} by the constant loops. It was then clear that a more direct account of 9.1 ought to involve some analogous map

$$A(X) \quad \overset{1}{+} \quad L(X);$$

where $L(X) = \overset{1}{+} X^{S^1}$, a map that ought to have a K -theoretic rather than a manifold-theoretic description. It is not hard to make such a map (the *trace*), using the methods of [16].

9.4 Remark This was generalized by Bokstedt. The trace map is reminiscent of the Dennis trace map from the algebraic K -theory of a ring to its Hochschild homology, and this observation pointed the way to the generalization. Since $A(BG)$ can be defined as the algebraic K -theory of a generalized ring which may be thought of as the group ring of G over the sphere spectrum, and since the Hochschild homology of a group ring $k[G]$ is the homology of $(BG)^{S^1}$ with coefficients in k , it was natural to imagine that should be a special case of a construction

$$K(R) \quad \overset{1}{+} \quad THH(R)$$

defined for reasonable ring spectra R , where $THH(R)$ is a kind of "Hochschild homology over the sphere spectrum". Bokstedt invented (and named) the object THH and defined the trace (originally for a class of generalized rings called *functors with smash product*). See [6].

The trace induces a map

$$\textcircled{y}A(Y) \xrightarrow{\textcircled{y}!} \textcircled{y}L(Y)$$

By 7.1 we have

$$\textcircled{y}L(Y) \cong \int_{k \in S^1} \textcircled{y}_+(Y; y)^{(S^1; k)} dk;$$

Using rotations of the circle to continuously identify $(S^1; k)$ with $(S^1; 1)$, where 1 is one point in S^1 , this last spectrum may be identified with

$$\int_{k \in S^1} \textcircled{y}_+(Y; y)^{(S^1; 1)} dk = \text{Map}(S^1_+; \textcircled{y}_+ Y);$$

and thus split into two factors $\textcircled{y}_+ Y \times \textcircled{y}_+ Y$. Projecting on the first factor and composing with \textcircled{y}_+ we get a map

$$\textcircled{y}A(Y) \xrightarrow{\textcircled{y}_+} \textcircled{y}_+ Y;$$

which is in fact an equivalence.

9.5 Remark It would have been very tedious to prove this last fact directly from 9.2. That would have meant combining the proofs of 9.2 and 9.3 and the construction of the trace and undoubtedly dealing with several different models for $A(X)$. Instead in [5] we took a shortcut, observing that the functors $\textcircled{y}A(Y)$ and $\textcircled{y}_+ Y$ are abstractly equivalent by 9.1, and then arguing by universal examples that a natural map between them must be an equivalence for all Y if this is so in some special cases where things can be checked.

It is possible, and very convenient, to use a homotopy fixed point construction to single out the factor of $\textcircled{y}L(Y)$ that is to correspond to $\textcircled{y}A(Y)$. Consider the obvious action of the circle group T on L . The trace can easily be made to factor through the homotopy fixed point spectrum

$$A(Y) \xrightarrow{f} L(Y)^{hT} \xrightarrow{g} L(Y);$$

and it was shown in [5] that the map

$$(9.6) \quad \textcircled{y}A(Y) \xrightarrow{g} (\textcircled{y}L(Y))^{hT}$$

resulting from this refined trace \sim is an equivalence. This form of 9.1 will be very useful for getting the higher derivatives.

We emphasize that the right hand side of 9.6 is not $\textcircled{y}((L(Y))^{hT})$, and that L^{hT} is not an analytic functor. There is a canonical map

$$\textcircled{y}((L(Y))^{hT}) \xrightarrow{h} (\textcircled{y}L(Y))^{hT};$$

because T acts continuously on the homotopy functor L ; but this map does not happen to be an equivalence.

9.7 Theorem For all $n \geq 1$ the (natural and symmetry-preserving) map

$$\partial_{y_1, \dots, y_n}^{(n)} A(Y) \neq (\partial_{y_1, \dots, y_n}^{(n)} L(Y))^{hT}$$

induced by \sim is an equivalence.

Proof In proving this we are allowed to ignore the symmetry; thus for this purpose we can get away with using 6.3 to view n^{th} derivatives as first derivatives of $(n - 1)^{st}$ derivatives.

The key point now is that the operations ∂_y and $()^{hT}$, which did not commute when applied to $L(Y)$, do commute when applied to $\partial_{y_2, \dots, y_n}^{(n-1)} L(Y)$. Once this is established we can argue by induction on n :

$$\partial_{y_1} \partial_{y_2, \dots, y_n}^{(n-1)} A(Y) = \partial_{y_1} ((\partial_{y_2, \dots, y_n}^{(n-1)} L(Y))^{hT}) = (\partial_{y_1} \partial_{y_2, \dots, y_n}^{(n-1)} L(Y))^{hT} :$$

What is needed to establish it is the equation

$$(9.8) \quad \partial_{y_2, \dots, y_n}^{(n-1)} L(Y) = \int_c \int_{k_2(S^1)^{(n-1)}} \frac{1}{+} (Y; y_2, \dots, y_n)^{(S^1; k_2, \dots, k_n)} dk$$

(an instance of 7.10), together with the observation that T is acting freely on $(S^1)^{(n-1)}$.

Now apply the principle given at the end of Section 5, taking G to be T , P to be $(S^1)^{(n-1)}$ and B to be the orbit space. It is clear from 9.6 that for a suitable system E , functorial in $(Y; y_2, \dots, y_n)$, we have

$$\partial_{y_2, \dots, y_n}^{(n-1)} L(Y) = \int_{p \in 2P} E_{(p)} dp$$

equivariantly and

$$\begin{aligned} \partial_y ((\int_{p \in 2P} E_{(p)} dp)^{hT}) &= \int_{b \in 2B} \partial_y E_b db \\ &= \int_{b \in 2B} \partial_y E_b db \\ &= (\int_{p \in 2P} \partial_y E_{(p)} dp)^{hT} \\ &= \int_{p \in 2P} \partial_y E_{(p)} dp)^{hT} : \end{aligned}$$

The differentiations under the integral are justified by the finiteness of B and of P . (This step fails when $n = 1$, basically because the appropriate B would then be the space BT , which is not so finite.) □

In the special case $Y = \ast$ the conclusion of 9.7 is that the n^{th} derivative of A is the homotopy fixed point spectrum for T acting on the S -dual of the one-point compactification of $(S^1)^{(n)}$, where $(S^1)^{(n)}$ has the obvious commuting actions of T and Σ_n . Since the T -action is free, the answer may be rewritten as the S -dual of the one-point compactification of $(S^1)^{(n)} = T$. As a Σ_n -space, $(S^1)^{(n)} = T$ is isomorphic to $\Sigma_n \wr_{C_n} (S^1 \wr_{C_n} V_{C_n})$, where the cyclic group C_n acts on S^1 as a subgroup of the rotations and acts linearly on the vector space V_{C_n} by the reduced regular representation. It follows that the n^{th} derivative of A at a one-point space is induced from C_n and the n^{th} layer of the Taylor tower is given by

$$(D_n A)(X) = \text{Map}(S^1_+ \wedge S^{V_{C_n}}; \Sigma^{-1}(X^{\wedge n}))^{C_n}.$$

9.9 Remark In the special case when X is a suspension ΣY this becomes $\text{Map}(S^1_+; \Sigma^{-1}(X^{\wedge n}))^{C_n}$. In fact, in that case the Taylor tower splits:

$$A(\Sigma Y) = A(\Sigma) \times \prod_{n \geq 1} \text{Map}(S^1_+; \Sigma^{-1}(X^{\wedge n}))^{C_n}$$

if Y is connected. A proof of this was sketched in [7] and corrected in [5]. This special case of the conclusion was used in the shortcut mentioned in 9.5.

9.10 Remark There is also a direct K -theoretic approach to all of this. Dundas and McCarthy [8] generalized 9.1 to (generalized) rings. Their statement is that the trace map induces an equivalence from "stable K -theory" to THH . This led in [13] to an extension of the main result of [5] to such generalized rings.

References

- [1] **G Arone**, *A generalization of Snaitch-type splitting*, Trans. AMS 351 (1999) 1123{1150
- [2] **G Z Arone, W G Dwyer**, *Partition complexes, Tits buildings, and symmetric products*, Proc. London Math. Soc. 3 (2001) 229{256
- [3] **G Z Arone, M Kankaanrinta**, *A functorial model for iterated Snaitch splitting with applications to calculus of functors*, from: "Stable and unstable homotopy", AMS 1{30 (1996)
- [4] **G Arone, M Mahowald**, *The Goodwillie tower of the identity functor and the unstable periodic homotopy of spheres*, Invent. Math. 135 (1999) 743{788
- [5] **M Bokstedt, G Carlsson, R Cohen, T Goodwillie, W-C Hsiang, I Madsen**, *On the Algebraic K -Theory of Simply Connected Spaces*, Duke Math. J. 84 (1996) 541{563

- [6] **M Bokstedt, W-C Hsiang, I Madsen**, *The cyclotomic trace and algebraic K-theory of spaces*, *Invent. Math.* (1993) 463{539
- [7] **G Carlsson, R Cohen, T Goodwillie, W-C Hsiang**, *The free loop space and the algebraic K-theory of spaces*, *K-theory* 1 (1987) 53{82
- [8] **B Dundas, R McCarthy**, *Stable K-theory and topological Hochschild homology*, *Annals of Math.* 140 (1994) 685{702
- [9] **T Goodwillie**, *Calculus I: The First Derivative of Pseudoisotopy Theory*, *K-Theory* 4 (1990) 1{27
- [10] **T Goodwillie**, *Calculus II: Analytic Functors*, *K-Theory* 5 (1992) 295{332
- [11] **B Johnson**, *The derivatives of homotopy theory*, *Trans. AMS* 347 (1995) 1295{1321
- [12] **L G Lewis, J P May, M Steinberger**, *Equivariant Stable Homotopy Theory*, *Lecture Notes in Mathematics* 1213, Springer-Verlag (1986)
- [13] **R McCarthy**, *Relative algebraic K-theory and topological cyclic homology*, *Acta Math.* 179 (1997) 197{222
- [14] **S Mac Lane**, *Categories for the Working Mathematician*, Springer-Verlag (1971)
- [15] **F Waldhausen**, *Algebraic K-theory of spaces*, *Lecture Notes in Mathematics* 1126, Springer-Verlag (1985) 318{419
- [16] **F Waldhausen**, *Algebraic K-theory of topological spaces, II*, *Lecture Notes in Mathematics* 763, Springer-Verlag (1979) 356{394
- [17] **G Whitehead**, *Elements of Homotopy Theory*, Springer-Verlag (1978)