

Geometry & Topology
 Volume 8 (2004) 1032–1042
 Published: 7 August 2004



Weighted L^2 -cohomology of Coxeter groups based on barycentric subdivisions

BORIS L OKUN

*Department of Mathematical Sciences
 University of Wisconsin–Milwaukee
 Milwaukee, WI 53201, USA*

Email: okun@uwm.edu

Abstract

Associated to any finite flag complex L there is a right-angled Coxeter group W_L and a contractible cubical complex Σ_L (the Davis complex) on which W_L acts properly and cocompactly, and such that the link of each vertex is L . It follows that if L is a generalized homology sphere, then Σ_L is a contractible homology manifold. We prove a generalized version of the Singer Conjecture (on the vanishing of the reduced weighted $L^2_{\mathfrak{q}}$ -cohomology above the middle dimension) for the right-angled Coxeter groups based on barycentric subdivisions in even dimensions. We also prove this conjecture for the groups based on the barycentric subdivision of the boundary complex of a simplex.

AMS Classification numbers Primary: 58G12

Secondary: 20F55, 57S30, 20F32, 20J05

Keywords: Coxeter group, aspherical manifold, barycentric subdivision, weighted L^2 -cohomology, Tomei manifold, Singer conjecture

Proposed: Benson Farb
 Seconded: Walter Neumann, Steven Ferry

Received: 15 March 2004
 Revised: 3 August 2004

1 Introduction

A construction of Davis ([1], [2], [4]), associates to any finite flag complex L , a “right-angled” Coxeter group W_L and a contractible cubical cell complex Σ_L on which W_L acts properly and cocompactly. W_L has the following presentation: the generators are the vertices of L , each generator has order 2, and two generators commute if they span an edge in L . The most important feature of this construction is that the link of each vertex of Σ_L is isomorphic to L . A simplicial complex L is a *generalized homology m -sphere* (for short, a GHS^m) if it is a homology m -manifold having the same homology as a standard sphere \mathbb{S}^m (the homology is with real coefficients.) It follows, that if L is a GHS^{n-1} , then Σ_L is a homology n -manifold.

If L is a simplicial complex, bL will denote the barycentric subdivision of L . bL is a flag simplicial complex. Let $\partial\Delta^n$ denote the boundary complex of the standard n -dimensional simplex.

We study a certain weighted L^2 -cohomology theory $L^2_{\mathbf{q}}\mathcal{H}^*$, described in [7], [5]. Suppose, for each vertex $v \in L$ we are given a positive real number q_v , and let \mathbf{q} denote the vector with components q_v . Given a minimal word $w = v_1 \dots v_n \in W_L$, let $\mathbf{q}^w = q_{v_1} \dots q_{v_n}$. For each W_L -orbit of cubes pick a representative σ_0 and let $w(\sigma) = w$ if $\sigma = w\sigma_0$. (The ambiguity in the choices will not matter in our discussion.) Let $L^2_{\mathbf{q}}C^i(\Sigma_L) = \{\sum c_{\sigma}\sigma \mid \sum c_{\sigma}^2 \mathbf{q}^{w(\sigma)} < \infty\}$ be the Hilbert space of infinite i -cochains, which are square-summable with respect to the weight \mathbf{q}^w . The usual coboundary operator d is then a bounded operator, and we define the reduced weighted $L^2_{\mathbf{q}}$ -cohomology to be $L^2_{\mathbf{q}}\mathcal{H}^i(\Sigma_L) = \text{Ker}(d^i) / \overline{\text{Im}(d^{i-1})}$. Similarly, one can define the reduced weighted $L^2_{\mathbf{q}}$ -homology, except, instead of the usual boundary operator one uses the adjoint of d . It follows from the Hodge decomposition that the resulting homology and cohomology spaces are naturally isomorphic. These spaces are Hilbert modules over the Hecke-von Neumann algebra $\mathcal{N}_{\mathbf{q}}$ (an appropriately completed Hecke algebra of W_L .) This allows us to introduce the weighted $L^2_{\mathbf{q}}$ Betti numbers — the dimension of $L^2_{\mathbf{q}}\mathcal{H}^i$ over $\mathcal{N}_{\mathbf{q}}$. If $\mathbf{q} = \mathbf{1} = (1, \dots, 1)$, we obtain the usual reduced L^2 -cohomology, and we omit the index \mathbf{q} . We write $\mathbf{q} \leq \mathbf{1}$, if each component of \mathbf{q} is ≤ 1 .

The following conjecture, attributed to Singer, goes back to 1970’s.

The Singer Conjecture If M^n is a closed aspherical manifold, then

$$L^2\mathcal{H}^i(\widetilde{M}^n) = 0 \text{ for all } i \neq n/2.$$

As explained in [5, Section 14], the appropriate generalization of the Singer Conjecture to the weighted case is the following conjecture:

The Generalized Singer Conjecture Suppose L is a flag GHS^{n-1} . Then $L_{\mathbf{q}}^2 \mathcal{H}^i(\Sigma_L) = 0$ for $i > n/2$ and $\mathbf{q} \leq \mathbf{1}$.

(Poincaré duality shows that for $\mathbf{q} = \mathbf{1}$ this conjecture implies the Singer Conjecture for Σ .)

This conjecture holds true for $n \leq 4$ by [5]. One of the main results of this paper is a proof of this conjecture for barycentric subdivisions in even dimensions. The proof uses a reduction to a very special case $L = b\partial\Delta^{2k-1}$.

We prove this case as Theorem 5.2. It turns out (Theorem 5.3), that this result implies the vanishing of the $L_{\mathbf{q}}^2$ -cohomology in a certain range for arbitrary right-angled Coxeter groups based on barycentric subdivisions. (For $\mathbf{q} = \mathbf{1}$, this implication is proved in [6].) In particular, it follows that the Generalized Singer Conjecture is true for all barycentric subdivisions in even dimensions (Theorem 5.4), and for $b\partial\Delta^n$ in all dimensions (Theorem 5.6).

This paper relies very heavily on [5]. In the inductive proofs we mostly omit the first steps, they are easy exercises using [5].

The author wishes to thank Mike Davis, Jan Dymara and Tadeusz Januszkiewicz for several illuminating discussions.

2 Vanishing conjectures

We will follow the notation from [5]. Given a flag complex L and a full subcomplex A , set:

$$\begin{aligned} \mathfrak{h}_i^{\mathbf{q}}(L) &= L^2 \mathcal{H}_i(\Sigma_L) \\ \mathfrak{h}_i^{\mathbf{q}}(A) &= L^2 \mathcal{H}_i(W_L \Sigma_A) \\ \mathfrak{h}_i^{\mathbf{q}}(L, A) &= L^2 \mathcal{H}_i(\Sigma_L, W_L \Sigma_A) \\ \mathfrak{b}_{\mathbf{q}}^i(L) &= \dim_{\mathcal{N}_{\mathbf{q}}}(\mathfrak{h}_i^{\mathbf{q}}(L)) \\ \mathfrak{b}_{\mathbf{q}}^i(L, A) &= \dim_{\mathcal{N}_{\mathbf{q}}}(\mathfrak{h}_i^{\mathbf{q}}(L, A)) \end{aligned}$$

The dimension of Σ_L is one greater than the dimension of L . Hence, $\mathfrak{b}_{\mathbf{q}}^i(L) = 0$ for $i > \dim L + 1$.

We will use the following three properties of $L_{\mathbf{q}}^2$ -homology.

Proposition 2.1 (See [5, Section 15])

The Mayer–Vietoris sequence *If $L = L_1 \cup L_2$ and $A = L_1 \cap L_2$, where L_1 and L_2 (and therefore, A) are full subcomplexes of L , then*

$$\rightarrow \mathfrak{h}_i^{\mathfrak{q}}(A) \rightarrow \mathfrak{h}_i^{\mathfrak{q}}(L_1) \oplus \mathfrak{h}_i^{\mathfrak{q}}(L_2) \rightarrow \mathfrak{h}_i^{\mathfrak{q}}(L) \rightarrow$$

is weakly exact.

The Künneth Formula *The Betti numbers of the join of two complexes are given by:*

$$b_{\mathfrak{q}}^k(L_1 * L_2) = \sum_{i+j=k} b_{\mathfrak{q}}^i(L_1) b_{\mathfrak{q}}^j(L_2).$$

Poincaré Duality *If L is a flag GHS^{n-1} , then $b_{\mathfrak{q}}^i(L) = b_{\mathfrak{q}-1}^{n-i}(L)$.*

If σ is a simplex in L , let L_{σ} denote the link of σ in L . To simplify notation we will write bL_v instead of $(bL)_v$ to denote the link of the vertex v in bL . Let \mathcal{C} be a class of GHS's closed under the operation of taking link of vertices, i.e. if $S \in \mathcal{C}$ and v is a vertex of S then $S_v \in \mathcal{C}$. Following Section 15 of [5] we consider several variations of the Generalized Singer Conjecture for the class \mathcal{C} .

I(n) *If $S \in \mathcal{C}$ and $\dim S = n - 1$, then $b_{\mathfrak{q}}^i(S) = 0$ for $i > n/2$ and $\mathfrak{q} \leq \mathbf{1}$.*

III'(2k + 1) *Let $S \in \mathcal{C}$ and $\dim S = 2k$. Let v be a vertex of S . Then the map $i_*: \mathfrak{h}_k^{\mathfrak{q}}(S_v) \rightarrow \mathfrak{h}_k^{\mathfrak{q}}(S)$, induced by the inclusion, is the zero homomorphism for $\mathfrak{q} \geq \mathbf{1}$.*

V(n) *Let $S \in \mathcal{C}$ and $\dim S = n - 1$. Let A be a full subcomplex of S .*

- *If $n = 2k$ is even, then $b_{\mathfrak{q}}^i(S, A) = 0$ for all $i > k$ and $\mathfrak{q} \leq \mathbf{1}$.*
- *If $n = 2k + 1$ is odd, then $b_{\mathfrak{q}}^i(A) = 0$ for all $i > k$ and $\mathfrak{q} \leq \mathbf{1}$.*

The argument in Section 16 of [5] goes through without changes if we consider only GHS's from a class \mathcal{C} to give the following:

Theorem 2.2 (Compare [5, Section 16]) *If we only consider GHS's from a class \mathcal{C} , then the following implications hold.*

- (1) $\mathbf{I}(2k + 1) \implies \mathbf{III}'(2k + 1)$.
- (2) $\mathbf{V}(n) \implies \mathbf{I}(n)$.
- (3) $\mathbf{V}(2k - 1) \implies \mathbf{V}(2k)$.
- (4) $[\mathbf{V}(2k) \text{ and } \mathbf{III}'(2k + 1)] \implies \mathbf{V}(2k + 1)$.

Let \mathcal{JD} denote the class of finite joins of the barycentric subdivisions of the boundary complexes of standard simplices:

$$\mathcal{JD} = \{b\partial\Delta^{n_1} * \dots * b\partial\Delta^{n_j}\}.$$

Lemma 2.3 *The class \mathcal{JD} is closed under the operation of taking link of vertices.*

Proof Let $S = b\partial\Delta^{n_1} * \dots * b\partial\Delta^{n_j}$ and $v \in S$. We can assume that $v \in b\partial\Delta^{n_1}$. Then $S_v = b\partial\Delta_v^{n_1} * b\partial\Delta^{n_2} * \dots * b\partial\Delta^{n_j} = b\partial\Delta^{\dim(v)} * b\partial\Delta^{n_1 - \dim(v) - 1} * b\partial\Delta^{n_2} * \dots * b\partial\Delta^{n_j}$, and therefore $S \in \mathcal{JD}$. □

Next, consider the following statement:

III''(2k+1) *Let v be a vertex of $b\partial\Delta^{2k+1}$. Then the map $i_*: \mathfrak{h}_k^{\mathfrak{q}}(b\partial\Delta_v^{2k+1}) \rightarrow \mathfrak{h}_k^{\mathfrak{q}}(b\partial\Delta^{2k+1})$, induced by the inclusion, is the zero homomorphism for $\mathfrak{q} \geq 1$.*

Lemma 2.4 **III''(2k+1) \implies III'(2k+1)** *for the class \mathcal{JD} .*

Proof By induction, we can assume that the lemma holds for all odd numbers $< 2k+1$. Then it follows from the Theorem 2.2 that **V(m)** and therefore **I(m)** hold for all $m < 2k+1$.

Let $S = b\partial\Delta^{n_1} * \dots * b\partial\Delta^{n_j}$ with $n_1 + \dots + n_j = 2k+1$ and $v \in S$. We assume that $v \in b\partial\Delta^{n_1}$. Then $S_v = b\partial\Delta_v^{n_1} * b\partial\Delta^{n_2} * \dots * b\partial\Delta^{n_j}$ and, by the Künneth formula, the map in question decomposes as the direct sum of maps of the form

$$(\mathfrak{h}_{k_1}^{\mathfrak{q}}(b\partial\Delta_v^{n_1}) \rightarrow \mathfrak{h}_{k_1}^{\mathfrak{q}}(b\partial\Delta^{n_1})) \otimes \bigotimes_{i=2}^j (\mathfrak{h}_{k_i}^{\mathfrak{q}}(b\partial\Delta^{n_i}) \rightarrow \mathfrak{h}_{k_i}^{\mathfrak{q}}(b\partial\Delta^{n_i}))$$

where $k_1 + \dots + k_j = k$. Since $n_1 + \dots + n_j = 2k+1$ it follows that $k_i < n_i/2$ for some index i . If $n_i < 2k+1$, then the range of the corresponding map in the above tensor product is 0 by **I(n_i)** and Poincaré duality, and therefore the tensor product map is 0. If $n_i = 2k+1$ then, in fact, $i = 1$ (the join is a trivial join) and the result follows from **III''(2k+1)**. □

Thus, it follows from Theorem 2.2, Lemmas 2.3 and 2.4, and induction on dimension, that in order to prove the Generalized Singer Conjecture for the class \mathcal{JD} all we need is to prove **III''(2k+1)**.

3 Removal of an odd-dimensional vertex

Let L be a simplicial complex and bL be its barycentric subdivision. The vertices of bL are naturally graded by "dimension": each vertex v of bL is the barycenter of a unique cell (which we still denote v) of the complex L , and we call the dimension of this cell the *dimension* of the vertex v . Let E_L denote the subcomplex of bL spanned by the even dimensional vertices. Let \mathcal{A}_L denote the set of full subcomplexes A of L containing E_L , which have the following property: if A contains a vertex of odd dimension $2j + 1$, then A contains all vertices of bL of dimensions $\leq 2j$. In other words, any such A can be obtained inductively from bL by repeated removal of an odd-dimensional vertex of the highest dimension.

If $L = \partial\Delta^n$ we will use the notation $E_n = E_L$ and $\mathcal{A}_n = \mathcal{A}_L$.

Lemma 3.1 *Assume $\mathbf{III}''(2m + 1)$ holds for $2m + 1 < n$. Then for any $(n - 1)$ -dimensional simplicial complex L and any complex $A \in \mathcal{A}_L$ we have:*

$$b_{\mathbf{q}}^i(A) = b_{\mathbf{q}}^i(bL) = 0 \text{ for } i > (n + 1)/2 \text{ and } \mathbf{q} \leq \mathbf{1}.$$

Proof By induction, we can assume that the lemma holds for all $m < n$. First, we claim that removal of odd-dimensional vertices does not change the homology above $(n + 1)/2$. Let $A \in \mathcal{A}_L$ and let $B = A - v$ where v is a vertex of the highest odd dimension of A . We let $\dim(v) = 2d - 1$, $1 \leq d \leq k$. We want to prove that $b_{\mathbf{q}}^i(A) = b_{\mathbf{q}}^i(B)$ for $i > (n + 1)/2$. Consider the Mayer-Vietoris sequence of the union $A = B \cup_{A_v} CA_v$:

$$\rightarrow \mathfrak{h}_i^{\mathbf{q}}(A_v) \rightarrow \mathfrak{h}_i^{\mathbf{q}}(B) \oplus \mathfrak{h}_i^{\mathbf{q}}(CA_v) \rightarrow \mathfrak{h}_i^{\mathbf{q}}(A) \rightarrow \mathfrak{h}_{i-1}^{\mathbf{q}}(A_v)$$

Suppose $i > (n + 1)/2$. Since $A_v = B \cap bL_v = B \cap (b\partial\Delta^{2d-1} * b(L_v))$, and since $B \in \mathcal{A}_L$, it follows, by construction, that A_v splits as the join:

$$A_v = b\partial\Delta^{2d-1} * A_1,$$

with $A_1 \in \mathcal{A}_{(L_v)}$. By inductive assumption the lemma holds for L_v , i.e. $b_{\mathbf{q}}^i(A_1) = b_{\mathbf{q}}^i(b(L_v)) = 0$ for $i > (n + 1)/2 - d$.

Since $\mathbf{III}''(2d - 1)$ holds by hypothesis, by Lemma 2.4 and Theorem 2.2, $\mathbf{I}(2d - 1)$ holds for the class \mathcal{JD} , and, thus, $b_{\mathbf{q}}^i(b\partial\Delta^{2d-1}) = 0$ for $i \geq d$.

Then, by the Künneth formula, $b_{\mathbf{q}}^{i-1}(A_v) = 0$ for $i - 1 \geq (n + 1)/2$, i.e. for $i > (n + 1)/2$. By [5, Proposition 15.2(d)], $b_{\mathbf{q}}^i(CA_v) = \frac{1}{q_v+1} b_{\mathbf{q}}^i(A_v)$. Therefore in the above sequence the terms corresponding to A_v and CA_v are 0, and

the claim follows. Then it follows by induction, that $b_{\mathbf{q}}^i(A) = b_{\mathbf{q}}^i(bL)$ for all $A \in \mathcal{A}_L$ and $i > (n + 1)/2$.

To prove the vanishing we note that, in particular, $b_{\mathbf{q}}^i(E_L) = b_{\mathbf{q}}^i(bL)$ for $i > (n + 1)/2$. Since E_L is spanned by the even-dimensional vertices of bL and since a simplex in bL has vertices of pairwise different dimensions, we have $\dim(E_L) = [(n + 1)/2] - 1$. Therefore, $b_{\mathbf{q}}^i(E_L) = 0$ for $i > (n + 1)/2$ and we have proved the lemma. \square

In the special case $L = \Delta^{2k+1}$ this lemma admits the following strengthening:

Lemma 3.2 *Let $n = 2k + 1$. Assume $\mathbf{III}''(2m + 1)$ holds for $2m + 1 < n$. Then for any complex $A \in \mathcal{A}_n$, $A \subset b\partial\Delta^n$, we have:*

$$b_{\mathbf{q}}^i(A) = b_{\mathbf{q}}^i(b\partial\Delta^n) \text{ for } i > k \text{ and } \mathbf{q} \leq \mathbf{1}.$$

Proof We proceed as in the previous proof. As before, we have $B = A - v$, $\dim(v) = 2d - 1$ and $A_v = b\partial\Delta^{2d-1} * A_1$, where now $A_1 \subset b\partial\Delta^{2k+1-2d}$. Therefore, the inductive assumption and the hypothesis on $\mathbf{III}''(2d - 1)$ imply that $b_i(A_1) = 0$ for $i > k + d$. The lemma follows as before. \square

As explained in [6], when $\mathbf{q} = \mathbf{1}$, the removal of the odd-dimensional vertex does not change homology in *all* dimensions. We record this result below.

Lemma 3.3 *Assume $\mathbf{III}''(2m + 1)$ holds for $2m + 1 < n$ and $\mathbf{q} = \mathbf{1}$. Then for any $(n - 1)$ -dimensional simplicial complex L and for any complex $A \in \mathcal{A}_L$, obtained by the repeated removal of highest odd-dimensional vertices, we have:*

$$b^*(A) = b^*(bL).$$

Proof Again we repeat the proof of Lemma 3.1. As before, we have the splitting $A_v = b\partial\Delta^{2d-1} * A_1$. The point now is that for $\mathbf{q} = \mathbf{1}$, $\mathbf{I}(2d - 1)$ and Poincaré duality imply $b^*(b\partial\Delta^{2d-1}) = 0$ and therefore $b^*(A_v) = 0$ by the Künneth formula. \square

4 Intersection form

Lemma 4.1 *Let L be a GHS^{2k} and let v be a vertex of L . Then the image of the restriction map on L^2 -cohomology $i^*: L^2\mathcal{H}^k(\Sigma_L) \rightarrow L^2\mathcal{H}^k(\Sigma_{L_v})$ is an isotropic subspace of the intersection form of Σ_{L_v} .*

Proof Note that the cup product of two L^2 -cocycles is an L^1 -cocycle. The intersection form is the result of evaluation of the cup product of two middle-dimensional cocycles on the fundamental class, which is L^∞ . Since Σ_{L_v} bounds a half-space in Σ_L , $i_*([\Sigma_{L_v}]) = 0$ in L^∞ -homology of Σ_L . Thus, if $a, b \in L^2\mathcal{H}^n(\Sigma_L)$, then $\langle i^*(a) \cup i^*(b), [\Sigma_{L_v}] \rangle = \langle a \cup b, i_*([\Sigma_{L_v}]) \rangle = 0$. \square

Lemma 4.2 *Let G be a group and let A be a bounded G -invariant (with respect to the diagonal action) non-degenerate bilinear form on a Hilbert submodule $M \subset \ell^2(G)$. Then A has no nontrivial G -invariant isotropic subspaces.*

Proof Let us consider the case $M = \ell^2(G)$ first. G -invariance and continuity of the form A implies that A is completely determined by its values $a_g = (g A 1)$, $g \in G$. It is convenient to think of the form as given by $(x A y) = \langle x, Ay \rangle$, where $\langle \cdot, \cdot \rangle$ is the inner product and $A = \sum_{g \in G} a_g g$ is a bounded G -equivariant operator on $\ell^2(G)$. Non-degeneracy of A means that $Ax = 0$ only if $x = 0$. A is the limit of the group ring elements, and Ax is the limit of the corresponding linear combinations of G -translates of x , i.e. $Ax = \lim \sum_{g \in G_n} a_g(gx)$, where G_n is some exhaustion of G by finite sets. It follows that if x belongs to G -invariant isotropic subspace, then Ax belongs to the closure of this subspace. Thus, we have $\langle Ax, Ax \rangle = (Ax A x) = 0$ by isotropy and continuity, therefore $x = 0$.

The case of general submodule $M \subset \ell^2(G)$ reduces to the above, since the bilinear form A can be extended to $\ell^2(G)$, for example, by taking the orthogonal sum $A \oplus \langle \cdot, \cdot \rangle$ of A on M and the inner product on the orthogonal complement of M . \square

5 Vanishing theorems

Our main technical results are the following two theorems.

Theorem 5.1 $\text{III}'(2k + 1)$ is true for all $k > 0$ and $\mathbf{q} = \mathbf{1}$.

Proof The proof is by induction on k . Suppose the theorem is true for all $m < k$.

Let v be a vertex of $b\partial\Delta^{2k+1}$. We need to show that the restriction map $i^*: \mathfrak{h}^k(b\partial\Delta^{2k+1}) \rightarrow \mathfrak{h}^k(b\partial\Delta_v^{2k+1})$ is the 0-map.

First let us suppose that v is a vertex of dimension 0, i.e. a vertex of Δ^{2k+1} .

Consider the action of the symmetric group \mathbf{S}_{2k+1} on Δ^{2k+1} which fixes the vertex v and permutes other vertices. This action gives a simplicial action of \mathbf{S}_{2k+1} on $b\partial\Delta^{2k+1}$ and therefore, after choosing a base point, lifts to a cubical action of \mathbf{S}_{2k+1} on $\Sigma_{b\partial\Delta^{2k+1}}$ stabilizing $\Sigma_{b\partial\Delta_v^{2k+1}}$. Let G' be the group of cubical automorphisms of $\Sigma_{b\partial\Delta^{2k+1}}$ generated by this action and the standard action of $W_{b\partial\Delta^{2k+1}}$, and let G be the orientation-preserving subgroup of G' . Similarly, let G'_v be the group of cubical automorphisms of $\Sigma_{b\partial\Delta_v^{2k+1}}$ generated by this action and the standard action of $W_{b\partial\Delta_v^{2k+1}}$, and let G_v be the orientation-preserving subgroup of G'_v .

We claim that, as a Hilbert G_v -module $L^2\mathcal{H}^k(\Sigma_{b\partial\Delta_v^{2k+1}})$, is a submodule of $\ell^2(G_v)$. Note that $b\partial\Delta_v^{2k+1}$ is naturally isomorphic to $b\partial\Delta^{2k}$.

Using the inductive assumption and Lemma 3.3, we can remove from $b\partial\Delta^{2k}$ all odd-dimensional vertices without changing the L^2 -cohomology: $\mathfrak{h}^*(E_{2k}) = \mathfrak{h}^*(b\partial\Delta^{2k})$. Since the action of \mathbf{S}_{2k+1} on $b\partial\Delta_v^{2k+1} = b\partial\Delta^{2k}$ preserves the dimension of the vertices, we have isomorphism $L^2\mathcal{H}^*(G_v\Sigma_{E_{2k}}) = L^2\mathcal{H}^*(\Sigma_{b\partial\Delta^{2k}})$ as Hilbert G_v -modules.

The complex E_{2k} is spanned by the even-dimensional vertices of $b\partial\Delta^{2k}$, which correspond to the proper subsets of vertices of Δ^{2k} of odd cardinality. Thus, the dimension of E_{2k} is $k - 1$, and its top-dimensional simplices are chains $v_0 < v_0v_1v_2 < \dots < v_0\dots v_{2k-2}$ of length k of distinct vertices of Δ^{2k} . Therefore \mathbf{S}_{2k+1} acts transitively on $(k - 1)$ -dimensional simplices of E_{2k} and it follows that G_v acts transitively on k -dimensional cubes of $G_v\Sigma_{E_{2k}}$. Therefore the space of k -cochains is a Hilbert G_v -submodule of $\ell^2(G_v)$, and the claim follows from the Hodge decomposition.

We have, by construction, $G_v = \text{Stab}_G(\Sigma_{b\partial\Delta_v^{2k+1}})$. Then the restriction map $i^*: L^2\mathcal{H}^k(\Sigma_{b\partial\Delta^{2k+1}}) \rightarrow L^2\mathcal{H}^k(\Sigma_{b\partial\Delta_v^{2k+1}})$ is G_v -equivariant and therefore its image is a G_v -invariant subspace of $L^2\mathcal{H}^k(\Sigma_{b\partial\Delta_v^{2k+1}})$. Since G_v acts preserving orientation, the intersection form is G_v -invariant. By Lemma 4.1 the image is isotropic, thus by Lemma 4.2 it is 0. Thus, the map $i^*: \mathfrak{h}^k(b\partial\Delta^{2k+1}) \rightarrow \mathfrak{h}^k(b\partial\Delta_v^{2k+1}) = L^2\mathcal{H}^k(W_{b\partial\Delta^{2k+1}}\Sigma_{b\partial\Delta_v^{2k+1}})$ is the 0-map.

For vertices of the other even dimensions the argument is similar. If $\dim(v) = 2d$, then its link is $b\partial\Delta^{2d} * b\partial\Delta^{2k-2d}$. Again, using Lemma 3.3, we remove, without changing the L^2 -cohomology, all odd-dimensional vertices from each factor to obtain $E_{2d} * E_{2k-2d}$. The group $\mathbf{S}_{2d+1} \times \mathbf{S}_{2k-2d+1}$ acts naturally on $b\partial\Delta^{2k+1}$ fixing the vertex v and stabilizing both the link and $E_{2d} * E_{2k-2d}$. This action is again transitive on the top-dimensional simplices of $E_{2d} * E_{2k-2d}$, and the rest of the argument goes through.

Finally, if v is an odd-dimensional vertex, $\dim(v) = 2d + 1$, then we have $b\partial\Delta_v^{2k+1} = b\partial\Delta^{2d+1} * b\partial\Delta^{2k-2d-1}$. The hypothesis on **III''** and Theorem 2.2 and Lemma 2.4 imply that both **I**($2d + 1$) and **I**($2k - 2d - 1$) hold. Therefore, by the Künneth formula $\mathfrak{h}^k(b\partial\Delta_v^{2k+1}) = 0$ in this case. \square

Theorem 5.2 *The Generalized Singer Conjecture holds true for $b\partial\Delta^{2k+1}$:*

$$b_{\mathbf{q}}^i(b\partial\Delta^{2k+1}) = 0 \text{ for } i > k \text{ and } \mathbf{q} \leq \mathbf{1}.$$

Proof We proceed by induction on k . Using the inductive assumption and Lemma 3.2, we can remove all odd-dimensional vertices without changing the weighted $L_{\mathbf{q}}^2$ -homology above k . Thus, since the remaining part E_{2k+1} is k -dimensional, the problem reduces to showing that $\mathfrak{h}_{k+1}^{\mathbf{q}}(E_{2k+1}) = 0$ for $\mathbf{q} \leq \mathbf{1}$. Since E_{2k+1} is k -dimensional, the natural map $\mathfrak{h}_{k+1}(E_{2k+1}) \rightarrow \mathfrak{h}_{k+1}^{\mathbf{q}}(E_{2k+1})$ is injective and the result follows from the Theorem 5.1. \square

Next, we list some consequences. Lemma 3.1 implies:

Theorem 5.3 *Let bL be the barycentric subdivision of an $(n-1)$ -dimensional simplicial complex L . Then*

$$b_{\mathbf{q}}^i(bL) = 0 \text{ for } i > (n + 1)/2 \text{ and } \mathbf{q} \leq \mathbf{1}.$$

Taking L to be a GHS^{2k-1} , we obtain:

Theorem 5.4 *The Generalized Singer Conjecture holds true for the barycentric subdivision of a GHS^{n-1} for all even n .*

For odd n we obtain a weaker statement:

Theorem 5.5 *Let bL be the barycentric subdivision of a GHS^{2k} . Then*

$$b_{\mathbf{q}}^i(bL) = 0 \text{ for } i > k + 1 \text{ and } \mathbf{q} \leq \mathbf{1}.$$

In particular,

$$b^i(bL) = 0 \text{ for } i \neq k, k + 1.$$

Specializing further, and combining with Theorem 5.2, we obtain:

Theorem 5.6 *The Generalized Singer Conjecture holds true for $b\partial\Delta^n$:*

$$b_{\mathbf{q}}^i(b\partial\Delta^n) = 0 \text{ for } i > n/2 \text{ and } \mathbf{q} \leq \mathbf{1},$$

and, therefore, for the class \mathcal{JD} .

Finally, let us mention an application of the above result to a more analytic object. Let T_n denote the space of all symmetric tridiagonal $(n+1) \times (n+1)$ -matrices with fixed generic eigenvalues, the so-called Tomei manifold. It is proved in [8] that T_n is an n -dimensional closed aspherical manifold.

Theorem 5.7 *The Singer Conjecture holds true for Tomei manifolds T_n .*

Proof The space T_n can be identified with a natural finite index orbifolal cover of $\Sigma_{b\partial\Delta^n}/W_{b\partial\Delta^n}$ [3]. Thus $\Sigma_{b\partial\Delta^n}$ is the universal cover of T^n , and the claim follows from the previous theorem. \square

References

- [1] **M W Davis**, *Groups generated by reflections and aspherical manifolds not covered by Euclidean space*, Ann. of Math. 117 (1983) 293–324 MathReview
- [2] **M W Davis**, *Coxeter groups and aspherical manifolds*, from: “Algebraic topology (Aarhus 1982)”, Lecture Notes in Math. 1051, Springer, New York (1984) 197–221 MathReview
- [3] **M W Davis**, *Some aspherical manifolds*, Duke Math. J. 55 (1987) 105–139 MathReview
- [4] **M W Davis**, *Nonpositive curvature and reflection groups*, from: “Handbook of Geometric Topology”, (R Daverman, R Sher, editors), Elsevier, Amsterdam (2002) 373–422 MathReview
- [5] **M W Davis**, **T Januszkiewicz**, **J Dymara**, **B L Okun**, *Weighted L^2 -cohomology of Coxeter groups*, arXiv:math.GT/0402377
- [6] **M W Davis**, **B L Okun**, *ℓ_2 -homology of Coxeter groups based on barycentric subdivisions*, Topol. Appl. 140 (2004) 197–202
- [7] **J Dymara**, *Thin buildings*, preprint
- [8] **C Tomei**, *The topology of isospectral manifolds of tridiagonal matrices*, Duke Math. J. 51 (1984) 981–996 MathReview