

BEYOND DELAUNAY SURFACES*

PETER A. DJONDJOROV, MARIANA TS. HADZHILAZOVA[†]
IVAÏLO M. MLADENOV[†] and VASSIL M. VASSILEV

*Institute of Mechanics, Bulgarian Academy of Sciences
Acad. G. Bonchev Str., Bl. 4, 1113 Sofia, Bulgaria*

[†]*Institute of Biophysics, Bulgarian Academy of Sciences
Acad. G. Bonchev Str., Bl. 21, 1113 Sofia, Bulgaria*

Abstract. An interesting class of axially symmetric surfaces, which generalizes Delaunay's unduloids and provides solutions of the shape equation is described in explicit parametric form. This class provide the first analytical examples of surfaces with periodic curvatures studied by K. Kenmotsu and leads to some unexpected relationships among Jacobian elliptic functions and their integrals.

1. Introduction

It is well-known that in aqueous solution, amphiphilic molecules (e.g., phospholipids) form spontaneously bilayers so that the hydrophilic heads of these molecules are located in both outer sides of the bilayer which are in contact with the liquid, while their hydrophobic tails remain at the interior. In many cases, the bilayer form a closed membrane, which is called a vesicle. Vesicles constitute well-defined and sufficiently simple model systems for studying basic physical properties of the more complex biological cells.

In 1973, Helfrich [3] had proposed the so-called **spontaneous curvature model** according to which the equilibrium shapes of a lipid vesicle are determined by the extremals of the curvature (shape) energy

$$\mathcal{F}_c = \frac{k_c}{2} \int_S (2H - \mathfrak{h})^2 dA + k_G \int_S K dA$$

*Reprinted from JGSP **18** (2010) 1–11.