

**A PRIORI BOUNDS FOR POSITIVE RADIAL SOLUTIONS OF QUASILINEAR EQUATIONS OF LANE–EMDEN TYPE**

SOOHYUN BAE

ABSTRACT. We consider the quasilinear equation  $\Delta_p u + K(|x|)u^q = 0$ , and present the proof of the local existence of positive radial solutions near 0 under suitable conditions on  $K$ . Moreover, we provide a priori estimates of positive radial solutions near  $\infty$  when  $r^{-\ell}K(r)$  for  $\ell \geq -p$  is bounded near  $\infty$ .

1. INTRODUCTION

We consider the equation

$$(1.1) \quad \Delta_p u + K(|x|)u^q = 0,$$

where  $\Delta_p u = \operatorname{div}(|\nabla u|^{p-2}\nabla u)$ ,  $n > p > 1$  and  $q > p - 1$ . Let  $r = |x|$  and  $\frac{d}{dr}u(r) = u_r(r)$ . Then, the radial version of (1.1) is

$$(1.2) \quad r^{1-n}(r^{n-1}|u_r|^{p-2}u_r)_r + K(r)u^q = 0.$$

For  $p = 2$ , the basic assumption of  $K$  for local solutions is (K):

- (i)  $K(r) \geq 0, \neq 0$ ;  $K(r)$  is continuous on  $(0, \infty)$ ;
- (ii)  $\int_0^r rK(r) dr < \infty$ , i.e.,  $rK(r)$  is integrable near 0.

Under condition (K), (1.2) with  $p = 2$  and  $u(0) = \alpha > 0$ , has a unique positive solution  $u_\alpha \in C^2(0, \varepsilon) \cap C[0, \varepsilon)$  for small  $\varepsilon > 0$ . In order to obtain local solutions (1.2) near 0, we assume (KP): (i) of (K), and for  $r > 0$  small,

$$\int_0^r t^{\frac{1-n}{p-1}} \left( \int_0^t s^{n-1} K(s) ds \right)^{\frac{1}{p-1}} dt < \infty.$$

For  $p = 2$ , this integrability is (ii) of (K). If  $K(r) = r^l$ , then it is easy to see that (KP) holds for  $l > -p$ . As a typical example, the equation

$$(1.3) \quad \Delta_p u + |x|^l u^q = 0$$

possesses a local radial solution  $\bar{u}_\alpha$  with  $\bar{u}_\alpha(0) = \alpha$  for each  $\alpha > 0$ , and has the scaling invariance:

$$(1.4) \quad \bar{u}_\alpha(r) = \alpha \bar{u}_1(\alpha^{\frac{1}{m}} r)$$

2020 *Mathematics Subject Classification*: primary 35J92; secondary 35B45.

*Key words and phrases*: quasilinear equation, positive solution, a priori bound.

This research was supported by the research fund of Hanbat National University in 2018.

Received August 26, 2022, accepted December 16, 2022. Editor Š. Nečasová.

DOI: 10.5817/AM2023-2-155

with  $m = \frac{p+l}{q-(p-1)}$ . Moreover, (1.3) has a singular solution which is invariant under the scaling in (1.4), the so-called self-similar solution. That is,

$$U(x) = L|x|^{-m},$$

where  $L$  is defined by

$$(1.5) \quad L = L(n, p, q, l) = [m^{p-1}(n - 1 - (m + 1)(p - 1))]^{\frac{1}{q-(p-1)}}.$$

This singular solution can be defined for  $l > -p$  and  $q > \frac{(p-1)(n+l)}{n-p}$  because  $n - 1 - (m + 1)(p - 1) > 0$ . Then, we observe the asymptotic self-similar behavior.

**Theorem 1.1.** *Let  $n > p > 1$  and  $q > \frac{(p-1)(n+l)}{n-p}$  with  $l > -p$ . If  $r^{-l}K(r) \rightarrow 1$  as  $r \rightarrow \infty$ , then any positive solution  $u$  of (1.2) near  $\infty$  satisfies one of the two asymptotic behavior: either*

$$(1.6) \quad \liminf_{r \rightarrow \infty} r^m u(r) \leq L \leq \limsup_{r \rightarrow \infty} r^m u(r) < \infty$$

with  $L = L(n, p, q, l)$  given by (1.5) or  $r^{(n-p)/(p-1)}u(r) \rightarrow C > 0$  as  $r \rightarrow \infty$ .

Moreover, (1.6) can be the asymptotic self-similarity

$$\lim_{r \rightarrow \infty} r^m u(r) = L.$$

In a forthcoming paper, we study entire solutions of (1.2) with this asymptotic behavior in a supercritical range.

**1.1. Lower bound.** The  $p$ -Laplace equation has the radial form

$$(1.7) \quad (|u_r|^{p-2}u_r)_r + \frac{n-1}{r}|u_r|^{p-2}u_r = 0,$$

where  $n > p > 1$ . Then, (1.7) possesses a solution  $|x|^{-\theta}$  with  $\theta = \frac{n-p}{p-1}$ . Let  $u$  be a positive radial solution satisfying the quasilinear inequality

$$(1.8) \quad r^{1-n}(r^{n-1}|u_r|^{p-2}u_r)_r = (|u_r|^{p-2}u_r)_r + \frac{n-1}{r}|u_r|^{p-2}u_r \leq 0.$$

If  $u_r(r_0) \leq 0$  for some  $r_0 > 0$ , then  $u_r(r) \leq 0$  for  $r > r_0$ . Hence,  $u$  is monotone near  $\infty$ . Assume  $u_r \leq 0$  for  $r \geq r_0$  with some  $r_0 > 1$ . Setting  $V(t) = r^\theta u(r)$  for  $t = \log r \geq t_0 = \log r_0$ , we see that  $g(t) = \theta V(t) - V'(t) = r^{\theta+1}(-u_r(r)) = r^{\frac{n-1}{p-1}}(-u_r(r))$  satisfies

$$\frac{d}{dt}(g^{p-1}(t)) = (n-1)g^{p-1}(t) + r^n[(-u_r)^{p-1}]_r \geq 0$$

for  $t \geq t_0$ . Hence,  $g$  is increasing for  $t \geq t_0$ . Then,  $V$  satisfies that for  $t > T \geq t_0$ ,

$$V'(t) - \theta V(t) \leq V'(T) - \theta V(T).$$

Suppose  $V'(T) < 0$ . Setting  $c = \theta V(T) - V'(T)$ , we have  $(e^{-\theta t}V(t))' \leq -ce^{-\theta t}$  and

$$V(t) \leq e^{\theta(t-T)}(V(T) - \frac{c}{\theta}) + \frac{c}{\theta} = e^{\theta(t-T)}\frac{V'(T)}{\theta} + \frac{c}{\theta}.$$

Hence,  $V$  has a finite zero. Therefore, in order for  $u$  to be positive near  $\infty$ ,  $V$  must be increasing and  $(r^\theta u(r))_r \geq 0$  near  $\infty$ . This is true obviously in the other case that  $u_r > 0$  near  $\infty$ .

**Lemma 1.2.** *Let  $n > p > 1$ . If  $u$  is a positive radial solution satisfying (1.8) near  $\infty$ , then  $r^{\frac{n-p}{p-1}}u(r)$  is increasing.*

Now, we classify positive solutions of (1.8) near  $\infty$  into two groups according to their behaviors. If  $r^{\frac{n-p}{p-1}}u$  converges to a positive constant at  $\infty$ , then we call  $u$  a fast decaying solution. Otherwise,  $u$  is a slowly decaying solution if  $r^{\frac{n-p}{p-1}}u(r) \rightarrow \infty$  as  $r \rightarrow \infty$ .

**1.2. Known results.** One of Liouville’s theorems related to  $p$ -Laplace equation is the nonexistence of nontrivial nonnegative solutions in  $W_{loc}^{1,p}(\mathbf{R}^n) \cap C(\mathbf{R}^n)$  to the following quasilinear inequality

$$-\Delta_p u \geq c|x|^l u^q$$

with  $c > 0$  and  $l > -p$ , when  $n > p > 1$  and

$$q \leq \frac{(p-1)(n+l)}{n-p}.$$

See [1, Theorem 3.3 (iii)]. For the existence of nontrivial solutions to

$$\Delta_p u + u^q = 0,$$

on  $\mathbf{R}^n$  with  $n > p > 1$  and  $q > p-1$ , it is necessary and sufficient that  $q \geq \frac{n(p-1)+p}{n-p}$  [6]. On the other hand, (1.3) with  $q = q_s := \frac{n(p-1)+p+p_l}{n-p}$  admits the one-parameter family of positive solutions given by

$$\bar{u}_\alpha(x) = \frac{\alpha}{(1 + \xi(\alpha^{\frac{p}{n-p}} |x|)^{\frac{p+l}{p-1}})^{\frac{n-p}{p+l}}}$$

with  $\xi = \xi_{p,n} = \frac{p-1}{(n-p)(n+l)^{1/(p-1)}}$  and  $\bar{u}_\alpha(0) = \alpha > 0$ . A radial solution  $u(x) = u(|x|)$  of (1.3) satisfies the equation

$$(1.9) \quad (|u_r|^{p-2}u_r)_r + \frac{n-1}{r}|u_r|^{p-2}u_r + r^l u^q = 0.$$

For  $l > -p$ , (1.9) with  $u(0) = \alpha > 0$ , has a unique positive solution  $u \in C^1(0, \epsilon) \cap C[0, \epsilon)$  for small  $\epsilon > 0$  such that  $|u_r|^{p-2}u_r \in C^1[0, \epsilon)$ . If  $q < q_s$ , then every local solution of (1.9) has a finite zero [2, 5]. In the opposite case  $q > q_s$ , every local solution of (1.9) is to be a slowly decaying solution [2, 3, 5].

## 2. LOCAL EXISTENCE

Let  $n \geq p > 1$ ,  $l > -p$  and  $q \geq p-1$ . First, in order to prove the local existence of positive radial solutions of (1.3), we consider the integral equation

$$u(r) = \alpha - \int_0^r t^{\frac{1-n}{p-1}} \left( \int_0^t s^{n-1+l} u^q(s) ds \right)^{\frac{1}{p-1}} dt$$

with  $\alpha > 0$ .

**2.1. Integral representation.** On a space

$$S = \{u \in C[0, \varepsilon] \mid 0 \leq u \leq \alpha\}$$

we study a nonlinear operator  $T$  from  $S$  to  $C[0, \varepsilon]$  by

$$T(u)(r) = \alpha - T_1(u)(r),$$

where

$$T_1(u)(r) = \int_0^r t^{\frac{1-n}{p-1}} \left( \int_0^t s^{n-1+l} u^q(s) ds \right)^{\frac{1}{p-1}} dt.$$

For  $\varepsilon > 0$  small enough,  $T_1$  satisfies that

$$0 \leq T_1 \leq \alpha^{\frac{q}{p-1}} \int_0^r t^{\frac{1-n}{p-1}} \left( \int_0^t s^{n-1+l} ds \right)^{\frac{1}{p-1}} dt \leq \left( \frac{\alpha^q}{n+l} \right)^{\frac{1}{p-1}} \frac{p-1}{p+l} \varepsilon^{\frac{p+l}{p-1}} \leq \alpha.$$

Hence,  $T(S) \subset S$ . Minkowski's inequality for  $p \geq 2$  shows that for  $u_1, u_2 \in S$ ,

$$\begin{aligned} \|T(u_2) - T(u_1)\| &\leq \int_0^r t^{\frac{1-n}{p-1}} \left( \int_0^t s^{n-1+l} |u_2^{\frac{q}{p-1}} - u_1^{\frac{q}{p-1}}|^{p-1} ds \right)^{\frac{1}{p-1}} dt \\ &\leq \frac{q}{p-1} \alpha^{\frac{q-(p-1)}{p-1}} \int_0^r t^{\frac{1-n}{p-1}} \left( \int_0^t s^{n-1+l} ds \right)^{\frac{1}{p-1}} dt \|u_2 - u_1\| \\ &= \frac{q}{p-1} \alpha^{\frac{q-(p-1)}{p-1}} \left( \frac{1}{n+l} \right)^{\frac{1}{p-1}} \frac{p-1}{p+l} \varepsilon^{\frac{p+l}{p-1}} \|u_2 - u_1\|. \end{aligned}$$

For  $1 < p < 2$ , we observe that for  $u_1, u_2 \in S$ ,

$$\begin{aligned} \|T(u_2) - T(u_1)\| &\leq \int_0^r t^{\frac{1-n}{p-1}} \frac{\alpha^{\frac{q(2-p)}{p-1}}}{p-1} \left( \int_0^t s^{n-1+l} ds \right)^{\frac{2-p}{p-1}} \left( \int_0^t s^{n-1+l} |u_2^q - u_1^q| ds \right) dt \\ &\leq \frac{q}{p-1} \alpha^{\frac{q-(p-1)}{p-1}} \int_0^r t^{\frac{1-n}{p-1}} \left( \int_0^t s^{n-1+l} ds \right)^{\frac{1}{p-1}} dt \|u_2 - u_1\| \\ &= \frac{q}{p-1} \alpha^{\frac{q-(p-1)}{p-1}} \left( \frac{1}{n+l} \right)^{\frac{1}{p-1}} \frac{p-1}{p+l} \varepsilon^{\frac{p+l}{p-1}} \|u_2 - u_1\|. \end{aligned}$$

Now, we assume that

$$\frac{p-1}{p+l} \max\left\{ \left( \frac{\alpha^q}{n+l} \right)^{\frac{1}{p-1}}, \frac{q}{p-1} \alpha^{\frac{q-(p-1)}{p-1}} \left( \frac{1}{n+l} \right)^{\frac{1}{p-1}} \right\} \varepsilon^{\frac{p+l}{p-1}} < \min\{\alpha, 1\}.$$

Then,  $T$  is a contraction mapping in  $S$  and thus  $T$  has a unique fixed point  $\bar{u}_\alpha$ .

Generally, we consider the integral equation under condition (KP),

$$u(r) = \alpha - \int_0^r t^{\frac{1-n}{p-1}} \left( \int_0^t s^{n-1} K(s) u^q(s) ds \right)^{\frac{1}{p-1}} dt.$$

Then, the integrability of (KP) shows in the same way the local existence of a positive solution  $u_\alpha$  with  $u_\alpha(0) = \alpha > 0$  to (1.2). Then, it is easy to see that there exists a sequence  $\{r_j\}$  going to 0 such that

$$(2.1) \quad \lim_{j \rightarrow \infty} r_j^{n-1} |u_r(r_j)|^{p-2} u_r(r_j) = 0,$$

and  $u_\alpha(r)$  is decreasing as long as  $u$  remains positive. Moreover,  $u_\alpha$  is strictly decreasing after  $K$  becomes positive.

**2.2. Fowler transform.** Let  $n > p > 1$  and  $q > \frac{(n+l)(p-1)}{n-p}$  with  $l > -p$ . Set  $m = \frac{p+l}{q-(p-1)}$ . Fowler transform  $V(t) = r^m u(r)$ ,  $t = \log r$ , of a positive solution to (1.2) satisfies

$$(2.2) \quad (p-1)(mV - V')^{p-2}(V'' - mV') - \xi(mV - V')^{p-1} + k(t)V^q = 0,$$

where  $\xi = n-1-(m+1)(p-1) = \frac{L^{q-(p-1)}}{m^{p-1}}$  with  $L$  given by (1.5), and  $k(t) = r^{-l}K(r)$ . Furthermore, if  $-r^{m+1}u_r(r) = mV - V' > 0$ , then (2.2) can be rewritten as

$$(p-1)(V'' - mV') - \xi(mV - V') = -\frac{k(t)V^q}{(mV - V')^{p-2}}$$

and

$$(p-1)V'' + aV' - \xi mV = -\frac{k(t)V^q}{(mV - V')^{p-2}},$$

where  $a = n-1-(2m+1)(p-1)$ . Setting  $b = \xi m = \frac{L^{q-(p-1)}}{m^{p-2}}$ , we have

$$(p-1)V'' + aV' - (b - \frac{k(t)V^{q-1}}{(mV - V')^{p-2}})V = 0.$$

That is,

$$(2.3) \quad (p-1)V'' + aV' - \frac{1}{m^{p-2}}L^{q-(p-1)}V + \frac{k(t)}{(mV - V')^{p-2}}V^q = 0,$$

which holds as long as the local solution remains positive.

### 3. A PRIORI ESTIMATES

In order to obtain upper bounds, we argue similarly as in Lemma 2.16, Lemma 2.20, Theorem 2.25 in [4].

**3.1. Upper bound.** Let  $n > p \geq -\ell$ . If  $u$  is a positive solution satisfying the inequality

$$(3.1) \quad (r^{n-1}|u_r|^{p-2}u_r)_r \leq -cr^{n-1+\ell}u^q$$

near  $\infty$  for some  $c > 0$ , then

$$(3.2) \quad r^{n-1}|u_r|^{p-2}u_r \leq r_0^{n-1}|u_r(r_0)|^{p-2}u_r(r_0) - c \int_{r_0}^r s^{n-1+\ell}u^q(s) ds$$

for  $r > r_0$ , if  $r_0$  is sufficiently large. Then, we may assume that  $u_r(r_0) \leq 0$ . Indeed, if  $u_r(r_0) > 0$ , then

$$r^{n-1}|u_r|^{p-2}u_r \leq r_0^{n-1}|u_r(r_0)|^{p-2}u_r(r_0) - cu^q(r_0)\frac{1}{n+\ell}(r^{n+\ell} - r_0^{n+\ell})$$

as long as  $u_r$  is positive. Hence,  $u_r$  is eventually negative. Therefore, (3.2) gives

$$r^{n-1}|u_r|^{p-2}u_r \leq -cu^q(r)\frac{1}{n+\ell}(r^{n+\ell} - r_0^{n+\ell})$$

and thus,

$$\frac{u_r}{u^{q/(p-1)}} \leq -c_1 r^{\frac{1+\ell}{p-1}}$$

for some  $c_1 > 0$ . Hence, we obtain

$$u(r) \leq \begin{cases} Cr^{-\frac{p+\ell}{q-(p-1)}} & \text{if } \ell > -p, \\ C(\log r)^{-\frac{p-1}{q-(p-1)}} & \text{if } \ell = -p \end{cases}$$

for some  $C > 0$ . Combining the a priori estimates and Lemma 1.2, we have the following assertion.

**Theorem 3.1.** *Let  $n > p \geq -\ell$  and  $q > \frac{(p-1)(n+\ell)}{n-p}$ . Then, every positive solution to (3.1) near  $\infty$  satisfies that*

$$C_1 r^{-\frac{p+\ell}{q-(p-1)}} \geq u(r) \geq C_2 r^{-\frac{n-p}{p-1}}$$

for  $\ell > -p$  and

$$C_1 (\log r)^{-\frac{p-1}{q-(p-1)}} \geq u(r) \geq C_2 r^{-\frac{n-p}{p-1}}$$

for  $\ell = -p$ .

In Theorem 3.1, we use the notation  $\ell$  instead of  $l$  to consider the case of  $\ell = -p$ . It is interesting to study the existence of positive entire solutions of (1.1) with the logarithmic asymptotic behavior at  $\infty$ .

**Lemma 3.2.** *Let  $q > \frac{(p-1)(n+l)}{n-p}$ . Assume  $K(r) = O(r^l)$  at  $\infty$  for some  $l > -p$ . If  $u$  is a positive solution to (3.1) near  $\infty$  and  $u(r) = O(r^{-m-\varepsilon})$  with some  $\varepsilon > 0$  at  $\infty$ , then  $u(r) = O(r^{\frac{p-n}{p-1}})$  at  $\infty$ .*

**Proof.** Integrating (1.2) over  $[r, \infty)$ , we obtain

$$u(r) = \int_r^\infty t^{\frac{1-n}{p-1}} \left( \int_0^t K(s)u^q(s)s^{n-1} ds \right)^{\frac{1}{p-1}} dt.$$

On the other hand, we have

$$\begin{aligned} \int_0^t K(s)u^q(s)s^{n-1} ds &\leq C + C \int_1^t s^{n-1+l-q(m+\varepsilon)} ds \\ &= \begin{cases} C + Ct^{n+l-q(m+\varepsilon)} & \text{if } n+l \neq q(m+\varepsilon), \\ C + C \log t & \text{if } n+l = q(m+\varepsilon). \end{cases} \end{aligned}$$

If  $n+l < q(m+\varepsilon)$ , we are done. If  $n+l \geq q(m+\varepsilon)$ , then

$$u(r) \leq \begin{cases} Cr^{\frac{p-n}{p-1}} + Cr^{\frac{p-n}{p-1}} (\log r)^{\frac{1}{p-1}} & \text{if } n+l = q(m+\varepsilon), \\ Cr^{\frac{p-n}{p-1}} + Cr^{\frac{p+l}{p-1} - \frac{q(m+\varepsilon)}{p-1}} & \text{if } p+l < q(m+\varepsilon) < n+l. \end{cases}$$

In case  $n+l = q(m+\varepsilon)$ , we replace  $\varepsilon$  by  $\frac{n-p}{p-1} - m - \delta$  in the above arguments, where  $\delta > 0$  is so small that  $\delta < \frac{n-p}{p-1} - m$ . Note that  $m < \frac{n-p}{p-1}$  iff  $q > \frac{(p-1)(n+l)}{n-p}$ .

$$u(r) \leq \begin{cases} Cr^{\frac{p-n}{p-1}} & \text{if } n+l = q(m+\varepsilon), \\ Cr^{\frac{p-n}{p-1}} + Cr^{\frac{p+l}{p-1} + \frac{q(p+l)}{(p-1)^2} - \frac{q^2(m+\varepsilon)}{(p-1)^2}} & \text{if } p+l < q(m+\varepsilon) < n+l. \end{cases}$$

In case  $q(m + \varepsilon) < n + l$ , we iterate this process to obtain

$$\begin{aligned} u(r) &\leq Cr^{\frac{p-n}{p-1}} + Cr^{\frac{p+l}{p-1}} \sum_{i=0}^{j-1} \left(\frac{q}{p-1}\right)^i - \frac{q^j(m+\varepsilon)}{(p-1)^j} \\ &= Cr^{\frac{p-n}{p-1}} + Cr^{-m-\varepsilon\left(\frac{q}{p-1}\right)^j} \end{aligned}$$

for any positive integer  $j$ . Since  $q > p - 1$ , we reach the conclusion after a finite number of iterations.  $\square$

**Lemma 3.3.** *Let  $q > \frac{(p-1)(n+l)}{n-p}$ . Assume  $K(r) = O(r^l)$  at  $\infty$  for some  $l > -p$ . If  $u(r) = o(r^{-m})$  at  $\infty$ , then  $(r^m u(r))_r < 0$  near  $\infty$ .*

**Proof.** Let  $V(t) = r^m u(r)$ ,  $t = \log r$ . Then,  $V$  satisfies (2.3). Suppose  $V'(T) = 0$  for some  $T$  near  $\infty$  and  $k(t)V^{q-(p-1)}(t) < m^{p-2}b$  for  $t \in [T, \infty)$ . Then,  $V''(T) > 0$  and  $V(t)$  is strictly increasing near  $T$  but for  $t > T$ . Since  $V \rightarrow 0$  at  $\infty$ , there exists  $T_1 > T$  such that  $V'(T_1) = 0$  and

$$V''(T_1) = \frac{1}{p-1} \left( b - \frac{1}{m^{p-2}} k(T_1) V^{q-(p-1)}(T_1) \right) V(T_1) \leq 0,$$

a contradiction.  $\square$

**Theorem 3.4.** *Let  $q > \frac{(p-1)(n+l)}{n-p}$ . Assume  $K(r) = O(r^l)$  at  $\infty$  for some  $l > -p$ . If  $u(r) = o(r^{-m})$  at  $\infty$ , then  $u(r) = O(r^{\frac{p-n}{p-1}})$  at  $\infty$ .*

**Proof.** Let  $\varphi(r) = r^m u(r)$ . Then,  $\varphi$  satisfies

$$\varphi_{rr} + \left(1 + \frac{a}{p-1}\right) \frac{1}{r} \varphi_r - \frac{b}{(p-1)r^2} \varphi + \frac{k}{(p-1)(m\varphi - r\varphi_r)^{p-2} r^2} \varphi^q = 0.$$

For  $\varepsilon > 0$ , define the elliptic operator

$$\mathcal{L}_\varepsilon \varphi = \Delta \varphi - \left[2m + (n-1) \frac{p-2}{p-1}\right] \frac{x \cdot \nabla \varphi}{|x|^2} - m \left( \frac{L^{q-(p-1)}}{m^{p-1}} - \varepsilon \right) \frac{\varphi}{|x|^2},$$

where  $\frac{L^{q-(p-1)}}{m^{p-1}} = n - 1 - (m+1)(p-1)$ . It follows from Lemma 3.3 that for any  $\varepsilon > 0$ , there exists  $R_\varepsilon > 0$  such that

$$\mathcal{L}_\varepsilon \varphi = m\varepsilon \frac{\varphi}{r^2} - \frac{k\varphi^q}{(p-1)r^2(m\varphi - r\varphi_r)^{p-2}} \geq \left(m\varepsilon - \frac{k\varphi^{q-(p-1)}}{(p-1)m^{p-2}}\right) \frac{\varphi}{r^2} \geq 0$$

in  $\mathbf{R}^n \setminus B_{R_\varepsilon}(0)$ . For  $0 < \varepsilon < n - 1 - (m+1)(p-1)$ , let  $\eta_\varepsilon(x) = |x|^{\sigma_\varepsilon}$  with  $\sigma_\varepsilon$  being the negative root of  $\sigma(\sigma-1) + (n-1-2m-(n-1)\frac{p-2}{p-1})\sigma - m\left(\frac{L^{q-(p-1)}}{m^{p-1}} - \varepsilon\right) = 0$ , i.e.,

$$\sigma_\varepsilon = \frac{1}{2} \left[ -\left(n-2-2m-(n-1)\frac{p-2}{p-1}\right) - \sqrt{D} \right],$$

where  $D = (n-1-2m-(n-1)\frac{p-2}{p-1})^2 + 4m\left(\frac{L^{q-(p-1)}}{m^{p-1}} - \varepsilon\right)$ . Setting  $C_\varepsilon = \varphi(R_\varepsilon)R_\varepsilon^{-\sigma_\varepsilon}$ , we see that  $\mathcal{L}_\varepsilon(\varphi - C_\varepsilon\eta_\varepsilon) \geq 0$  in  $\mathbf{R}^n \setminus B_{R_\varepsilon}(0)$  and  $\varphi(R_\varepsilon) = C_\varepsilon\eta_\varepsilon(R_\varepsilon)$ ,  $\varphi - C_\varepsilon\eta_\varepsilon \rightarrow 0$  as  $r \rightarrow \infty$ . Then, the maximum principle implies that  $\varphi - C_\varepsilon\eta_\varepsilon \leq 0$  in  $\mathbf{R}^n \setminus B_{R_\varepsilon}(0)$ . Hence,  $\varphi(r) \leq C_\varepsilon\eta_\varepsilon(r)$  at  $\infty$ . Then, Lemma 3.2 implies the conclusion.  $\square$

**Proof of Theorem 1.1.** When  $k(t) = r^{-l}K(r) \rightarrow 1$  as  $t = \log r \rightarrow +\infty$ , it follows from Theorem 3.1 and (2.3) that slowly decaying solutions satisfy

$$\liminf_{r \rightarrow \infty} r^m u(r) \leq L \leq \limsup_{r \rightarrow \infty} r^m u(r) < \infty.$$

Indeed, at every local minimum (maximum) point of  $V(t) = r^m u(r)$ ,  $V$  satisfies

$$\frac{1}{m^{p-2}} L^{q-(p-1)} V \geq (\leq) \frac{k(t)}{(mV)^{p-2}} V^q.$$

If  $V$  is monotonically increasing near  $+\infty$ , then it is easy to see that  $V \rightarrow L$  as  $t \rightarrow +\infty$  by (2.3). If  $V$  is monotonically decreasing and  $V \rightarrow 0$ , then it follows from Lemma 1.2 and Theorem 3.4 that  $r^{\frac{n-p}{p-1}} u(r) \rightarrow C$  for some  $C > 0$ .  $\square$

## REFERENCES

- [1] Bidaut-Véron, M.-F., Pohozaev, S., *Nonexistence results and estimates for some nonlinear elliptic problems*, J. Anal. Math. **84** (2001), 1–49.
- [2] Kawano, N., Ni, W.-M., Yotsutani, S., *A generalized Pohozaev identity and its applications*, J. Math. Soc. Japan **42** (1990), 541–564.
- [3] Kawano, N., Yanagida, E., Yotsutani, S., *Structure theorems for positive radial solutions to  $\operatorname{div}(|Du|^{m-2} Du) + K(|x|)u^q = 0$  in  $\mathbf{R}^n$* , J. Math. Soc. Japan **45** (1993), 719–742.
- [4] Li, Y., Ni, W.-M., *On conformal scalar curvature equation in  $\mathbf{R}^n$* , Duke Math. J. **57** (1988), 895–924.
- [5] Ni, W.-M., Serrin, J., *Existence and non-existence theorems for ground states of quasilinear partial differential equations: The anomalous case*, Atti Convegna Lincei **77** (1986), 231–257.
- [6] Serrin, J., Zou, H., *Cauchy-Liouville and universal boundedness theorems for quasilinear elliptic equations and inequalities*, Acta Math. **189** (2002), 79–142.

DEPARTMENT OF MATHEMATICAL SCIENCES, HANBAT NATIONAL UNIVERSITY,  
 DAEJEON 34158, REPUBLIC OF KOREA  
*E-mail*: shbae@hanbat.ac.kr